



**QUEEN'S
UNIVERSITY
BELFAST**

Microbial mediated arsenic biotransformation in wetlands

Zhang , S.-Y., Williams, P. N., Luo, J., & Zhu, Y.-G. (2017). Microbial mediated arsenic biotransformation in wetlands. *Frontiers of Environmental Science & Engineering*, 11(1). <https://doi.org/10.1007/s11783-017-0893-y>

Published in:
Frontiers of Environmental Science & Engineering

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© 2016 Higher Education Press and Springer-Verlag Berlin Heidelberg.
This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access
This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

1 **Microbial mediated arsenic biotransformation in wetlands**
2 **Si-Yu Zhang^{1, 2}, Paul N. Williams³, Jinming Luo⁴, Yong-Guan Zhu (*)^{1, 5}**
3
4 1 State Key Laboratory of Urban and Regional Ecology, Research Center for
5 Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China
6 2 State Key Joint Laboratory of Environment Simulation and Pollution Control,
7 School of Environment, Beijing Normal University, Beijing 100875, China
8 3 Institute for Global Food Security, School of Biological Sciences, Queen's
9 University Belfast, Belfast, BT9 7BN, UK
10 4 Key Laboratory of Drinking Water Science and Technology, Research Center for
11 Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085.
12 5 Key Laboratory of Urban Environment and Health, Institute of Urban Environment,
13 Chinese Academy of Sciences, Xiamen 361021, China.
14 * *Corresponding author*

Abstract

Arsenic (As) is a pervasive environmental toxin and carcinogenic metalloid. It ranks at the top of the US priority List of Hazardous Substances and causes worldwide human health problems. Wetlands, including natural and artificial ecosystems (i.e. paddy soils) are highly susceptible to As enrichment; acting not only as repositories for water but a host of other elemental/chemical moieties. While macro-scale processes (physical and geological) supply As to wetlands, it is the micro-scale biogeochemistry that regulates the fluxes of As and other trace elements from the semi-terrestrial to neighboring plant/aquatic/atmospheric compartments. Among these fine-scale events, microbial mediated As biotransformations contribute most to the element's changing forms, acting as the 'switch' in defining a wetland as either a source or sink of As. Much of our understanding of these important microbial catalyzed reactions follows relatively recent scientific discoveries. Here we document some of these key advances, with focuses on the implications that wetlands and their microbial mediated transformation pathways have on the global As cycle, the chemistries of microbial mediated As oxidation, reduction and methylation, and future research priorities areas.

Keywords

Arsenic, wetland, microbes, switch

1 **1 Introduction**

2 Arsenic (As) exists in four oxidation states (-III, 0, +III, and +V), and is present in
3 both inorganic and organic forms. Commonly found in highly auriferous regions, As
4 also associates with the mineral ores of other metals such as iron, copper, and lead [1].
5 Speciation in addition to concentration is fundamental to the understanding of As
6 toxicity, mobility and fate in the environment [2]. In the biosphere, arsenate (As^{V}) and
7 arsenite (As^{III}) are the most abundant inorganic As species. Under aerobic conditions,
8 As^{V} dominates, is typically strongly absorbed to iron oxy-hydroxide minerals, and
9 exhibits limited mobility/bioavailability [3]. Conversely, under anaerobiosis, a shift in As
10 speciation to the trivalent form means the element is both more toxic and inherently
11 more labile [3].

12 Common organic As species found in our bodies as well as in waters, soils, feeds
13 and foods include monomethylarsenate (MMAs^{V}), dimethylarsenate (DMAs^{V}),
14 trimethylarsine oxide (TMAOs), arsenosugars and arsenobetaine (AsB) (Table 1).
15 They can be introduced into the environment due to microbial mediated
16 biomethylation [4] and via anthropogenic activities, such as pesticide/fertilizer
17 applications. Historically, the organic arsenical Roxarsone (2-nitrophenol-4-arsonic
18 acid), has been added to poultry feeds as a growth promoter, however this has been
19 largely phased out due to both animal and environmental health concerns [5]. Most,
20 but not all, of the organic As species are considerably less toxic compared to their
21 inorganic As counterparts. Typically, they are found **with** high concentrations in
22 marine organisms. Arsenosugars constitute the main organo-As species found in algae,

1 while in other marine dwellers, from lobsters to white fish, the toxin is packaged as
2 the inert and very stable/unreactive AsB [6]. In freshwaters and soils/sediments,
3 generally MMAs^V, DMAs^V and volatile TMAOs are the most commonly detected
4 organic As species [7].

5 As the interface between terrestrial and aquatic ecosystems, wetlands occupy
6 between 5.3 and 12.8 million km² of the earth's surface [8]. Defined by their
7 inundation by water and constant transition between wet and dry states, oxygen
8 availability is typically very heterogeneous resulting in regions of water and sediment
9 where anaerobism is commonplace [9]. Because of their wide distribution, high
10 productivity, and rich biodiversity, wetlands have been viewed as having a major
11 influence on many global biogeochemistries, including the carbon and nitrogen cycles
12 [10, 11] in addition to impacting on As transfer [2].

13 Owing to their depositional characteristics, wetlands are also highly susceptible
14 to contamination by heavy metals and other toxic trace elements. Macro-scale
15 processes (physical and geological events), which include anthropogenic activities,
16 wet, dry deposition, erosion, volcanism, and weathering are essential in determining
17 As loading in these environments (Figure 1). Even seemingly 'pristine' wetland
18 habitats/ecosystems, with Yellowstone National Park being a prime example [12], can
19 be 'naturally' enriched in As. However, anthropogenic activities such as mining,
20 smelting, urban waste and wastewater discharges, fertilizer and pesticide use also
21 contribute significantly to the build-up of As stores in wetland [13]. Often,
22 disconnected from the surrounding geology, these human induced redistributions of

1 As can be the hardest to predict and hence pose perhaps the greatest hazard to human
2 health.

3 Despite the importance of macro-scale processes in determining As loading into
4 wetlands, it is micro-scale biogeochemistry, which includes desorption, dissolution
5 [14] and microbial mediated As^V reduction, As^{III} oxidation and methylation [2] that
6 mainly regulate the fluxes of As from the semi-terrestrial to either aquatic or
7 atmospheric compartments (Figure 1). Recent advances in molecular methods have
8 enabled great leaps in our understanding of microbial mediated As biotransformations.
9 This review summarizes the role of wetland microbiology within the global As cycle,
10 and the distribution and behavior of As in water submerged soils and sediments.
11 Moreover, it provides a detailed overview of the key transformation **biochemistries**,
12 ranking the events in order of priority and discussing their interactions within a
13 chemically complex and heterogeneous localized environment. Finally, an outlook for
14 future research areas and introduction to emerging new technologies for measuring As
15 cycling in wetlands is presented.

16

17 **2 Distribution and behavior of As in wetlands**

18 **2.1 Overview**

19 Wetlands can be broadly classified as either '*natural*', which includes marine, **coastal**
20 zone and inland wetlands, or '*artificial*' wetlands such as paddy soils. Although, the
21 true state of most systems is perhaps as a mixture/intermediary of the two. The
22 definition of what constitutes, as a 'contaminated sediment' is similarly quite open to

1 interpretation, as the acceptable quality criteria for As in wetland soils varies between
2 countries (Table 2). For example, the Canadian Council of Ministers of the
3 Environment (CCME), have set interim quality guidelines (ISQGs) for
4 marine/estuarine sites at 7.24 mg kg^{-1} dry weight [15], whereas trigger thresholds in
5 China are marked slightly higher, so a site is only given contamination status when
6 concentrations of 20 mg kg^{-1} are breached [16]. An alternative method to evaluate the
7 quality of a wetland, is to consider the As concentration in extractable waters.
8 Recently reported incidences where detected As in overlying surface waters exceeded
9 the US Environmental Protection Agency's (EPA) maximum containment level of As
10 in drinking water ($10 \text{ } \mu\text{g L}^{-1}$) are summarized in Table 3.

11 Perhaps the focal point of As contamination in wetlands, where to date most of
12 the scientific research attention has been directed lies in southeast Asia [17-21]. Here
13 the As source originally derived from eroded Himalayan rock, now resides as buried
14 lens of sediment making up the Bay of Bengal Delta. These near-surface alluvia are
15 the cause of As release to aquifers, and threaten tens of millions of people consuming
16 these waters [22]. In samples collected from fifteen selected Bhaluka wetlands in
17 Bangladesh, the concentration of As in the waters ranged from 7 to $80 \text{ } \mu\text{g L}^{-1}$, with 93%
18 found exceeding WHO recommended permissible limits [23]. For further information
19 on the Bangladesh As crisis, refer to reviews by Meharg *et al.* [24] and Meharg &
20 Zhao [25].

21 Paddy field contamination in the region is also a serious problem, here high soil
22 As concentrations are linked to the quality of the groundwaters used for irrigation [20].

1 This problem was uncovered in 2003, when seventy-one soils collected from three
2 districts, revealed a range of As concentrations in soils spanning 3.1 mg kg⁻¹ (*baseline*)
3 up to 42.5 mg kg⁻¹ [20]. While a later study by Stroud *et al.* [26], reported an even
4 greater variation in total As concentrations in twelve paddy soils from Bangladesh,
5 again similar baselines of 4 mg kg⁻¹, but this time the inputs of As from irrigation
6 caused soils to reach 138 mg kg⁻¹. This highlights the ease in which As can
7 accumulate in agricultural-wetland soils, which then leads to the direct transfer to
8 food supply chains [20, 27].

9 China, meanwhile, is another country that has extensively studied the problem of
10 As build-up in wetlands [28-31]. Here though the main inputs derive not from
11 groundwaters, but instead take many different forms/guises. Although, often difficult
12 to characterize due to the mixing of multiple and varying sources, they do share a
13 commonality, in that inputs have intensified as the country had modernized. For
14 example, freshwater and salt marsh sediments in the Yellow river delta, before the
15 Xiaolangdi reservoir was built, were already As enriched, recording values of 30 mg
16 kg⁻¹. However, after it was constructed, changes to the sediment flows coupled with
17 increased human settlement and activity in the area, resulted in further As deposition
18 taking the concentration up to 45 mg kg⁻¹ [32], equivalent to a 4.5-fold increase from
19 baseline levels for soils in the Yangtze river delta [33].

20 Like Bangladesh, paddy field contamination is also a topic of importance in
21 China, with numerous studies highlighting this as a prominent issue, see Table 3
22 [28-31, 34]. However, total As concentrations only tell part of the story. Here As

1 species are essential to understand the bioavailability and uptake of As in rice. Based
2 on soil pore waters collected from Bangladesh and Chinese paddy soils, As^{III} is the
3 dominant species, accounting for 66-100% of the total As [26]. Methylated As species
4 (DMAs^V and MMAs^V) are also commonly detected in rice soils [34]. Although the
5 uptake of organic As into rice is not as rapid as As^{III}, once inside roots they quickly
6 translocate to the above-ground tissues [25]. Volatile TMAs^{III} has also been recently
7 detected in field studies of paddy soils from Spain and Bangladesh, albeit only at
8 trace-concentrations [35].

9 Far from just being confined to Asia, the problem of As contamination in
10 **wetlands** is a feature on other continents as well. For example, concentrations of total
11 As in twenty-five surface sediments (0-4 cm) collected from wetlands in
12 Massachusetts in the U.S., ranged from 20 to 2100 mg kg⁻¹ [36]. In southwest Spain,
13 the contamination by As in the Guadalquivir marshes, arising from the mine-tailing
14 spill in Aznalcollar, led to severe wetland soil pollution, with far reaching biological
15 effects for local wildlife, a subject covered in depth by a number of related
16 publications (Table 3) [37-39].

17

18 **2.2 Biogeochemistry of As transformation and transport in wetlands**

19 The predominant form of As in soil prior to flooding is As^V. Bound commonly to iron
20 (oxy)hydroxides, due in equal measure to it's high affinity for the species and
21 abundance in most soils, the lability and subsequently toxicity of As^V is relatively
22 low [40]. Saturating the system with water causes soil redox potentials to rapidly

1 decrease, in response to the depletion of electron acceptors, such as oxygen, nitrate,
2 manganese oxides and iron oxides/hydroxides. When Eh drops to below -200 mV
3 porewater As^V concentrations decline as the species is transformed to As^{III} [25]. In the
4 short-term, buffering can occur, because in these highly reducing conditions iron
5 oxides/hydroxides are solubilized, releasing As^V. However, As^{III} development will
6 dominate overtime [25, 41, 42]. This phenomenon can be further enhanced, by the
7 direct reduction of soil reservoirs of As^V. Due to As^{III} being less strongly bound to soil
8 particle surfaces than As^V, it partitions more readily into solution phase, increasing the
9 overall bioavailability of the toxin [40, 43].

10 Organic carbon (OC) has also been shown to mobilize As in aquifers in south
11 Bangladesh [44], while high concentrations of dissolved organic matter (DOM) in soil
12 porewaters can compete with surface absorbed As^V and As^{III} displacing them into
13 solution [21, 25]. Other important chemistries that control As transport, include the
14 grouping of interactions that As has with sulfur rich minerals [45]. Under highly
15 reducing conditions, and in the presence of dissolved sulfide, As^{III} can form stable
16 complexes, for example, orpiment (As₂S₃) or realgar (As₄S₄) [36]. Immobilizing As
17 by controlling redox conditions, to favor As-S precipitation, is a practice used in
18 constructed wetlands for reducing As bioavailability [46]. However, the sites still
19 require careful management, as oxidation of the sediment can result in the dissolution
20 of sulfides and re-mobilization of As^{III} back into the environment [47, 48].

3 Mechanisms of microbial mediated As biotransformation in wetlands

Compared to chemical change, microbial mediated As redox reactions occur far more rapidly [49]. Study has showed As^V reduction in soils to be significantly suppresses when γ -irradiated, highlighting the dominant role the microbiota play in this in solum As transfer step [40]. Microbial activity can be increased/simulated by enhancing the availability of OC sources, a response that differs between species/communities. This in turn contributes to different rates and forms of As biotransformation, impacting on both inorganic and methylated As trends [28]. Indeed, most methylated As species detected in wetlands are originally derived from microbial mediated As reactions [31, 35], because plants are not efficient at methylating inorganic As [50]. These different As species with their ranging properties/characteristics define the manner that As is transferred back and forth from the main land and aquatic stores [51-54]. Some volatile As is released into the atmosphere, in the form of AsH₃, CH₃AsH₂ (MeAsH₂), (CH₃)₂AsH (Me₂AsH), and (CH₃)₃As (TMA^{III}As). However compared to the land and water exchanges the fluxes are relatively minor [28, 35].

The evolution of such effective and diverse microbial mediated As biotransformation systems arose because they protect against As toxicity. The pathways of how microbes deal with As have been extensively studied in the last couple of decades, revealing various genes and enzymes responsible for As biotransformation and its biotransportation out of cells [50, 55-57]. The main systems used in As biotransformation by microbes in wetlands are further summarized below.

3.1 As^{III} oxidation in wetlands

Rapid As^{III} oxidation in the environment is mediated by metabolic microbial processes [1, 58], which are catalyzed by As^{III} oxidase (Aio). This enzyme is encoded by *aioA* and *aioB* genes for the two subunits AioA (AoxB) and AioB (AoxA) [55, 59]. Microbial mediated As^{III} oxidation is considered as one of the primary As detoxification mechanisms for microbes because it can oxidize As^{III} to the less toxic As^V [55]. The high abundance of *aioA* genes in paddy soils under flooding conditions further suggests the importance of this pathway [31, 60]. As^{III} oxidizing bacteria in paddy soils mainly assign to the following family groupings: *Phyllobacteriaceae*, *Bradyrhizobiaceae*, *Methylobacteriaceae*, *Rhizobiales*, *Burkholderiales* and *Comamonadaceae* [60, 61].

In other types of wetland, for example coastal sediments, a high diversity of *aioA*-like genes have also been detected. The most abundant *aioA*-like genes derived from *Roseobacter litoralis* Och 149 [62]. As^{III} oxidizing bacteria isolated from natural and constructed wetlands in the Republic of Korea showed that they were all able to grow in the presence of high concentrations of As^{III} (10 mM). The bacterium identified as *Pseudomonas stutzeri* strain GIST-BDAN2 showed an especially high activity of As^{III} oxidation (completely oxidized 1 mM As^{III} to As^V within 25-30 h) and possessed both the *aoxB* and *aoxR* genes [63]. Three As^{III} oxidizing bacteria including *Agrobacterium tumefaciens*, *Pseudomonas fluorescens*, and *Variovorax paradoxus*-like organisms were also isolated from the same Madison River Valley soils that also supported large populations of As^V reducing bacteria [64]. This

1 highlights the close overlay of the microbial catalyzed oxidation and reduction
2 pathways, and how this dichotomy functions as the principal 'switch' which controls
3 As fate in a specific site or zone within a site.

5 **3.2 As^V reduction in wetlands**

6 Microbial mediated As^V reduction segregates into two functional schemes, the
7 respiratory and the detoxification pathway. The respiratory pathway is catalyzed by
8 the As^V respiratory reductase (ArrA) complex, which consists of a large catalytic
9 subunit (ArrA) and a small subunit (ArrB) [65-67]. It can mediate anaerobic
10 dissimilatory As^V reduction and couple this with energy production [68-70]. A more
11 universal system of As^V detoxification reduction, is catalyzed by the cytoplasmic As^V
12 reductase (ArsC), which is present in both aerobic and anaerobic microbes [71]. The
13 *arsC* gene occurs in the *ars* operons next to the *arsB* gene that functions as an As^{III}
14 membrane pump in most bacteria [56], controlling cytoplasmic As^V reduction, and
15 thereafter As^{III} efflux.

16 In wetland sediments, microbial mediated As^V reduction is considered to be a
17 crucial mechanism controlling As^V mobilization [72], and contributing factor leading
18 to As contamination of ground water, as has happened in south and southeast Asia [22,
19 73]. Due to the anaerobic conditions encountered in wetlands, dissimilatory As^V
20 respiring bacteria are quite commonplace. The dissimilatory As^V reduction gene *arrA*
21 has been detected in various wetlands, including paddy soils [60, 61], coastal
22 sediments from south China [62], and estuarine sediments from Chesapeake Bay in

1 the U.S. [74]. Zhang *et al.* [34] revealed that *arrA* sequences detected in paddy soils
2 were analogous to those found in *Geobacter* species, which have been frequently
3 found in As rich sediments before [72, 75]. Similarly, *Geobacter* are prominent in
4 Japanese paddy soil [54]. Observations of *Geobacter* species OR-1 have revealed that
5 in addition to As, ferrihydrite is also being used as an electron acceptor, thus
6 catalyzing the dissolution of As from As^V-absorbed ferrihydrite, by promoting ferrous
7 iron formation [54]. Moreover, As K-edge X-ray absorption near-edge structure
8 analysis demonstrated the OR-1 can also reduce soil/solid bound As^V directly. In
9 addition to *Geobacter*-related bacteria, other microbiota, such as *Shewanella* species,
10 that utilize the coupling/uncoupling of iron and As reduction as a means of acquiring
11 energy, are also capable of dissimilatory As^V reduction [76, 77].

12 Similar to *arrA*, *arsC* has also been frequently detected in paddy soils [60, 61]
13 and coastal sediments [62]. The As^V reducing microbes with the most active
14 detoxification pathways in paddy soils mostly belonged to typically rhizospheric
15 bacteria groups, such as *Rhizobiales* and *Pseudomonadales* [61]. Study of pure culture
16 bacteria from Madison River Valley soils revealed five isolates, *Agrobacterium*
17 *tumefaciens*, *Flavobacterium* sp., *Microbacterium* sp. and two *Arthrobacter* sp.-like
18 organisms capable of rapidly reducing As^V under aerobic conditions, with the later
19 found to possess new putative *arsC* genes [64]. Seventeen As-resistant bacteria
20 isolated from Mandovi and Zuari estuarine water systems revealed *arsA* (ATPase),
21 *arsB* and *arsC* genes on their plasmid DNA. While, *arsC* genes were individually
22 detected in thirteen bacterial isolates, including the genera's *Halomonas* and

1 *Acinetobacter* [78]. Using genome sequencing of *Bacillus* bacterium isolated from
2 Andean wetlands in northern Chile, the *arsC* gene function was found as
3 detoxification As^V reduction [79]. Together with the As^V respiring bacteria, these
4 detoxification As^V reducing microbes catalyze the reduction of both the soil absorbed
5 As^V and dissolved As^V in solution into As^{III}.

6

7 **3.3 As^{III} methylation and volatilization in wetlands**

8 Organic As is introduced primarily into the environment via the microbial catalyzed
9 methylation of As^{III}. The As^{III} S-adenosylme-thionine methyltransferase (ArsM),
10 which is responsible for As^{III} methylation has recently been found to be common to
11 many different microbes [12, 80-82]. It is encoded by *arsM* genes, and **catalyzes** the
12 generation of less toxic organic As species, such as MMAs^V, DMAs^V and TMAO.
13 The volatile As gas including MeAsH₂, Me₂AsH and TMAAs^{III} could also be generated
14 during the biomethylation processes. To date, TMAAs^{III} is the most commonly detected
15 volatile As species in the natural environment [28, 35] and under *in vitro* conditions
16 during pure culture [81-83].

17 In microcosm experiments using As contaminated paddy soils, Huang *et al.* [28]
18 and Jia *et al.* [29] found evidence of both microbial mediated As^{III} methylation and
19 volatilization. Recently discovered universal primers for the amplification of *arsM*
20 genes have uncovered *arsM* sequences as being widespread in numerous paddy soils
21 [29, 61]. Continued investigation reveals these *arsM* sequences mainly derive from
22 the phyla *Gemmatimonadales*, *Firmicutes*, *Actinobacteria*, and *Proteobacteria* and the

1 domain *Archaea* [61]. By applying a metagenomic approach, Cai *et al.* [62] also
2 found *arsM* genes in coastal sediments, but the dominant bacteria species involved in
3 As^{III} methylation were not identified in this study. Furthermore the ability to
4 methylate and volatilize As^{III} has been discovered in three cyanobacteria species that
5 are common in paddy soils [82], and a small free-living eukaryote *Ostreococcus tauri*
6 found in coastal waters [83]. In general, to the best of our knowledge, the study of
7 As^{III} methylation and volatilization in wetlands is still rather limited, but given the
8 rapid advance in our understanding of the importance of this aspect of the As
9 biogeochemical cycle, this is an area that will likely be of increasing interest in the
10 future.

11

12 **3.4 Environmental factors affecting As biotransformation microbes**

13 Various environmental factors affecting the diversity, behavior and metabolism of As
14 biotransformation microbes have been observed in wetlands. One of the most
15 important is the As concentration of the soils/sediments with As biotransformation
16 microbe abundance found to correlate positively with As concentration in paddy soils
17 [61]. Similarly, *arsM* gene abundance and the concentration of methylated As in
18 paddy soil solutions also exhibit a strong positive correlation [29]. The amount and
19 form of OM in wetlands is also key, for example, several experiments carried out in
20 microcosms using paddy soils have demonstrated that the amendment of OM
21 promotes the activity of As^{III} oxidizing bacteria [28]. The application of rice straw can
22 increase the community diversity of As^{III} oxidizing bacteria [60], and the activity of

1 As^{III} methylation microbes in paddy soils [28, 29]. Straw also has an impact on As^V
2 detoxification reducing microbes, acting to moderately simulate their activity, but
3 greatly enhance the total abundance of As^V reducers [28].

4 The concentration of SO₄²⁻ in paddy soils strongly influences microbial
5 community compositions, modifying both As^{III} oxidation and As^V detoxification
6 reduction activity. It is postulated that this is due to the possible sharing of microbial
7 groups of sulfur oxidation and As redox [61]. The microbial community composition
8 of bacteria/archaea involved in As biotransformations is also particularly sensitive to
9 iron concentrations. This is reasonable, especially considering that the
10 absorption-desorption dynamics of iron /oxyhydroxides and As^V influence the
11 concentration and species of bioavailable As in wetlands [41, 42]. For the microbial
12 communities carrying out As^{III} methylation, it is the available NO₃⁻-N, NH₄⁺-N, that
13 appears to be the most sensitive environmental parameter influencing community
14 composition [61]. Other factors, such as pH, redox potential, alkalinity, temperature
15 and dissolved oxygen, also contribute and influence As biotransformation microbial
16 abundance and activity [46]. However, investigation into this topic is still in its
17 infancy. Much remains to be done to improve our understanding of how we can
18 optimize the function of microbial mediated As biotransformations by changing the
19 management of local wetland environments. A significant hurdle, blocking this goal
20 has been the lack of suitable methods to observe and measure the dynamic
21 environmental conditions, at scales appropriate to the biota driving the change.
22 However, with the development of new multi-parameter, visualization tools such as

diffusive gradients in thin films (DGT)/planar optode sandwich sensors [84, 85] and advanced radiography and isotope imaging, *in situ* mapping of As and other key elements is now possible. Employing these technologies concurrently with 2D enzyme activity plots (zymography) and microbial imaging, i.e. fluorescence *in situ* hybridization (FISH) or oligonucleotides labeled probes with light emitting chromophores [86], offers further exciting possibilities to unravel the complex environment-biota interactions impacting on wetland As release.

4 Concluding remarks and perspectives

Microbes act as the ‘switch’ turning a wetland from an As sink to source through As^V reduction, As^{III} oxidation, As^{III} methylation, and As volatilization processes (Figure 1). As^V reducing bacteria/archaea can enhance As mobilization from sediments to water [54, 75], while As^{III} oxidizing microbes act to decrease As mobility and bioavailability [3]. The two groups of As^V reducers and As^{III} oxidizers act in tandem, controlling the direction/path of the inorganic As cycle. While wetland As^{III} methylators create organic As species (MMAs^V, DMAs^V, TMAs^{III}), they also generate volatile As, promoting its release into the atmosphere [2].

Considering the key role wetlands play in delivering essential ecosystems services and the large area they occupy, the microbial mediated As biotransformations they host/support remain indispensable components of a ‘safe’ global As biogeochemical cycle, one that functions in the best interests of humankind. Failure to live in balance with As’ complex biogeochemistry can have devastating consequences,

1 from the release of As from near-surface wetland sediments to underground drinking
2 water supplies [72], to its transfer into rice plants [41].

3 Better management of wetland resources for optimized microbial control of As
4 could potentially yield huge gains in public and environmental health. However more
5 research is needed to improve our understanding of how to make best use of these
6 microbial mediated reactions. We recommend research attention is immediately
7 directed at gaps in our understanding of wetland As cycling, such as microbial
8 organic-As degradation, as the literature available on this topic is scant. Furthermore,
9 continued improvements to the technologies available to study *in situ*, both the
10 chemistry and biology of the mercurial wetland environment is needed.

11

12 **Acknowledgements**

13 The authors declare no conflicts of interest and financial disclosures. We would like to
14 thank the Chinese Academy of Sciences President's International Fellowship Initiative
15 (CAS-PIFI 2016VEC001) and China Postdoctoral Science Foundation (No.
16 212400241).

17

References

1. Oremland R S, J F Stolz. The ecology of arsenic. *Science*, 2003, 300(5621): 939-944.
2. Zhu Y-G, M Yoshinaga, F-J Zhao, B P Rosen. Earth abides arsenic biotransformations. *Annual Review of Earth and Planetary Sciences*, 2014, 42(0): 443-467.
3. Bhattacharya P, A H Welch, K G Stollenwerk, M J McLaughlin, J Bundschuh, G Panaullah. Arsenic in the environment: Biology and Chemistry. *Science of the Total Environment*, 2007, 379(2-3): 109-120.
4. Bentley R, T G Chasteen. Microbial methylation of metalloids: arsenic, antimony, and bismuth. *Microbiology and Molecular Biology Reviews*, 2002, 66(2): 250-271.
5. Silbergeld E K, K Nachman. The environmental and public health risks associated with arsenical use in animal feeds. *Annals of the New York Academy of Sciences*, 2008, 1140(1): 346-357.
6. Murray L A, A Raab, I L Marr, J Feldmann. Biotransformation of arsenate to arsenosugars by *Chlorella vulgaris*. *Applied organometallic chemistry*, 2003, 17(9): 669-674.
7. Moore J W, S Ramamoorthy. Heavy metals in natural waters: applied monitoring and impact assessment. 2012: Springer Science & Business Media.
8. Zedler J B, S Kercher. Wetland resources: status, trends, ecosystem services, and restorability. *Annu. Rev. Environ. Resour.*, 2005, 30: 39-74.
9. Keddy P A. Wetland ecology: principles and conservation. 2010: Cambridge University Press.
10. Chmura G L, S C Anisfeld, D R Cahoon, J C Lynch. Global carbon sequestration in tidal, saline wetland soils. *Global biogeochemical cycles*, 2003, 17(4).
11. Wang S, Y Wang, X Feng, L Zhai, G Zhu. Quantitative analyses of ammonia-oxidizing Archaea and bacteria in the sediments of four nitrogen-rich wetlands in China. *Applied microbiology and biotechnology*, 2011, 90(2): 779-787.
12. Qin J, C R Lehr, C Yuan, X C Le, T R McDermott, B P Rosen. Biotransformation of arsenic by a Yellowstone thermoacidophilic eukaryotic alga. *Proceedings of the National Academy of Sciences*, 2009, 106(13): 5213-5217.
13. Bhakta J N, Y Munekage. Spatial distribution and contamination status of arsenic, cadmium and lead in some coastal shrimp (*Macrobrachium rosenbergii*) farming ponds of Viet Nam. *Pacific Journal of Science and Technology*, 2009, 11: 606-15.
14. Wang S, C N Mulligan. Occurrence of arsenic contamination in Canada: sources, behavior and distribution. *Science of the Total Environment*, 2006, 366(2-3): 701-721.
15. ENVIRONMENT C C O M O T. Canadian Sediment Quality Guidelines for the Protection of Aquatic Life: Summary Tables. *Canadian Environmental Quality Guidelines*, 1999. 2001, Canadian Council of Ministers of the Environment Winnipeg.
16. Li Q, Z Wu, B Chu, N Zhang, S Cai, J Fang. Heavy metals in coastal wetland sediments of the Pearl River Estuary, China. *Environmental Pollution*, 2007, 149(2): 158-164.
17. Huq S I, A Rahman, N Sultana, R Naidu. Extent and severity of arsenic contamination in soils of Bangladesh. *Fate of arsenic in the environment*. Dhaka: Bangladesh University of Engineering and Technology, 2003: 69-84.
18. Huq S I, J U M Shoaib. Soils and humans. in *The Soils of Bangladesh*. 2013, Springer. p. 125-129.
19. Lu Y, E E Adomako, A Solaiman, M R Islam, C Deacon, P Williams, G Rahman, A A Meharg.

- 1 Baseline soil variation is a major factor in arsenic accumulation in Bengal Delta paddy rice.
2 Environmental science & technology, 2009, 43(6): 1724-1729.
- 3 20. Meharg A A,M M Rahman. Arsenic contamination of Bangladesh paddy field soils:
4 implications for rice contribution to arsenic consumption. Environmental Science &
5 Technology, 2003, 37(2): 229-234.
- 6 21. Williams P N, H Zhang, W Davison, A A Meharg, M Hossain, G J Norton, H Brammer,M R
7 Islam. Organic Matter Solid Phase Interactions Are Critical for Predicting Arsenic Release
8 and Plant Uptake in Bangladesh Paddy Soils. Environmental science & technology, 2011,
9 45(14): 6080-6087.
- 10 22. Polizzotto M L, B D Kocar, S G Benner, M Sampson,S Fendorf. Near-surface wetland
11 sediments as a source of arsenic release to ground water in Asia. Nature, 2008, 454(7203):
12 505-508.
- 13 23. Alam M, M Ali, N A Al-Harbi,T R Choudhury. Contamination status of arsenic, lead, and
14 cadmium of different wetland waters. Toxicological & Environmental Chemistry, 2011, 93(10):
15 1934-1945.
- 16 24. Meharg A. Venomous earth: How arsenic caused the world's worst mass poisoning. 2005.
- 17 25. Meharg A A,F-J Zhao. Biogeochemistry of Arsenic in Paddy Environments. in Arsenic & Rice.
18 2012, Springer. p. 71-101.
- 19 26. Stroud J L, M A Khan, G J Norton, M R Islam, T Dasgupta, Y-G Zhu, A H Price, A A Meharg,
20 S P McGrath,F-J Zhao. Assessing the labile arsenic pool in contaminated paddy soils by
21 isotopic dilution techniques and simple extractions. Environmental science & technology,
22 2011, 45(10): 4262-4269.
- 23 27. Williams P N, A Villada, C Deacon, A Raab, J Figuerola, A J Green, J Feldmann,A A Meharg.
24 Greatly enhanced arsenic shoot assimilation in rice leads to elevated grain levels compared to
25 wheat and barley. Environmental Science & Technology, 2007, 41(19): 6854-6859.
- 26 28. Huang H, Y Jia, G X Sun,Y G Zhu. Arsenic speciation and volatilization from flooded paddy
27 soils amended with different organic matters. Environmental science & technology, 2012,
28 46(4): 2163-2168.
- 29 29. Jia Y, H Huang, M Zhong, F-H Wang, L-M Zhang,Y-G Zhu. Microbial arsenic methylation in
30 soil and rice rhizosphere. Environmental science & technology, 2013, 47(7): 3141-3148.
- 31 30. Williams P N, M Lei, G Sun, Q Huang, Y Lu, C Deacon, A A Meharg,Y-G Zhu. Occurrence
32 and partitioning of cadmium, arsenic and lead in mine impacted paddy rice: Hunan, China.
33 Environmental science & technology, 2009, 43(3): 637-642.
- 34 31. Zhao F-J, E Harris, J Yan, J Ma, L Wu, W Liu, S P McGrath, J Zhou,Y-G Zhu. Arsenic
35 methylation in soils and its relationship with microbial *arsM* abundance and diversity, and As
36 speciation in rice. Environmental science & technology, 2013, 47(13): 7147-7154.
- 37 32. Bai J, R Xiao, K Zhang,H Gao. Arsenic and heavy metal pollution in wetland soils from tidal
38 freshwater and salt marshes before and after the flow-sediment regulation regime in the
39 Yellow River Delta, China. Journal of Hydrology, 2012, 450: 244-253.
- 40 33. Gorenc S, R Kostaschuk,Z Chen. Spatial variations in heavy metals on tidal flats in the
41 Yangtze Estuary, China. Environmental Geology, 2004, 45(8): 1101-1108.
- 42 34. Zhang S-Y, F-J Zhao, G-X Sun, J-Q Su, X-R Yang, H Li,Y-G Zhu. Diversity and abundance of
43 arsenic biotransformation genes in paddy soils from Southern China. Environmental science
44 & technology, 2015, 49(7): 4138-4146.

- 1 35. Mestrot A, J Feldmann, E M Krupp, M S Hossain, G Roman-Ross,A A Meharg. Field fluxes
2 and speciation of arsines emanating from soils. *Environmental science & technology*, 2011,
3 45(5): 1798-1804.
- 4 36. Wilkin R T,R G Ford. Arsenic solid-phase partitioning in reducing sediments of a
5 contaminated wetland. *Chemical Geology*, 2006, 228(1): 156-174.
- 6 37. Grimalt J O, M Ferrer,E Macpherson. The mine tailing accident in Aznalcollar. *Science of the*
7 *Total Environment*, 1999, 242(1): 3-11.
- 8 38. Kraus U,J Wiegand. Long-term effects of the Aznalcóllar mine spill—heavy metal content and
9 mobility in soils and sediments of the Guadiamar river valley (SW Spain). *Science of the Total*
10 *Environment*, 2006, 367(2): 855-871.
- 11 39. Mateo R, M A Taggart, A J Green, C Cristòfol, A Ramis, H Lefranc, J Figuerola,A A Meharg.
12 Altered porphyrin excretion and histopathology of greylag geese (*Anser anser*) exposed to soil
13 contaminated with lead and arsenic in the Guadalquivir Marshes, southwestern Spain.
14 *Environmental Toxicology and Chemistry*, 2006, 25(1): 203-212.
- 15 40. Yamaguchi N, T Nakamura, D Dong, Y Takahashi, S Amachi,T Makino. Arsenic release from
16 flooded paddy soils is influenced by speciation, Eh, pH, and iron dissolution. *Chemosphere*,
17 2011, 83(7): 925-932.
- 18 41. Xu X, S McGrath, A Meharg,F Zhao. Growing rice aerobically markedly decreases arsenic
19 accumulation. *Environmental science & technology*, 2008, 42(15): 5574-5579.
- 20 42. Cummings D E, F Caccavo, S Fendorf,R F Rosenzweig. Arsenic mobilization by the
21 dissimilatory Fe (III)-reducing bacterium *Shewanella alga* BrY. *Environmental Science &*
22 *Technology*, 1999, 33(5): 723-729.
- 23 43. Takahashi Y, R Minamikawa, K H Hattori, K Kurishima, N Kihou,K Yuita. Arsenic behavior
24 in paddy fields during the cycle of flooded and non-flooded periods. *Environmental science &*
25 *technology*, 2004, 38(4): 1038-1044.
- 26 44. Harvey C F, C H Swartz, A Badruzzaman, N Keon-Blute, W Yu, M A Ali, J Jay, R Beckie, V
27 Niedan,D Brabander. Arsenic mobility and groundwater extraction in Bangladesh. *Science*,
28 2002, 298(5598): 1602-1606.
- 29 45. Bostick B C, C Chen,S Fendorf. Arsenite retention mechanisms within estuarine sediments of
30 Pescadero, CA. *Environmental science & technology*, 2004, 38(12): 3299-3304.
- 31 46. Lizama K, T D Fletcher,G Sun. Removal processes for arsenic in constructed wetlands.
32 *Chemosphere*, 2011, 84(8): 1032-1043.
- 33 47. Morse J W. Interactions of trace metals with authigenic sulfide minerals: implications for their
34 bioavailability. *Marine Chemistry*, 1994, 46(1): 1-6.
- 35 48. Saulnier I,A Mucci. Trace metal remobilization following the resuspension of estuarine
36 sediments: Saguenay Fjord, Canada. *Applied Geochemistry*, 2000, 15(2): 191-210.
- 37 49. Kirk G. The biogeochemistry of submerged soils. 2004: John Wiley & Sons.
- 38 50. Ye J, C Rensing, B P Rosen,Y G Zhu. Arsenic biomethylation by photosynthetic organisms.
39 *Trends in Plant Science*, 2012, 17(3): 155-162.
- 40 51. Maguffin S C, M F Kirk, A R Daigle, S R Hinkle,Q Jin. Substantial contribution of
41 biomethylation to aquifer arsenic cycling. *Nature Geoscience*, 2015.
- 42 52. Drahota P, L Falteisek, A Redlich, J Rohovec, T Matoušek,I Čepička. Microbial effects on the
43 release and attenuation of arsenic in the shallow subsurface of a natural geochemical anomaly.
44 *Environmental Pollution*, 2013, 180: 84-91.



- 1 53. Mumford A C, J L Barringer, W M Benzel, P A Reilly,L Young. Microbial transformations of
2 arsenic: mobilization from glauconitic sediments to water. Water research, 2012, 46(9):
3 2859-2868.
- 4 54. Ohtsuka T, N Yamaguchi, T Makino, K Sakurai, K Kimura, K Kudo, E Homma, D T Dong,S
5 Amachi. Arsenic dissolution from Japanese paddy soil by a dissimilatory arsenate-reducing
6 bacterium *Geobacter* sp. OR-1. Environmental science & technology, 2013, 47(12):
7 6263-6271.
- 8 55. Slyemi D,V Bonnefoy. How prokaryotes deal with arsenic†. Environmental microbiology
9 reports, 2012, 4(6): 571-586.
- 10 56. Silver S,L T Phung. Genes and enzymes involved in bacterial oxidation and reduction of
11 inorganic arsenic. Applied and environmental microbiology, 2005, 71(2): 599-608.
- 12 57. Rosen B P,Z Liu. Transport pathways for arsenic and selenium: a minireview. Environment
13 International, 2009, 35(3): 512-515.
- 14 58. Mukhopadhyay R, B P Rosen, L T Phung,S Silver. Microbial arsenic: from geocycles to genes
15 and enzymes. FEMS Microbiology Reviews, 2002, 26(3): 311-325.
- 16 59. Stolz J F, P Basu, J M Santini,R S Oremland. Arsenic and Selenium in Microbial Metabolism*.
17 Annu. Rev. Microbiol., 2006, 60: 107-130.
- 18 60. Jia Y, H Huang, Z Chen,Y-G Zhu. Arsenic uptake by rice is influenced by microbe-mediated
19 arsenic redox changes in the rhizosphere. Environmental science & technology, 2014, 48(2):
20 1001-1007.
- 21 61. Zhang S-Y, F Zhao, G Sun, J Su, X Yang, H Li,Y-G Zhu. Diversity and abundance of arsenic
22 biotransformation genes in paddy soils from Southern China. Environmental science &
23 technology, 2015.
- 24 62. Cai L, K Yu, Y Yang, B-w Chen, X-d Li,T Zhang. Metagenomic exploration reveals high
25 levels of microbial arsenic metabolism genes in activated sludge and coastal sediments.
26 Applied microbiology and biotechnology, 2013, 97(21): 9579-9588.
- 27 63. Chang J-S, I-H Yoon, J-H Lee, K-R Kim, J An,K-W Kim. Arsenic detoxification potential of
28 aox genes in arsenite-oxidizing bacteria isolated from natural and constructed wetlands in the
29 Republic of Korea. Environmental geochemistry and health, 2010, 32(2): 95-105.
- 30 64. Macur R E, C R Jackson, L M Botero, T R Mcdermott,W P Inskeep. Bacterial populations
31 associated with the oxidation and reduction of arsenic in an unsaturated soil. Environmental
32 science & technology, 2004, 38(1): 104-111.
- 33 65. Afkar E, J Lisak, C Saltikov, P Basu, R S Oremland,J F Stolz. The respiratory arsenate
34 reductase from *Bacillus selenitireducens* strain MLS10. FEMS microbiology letters, 2003,
35 226(1): 107-112.
- 36 66. Saltikov C W,D K Newman. Genetic identification of a respiratory arsenate reductase.
37 Proceedings of the National Academy of Sciences, 2003, 100(19): 10983-10988.
- 38 67. Krafft T,J M Macy. Purification and characterization of the respiratory arsenate reductase of
39 *Chrysiogenes arsenatis*. European Journal of Biochemistry, 1998, 255(3): 647-653.
- 40 68. van Lis R, W Nitschke, S Duval,B Schoepp-Cothenet. Arsenics as bioenergetic substrates.
41 Biochimica et Biophysica Acta (BBA)-Bioenergetics, 2013, 1827(2): 176-188.
- 42 69. Malasarn D, W Saltikov, K M Campbell, J M Santini, J G Hering,D K Newman. *arrA* is a
43 reliable marker for As(V) respiration. Science, 2004, 306(5695): 455-455.
- 44 70. Hoeft S E, T R Kulp, J F Stolz, J T Hollibaugh,R S Oremland. Dissimilatory arsenate

- 1 reduction with sulfide as electron donor: Experiments with mono lake water and isolation of
- 2 strain MLMS-1, a chemoautotrophic arsenate respirer. *Applied and Environmental*
- 3 *Microbiology*, 2004, 70(5): 2741-2747.
- 4 71. Bhattacharjee H,B P Rosen. Arsenic metabolism in prokaryotic and eukaryotic microbes.
- 5 *Molecular microbiology of heavy metals.*, ed. S.S. Nies DH. Vol. 6. 2007, Heidelberg,
- 6 Germany: Springer. 371-406.
- 7 72. Islam F S, A G Gault, C Boothman, D A Polya, J M Charnock, D Chatterjee,J R Lloyd. Role
- 8 of metal-reducing bacteria in arsenic release from Bengal delta sediments. *Nature*, 2004,
- 9 430(6995): 68-71.
- 10 73. Fendorf S, H A Michael,A van Geen. Spatial and temporal variations of groundwater arsenic
- 11 in South and Southeast Asia. *Science*, 2010, 328(5982): 1123-1127.
- 12 74. Song B, E Chyun, P R Jaffé,B B Ward. Molecular methods to detect and monitor dissimilatory
- 13 arsenate - respiring bacteria (DARB) in sediments. *Fems Microbiology Ecology*, 2009, 68(1):
- 14 108-117.
- 15 75. Héry M, B Van Dongen, F Gill, D Mondal, D Vaughan, R Pancost, D Polya,J Lloyd. Arsenic
- 16 release and attenuation in low organic carbon aquifer sediments from West Bengal.
- 17 *Geobiology*, 2010, 8(2): 155-168.
- 18 76. Oremland R S,J F Stolz. Arsenic, microbes and contaminated aquifers. *Trends in microbiology*,
- 19 2005, 13(2): 45-49.
- 20 77. Tufano K J, C Reyes, C W Saltikov,S Fendorf. Reductive processes controlling arsenic
- 21 retention: revealing the relative importance of iron and arsenic reduction. *Environmental*
- 22 *science & technology*, 2008, 42(22): 8283-8289.
- 23 78. Sunita M S L, S Prashant, P B Chari, S N Rao, P Balaravi,P K Kishor. Molecular identification
- 24 of arsenic-resistant estuarine bacteria and characterization of their ars genotype.
- 25 *Ecotoxicology*, 2012, 21(1): 202-212.
- 26 79. Vilo C, A Galetovic, J E Araya, B Gómez-Silva,Q Dong. Draft genome sequence of a *Bacillus*
- 27 bacterium from the Atacama Desert wetlands metagenome. *Genome announcements*, 2015,
- 28 3(4): e00955-15.
- 29 80. Qin J, B P Rosen, Y Zhang, G Wang, S Franke,C Rensing. Arsenic detoxification and
- 30 evolution of trimethylarsine gas by a microbial arsenite S-adenosylmethionine
- 31 methyltransferase. *Proceedings of the National Academy of Sciences*, 2006, 103(7):
- 32 2075-2080.
- 33 81. Wang P-P, G-X Sun,Y-G Zhu. Identification and characterization of arsenite methyltransferase
- 34 from an archaeon, *Methanosarcina acetivorans* C2A. *Environmental science & technology*,
- 35 2014, 48(21): 12706-12713.
- 36 82. Yin X, J Chen, J Qin, G Sun, B Rosen,Y Zhu. Biotransformation and volatilization of arsenic
- 37 by three photosynthetic cyanobacteria. *Plant Physiology*, 2011, 156(3): 1631-1638.
- 38 83. Zhang S-Y, G-X Sun, X-X Yin, C Rensing,Y-G Zhu. Biomethylation and volatilization of
- 39 arsenic by the marine microalgae *Ostreococcus tauri*. *Chemosphere*, 2013.
- 40 84. Williams P N, J Santner, M Larsen, N J Lehto, E Oburger, W Wenzel, R N Glud, W Davison,H
- 41 Zhang. Localized flux maxima of arsenic, lead, and iron around root apices in flooded lowland
- 42 rice. *Environmental science & technology*, 2014, 48(15): 8498-8506.
- 43 85. Guan D-X, P N Williams, J Luo, J-L Zheng, H-C Xu, C Cai,L Q Ma. Novel precipitated
- 44 zirconia-based DGT technique for high-resolution imaging of oxyanions in waters and

- 1 sediments. Environmental science & technology, 2015, 49(6): 3653-3661.
- 2 86. Oburger E,H Schmidt. New methods to unravel rhizosphere processes. Trends in plant science,
3 2016, 21(3): 243-255.
- 4 87. Chapman P M, F Wang, C Janssen, G Persoone,H E Allen. Ecotoxicology of metals in aquatic
5 sediments: binding and release, bioavailability, risk assessment, and remediation. Canadian
6 Journal of Fisheries and Aquatic Sciences, 1998, 55(10): 2221-2243.
- 7 16. National Standard of PR China, 2002. National Standard of PR China Marine
8 Sediment Quality (GB 18668-2002) Standards Press of China, Beijing (2002) (in
9 Chinese).

10

1 Table 1. Structure of prevalent As species in the environment.

Species	Structure of As speciation
Arsenite, As ^{III}	$\begin{array}{c} \text{HO}-\text{As}-\text{OH} \\ \\ \text{OH} \end{array}$
Aesenate, As ^V	$\begin{array}{c} \text{O} \\ \\ \text{HO}-\text{As}-\text{OH} \\ \\ \text{OH} \end{array}$
Methylarsonate, MMAs ^V	$\begin{array}{c} \text{O} \\ \\ \text{HO}-\text{As}-\text{OH} \\ \\ \text{CH}_3 \end{array}$
Dimethylarsinate, DMAs ^V	$\begin{array}{c} \text{O} \\ \\ \text{H}_3\text{C}-\text{As}-\text{OH} \\ \\ \text{CH}_3 \end{array}$
Thrimethylarsine oxide, TMAO	$\begin{array}{c} \text{O} \\ \\ \text{H}_3\text{C}-\text{As}-\text{CH}_3 \\ \\ \text{CH}_3 \end{array}$
Aesenosugars	
Arsenobetaine, AsB	

2

3

Table 2. The quality guidelines for As contamination in wetlands from different countries.

Sediment quality guidelines in various countries	Countries	Level	Arsenic concentration (mg kg ⁻¹)	References
Hongkong ISQVs	China	ISQV-low	8.2	[87]
		ISQV-high	70	
Sediment Quality Criteria	China	Class I	20	[16]
		Class II	65	
Canadian Environmental Quality Guideline	Canada	ISQGs	7.24	[15]
		PEL	41.6	

ISQV, interim sediment quality value; ISQGs, interim sediment quality guidelines; PEL: probable effect level.

Table 3. Summary of recently detected arsenic contaminated wetlands.

Wetland types	Country	Sampling Location	As concentration (mg kg ⁻¹ / µg L)	References
Coastal wetlands	China	Yellow River delta	38	[32]
			45	
		Yangtze River delta	10	[33]
Inland wetlands	U.S.	Massachusetts	20-2100	[36]
	Spain	Guadalquivir	20	[38]
		Fatehpur	72-114	
Paddy soils	Bangladesh	Dhumrakandi	62-138	[26]
		Paranpur	73-77	
		Faridpur	34	
	India	De Ganga	17	[61]
	China	Chenzhou	60	

		Qiyang	79	
		Anqing	19	
		Jiaxing	20	
		Yingtian	16	
		Jingzhou	19	
		Changde	16	
		Jiangmen	25	
		Guilin	21	
		Guiyang	21	
		Zhanjiang	18	
Wetland waters	Bangladesh	Malabar	60	[23]
		Gerajan	80	
		Dohuria	11-14	

Behi	11
Porabait	26
Barakhaillah	20
Jora	11
Uhila	18
Barakuri	11
Jerukuri	11
Bigaira	11

Figure caption

Figure 1. Sources of As introduced to wetland, microbial mediated As biotransformation in wetland, and genes responsible for As^V respiratory reduction, As^V detoxification reduction, As^{III} oxidation, and As^{III} methylation.