



**QUEEN'S
UNIVERSITY
BELFAST**

Social capital, deprivation and self-rated health: Does reporting heterogeneity play a role? Results from the English Longitudinal Study of Ageing

O'Doherty, M. G., French, D., Steptoe, A., & Kee, F. (2017). Social capital, deprivation and self-rated health: Does reporting heterogeneity play a role? Results from the English Longitudinal Study of Ageing. *Social Science & Medicine*, 179, 191. Advance online publication. <https://doi.org/10.1016/j.socscimed.2017.03.006>

Published in:
Social Science & Medicine

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2017 Elsevier Ltd. All rights reserved.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>, which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

1 Social capital, deprivation and self-rated health: does
2 reporting heterogeneity play a role? Results from the English
3 Longitudinal Study of Ageing

4
5 **Abstract**

6 Self-rated health (SRH) is commonly assessed in large surveys, though responses can be
7 influenced by different individuals' perceptions of and beliefs about health. Therefore,
8 instead of providing evidence of 'true' health disparities across groups, findings may actually
9 reflect reporting heterogeneity.

10 Using data from participants aged 50 years and older from the English Longitudinal
11 Study of Ageing (ELSA) Wave 3 (2006/07; participation rate =73%), associations between
12 three dimensions of social capital (local area & trust, social support and social networks),
13 deprivation and SRH were examined using the vignette methodology in 2341 individuals
14 who completed both the self-report and at least one of the 18 vignettes. Analysis employed
15 a hierarchical probit model (HOPIT).

16 Individuals expressing low local area & trust social capital (beta= -0.276, p<0.001)
17 and those with poor social networks (beta= -0.280, p<0.001) were more likely to report poor
18 SRH in HOPIT models accounting for reporting heterogeneity, but unadjusted ordered probit
19 analyses still correctly show a negative relationship between low local area & trust social
20 capital (beta= -0.243, p<0.001) and those with poor social networks (beta= -0.210, p<0.01),
21 though they somewhat tend to underestimate its strength. Neither social support nor
22 deprivation appeared to have any effect on SRH regardless of reporting heterogeneity.

23 Anchoring vignettes offer a relatively uncomplicated and cost-effective way of
24 identifying and correcting for reporting heterogeneity to improve comparative validity of
25 self-report measures of health. This analysis underlines the need for caution when using
26 unadjusted self-reported measures to study the effects of social capital on health.

27

28 **Keywords**

29 Social capital; self-rated health; ageing; vignettes; health disparities; deprivation

30 **Word count:** 7988

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47 **Introduction**

48 Empirical evidence has consistently demonstrated a relationship between social
49 capital and self-rated health (SRH) (Chen and Meng, 2015; Giordano et al., 2012; Kawachi et
50 al., 1999; Kawachi and Berkman, 2014; Koutsogeorgou et al., 2015; Nieminen et al., 2013,
51 2010), but because there is no 'gold standard' of how to measure social capital, the strength
52 of the association is uncertain. A simple definition of social capital is: the “resources that are
53 accessed by individuals as a result of their membership of a network or a group” (Kawachi
54 and Berkman, 2014). While there is a debate around the conceptualisation of social capital
55 (Kawachi et al., 2004; Poortinga, 2006; Szreter and Woolcock, 2004) most agree that it is
56 multidimensional and that it carries different interpretations depending on who defines it
57 and on their disciplinary traditions.

58 Social capital has been suggested to improve health through norms and attitudes
59 that influence healthy behaviours, and psychosocial networks that increase access to health
60 care and mechanisms that enhance self-esteem (Kawachi et al., 1999; Kawachi and
61 Berkman, 2014; Lindström, 2008). Conversely, social capital can also have a negative impact
62 on health, including the promotion (but also cessation) of risky behaviours (e.g. smoking),
63 exchanging wrong information, the exclusion of ‘outsiders’, and downward-levelling norms
64 (Burt, 1992; Campos-Matos et al., 2016; Christakis and Fowler, 2008; Kawachi and Berkman,
65 2014; Rosenquist et al., 2011). Interventions have been conducted to evaluate social
66 capital’s impact on health with varying success (Coll-Planas et al., 2016).

67 While the literature is vast on the health effects of individual level disadvantage,
68 area level deprivation can affect health by increasing an individual’s sense of being deprived
69 of status, resulting in frustration, shame and stress, which in turn may lead to adverse
70 health consequences. On the other hand areas which are least deprived may be wealthier

71 and thus have more local facilities and resources which can have a positive impact on health
72 (Glymour et al., 2014; Zhang et al., 2013). However, rather than using instruments designed
73 for a specific purpose (Sánchez-Santos et al., 2013), different methods to measure
74 deprivation at area-level have been employed, making the strength of associations
75 uncertain. The Index of Multiple Deprivation (IMD) is a measure of deprivation at the lower
76 super output area (LSOA) which has been used in the UK since 2000 and is an instrument
77 specifically designed for such a purpose (Noble et al., 2004). It is based on the idea of
78 distinct dimensions of deprivation which can be recognised and measured separately, but
79 combined into an overall score.

80 To measure health in large cohort surveys, it is common that a subjective measure
81 based on self-report is employed, preferably in combination with the use of objective
82 measures. However, the latter may be too expensive to implement in large population
83 surveys. Nevertheless SRH has been shown to have robust associations with “hard”
84 outcomes such as mortality (Barger et al., 2016). Many international cohort studies have
85 employed measures of SRH collected sequentially over time, including the English
86 Longitudinal Study of Ageing (ELSA) (Stephens et al., 2013). However, the primary issue with
87 using such self-reports alone is that different individuals may have different beliefs and
88 perceptions about the concept of health. The comparability of self-reported information can
89 vary across social groups (within countries) or across countries because of: unequal access
90 to medical providers or health information; diagnosis avoidance (inadvertent or intentional
91 avoidance of medical screening/testing); or interpersonal incomparability across groups if
92 they use different reference groups or interpret questions or concepts differently (Burgard
93 and Chen, 2014). Researchers also usually have little insight as to what individuals are
94 actually thinking of when they assess their health (Au and Johnston, 2014). Therefore,

95 instead of providing evidence of 'true' health disparities, findings may actually reflect
96 *reporting heterogeneity*.

97 To help overcome the problems of interpersonal incomparability of subjective
98 measures, such as self-reports, King et al. 2004 proposed a technique using anchoring
99 vignettes(King et al., 2004). The vignettes were presented as a way to alleviate problems
100 which occur when different groups of participants understand and use the Likert scales for
101 self-reports in different ways (e.g. 1=very bad health to 5=very good health). This
102 heterogeneity in reporting styles is also known as differential item functioning (DIF).
103 Graphically, this problem is illustrated in Figure 1. Previous research has been conducted
104 into the use of anchoring vignettes to assess group differences in SRH (Au and Lorgelly,
105 2014; Grol-Prokopczyk et al., 2015, 2011; Peracchi and Rossetti, 2012; Xu and Xie, 2015), but
106 to our knowledge, only one so far has specifically used the anchoring vignettes technique to
107 improve comparability of SRH and social capital (Chen and Meng, 2015).

108 As outlined above, studies have demonstrated associations between social capital
109 and SRH. However, these studies did not take into account the possibility of reporting
110 heterogeneity distorting SRH disparities associated with social capital. Therefore, using
111 nationally representative data, we aim to better estimate the relationship between social
112 capital and SRH among English adults aged 50 years and older. By improving the
113 interpersonal comparability of SRH, we can conduct simulations to illustrate the potential
114 magnitude of the effect of reporting heterogeneity in estimating the distribution of SRH
115 from self-reported survey data.

116

117 **Methods**

118 *Population*

119 ELSA is a panel study of a representative cohort of men and women living in England
120 aged 50 years and older, and their partners of any age. It was designed as a sister study to
121 the Health and Retirement Study in the US and follows many of the same principles. The
122 study commenced in 2002, and the sample has been followed up every two years using
123 computer-assisted personal interviews and self-completion questionnaires, with an
124 additional nurse visit for the assessment of biomarkers every four years (main interview).
125 More detailed information on the design of ELSA can be found elsewhere(Steptoe et al.,
126 2013). Data for this current study is from ELSA Wave 3 (2006/07). The participation rate in
127 Wave 3 was 73% (total individual respondents to wave 3 divided by total individuals eligible
128 for wave 3). After excluding partners aged <50 years (n=428), 9343 main interviews were
129 completed. 2341 individuals also completed a module on self-completion health vignettes
130 (at least one of the 18 vignettes answered; covering the health domains pain, sleeping,
131 mobility, memory, breathing and depression) and a self-rated health question using a similar
132 five-point Likert scale.

133

134 *Self-rated health assessment*

135 SRH was collected during the main interview. Individuals were asked to rate their
136 own general health on a five-point Likert scale ('Would you say your health is...') which was
137 reverse coded to be increasing in good health (1= very poor to 5=very good).

138

139 *Social capital assessment*

140 The framework adopted by the Office for National Statistics (ONS) (Siegler, 2014)
141 and introduced by the Organisation for Economic Co-operation and Development (OECD)
142 (Scrivens and Smith, 2013) was used as a basis to select 21 different variables within ELSA

143 that could be used to represent social capital. In this framework, there are four different
144 aspects of social capital: [1] personal relationships; [2] social network support; [3] civic
145 engagement; and [4] trust and cooperative norms. Of the 21 variables selected, only two
146 mapped onto 'civic engagement' (member of at least one organisation, club or society and
147 voluntary work). Therefore, the 21 variables were reduced to three social capital dimensions
148 using factor analysis. The three dimensions were [1] local area & trust, [2] social support and
149 [3] social networks. Factor loadings ≥ 0.3 were retained. The factor loadings and dimensions
150 are outlined in Table S1 in the Online Supplementary Material.

151 The three dimensions created align well with the ONS framework. [1] "Local area &
152 trust" matches the concepts of trust and cooperative norms or shared values that shape the
153 way people behave towards each other and as members of society (nine factors; all found in
154 the same section of ELSA questionnaire - local area), [2] "social support" is closely related to
155 the level of resources or support that a person can draw from in their personal relationships
156 (six factors; all in reference to spouse/partner, children, family and friends), and [3] "social
157 networks" incorporates aspects of both "personal relationships" and "civic engagement" (six
158 factors). It includes variables which refer to both the structure and nature of people's
159 personal relationships (number of close relationships, meet ups/communication) and the
160 actions and behaviours that can be seen as contributing positively to the collective life of a
161 community or society (member of an organisation and volunteering). The composite
162 reliability {a test of internal consistency - measures the overall reliability of a collection of
163 heterogeneous but similar items} (Colwell, 2016) of the dimensions of social capital were
164 0.84 (local area & trust), 0.81 (social support) and 0.66 (social networks).

165 The three factor scores were divided into quintiles. The top quintile included persons
166 with high levels of social capital with regard to the dimension in question. Respectively, the

167 bottom quintile included those with the least social capital in that dimension. For the
168 purposes of this analysis, we created two separate dichotomised variables for each
169 dimension. High social capital coded as 1=top quintile and 0=bottom four quintiles. Low
170 social capital coded as 1=bottom quintile and 0=top four quintiles.

171

172 *Deprivation assessment*

173 ELSA deprivation data was obtained separately via an application process which was
174 approved by the NatCen Data Release Panel. IMD2004 is a measure of multiple deprivation
175 at the lower super output area (LSOA) (Noble et al., 2004). IMD2004 is based on the idea of
176 distinct dimensions of deprivation which can be recognised and measured separately.
177 People may be counted as deprived in one or more of the domains depending on the
178 number of types of deprivation that they experience. IMD2004 is conceptualised as a
179 weighted area level aggregation of these specific dimensions of deprivation: [1] income
180 deprivation; [2] employment deprivation; [3] health deprivation and disability; [4]
181 education, skills and training deprivation; [5] barriers to housing and services; [6] living
182 environment deprivation; [7] crime. Each dimension index consists of a score which is then
183 ranked. The higher the score, the more deprived is the LSOA. The IMD2004 scores were
184 provided from NatCen as quintiles. The top quintile included persons who were most
185 deprived. Respectively, the bottom quintile included those who were least deprived. For the
186 purposes of this analysis, we created two separate dichotomised variables. Most deprived
187 coded as 1=top quintile and 0=bottom four quintiles. Least deprived coded as 1=bottom
188 quintile and 0=top four quintiles.

189

190 *Vignette assessment*

191 The 18 vignettes within the health self-completion questionnaire are outlined in the
192 Online Supplementary Material. They cover several different health domains, including pain,
193 sleeping, mobility, memory, breathing and depression (three vignettes each). Briefly,
194 individuals were asked to rate the health limitations of various hypothetical persons who
195 experience different circumstances related to health on a five-point Likert scale. Possible
196 responses, once reverse coded to be increasing in good health, ranged from 1= extreme
197 health problem to 5=no health problem. Individuals were asked to assume that each of the
198 hypothetical persons had the same age and background as their own. Anchoring vignettes
199 are designed to take into account the fact that people of different countries, sex, age bands
200 and socio-economic groups may rate similar circumstances differently. Further detailed
201 information on anchoring vignettes can be found elsewhere (Jones et al., 2013).

202

203 *Covariates assessment*

204 Health behaviours and other covariates were recorded during the main interviews.
205 Four health behaviours included smoking, alcohol, physical activity and sleeping. Smoking
206 status was coded as current vs. not current smoker. Alcohol frequency, but not consumption
207 volume was available and was coded as low/moderate (once or twice per week to once or
208 twice per year), high (almost every day to three or four times per week) and abstainer (not
209 at all in last 12 months). Physical activity was coded as active (moderate to high physical
210 activity) vs. low/sedentary. Sleeping was coded as restless during past week vs. not restless.

211 Other covariates included age, sex, living arrangements, education and household
212 income. Age was classified into four categories: 50-59, 60-69, 70-79 and 80+ years old.
213 Living arrangements were classified into two categories: living alone and
214 cohabiting/married. Education was classified into three categories: basic (no/basic

215 qualifications), secondary (higher education but below a degree), and higher (degree or
216 above). Income was included as a continuous variable which was based on the sum of
217 employment, state benefit, state and private pension, asset, and other income; each
218 member of the benefit unit was assigned the total benefit unit level income. The OECD
219 equivalence scale was used (assigned a weight of 1 to the household head, 0.5 to second
220 adults and dependent children aged 14 and over and a weight of 0.3 to children under 14
221 years of age) (39) and total income was scaled by a factor of £1000.

222

223 **Statistical methods**

224 All statistical analysis was performed using STATA IC V.13.1 (StataCorp, Texas, USA).

225 A standard ordered probit model was used as a baseline model with which we could
226 compare our more flexible specification which does not impose the assumption of reporting
227 homogeneity, to assess the extent to which this assumption biases the estimated health
228 effects.

229 Individuals rated the vignettes describing the hypothetical cases similar to how they
230 rated their own SRH. As they represent fixed levels of health, individual variation in vignette
231 ratings characterise reporting heterogeneity (DIF). This 'external' vignette information can
232 therefore be used to model the cut-points on the Likert scale (which are assumed fixed in
233 the ordered probit model) as functions of the individual's characteristics. These cut-points
234 can then be used to purge reporting heterogeneity from the SRH, making it possible to
235 identify 'true' health effects. This is achieved through the use of a hierarchical probit model
236 (HOPIT).

237 The HOPIT model has two components. The vignette component models the cut-
238 points as functions of the covariates allowing for reporting heterogeneity. To relax the

239 restriction of parallel cut-point shift (covariates affect all cut-points by the same magnitude)
240 in this component, a generalised ordered probit model is used. The health component
241 represents the relationship between SRH and covariates, with the cut-points determined by
242 the vignette component, linking individual's SRH to the observed severity categories.
243 Further detailed information on these models can be found elsewhere (Jones et al., 2013).
244 The reference scale used in the HOPIT approach is arbitrary as it is the group represented by
245 the omitted categories in the generalised ordered probit. By applying any reference scale of
246 interest, it is possible to conduct simulations to illustrate the potential magnitude of the
247 effect of reporting heterogeneity in estimating the distribution of SRH. This is achieved by
248 reclassifying all responses and making them consistent with that scale (Heiland and Yin,
249 2015). For example, the predicted distribution of health categories can be visualised by
250 applying a HOPIT correction for self-reporting heterogeneity *and* reclassifying all responses
251 in accordance with the inferred response scales of people with either high or low social
252 capital, for each dimension of interest.

253

254 **Results**

255 Table 1 represents descriptive statistics for sociodemographic variables, health behaviours
256 and vignette ratings for the whole sample, and for the sub-groups of high and low social
257 capital within each of the three dimensions and deprivation. The main results are for all
258 individuals who answered at least one of the 18 health vignettes and the SRH question
259 (whole sample; $n_{\max}=2,341$ individuals contributing a maximum of 42,138 observations
260 [person-vignettes]). Those individuals with high social capital in any of the dimensions were
261 generally older (not statistically significant for social networks), married females. Those with
262 good social networks tended to be highly educated ($P<0.01$) whereas the opposite was seen

263 in those with good social support ($P < 0.001$). As for health behaviours, those with low local
264 area & trust social capital and good social networks tended to be non-smokers ($P = 0.01$ and
265 $P < 0.001$, respectively). Those with good social networks were more likely to be drinkers
266 ($P = 0.02$). Those with high local area & trust social capital and good social networks reported
267 higher physical activity ($P = 0.05$ and $P < 0.001$, respectively). Individuals with high social
268 capital in any of the dimensions reported better sleep (borderline statistically significant for
269 social networks, $P = 0.06$). Those with high social capital in any of the dimensions also rated
270 their SRH higher. Individuals who were least deprived tended to be married and more highly
271 educated (both $P < 0.001$). They tended not to be current smokers but drank alcohol more
272 frequently (both $P < 0.001$). However, they were more physically active ($P < 0.001$) and
273 reported better sleep ($P < 0.01$). They rate their own SRH higher than the most deprived.

274 Table 2 compares the estimated coefficients in the latent health index implied by the
275 different specifications of the ordered probit model and HOPIT. Two different models are
276 shown: Model 1 includes all our dimensions of social capital (local area and trust, social
277 support and social networks) and deprivation simultaneously with age and sex, and Model 2
278 is similar to Model 1 except it also includes the sociodemographic and health behaviours. A
279 model including only one dimension of social capital (e.g. social support only) or deprivation
280 at a time, along with age and sex and the vignette dummies was also derived, though the
281 results were similar to those seen in model 1 (data not shown). For direct comparisons to be
282 made between the two specifications, the scale of the estimated sigma in the HOPIT needs
283 to be close to 1 because the scale in the ordered probit is normalised to 1, while it is
284 estimated (up to the normalisation of scale in the vignette component) in the HOPIT. The
285 estimated sigma in this analysis for model 1 was 1.16, but was 1.02 in model 2. Therefore,
286 making direct comparisons between the two specifications in the fully adjusted model

287 (model 2) is not problematic, but caution needs to be taken when making direct
288 comparisons between the two specifications in model 1. Thus, the following results are in
289 reference to model 2. The ordered probit is the most restricted specification that disregards
290 any reporting heterogeneity. Individuals expressing low local area & trust social capital (beta
291 = -0.243, $p < 0.001$) and those with poor social networks (beta = -0.210, $p < 0.01$) were more
292 likely to report poorer SRH. These findings remained evident when allowing for non-parallel
293 cut-point shift (HOPIT), though ignoring reporting heterogeneity tended to marginally
294 underestimate the detrimental effect on SRH of having low local area & trust social capital
295 (Beta = -0.276, $p < 0.001$) and poor social networks (Beta = -0.280, $p < 0.001$). Post-estimation
296 tests (using “suest” command in STATA which tests for intra-model and cross-model
297 hypotheses) between the betas in the ordered probit versus the HOPIT model were not
298 statistically significant for low local area & trust ($p=0.61$) or poor social networks ($p=0.28$).

299 Neither social support nor deprivation appeared to have any effect on SRH
300 regardless of DIF. However, in model 1, a significant negative coefficient for low social
301 support (beta = -0.209, $p < 0.01$) and for the most deprived group (beta = -0.186, $p < 0.05$),
302 and a significant positive coefficient for the least deprived (beta = 0.308, $p < 0.001$) lost
303 statistical significance once adjusted for sociodemographic variables and health behaviours
304 (model 2). Males reported poorer SRH in both the ordered probit (beta = -0.200, $p < 0.001$)
305 and HOPIT models (beta = -0.261, $p < 0.001$). The age categories were negatively associated
306 with SRH (model 2) and these effects remained non-significant, except for age 70-79 which
307 became statistically significant when reporting heterogeneity was accounted for (beta = -
308 0.185, $p < 0.05$).

309 Table S2 in the Online Supplementary Material compares the estimated coefficients
310 of the ordered probit model and HOPIT for each of the six domains of health that were also

311 asked in the self-completion questionnaire. The same covariates used in Model 2, Table 2
312 were used. Overall, these individual results align well with our main analysis using SRH to
313 represent an overall indicator of general health; the negative effect of low social capital is
314 greater for the majority of the six health domains and all three measures of social capital
315 when reporting heterogeneity is accounted for. Post-estimation tests between the betas for
316 social capital in the ordered probit versus the HOPIT models were statistically significant at
317 the 5% level across three domains of health and statistically significant at the 10% level
318 across four domains (emboldened in Table S2).

319 The response scales inferred from vignette classifications made by respondents of
320 high and low social capital within each of our three dimensions, and deprivation, can be
321 useful to researchers who rely on self-reported measures. Table S3 in the Online
322 Supplementary Material shows the results of the generalised ordered probit model of
323 individuals' rating of the vignettes' health (vignette component of the HOPIT model). This
324 model accommodates the potential for a non-parallel cut-point shift, allowing the covariates
325 to affect each of the cut-points differently. The coefficients vary considerably across cut-
326 points, and in many cases, the effects are not monotonic. Two model specifications were
327 performed similar to Table 2. A positive coefficient implies a rightwards shift in the cut-
328 point, suggesting that, on average, individuals from the corresponding group characterize
329 the health problems presented in the vignette as more severe. Likewise, a negative
330 coefficient implies a leftwards shift in the cut-point.

331 Figure 2 displays simulations to illustrate the potential magnitude of the effect of
332 reporting heterogeneity in estimating the distribution of SRH. The top graph shows the
333 empirical (unadjusted) distribution of SRH among ELSA participants aged 50 years and older
334 who answered at least one of the 18 health vignettes. The second graph represents a

335 predicted distribution of SRH using HOPIT procedures without any correction for reporting
336 heterogeneity (similar to an ordered probit model). In the latent (own) health index, the
337 same set of covariates as in Model 2 were included (age, sex, sociodemographic variables
338 and health behaviours).

339 By applying any reference scale of interest in the HOPIT specification, we can
340 reclassify all responses and make them consistent with that response scale. The bottom
341 graphs in Figure 2 display the predicted distribution of SRH after applying a HOPIT correction
342 for reporting heterogeneity and reclassifying all responses in accordance with the reference
343 scale of interest: response scales of high and low social capital within each of our three
344 dimensions or in accordance with the response scales of least/most deprived. The predicted
345 distributions are consistent with the findings reported in Table S2 and differ mainly from the
346 second graph at the threshold good vs. very good health. For example, when the scales
347 inferred for the groups with poor social networks were used, the predicted distributions
348 were more concentrated at the category “very good health”, consistent with Model 2, Table
349 S2. They have a lower threshold to what constitutes very good health compared to those
350 with good social networks.

351

352 **Discussion**

353 SRH is a subjective measure often used as an indicator of general health in large
354 cohort studies, and is regarded as a robust predictive measure of mortality, morbidity &
355 physical functioning (Grol-Prokopczyk et al., 2011). The conceptual framework for health
356 supports the view that it is best represented as a multidimensional set of domains (Salomon
357 et al., 2003). The World Health Organisation (WHO) developed a set of core health domains
358 that best describe different aspects of health status directly (Sadana et al., 2002; Salomon et

359 al., 2003). The 18 vignettes used in ELSA covered six health domains: pain, sleeping,
360 mobility, memory, breathing and depression, which are included as part of the WHO's core
361 domains of health. There were three vignettes per domain listing the health condition (e.g.
362 pain) in increasing severity. Furthermore, studies have consistently shown associations
363 between poor SRH and physical health (pain, sleeping, breathing and mobility), whether or
364 not it relates to limitations (e.g. our vignettes ask about health problems/limitations), and
365 mental health (Borim et al., 2014; Chang-Quan et al., 2010; Latham and Peek, 2013).
366 Therefore, it was felt that this set of six health domains covered by the vignettes would be
367 sufficiently exhaustive to capture the most common dimensions of SRH in our main analysis.

368 What our results show is that low local area & trust and poor social networks are
369 associated with poorer SRH in HOPIT models accounting for reporting heterogeneity, but
370 while ordered probit analyses still correctly show a negative relationship between these
371 social capital dimensions and SRH, they somewhat underestimate its strength. Moreover,
372 our simulations illustrate the potential magnitude of reporting heterogeneity in estimating
373 the distribution of SRH by demonstrating the impact of different response scales. In
374 particular, the distribution at the cut-point good vs. very good health tended to differ
375 (across social capital and deprivation categories) after applying a HOPIT correction for self-
376 reporting heterogeneity and reclassifying all responses in accordance with the chosen scale.
377 Also, the bad and very bad self-reported health distribution was greatly diminished after
378 applying the HOPIT correction and reclassification. Our analysis highlights the caution that
379 needs exercised when using unadjusted self-reported measures to study the effects of social
380 capital and deprivation on health.

381 Social capital, as highlighted in the introduction, is a multidimensional concept which
382 can have both positive and negative effects on health. When we applied the most flexible

383 model incorporating the hypothetical health vignettes and accommodating for non-parallel
384 cut-point shift (HOPIT) arising from reporting heterogeneity, we demonstrated that those
385 individuals with low local area & trust social capital and poor social networks were less likely
386 to report good health. These results are consistent with the literature.

387 Having a higher opinion of your local area and a greater sense of trust, and belonging
388 to broader social networks can bring certain benefits and resources that would not
389 otherwise be available. These resources are not all necessarily at an individual level but can
390 be garnered via the group-level dynamics within such environments (Kawachi and Berkman,
391 2014). Individual health benefits secured by virtue of membership include social support
392 (exchange of affective support), social influence (promotion of healthy behaviours), social
393 control (status and rewards) and social participation (opportunities to learn new skills, self-
394 esteem and promotion of belongingness). Additional benefits to health include access to
395 material resources such as health services, job opportunities and finance (Eriksson, 2011;
396 Kawachi et al., 1999; Kawachi and Berkman, 2014; Lindström, 2008). Collective health
397 benefits secured through norms and collective efficacy include trust, solidarity and
398 reciprocity, which promotes a health-enabling environment through attitudes that influence
399 healthy behaviours, diffusion of knowledge and information (social contagion) and the
400 potential to influence political and community decisions/resources (Eriksson, 2011; Kawachi
401 and Berkman, 2014). The availability and distribution of such resources will have an impact
402 on how individual's not only rate their own SRH, but may modify their judgement of what
403 constitutes good and bad health in the hypothetical vignettes.

404 Contrary to previous research which has found detrimental health effects of
405 deprivation (Diez Roux and Mair, 2010; Roux et al., 2001; Stafford and Marmot, 2003), the
406 current study found no evidence of an effect on SRH of living in a deprived neighbourhood

407 in both the ordered probit and HOPIT model specifications when adjusted for
408 sociodemographic variables and health behaviours. This could be due, at least in part, to
409 how deprivation is measured. The conceptual framework behind IMD2004 uses LSOA data
410 to construct an aggregate area based score and is agnostic with respect to the causes of
411 deprivation. Therefore, a LSOA scored as relatively deprived by the index may contain large
412 numbers of people who are not deprived, and conversely, LSOA which are relatively less
413 deprived might contain people experiencing significant disadvantage. Nevertheless we
414 acknowledge that both area level and individual level attributes contribute to deprivation
415 and we may not have been able to fully separate their effects. Therefore, caution is
416 warranted when interpreting our findings for deprivation, especially as the data is cross-
417 sectional in nature.

418 We originally hypothesised that relying on SRH alone without accounting for
419 reporting heterogeneity would underestimate the detrimental effect of low social capital on
420 SRH. Unadjusted ordered probit analyses still correctly demonstrated a negative
421 relationship between some of the social capital dimensions and SRH, though they somewhat
422 underestimated its strength (Table 2). It was also hypothesised that those with low social
423 capital might use lower response thresholds for what constitutes a health problem when
424 responding to the hypothetical vignettes (Table S3). Our simulations illustrate the impact of
425 these response thresholds on standard measures of SRH when reclassifying all responses in
426 accordance with high and low social capital (for each dimension) and deprivation.

427 Overall, this study demonstrates the importance of accounting for reporting
428 heterogeneity when conducting comparative studies, either between sub-groups or across
429 whole countries. Anchoring vignettes offer a relatively uncomplicated and cost-effective
430 way of identifying and correcting for DIF to improve comparative validity of self-reported

431 measures such as SRH. Future research is needed to improve vignette methodology while
432 retaining its simplicity with respect to survey operation and anchoring performance,
433 especially with large scale population surveys in which resources are limited.

434

435 **Strengths and Limitations**

436 Anchoring vignettes have a number of advantages over earlier methods of
437 identifying and correcting for DIF. They are less error-prone and can both identify DIF and
438 statistically correct for it (HOPIT); they are relatively cheap to implement in that they only
439 require a small number of additional survey items and be given to a proportion of the whole
440 sample; and they may allow a means of improving comparative validity of self-reported
441 measures. Health vignettes thus have the potential to serve a valuable role in health
442 research, enabling more accurate empirical work and more rigorous honing of theory (Grol-
443 Prokopczyk et al., 2011). However, the use of anchoring vignettes comes with potential
444 limitations. The assumptions of vignette equivalence and response consistency may not
445 always hold true in the HOPIT models. For example, given the complex multidimensional
446 nature of health, vignette descriptions are likely to be incomplete, and individuals may call
447 upon their own experience to impute the missing information (lack of vignette equivalence)
448 (van Soest et al., 2011). Similarly, individuals may report their own situation with a certain
449 strategic consideration that is absent from vignette assessment (failure of response
450 consistency) (d’Uva et al., 2011). The precise wording of the cut-points used in the current
451 study between the SRH and the health vignettes varied somewhat though it was generally
452 thought to impart the same understanding (e.g. ‘no health problem’ in vignette equivalent
453 to ‘very good health’ in the SRH). A few researchers have attempted to test these
454 assumptions separately (d’Uva et al., 2011; Grol-Prokopczyk et al., 2015), but rigorous tests

455 of these assumptions require extra data such as valid and reliable objective health
456 measures, which were not collected at Wave 3 of ELSA. Additionally, as with SRH, there may
457 be reporting heterogeneity in the social capital variables, but to our knowledge no
458 nationally representative study, including ELSA, has developed social capital vignettes.
459 Therefore, we could not take into account reporting heterogeneity in these measures.
460 However, the present study is a first step towards a better understanding of the effects of
461 reporting heterogeneity and the utility of anchoring vignettes in survey data on the social
462 capital and deprivation disparities in health. Other limitations of the current study are the
463 potential for unmeasured covariates and residual confounding and the fact that the health
464 vignettes module was only completed once at Wave 3 (2006/07). Therefore, we could not
465 analyse vignettes longitudinally, incorporating changes in perceptions and reporting of
466 health into the models. However, with 18 vignettes in total, covering six different health
467 domains, we have a very comprehensive data set in a large, representative sample of
468 individuals aged 50 years and older throughout England.

469

470 **Avenues for Future Research**

471 These results may be more indicative of 'true' health disparities or may be the result of
472 diverging 'attitudes' between social capital groupings. Overall, policy solutions require an
473 overarching approach by addressing the social determinants of health that are inclusive of
474 all sectors of the community. High quality research is required to identify how best to tackle
475 health inequalities and policy solutions for each group might be quite different.

476

477 **References**

478 Au, N., Johnston, D.W., 2014. Self-assessed health: What does it mean and what does it

479 hide? Soc. Sci. Med. 121, 21–28. doi:10.1016/j.socscimed.2014.10.007

480 Au, N., Lorgelly, P.K., 2014. Anchoring vignettes for health comparisons: an analysis of
481 response consistency. Qual. Life Res. 23, 1721–31. doi:10.1007/s11136-013-0615-2

482 Barger, S.D., Cribbet, M.R., Muldoon, M.F., 2016. Participant-Reported Health Status
483 Predicts Cardiovascular and All-Cause Mortality Independent of Established and
484 Nontraditional Biomarkers: Evidence From a Representative US Sample. J. Am. Heart
485 Assoc. 5. doi:10.1161/JAHA.116.003741

486 Borim, F.S.A., Neri, A.L., Francisco, P.M.S.B., Barros, M.B. de A., 2014. Dimensions of self-
487 rated health in older adults. Rev. Saude Publica 48, 714–22. doi:10.1590/s0034-
488 8910.2014048005243

489 Burgard, S.A., Chen, P. V., 2014. Challenges of health measurement in studies of health
490 disparities. Soc. Sci. Med. 106, 143–150. doi:10.1016/j.socscimed.2014.01.045

491 Burt, R., 1992. Social capital: it's origins and applications in modern sociology. Annu. Rev.
492 Sociol. 24, 1–24.

493 Campos-Matos, I., Subramanian, S. V, Kawachi, I., 2016. The “dark side” of social capital:
494 trust and self-rated health in European countries. Eur. J. Public Health 26, 90–5.
495 doi:10.1093/eurpub/ckv089

496 Chang-Quan, H., Xue-Mei, Z., Bi-Rong, D., Zhen-Chan, L., Ji-Rong, Y., Qing-Xiu, L., 2010.
497 Health status and risk for depression among the elderly: a meta-analysis of published
498 literature. Age Ageing 39, 23–30. doi:10.1093/ageing/afp187

499 Chen, H., Meng, T., 2015. Bonding, bridging, and linking social capital and self-rated health
500 among Chinese adults: Use of the anchoring vignettes technique. PLoS One 10,
501 e0142300. doi:10.1371/journal.pone.0142300

502 Christakis, N.A., Fowler, J.H., 2008. The collective dynamics of smoking in a large social

503 network. *N. Engl. J. Med.* 358, 2249–58. doi:10.1056/NEJMsa0706154

504 Coll-Planas, L., Nyqvist, F., Puig, T., Urrútia, G., Solà, I., Monteserín, R., 2016. Social capital
505 interventions targeting older people and their impact on health: a systematic review. *J.*
506 *Epidemiol. Community Health* jech-2016-208131. doi:10.1136/JECH-2016-208131

507 Colwell, S., 2016. The composite reliability calculator. doi:10.13140/RG.2.1.4298.088

508 d’Uva, T.B., Lindeboom, M., O’Donnell, O., van Doorslaer, E., 2011. Slipping Anchor? Testing
509 the Vignettes Approach to Identification and Correction of Reporting Heterogeneity. *J.*
510 *Hum. Resour.* 46, 875–906.

511 Diez Roux, A. V., Mair, C., 2010. Neighborhoods and health. *Ann. N. Y. Acad. Sci.* 1186, 125–
512 145. doi:10.1111/j.1749-6632.2009.05333.x

513 Eriksson, M., 2011. Social capital and health--implications for health promotion. *Glob.*
514 *Health Action* 4, 5611. doi:10.3402/gha.v4i0.5611

515 Giordano, G.N., Björk, J., Lindström, M., 2012. Social capital and self-rated health – A study
516 of temporal (causal) relationships. *Soc. Sci. Med.* 75, 340–348.
517 doi:10.1016/j.socscimed.2012.03.011

518 Glymour, M., Avendano, M., Kawachi, I., 2014. Socioeconomic status and health, in:
519 Berkman, L., Kawachi, I., Glymour, M. (Eds.), *Social Epidemiology*. Oxford University
520 Press, New York, pp. 17–62.

521 Grol-Prokopczyk, H., Freese, J., Hauser, R.M., 2011. Using anchoring vignettes to assess
522 group differences in general self-rated health. *J. Health Soc. Behav.* 52, 246–61.
523 doi:10.1177/0022146510396713

524 Grol-Prokopczyk, H., Verdes-Tennant, E., McEniry, M., Ispány, M., 2015. Promises and
525 Pitfalls of Anchoring Vignettes in Health Survey Research. *Demography* 52, 1703–28.
526 doi:10.1007/s13524-015-0422-1

527 Hageaars, A., de Vos, K., Zaidi, M., 1994. Poverty Statistics in the Late 1980s: Research
528 Based on Micro-data. Eurostat, Luxembourg.

529 Heiland, F., Yin, N., 2015. Anchoring vignettes in the Health and Retirement Study: how do
530 medical professionals and disability recipients characterize the severity of work
531 limitations? PLoS One 10, e0126218. doi:10.1371/journal.pone.0126218

532 Jones, A.M., Rice, N., Bago D’Uva, T., Balia, S., 2013. Reporting heterogeneity in health, in:
533 Applied Health Economics. Routledge, New York, pp. 69–105.

534 Kawachi, I., Berkman, L.F., 2014. Social capital, social cohesion, and health, in: Berkman, L.F.,
535 Kawachi, I., Glymour, M.M. (Eds.), Social Epidemiology. Oxford University Press, New
536 York, pp. 290–319.

537 Kawachi, I., Kennedy, B.P., Glass, R., 1999. Social capital and self-rated health: a contextual
538 analysis. Am. J. Public Health 89, 1187–93.

539 Kawachi, I., Kim, D., Coutts, A., Subramanian, S., 2004. Commentary: Reconciling the three
540 accounts of social capital. Int. J. Epidemiol. 33, 682–690. doi:10.1093/ije/dyh177

541 King, G., Murray, C., Salomon, J., Tandon, A., 2004. Enhancing the Validity and Cross-cultural
542 Comparability of Measurement in Survey Research. Am. Polit. Sci. Rev. 98, 191–207.

543 Koutsogeorgou, E., Nyqvist, F., Nygard, M., Cerniauskaite, M., Quintas, R., Raggi, A.,
544 Leonardi, M., 2015. Social capital and self-rated health among older adults: a
545 comparative analysis of Finland, Poland and Spain. Ageing Soc. 35, 653–667.
546 doi:10.1017/S0144686X13001025

547 Latham, K., Peek, C.W., 2013. Self-rated health and morbidity onset among late midlife U.S.
548 adults. J. Gerontol. B. Psychol. Sci. Soc. Sci. 68, 107–16. doi:10.1093/geronb/gbs104

549 Lindström, M., 2008. Social Capital and Health-Related Behaviors, in: Kawaci, I.,
550 Subramanian, S., Kim, D. (Eds.), Social Capital and Health. Springer, New York, pp. 215–

551 238. doi:10.1007/978-0-387-71311-3_10

552 Nieminen, T., Martelin, T., Koskinen, S., Aro, H., Alanen, E., Hyypä, M.T., 2010. Social
553 capital as a determinant of self-rated health and psychological well-being. *Int. J. Public*
554 *Health* 55, 531–42. doi:10.1007/s00038-010-0138-3

555 Nieminen, T., Prättälä, R., Martelin, T., Härkänen, T., Hyypä, M.T., Alanen, E., Koskinen, S.,
556 2013. Social capital, health behaviours and health: a population-based associational
557 study. *BMC Public Health* 13, 613. doi:10.1186/1471-2458-13-613

558 Noble, M., Wright, G., Dibben, C., Smith, G., McLennan, D., Anttila, C., Barnes, H., Mokhtar,
559 C., Noble, S., Avenell, D., Gardner, J., Covizzi, I., Lloyd, M., 2004. *The English Indices of*
560 *Deprivation 2004*. London.

561 Peracchi, F., Rossetti, C., 2012. Heterogeneity in health responses and anchoring vignettes.
562 *Empir. Econ.* 42, 513–538. doi:10.1007/s00181-011-0530-8

563 Poortinga, W., 2006. Social capital: An individual or collective resource for health? *Soc. Sci.*
564 *Med.* 62, 292–302. doi:10.1016/j.socscimed.2005.06.008

565 Rosenquist, J.N., Fowler, J.H., Christakis, N.A., 2011. Social network determinants of
566 depression. *Mol. Psychiatry* 16, 273–81. doi:10.1038/mp.2010.13

567 Roux, A.V.D., Merkin, S.S., Arnett, D., Chambless, L., Massing, M., Nieto, F.J., Sorlie, P., Szklo,
568 M., Tyroler, H.A., Watson, R.L., 2001. Neighborhood of Residence and Incidence of
569 Coronary Heart Disease. *N. Engl. J. Med.* 345, 99–106.
570 doi:10.1056/NEJM200107123450205

571 Sadana, R., Tandon, A., Murray, C., Serdobova, I., Cao, Y., Xie, W., Chatterji, S., Ustün, B.,
572 2002. Describing population health in six domains: comparable results from 66
573 household surveys (No. GPE discussion paper no. 43).

574 Salomon, J., Mathers, C., Chatterji, S., Sadana, R., Üstün, T., Murray, C., 2003. Quantifying

575 individual levels of health: definitions, concepts and measurement issues., in: Murray,
576 C., Evans, D. (Eds.), Health Systems Performance: Assessment Debates, Methods and
577 Empiricisms. WHO, Geneva, pp. 301–318.

578 Sánchez-Santos, M.T., Mesa-Frias, M., Choi, M., Nüesch, E., Asunsolo-Del Barco, A., Amuzu,
579 A., Smith, G.D., Ebrahim, S., Prieto-Merino, D., Casas, J.P., 2013. Area-level deprivation
580 and overall and cause-specific mortality: 12 years' observation on British women and
581 systematic review of prospective studies. PLoS One 8, e72656.
582 doi:10.1371/journal.pone.0072656

583 Scrivens, K., Smith, C., 2013. Four Interpretations of Social Capital. OECD Publishing.
584 doi:10.1787/5JZBCX010WMT-EN

585 Siegler, V., 2014. Measuring Social Capital.

586 Stafford, M., Marmot, M., 2003. Neighbourhood deprivation and health: does it affect us all
587 equally? Int. J. Epidemiol. 32, 357–66. doi:10.1093/IJE/DYG084

588 Steptoe, A., Breeze, E., Banks, J., Nazroo, J., 2013. Cohort profile: the English longitudinal
589 study of ageing. Int. J. Epidemiol. 42, 1640–8. doi:10.1093/ije/dys168

590 Szreter, S., Woolcock, M., 2004. Health by association? Social capital, social theory, and the
591 political economy of public health. Int. J. Epidemiol. 33, 650–67.
592 doi:10.1093/ije/dyh013

593 van Soest, A., Delaney, L., Harmon, C., Kapteyn, A., Smith, J.P., 2011. Validating the use of
594 anchoring vignettes for the correction of response scale differences in subjective
595 questions. J. R. Stat. Soc. Ser. A (Statistics Soc. 174, 575–595. doi:10.1111/j.1467-
596 985X.2011.00694.x

597 Xu, H., Xie, Y., 2015. Assessing the Effectiveness of Anchoring Vignettes in Bias Reduction for
598 Socioeconomic Disparities in Self-rated Health among Chinese Adults. Sociol.

599 Methodol. 81175015599808. doi:10.1177/0081175015599808

600 Zhang, X., Cook, P.A., Lisboa, P.J., Jarman, I.H., Bellis, M.A., 2013. The effects of deprivation
601 and relative deprivation on self-reported morbidity in England: an area-level ecological
602 study. *Int. J. Health Geogr.* 12, 5. doi:10.1186/1476-072X-12-5

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623 **Figure 1.** Reporting of health across two groups illustrating reporting heterogeneity

624

625 A hypothetical vignette person with the same objective degree of health (represented by
626 the dotted vertical line) is classified as having an extreme health problem by individuals with
627 high social capital, while individuals with low social capital may characterise the same
628 person as having a severe health problem.

629

630 **Figure 2.** Simulations illustrating the potential magnitude of the effect of reporting
631 heterogeneity in estimating the distribution of health problem severity from self-reported
632 survey data using the ELSA Wave 3 (2006/07) cohort, men and women aged 50 years and
633 older

634

635 “Empirical distribution” refers to the distribution of self-rated health among the whole
636 sample who have answered at least one of the 18 health vignettes and the self-report.

637 “Estimated distribution without adjusting for reporting heterogeneity” refers to the
638 distribution of self-rated health estimated using the HOPIT procedure but without adjusting
639 reporting heterogeneity, which is similar to an ordered probit model. “Reclassification using
640 high social capital/least deprived” and “Reclassification using low social capital/most
641 deprived” refers to the distribution of self-rated health adjusted for reporting heterogeneity
642 in accordance with the estimated scales (based on Model 2, Table S2) for high and low social
643 capital with regard to the dimension in question/least and most deprived.

Table 1 Descriptive statistics for analytic sample in the ELSA Wave 3 (2006/07) cohort, men and women aged 50 years and older^a

	Local area & trust						Social support				Social networks				Index of Multiple Deprivation 2004			
	Whole sample		High ^b		Low ^b		High		Low		High		Low		Least deprived ^c		Most deprived ^c	
	<i>(n_{max} = 2341)</i> <i>(obs_{max} = 42,138)</i>		<i>(n_{max} = 418)</i> <i>(obs_{max} = 7524)</i>		<i>(n_{max} = 403)</i> <i>(obs_{max} = 7254)</i>		<i>(n_{max} = 417)</i> <i>(obs_{max} = 7506)</i>		<i>(n_{max} = 431)</i> <i>(obs_{max} = 7758)</i>		<i>(n_{max} = 408)</i> <i>(obs_{max} = 7344)</i>		<i>(n_{max} = 410)</i> <i>(obs_{max} = 7380)</i>		<i>(n_{max} = 616)</i> <i>(obs_{max} = 11,088)</i>		<i>(n_{max} = 285)</i> <i>(obs_{max} = 5130)</i>	
	Percent / Mean	SD	Percent / Mean	SD	Percent / Mean	SD	Percent / Mean	SD	Percent / Mean	SD	Percent / Mean	SD	Percent / Mean	SD	Percent / Mean	SD	Percent / Mean	SD
Age	65.40	10.10	66.74	10.22	62.96	9.80	66.92	10.12	63.21	9.78	65.50	9.30	64.81	10.30	65.18	10.12	65.66	10.04
50-59	0.35	0.48	0.32	0.47	0.45	0.50	0.30	0.46	0.44	0.50	0.33	0.47	0.39	0.49	0.36	0.48	0.32	0.47
60-69	0.30	0.46	0.27	0.44	0.28	0.45	0.28	0.45	0.30	0.46	0.31	0.46	0.29	0.45	0.30	0.46	0.31	0.46
70-79	0.24	0.43	0.29	0.46	0.20	0.40	0.29	0.46	0.17	0.38	0.27	0.45	0.22	0.42	0.23	0.42	0.27	0.45
80+	0.11	0.31	0.12	0.32	0.07	0.25	0.12	0.33	0.09	0.28	0.08	0.28	0.10	0.31	0.11	0.31	0.09	0.29
Male	0.44	0.50	0.40	0.49	0.48	0.50	0.40	0.49	0.49	0.50	0.38	0.49	0.54	0.50	0.45	0.50	0.41	0.49
Living arrangements																		
Living alone	0.28	0.45	0.27	0.44	0.33	0.47	0.25	0.43	0.37	0.48	0.25	0.43	0.33	0.47	0.22	0.41	0.44	0.50
Cohabit/married	0.72	0.45	0.73	0.44	0.67	0.47	0.75	0.43	0.63	0.48	0.75	0.43	0.67	0.47	0.78	0.41	0.56	0.50
Education^d																		
Basic	0.43	0.49	0.40	0.49	0.42	0.49	0.53	0.50	0.32	0.47	0.27	0.44	0.45	0.50	0.30	0.46	0.64	0.48
Secondary	0.40	0.49	0.45	0.50	0.38	0.49	0.39	0.49	0.42	0.49	0.49	0.50	0.39	0.49	0.47	0.50	0.27	0.45
Higher	0.17	0.38	0.15	0.36	0.19	0.39	0.08	0.27	0.26	0.44	0.24	0.43	0.16	0.37	0.23	0.42	0.08	0.28
Current smoker	0.14	0.35	0.12	0.33	0.18	0.39	0.18	0.38	0.14	0.34	0.06	0.24	0.20	0.40	0.09	0.28	0.28	0.45
Drinking frequency^e																		
Low / moderate	0.53	0.50	0.53	0.50	0.55	0.50	0.54	0.50	0.51	0.50	0.54	0.50	0.51	0.50	0.51	0.50	0.61	0.49
High	0.35	0.48	0.35	0.48	0.32	0.47	0.33	0.47	0.38	0.49	0.38	0.48	0.35	0.48	0.42	0.49	0.21	0.41
Abstainer	0.12	0.32	0.12	0.33	0.13	0.34	0.13	0.33	0.11	0.31	0.08	0.27	0.13	0.34	0.07	0.26	0.19	0.39
Physical activity; active	0.72	0.45	0.73	0.44	0.67	0.47	0.69	0.46	0.73	0.44	0.81	0.39	0.65	0.48	0.80	0.40	0.54	0.50
No complaint sleeping	0.59	0.49	0.66	0.47	0.50	0.50	0.62	0.49	0.53	0.50	0.65	0.48	0.59	0.49	0.64	0.48	0.55	0.50
Self-rated health ^f	3.89	0.88	4.00	0.83	3.66	0.88	3.92	0.91	3.84	0.85	4.04	0.80	3.68	0.96	4.05	0.79	3.52	0.97
Vignette: Pain	3.04	1.06	3.02	1.06	3.03	1.04	3.01	1.08	3.10	1.01	2.99	1.05	3.05	1.06	3.05	1.04	2.98	1.07
Vignette: Sleep	2.59	0.85	2.54	0.82	2.59	0.85	2.60	0.83	2.65	0.82	2.50	0.81	2.62	0.87	2.57	0.81	2.58	0.85
Vignette: Mobility	2.68	1.00	2.66	0.99	2.66	0.98	2.62	1.00	2.72	0.98	2.63	0.99	2.66	0.97	2.66	0.98	2.66	0.97
Vignette: Memory	3.11	1.00	3.10	1.02	3.12	0.98	3.07	1.02	3.18	0.98	3.14	1.00	3.09	0.98	3.15	0.99	3.05	0.97
Vignette: Breathing	2.17	0.95	2.17	0.94	2.18	0.93	2.17	0.98	2.17	0.92	2.13	0.90	2.19	0.96	2.16	0.92	2.15	0.89
Vignette: Depression	2.60	1.04	2.58	1.05	2.59	1.02	2.59	1.07	2.62	1.02	2.55	1.03	2.65	1.05	2.55	1.01	2.55	1.01

^a Excludes those aged <50 years, did not participate in health self-completion questionnaire (no vignette responses)) or no self-rated health reported

^b High = top quintile of factor-analysis score for social capital dimension; Low = bottom quintile of factor-analysis score for social capital dimension

^c Least = bottom quintile of IMD2004; Most = top quintile of IMD2004

^d Basic = no/basic qualifications; Secondary = higher education but below a degree; Higher = degree or above

^e Low/moderate = 2 times per week or less; High = 3+ times per week; Abstainer = no times in previous 12 months

^f Five-point Likert scale (1 = very bad/extreme problems to 5 = very good/no problems)

Table 2 Ordered probit and HOPIT regressions of self-rated health in the ELSA Wave 3 (2006/07) cohort, men and women aged 50 years and older^a

	Ordered probit		HOPIT		Ordered probit		HOPIT	
	β	SE	β	SE	β	SE	β	SE
	Model 1				Model 2			
Local area & trust								
High ^b	-0.012	(0.064)	-0.005	(0.078)	0.028	(0.067)	0.055	(0.073)
Low ^b	-0.363***	(0.065)	-0.433***	(0.078)	-0.243***	(0.068)	-0.276***	(0.073)
Social support								
High	-0.040	(0.066)	-0.147	(0.080)	0.024	(0.070)	-0.063	(0.076)
Low	-0.099	(0.063)	-0.209**	(0.076)	-0.027	(0.066)	-0.119	(0.071)
Social networks								
High	0.034	(0.065)	0.088	(0.080)	-0.044	(0.069)	0.003	(0.076)
Low	-0.272***	(0.064)	-0.407***	(0.076)	-0.210**	(0.067)	-0.280***	(0.071)
Deprivation								
Least deprived ^c	0.154**	(0.056)	0.308***	(0.068)	0.003	(0.059)	0.110	(0.064)
Most deprived ^c	-0.275***	(0.080)	-0.186*	(0.096)	-0.047	(0.086)	0.155	(0.092)
Age 60-69	-0.156**	(0.060)	-0.265***	(0.072)	-0.066	(0.063)	-0.133	(0.069)
Age 70-79	-0.371***	(0.064)	-0.520***	(0.077)	-0.130	(0.072)	-0.185*	(0.078)
Age 80+	-0.542***	(0.089)	-0.674***	(0.108)	-0.099	(0.106)	-0.131	(0.114)
Male	-0.040	(0.049)	-0.083	(0.059)	-0.200***	(0.053)	-0.261***	(0.057)
Vignette dummies	No		Yes		No		Yes	
Socio-demographic dummies	No		No		Yes		Yes	
Health dummies	No		No		Yes		Yes	
<i>N</i>	2046		2046		1926		1926	

^a Excludes those aged <50 years, did not participate in health self-completion questionnaire (no vignette responses) or no self-rated health reported

^b High = top quintile of factor-analysis score for social capital dimension; Low = bottom quintile of factor-analysis score for social capital dimension

^c Least = bottom quintile of IMD2004; Most = top quintile of IMD2004

Model 1: All dimensions of social capital (local area & trust, social support and social networks) & IMD2014 simultaneously

Model 2: All dimensions of social capital (local area & trust, social support and social networks), IMD2014, the socio-demographic covariates (education, living arrangements and income), and health behaviours (smoking, alcohol, physical activity and sleep) simultaneously

* p<0.05, ** p<0.01, *** p<0.001

Figure 1: 1.5 columns




Figure 2: 2 columns

