



**QUEEN'S
UNIVERSITY
BELFAST**

Metformin use and survival from lung cancer: a population-based cohort study

McMenamin, U., Cardwell, C., Hughes, C., & Murray, L. (2016). Metformin use and survival from lung cancer: a population-based cohort study. *Lung Cancer*, 94, 35-39. <https://doi.org/10.1016/j.lungcan.2016.01.012>

Published in:
Lung Cancer

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright © 2016 Elsevier Ireland Ltd. This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

Metformin use and survival from lung cancer: a population-based cohort study

Authors:

Úna C. Mc Menamin¹, Chris R. Cardwell¹, Carmel M. Hughes², Liam M. Murray^{1,3}.

Affiliations:

¹Cancer Epidemiology and Health Services Research Group, Centre for Public Health, Queen's University Belfast, Belfast, Northern Ireland.

²School of Pharmacy, Queen's University Belfast, Northern Ireland, BT9 7BL.

³Centre of Excellence for Public Health (NI), Centre for Public Health, Queen's University Belfast, Belfast, Northern Ireland, UK.

Corresponding Author:

Dr Úna Mc Menamin

Mail: Centre for Public Health, Institute of Clinical Sciences Block B, Queen's University Belfast, Royal Victoria Hospital, Grosvenor Road, Belfast, BT12 6BA, Northern Ireland

Phone: +44(0)2890633956.

Fax: +44(0)2890635900

E-mail: u.mcmenamin@qub.ac.uk

Abstract word count: 222

Manuscript word count: 1,472

Number of tables and figures: 2

Abbreviations: BMI, body mass index; CIs, confidence intervals; CPRD, Clinical Practice Research Datalink; DDD, daily defined dose; GP, general practitioner; HR, hazard ratio; NCDR, National cancer Data Repository; ONS, Office of National Statistics; UK, United Kingdom; WHO, World Health Organisation.

Abstract

Preclinical evidence suggests that metformin, a widely prescribed anti-diabetic drug, may inhibit lung cancer progression. We investigated whether metformin use was associated with decreased risk of cancer-specific mortality in lung cancer patients. This study included newly diagnosed lung cancer patients (identified from English National Cancer Data Repository, 1998-2009) with type 2 diabetes (based on UK Clinical Practice Research Datalink prescriptions and diagnosis records). Lung cancer deaths occurring up to 2012 were identified using Office of National Statistics mortality data and the association between metformin use (before and after diagnosis) and risk of lung cancer-specific mortality was calculated using Cox regression models. In analysis of 533 patients, we found a weak non-significant reduction in lung cancer-specific mortality with metformin use after diagnosis (adjusted HR, 0.86; 95% CI, 0.68-1.09). No association was evident for metformin use before diagnosis and cancer-specific mortality in analysis of 1,350 patients (adjusted HR, 0.97; 95% CI, 0.86, 1.11). Associations were similar by duration of use. In addition, after adjustment for potential confounders, there was little evidence of an association between the use of other anti-diabetic medications (either before or after diagnosis) and lung cancer-specific mortality; including sulfonylureas, insulin or other anti-diabetic medications (such as thiazolidinediones). Overall, the results from this population-based study provide little evidence of a protective association between metformin use and cancer mortality in lung cancer patients.

Keywords: metformin; lung cancer survival; pharmacoepidemiology; CPRD

1. Introduction

Emerging evidence from preclinical studies suggests a potential antitumour effect for metformin, a biguanide derivative commonly prescribed to treat type 2 diabetes, possibly due to the activation of AMP-activated protein kinase, leading to the inhibition of the mTOR signalling pathway [1]. Metformin use has been associated with improved survival in patients with diabetes with breast [2], colorectal [3] and prostate cancer [4]; however there are limited data in lung cancer patients. Constitutive mTOR signaling plays an important role in the development of lung cancer and *in vitro* and animal studies have reported antiproliferative effects [5] and proapoptotic effects [6], as well as the inhibition of lung tumour growth [7]. Few epidemiological studies have assessed the influence of metformin on lung cancer-specific outcomes after diagnosis. Some, but not all [8] have reported improvements in progression-free survival [9] and disease-free survival [10,11] but these investigations were limited by small sample sizes [8-11], drug exposure determined at limited numbers of time-points [8-11] and a lack of adjustment for important baseline patient characteristics [9]. Other studies have investigated metformin use and overall survival [12-14], but these results could reflect mortality from non-cancer causes.

In this population-based study, we investigate for the first time the impact of metformin use, either before or after diagnosis, on lung cancer-specific survival. Secondary analysis investigated the influence of other anti-diabetic medications.

2. Materials and Methods

2.1 Study design

As described previously [15], we utilised record linkages between the English National Cancer Data Repository (NCDR) and the United Kingdom (UK) Clinical Practice Research Datalink (CPRD) to identify lung cancer patients diagnosed between 1998 and 2009. Office of National Statistics (ONS) mortality data provided deaths up to 2012. We restricted the cohort to those with a prior diagnosis of type 2 diabetes (based on a GP diagnosis of diabetes or at least one anti-diabetic medication prescription). Patients with type 1 diabetes were excluded and were defined as those with a diagnosis of type 1 diabetes in addition to at least one prescription for insulin before their cancer diagnosis. Patients were also excluded if they had a previous cancer (with the exception of non-melanoma skin cancers) or if their diagnosis date preceded CPRD research quality records. Ethical approval for all observational studies conducted using CPRD data has been obtained from a multicentre research ethics committee.

2.2 Exposure data

Data on post-diagnostic anti-diabetic medication usage was determined from general practitioner (GP) prescribing records contained within the CPRD according to the British National Formulary (BNF) [16] including metformin, sulfonylureas, insulin or other anti-diabetic medications (including thiazolidinediones).

2.3 Covariates

The NCDR provided clinical data on tumour histology and receipt of cancer treatments including surgery, chemotherapy and radiotherapy. Information on lifestyle factors before diagnosis including smoking, alcohol and BMI was obtained from CPRD records. Comorbidities prior to diagnosis were derived from GP records and included those within a

recent adaptation of the Charlson Comorbidity index [17]. A measure of deprivation was available from CPRD records [18]. Other medications including low-dose aspirin and statins were determined from GP-prescription records. Haemoglobin A1c (HbA1c) levels in the year prior to diagnosis were obtained from GP records.

2.4 Statistical analysis

2.4.1 Statistical analysis for anti-diabetic medication use after diagnosis

The analysis of drug use after diagnosis used Cox regression models to produce unadjusted and adjusted hazard ratios (HR) and 95% confidence intervals (CI) for the association between anti-diabetic medication usage and lung cancer-specific mortality. Patients that died in the first 6 months after diagnosis were removed as drug use during this time is unlikely to exert an effect on cancer death. Drug use was treated as time-varying [19], and was lagged by 6 months to remove prescriptions immediately prior to death (as these may reflect end-of-life care). Dose-response relationships were investigated for metformin use by increasing number of prescriptions and defined daily doses (DDDs), as defined by the WHO [20]. Sub-group analyses for metformin use and lung cancer mortality were carried out by pre-diagnostic drug use and tumour histology. Separate sensitivity analyses additionally adjusted for smoking and HbA1c level (among individuals with available information) and type 2 diabetes duration (among individuals with at least 5 years of records before diagnosis).

2.4.2 Statistical analysis for anti-diabetic medication use before diagnosis

Cox regression models were utilised to calculate unadjusted and adjusted HRs and 95% CIs for the association between anti-diabetic medication usage in the year prior to diagnosis and lung

cancer-specific mortality. Patients with less than 1 year of CPRD records prior to diagnosis were removed. Similarly, dose-response relationships for metformin use were examined. Sensitivity analyses additionally adjusted for smoking, HbA1c level and type 2 diabetes duration.

3. Results

3.1 Patient cohort

A total of 1,443 lung cancer patients with diabetes were identified. The analysis of anti-diabetic medication use after diagnosis included 533 patients, after 910 patients with less than 6 months follow-up were excluded. The analysis of anti-diabetic medication use before diagnosis included 1,350 patients, after 93 patients with less than 1 year of records prior to diagnosis were excluded. Patient characteristics by metformin use are listed in Table 1. Users were more likely to be diagnosed more recently, be younger, be obese, to have had surgery and chemotherapy and to be users of statins and other anti-diabetic medications (except insulin). Some comorbidities were less common in users of metformin after diagnosis, Table 1.

3.2 Association between anti-diabetic medication use after diagnosis and survival

The results for the association between anti-diabetic medication use after diagnosis and survival are presented in Table 2. Metformin use after diagnosis was weakly associated with a reduction in lung cancer death, however, after adjustment for potential confounders, results attenuated (adjusted HR 0.86, 95% CI 0.68-1.09) and there was no significant dose-response relationship by increasing number of prescriptions or DDDs. In sub-group analyses, results were similar after stratification by pre-diagnostic metformin use (non-users: adjusted HR 0.71, 95% CI 0.45-1.13; users: adjusted HR 0.83, 95% CI 0.41-1.69, p for interaction=0.52) and tumour histology (small cell lung cancer: adjusted HR 0.61, 95% CI 0.23-1.60; non-small cell

lung cancer: adjusted HR 0.85, 95% CI 0.63-1.14), p for interaction=0.87). Additional adjustment for other factors did not materially alter effect estimates, including smoking (adjusted HR 0.86, 95% CI 0.68-1.10), HbA1c level (adjusted HR 0.88, 0.68-1.13) and diabetes duration (adjusted HR 0.82, 95% CI 0.62-1.10).

There was no evidence of an association for sulfonylurea or insulin use. There was a suggestion of an increased risk of cancer mortality with the use of other anti-diabetic medication (including thiazolidinediones); but analysis was not significant and based on only 45 users (adjusted HR 1.42, 95% CI 0.94-1.13).

3.3 Association between anti-diabetic medication use before diagnosis and survival

No association between metformin use prior to diagnosis and lung cancer-specific mortality was observed and no dose-response association was apparent, Table 2. The findings remained largely unchanged after additional adjustment for smoking, HbA1c level and type 2 diabetes duration (data not shown). Sulfonylurea, insulin and other anti-diabetic medication use were not associated with cancer-specific mortality, Table 2.

3. Discussion

Using population-based data, we observed a 14% non-significant reduction in cancer-specific mortality with metformin use after diagnosis in a large cohort of lung cancer patients with pre-existing type 2 diabetes, however this reduction was weak in magnitude and there was no clear evidence of a dose-response association.

No study has investigated the influence of metformin use on cancer-specific survival in a cohort of lung cancer patients. Inconsistent with our findings, two previous studies reported significant reductions in all-cause mortality in users of metformin before [13] or after [14] lung cancer

diagnosis. A further study reported a shorter overall survival in pre-diagnostic metformin users compared to non-users [12]. Unlike our investigation, previous studies did not investigate dose-dependent relationships [12-14] and metformin use after diagnosis was time-fixed [14], thus potentially introducing immortal time bias. Moreover, results from studies investigating all-cause mortality could be reflective of non-cancer mortality.

Strengths of this study included an unselected lung cancer cohort and record linkages between NCDR and ONS data, allowing for robust verification of cancer diagnoses and deaths. High-quality GP prescribing data [21] eliminated any recall bias and allowed for temporal associations to be examined. Furthermore, we provided an inclusive and simultaneous examination of all anti-diabetic medications and risk of lung cancer-specific mortality. The present study is the largest to investigate metformin use and outcomes after lung cancer diagnosis; however, a lack of statistical power in sub-group analyses is possible. Although we did not have information on medication compliance, it is unlikely that this is affecting our results because similar findings were observed in dose-response analyses. We were also unable to adjust for cancer stage, thus potentially introducing residual confounding.

4. Conclusion

In conclusion, the results from this novel population-based study provide little evidence that metformin use is associated with increased lung cancer survival. Further investigation in larger sample sizes is required to verify these findings.

Conflict of interest statement: The authors declared no conflict of interest.

Financial support: CRC and ÚMM were supported by a Health and Social Care Research and Development, Public Health Agency, Northern Ireland, funded UK NIHR Career Development Fellowship, which also funded access to the CPRD dataset.

Acknowledgments: This study is based in part on data from the Clinical Practice Research Datalink obtained under licence from the UK Medicines and Healthcare products Regulatory Agency. However, the interpretation and conclusions contained in this study are those of the authors alone.

References

1. Dowling RJO, Goodwin PJ, Stambolic V. Understanding the benefit of metformin use in cancer treatment. *BMC Med* 2011; 9:33.
2. Vissers PAJ, Cardwell CR, van de Poll-Franse LV, Young IS, Pouwer F, Murray LJ. The association between glucose-lowering drug use and mortality among breast cancer patients with type 2 diabetes. *Breast Cancer Res Treat* 2015;150:427-37.
3. Garrett CR, Hassabo HM, Bhadkamkar NA, Wen S, Baladandavuthapain V, Kee BK, Eng C, Hassan MM. Survival advantage observed with the use of metformin in patients with type II diabetes and colorectal cancer. *Br J Cancer* 2012;106:1374-8.
4. Margel D, Urbach DR, Lipscombe LL, Bell CM, Kulkarni G, Qustin PC, Fleshner N. Metformin use and all-cause and prostate cancer-specific mortality among men with diabetes. *J Clin Oncol* 2013;31:3069-75.
5. Storozhuk Y, Hopmans SN, Sanli T, Barron C, Tsiani E, Cutz JC, Pond G, Wright J, Singh G, Tsakiridis T. Metformin inhibits growth and enhances radiation response of non-small cell lung cancer (NSCLC) through ATM and AMPK. *Br J Cancer* 2013; 108:2021-31.
6. Wu N, Gu C, Gu H, Hu H, Han Y, Li Q. Metformin induces apoptosis of lung cancer cells through activating JNK/p38 MAPK pathway and GADD153. *Neoplasma* 2011;58:482-490.
7. Memmott RM, Mercado JR, Maier CR, Kawabata S, Fox SD, Dennis PA. Metformin prevents tobacco carcinogen--induced lung tumorigenesis. *Cancer Prev Res* 2010; 3:1066-76.

8. Ahmed I, Ferro A, Cohler A, Langenfeld J, Surakanti SG, Aisner J, Zou W, Haffty BG, Jabbour SK. Impact of metformin use on survival in locally-advanced, inoperable non-small cell lung cancer treated with definitive chemoradiation. *J Thorac Dis* 2015;7:346-55.
9. Tan BX, Yao WX, Ge J, Peng XC, Du XB, Zhang R, Yao B, Xie K, Li LH, Dong H, Gao F, Zhao F, Hou JM, Su JM, Liu JY. Prognostic influence of metformin as first-line chemotherapy for advanced nonsmall cell lung cancer in patients with type 2 diabetes. *Cancer* 2011; 117:5103-11.
10. Xu T, Liang, G, Yang, L, Zhang F. Prognosis of small cell lung cancer patients with diabetes treated with metformin. *Clin Transl Oncol* 2015 [Epub ahead of print].
11. Kong F, Gao F, Liu H, Chen L, Zheng R, Yu J, Li X, Liu G . Metformin use improves the survival of diabetic combined small-cell lung cancer patients. *Tumour Biol* 2015 [Epub ahead of print].
12. Mazzone PJ, Rai H, Beukemann M, Xu M, Jain A, Sasidhar M. The effect of metformin and thiazolidinedione use on lung cancer in diabetics. *BMC Cancer* 2012;12:410.
13. Lin JJ, Gallagher EJ, Sigel K, Mhango G, Galsky MD, Smith CB, LeRoith D, Wisnivesky JP. Survival of patients with stage IV lung cancer with diabetes treated with metformin. *Am J Respir Crit Care Med* 2015; 191:448-54.
14. Currie CJ, Poole CD, Jenkins-Jones S, Gale EA, Johnson JA, Morgan CL. Mortality after incident cancer in people with and without type 2 diabetes: impact of metformin on survival. *Diabetes Care*. 2012; 35:299-304.

15. Cardwell CR, Mc Menamin Ú, Hughes CM, Murray LJ. Statin use and survival from lung cancer: a population-based cohort study. *Cancer Epidemiol Biomarkers Prev* 2015; 24:833-41.
16. British National Formulary 66th ed. 2013. Joint Formulary Committee. London, UK: BMJ Group and Pharmaceutical Press.
17. Khan NF, Perera R, Harper S, Rose PW. Adaptation and validation of the Charlson Index for Read/OXMIS coded databases. *BMC Fam Pract* 2010; 11:1.
18. Noble M, Wright G, Dibben C, et al. Indices of deprivation 2004: Report to the office of the deputy prime minister. London, UK, Neighb Renew Unit. 2004.
19. Lévesque LE, Hanley JA, Kezouh A, Suissa S. Problem of immortal time bias in cohort studies: example using statins for preventing progression of diabetes. *Br Med J* 2010; 340:907-911.
20. World Health Organisation: World Health Organisation Collaborating Centre for Drug Statistics Methodology. Available from: <http://www.whocc.no/>.
21. Jick H, Jick SS D LE. Validation of information recorded on general practitioner based computerised data resource in the United Kingdom. *Br Med J* 1991;302:766–8.

Table 1. Characteristics of lung cancer patients by metformin use

Characteristics	Metformin use in year prior to diagnosis ^a		Metformin use in year after diagnosis ^b		Metformin use after diagnosis ^b	
	User n (%)	Non-user n (%)	User n (%)	Non-user n (%)	User n (%)	Non-user n (%)
	(n=1,350)		(n=533)		(n=533)	
Year of diagnosis: 1998-2000	45 (7)	108 (16)	25 (9)	39 (16)	23 (9)	41 (15)
2001-2003	113 (17)	163 (24)	33 (12)	57 (23)	29 (12)	61 (22)
2004-2006	175 (26)	191 (28)	84 (30)	58 (23)	71 (28)	71 (25)
2007-2009	331 (50)	224 (33)	141 (50)	96 (38)	128 (51)	109 (39)
Age at diagnosis: < 50	1 (0)	8 (1)	0 (0)	7 (3)	0 (0)	7 (2)
50-59	57 (9)	57 (9)	33 (12)	12 (5)	30 (12)	15 (5)
60-69	183 (28)	183 (28)	90 (32)	67 (27)	81 (32)	76 (27)
70-79	286 (43)	286 (43)	116 (41)	109 (44)	104 (41)	121 (43)
80-89	130 (20)	130 (20)	43 (15)	54 (22)	35 (14)	62 (22)
≥ 90	7 (1)	7 (1)	1 (0)	1 (0)	1 (0)	1 (0)
Gender: Males	437 (66)	445 (65)	190 (67)	153 (61)	174 (64)	169 (65)
Treatment within 6 months of cancer diagnosis						
Surgery ^c	77 (15)	47 (9)	71 (32)	26 (13)	67 (33)	30 (14)
Chemotherapy	153 (23)	100 (15)	100 (35)	69 (28)	90 (36)	79 (28)
Radiotherapy	182 (27)	196 (29)	103 (36)	100 (40)	88 (35)	115 (41)
Histology: Non-small cell	371 (56)	367 (54)	206 (73)	159 (64)	184 (73)	181 (64)
Small cell	86 (13)	76 (11)	38 (13)	31 (12)	33 (13)	36 (13)
Missing	207 (31)	243 (35)	39 (14)	60 (24)	34 (14)	65 (23)
Smoking status prior to cancer diagnosis						
Non-smoker	101 (15.2)	99 (14)	41 (15)	35 (14)	37 (15)	39 (14)
Ex-smoker	331 (50)	324 (47)	143 (51)	121 (48)	127 (51)	137 (49)
Current smoker	220 (33)	224 (33)	92 (32)	82 (33)	80 (32)	94 (33)
Missing	12 (2)	39 (6)	7 (2)	12 (5)	7 (3)	12 (4)
Alcohol consumption prior to diagnosis						
Never	174 (26)	138 (20)	68 (24)	54 (22)	61 (24)	61 (22)
Ever	449 (68)	466 (68)	198 (70)	171 (68)	175 (70)	194 (69)
Missing	41 (6)	82 (12)	17 (6)	25 (10)	15 (6)	27 (10)
BMI (kg/m²) prior to diagnosis: mean (sd)						
Underweight (<18.5)	11 (2)	20 (3)	4 (1)	8 (3)	4 (2)	8 (3)
Normal (18.5 to 25)	174 (26)	211 (31)	58 (21)	68 (27)	52 (21)	74 (26)
Overweight (25-30)	236 (36)	239 (35)	103 (36)	95 (38)	88 (35)	110 (39)
Obese (>30)	219 (33)	156 (23)	106 (38)	67 (27)	96 (38)	77 (27)
Missing	24 (4)	60 (9)	12 (4)	12 (5)	11 (4)	13 (5)
Deprivation fifth: 1 st (least deprived)	102 (15)	102 (15)	52 (18)	36 (14)	45 (18)	43 (15)
2 nd	132 (20)	141 (21)	53 (19)	60 (24)	49 (20)	64 (23)
3 rd	120 (18)	143 (21)	50 (18)	52 (21)	44 (18)	58 (21)
4 th	150 (23)	154 (23)	57 (20)	50 (20)	49 (20)	58 (21)
5 th (most deprived)	158 (24)	144 (21)	68 (24)	51 (20)	61 (24)	58 (21)
Missing	2 (0)	2 (0)	3 (1)	1 (0)	3 (1)	1 (0)
Comorbidity prior to cancer diagnosis						
Cerebrovascular disease	79 (12)	97 (14)	22 (8)	37 (15)	19 (8)	40 (14)
Chronic pulmonary disease	174 (26)	199 (29)	68 (24)	81 (32)	64 (26)	85 (30)
Congestive heart disease	60 (9)	80 (12)	16 (6)	26 (10)	13 (5)	29 (10)
Myocardial infarction	103 (16)	106 (16)	36 (13)	38 (15)	34 (14)	40 (14)
Peptic ulcer disease	49 (7)	53 (8)	24 (9)	19 (8)	18 (7)	25 (9)
Peripheral vascular disease	129 (19)	115 (17)	43 (15)	35 (14)	39 (16)	39 (14)
Renal disease	91 (14)	118 (17)	30 (11)	49 (20)	27 (11)	52 (18)
Other medication use after diagnosis						
Low-dose aspirin use ^d	371 (56)	350 (51)	157 (56)	123 (49)	136 (54)	106 (38)
Statin use ^d	461 (69)	328 (48)	201 (71)	130 (52)	184 (73)	117 (42)
Anti-diabetic medication after diagnosis						
Sulfonylureas ^d	328 (49)	236 (34)	114 (40)	92 (37)	126 (50)	84 (30)
Insulin ^d	80 (12)	102 (15)	27 (10)	46 (18)	41 (16)	51 (18)
Other ADDs ^d	94 (14)	30 (4)	37 (13)	13 (5)	30 (12)	15 (5)
HbA1c level in year prior to diagnosis						
≤6.5%	161 (24)	223 (34)	62 (22)	87 (35)	54 (22)	95 (34)
>6.5%	478 (72)	340 (50)	206 (73)	127 (51)	184 (73)	149 (53)
Missing	25 (4)	113 (17)	15 (5)	36 (14)	13 (5)	38 (14)

^a Analysis includes lung cancer patients who have more than 1 year of records prior to diagnosis. ^b Analysis includes lung cancer patients who live more than 6 months after diagnosis. ^c Excluding cancer patients from Thames Registry as surgery information not available. ^d Other medication use after diagnosis for metformin use after diagnosis columns, other medication use in year prior to diagnosis for metformin use in year prior to diagnosis columns. ADD= anti-diabetic drug

Table 2: Association between anti-diabetic medication usage and cancer-specific mortality in lung cancer patients with type 2 diabetes

	Users			Non-users			Unadjusted HR (95% CI)	P	Adjusted HR (95% CI)	P
	Cancer-specific deaths	All patients	Person years	Cancer-specific deaths	All patients	Person years				
ADD use after diagnosis^{a,b}							(n=533)		(n=533)	
Metformin use ^c	149	251	386	211	282	305	0.80 (0.66-1.00)	0.05	0.86 (0.68-1.09)	0.21
Metformin use 1-11 prescriptions ^d	119	159	217	211	282	305	0.82 (0.65-1.03)	0.09	0.87 (0.68-1.11)	0.26
Metformin use ≥12 prescriptions ^d	30	92	169				0.75 (0.47-1.17)	0.20	0.82 (0.51-1.30)	0.40
Metformin use 1-365 DDDs ^d	121	172	236	211	282	305	0.79 (0.63-0.99)	0.05	0.84 (0.65-1.07)	0.16
Metformin use ≥365 DDDs ^d	28	79	150				0.91 (0.56-1.48)	0.71	1.01 (0.62-1.67)	0.96
Sulfonylurea use ^c	132	210	278	228	323	412	1.10 (0.88-1.37)	0.42	1.10 (0.87-1.40)	0.41
Insulin use ^c	55	92	137	305	441	554	1.02 (0.75-1.34)	0.96	1.06 (0.78-1.45)	0.70
Other ADD use ^c	30	45	54	330	488	637	1.21 (0.83-1.76)	0.32	1.42 (0.94-1.13)	0.09
ADD use in year prior to diagnosis^{e,f}							(n=1,350)		(n=1,350)	
Metformin use	514	664	536	558	686	505	0.89 (0.79-1.00)	0.06	0.97 (0.86-1.11)	0.67
Metformin use 1-11 prescriptions	375	483	403	558	686	505	0.88 (0.77-1.00)	0.05	0.95 (0.83-1.10)	0.51
Metformin use ≥12 prescriptions	139	181	133				0.92 (0.76-1.11)	0.38	1.03 (0.85-1.25)	0.77
Metformin use 1-365 DDDs	455	577	467	558	686	505	0.91 (0.80-1.02)	0.12	0.98 (0.86-1.12)	0.78
Metformin use ≥365 DDDs	59	87	69				0.78 (0.60-1.02)	0.07	0.90 (0.68-1.19)	0.47
Sulfonylurea use	453	564	389	619	786	651	1.11 (0.99-1.26)	0.08	1.07 (0.94-1.21)	0.31
Insulin use	143	182	148	929	1,168	892	0.93 (0.78-1.11)	0.40	1.00 (0.84-1.21)	0.97
Other ADD use	99	124	103	973	1,226	938	0.91 (0.74-1.13)	0.40	0.98 (0.79-1.21)	0.84

^a Analysis includes lung cancer patients who live more than 6 months after diagnosis

^b Adjusted analyses controlled for year of diagnosis, age at diagnosis, sex, radiotherapy within 6 months, chemotherapy within 6 months, comorbidities (prior to diagnosis, including cerebrovascular disease, chronic pulmonary disease, congestive heart disease, diabetes, myocardial infarction, peptic ulcer disease, peripheral vascular disease, renal disease), other anti-diabetic medication use (after diagnosis, as time varying covariates, including metformin, sulfonylureas, insulin and other ADDs), other medication use (after diagnosis, as time varying covariates, specifically low-dose aspirin and statins) and deprivation (in fifths).

^c Medication use modelled as a time varying covariate with an individual considered a non-user prior to 6 months after first medication usage and a user after this time, excludes deaths in the first 6 months after cancer diagnosis.

^d Medication use modelled as a time varying covariate with an individual considered a non-user prior to 6 months after first medication usage, a user of 0 to 12 prescriptions from 6 months after first prescription to 6 months after 12th prescription (or 365 DDDs) and a greater user after this time, excludes deaths in the 6 months after cancer diagnosis.

^e Analysis includes lung cancer patients who have more than 1 year of records prior to diagnosis

^f Adjusted analyses controlled for year of diagnosis, age at diagnosis, sex, comorbidities (prior to diagnosis, including cerebrovascular disease, chronic pulmonary disease, congestive heart disease, diabetes, myocardial infarction, peptic ulcer disease, peripheral vascular disease, renal disease), other anti-diabetic medication use (in year prior to diagnosis, including metformin, sulfonylureas, insulin and other ADDs), other medication use (in year prior to diagnosis, specifically low-dose aspirin and statins) and deprivation (in fifths).
ADD= anti-diabetic drug