



**QUEEN'S
UNIVERSITY
BELFAST**

Expression pattern of arenicins - the antimicrobial peptides of polychaete *Arenicola marina*

Maltseva, A., Kotenko, O., Kokryakov, V., Starunov, I., & Krasnodembskaya, A. (2014). Expression pattern of arenicins - the antimicrobial peptides of polychaete *Arenicola marina*. *Frontiers in Physiology*, Article 120063. <https://doi.org/10.3389/fphys.2014.00497>

Published in:
Frontiers in Physiology

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2014 Maltseva, Kotenko, Kokryakov, Starunov and Krasnodembskaya. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

Expression pattern of arenicins - the antimicrobial peptides of polychaete *Arenicola marina*

Arina L. Maltseva, Olga N. Kotenko, Vladimir N. Kokryakov, Viktor V. Starunov and Anna D. Krasnodembskaya

Journal Name:	Frontiers in Physiology
ISSN:	1664-042X
Article type:	Original Research Article
Received on:	24 Sep 2014
Accepted on:	02 Dec 2014
Provisional PDF published on:	02 Dec 2014
www.frontiersin.org:	www.frontiersin.org
Citation:	Maltseva AL, Kotenko ON, Kokryakov VN, Starunov VV and Krasnodembskaya AD(2014) Expression pattern of arenicins - the antimicrobial peptides of polychaete <i>Arenicola marina</i> . <i>Front. Physiol.</i> 5:497. doi:10.3389/fphys.2014.00497
Copyright statement:	© 2014 Maltseva, Kotenko, Kokryakov, Starunov and Krasnodembskaya. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY) . The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after rigorous peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

Expression pattern of arenicins - the antimicrobial peptides of polychaete *Arenicola marina*

Arina L. Maltseva^{*1}, Olga N. Kotenko¹, Vladimir N. Kokryakov^{2,3}, Viktor V. Starunov¹, Anna D. Krasnodembskaya⁴

¹ - Department of Invertebrate Zoology, St-Petersburg State University, St-Petersburg, Russia

² - Department of Biochemistry St-Petersburg State University, St-Petersburg, Russia

³ - Department of General Pathology and Pathophysiology, Institute of Experimental Medicine, Russian Academy of Medical Sciences, Saint Petersburg, Russia.

⁴ - Centre for Infection and Immunity, School of Medicine, Dentistry and Biomedical Sciences, Queens University Belfast, UK

* Corresponding author:

Arina L. Maltseva

St-Petersburg State University

Department of Invertebrate Zoology

Universitetskaya 7/9

St-Petersburg, 199034, Russia

a.maltseva@bio.spbu.ru

Abstract

Immune responses of invertebrate animals are mediated through innate mechanisms, among which production of antimicrobial peptides play an important role. Although evolutionary Polychaetes represent an interesting group closely related to a putative common ancestor of other coelomates, their immune mechanisms still remain scarcely investigated. Previously our group has identified arenicins - new antimicrobial peptides of the lugworm *Arenicola marina*, since then these peptides were thoroughly characterized in terms of their structure and inhibitory potential. In the present study we addressed the question of the physiological functions of arenicins in the lugworm body. Using molecular and immunocytochemical methods we demonstrated that arenicins are expressed in the wide range of the lugworm tissues - coelomocytes, body wall, extravasal tissue and the gut. The expression of arenicins is constitutive and does not depend on stimulation of various infectious stimuli. Most intensively arenicins are produced by mature coelomocytes where they function as killing agents inside the phagolysosome. In the gut and the body wall epithelia arenicins are released from producing cells via secretion as they are found both inside the epithelial cells and in the contents of the cuticle. Collectively our study showed that arenicins are found in different body compartments responsible for providing a first line of defence against infections, which implies their important role as key components of both epithelial and systemic branches of host defence.

Key words: antimicrobial peptides; invertebrate immunity; coelomocytes; *Arenicola marina*; annelid.

Introduction

Antimicrobial peptides (AMPs) are relatively small (not exceeding 100 amino acids) usually cationic polypeptidic molecules with prominent inhibitory potential against various microbial pathogens. It is well recognized that AMPs constitute an important component of the innate immunity, having a role in both effector and regulatory functions. The fact that AMPs are present in a wide range of organisms including plants, vertebrate and invertebrate animals, protists and prokaryotes (rev. in Boman, 2003; Reddy et al., 2004; Yeaman & Bayers, 2006; Otero-González et al., 2010; Pasupuleti et al., 2012) points to their importance for host defence against infections throughout the evolutionary process.

1 The wide distribution of AMPs makes them an attractive object for comparative immunology. Such
2 studies can provide insights into the mechanisms underpinning the evolution of innate immune
3 responses and also lead to identification of new effective molecular structures which could be
4 developed as new therapeutics. AMPs of invertebrate animals are of particular importance, as their
5 host defence against infections relies entirely on the mechanisms of innate immunity. Historically
6 insects represent a group of invertebrate animals where the immune system has drawn intensive
7 interest of investigators and consequently their AMPs are the most well characterized (Bulet et al.,
8 1999; Otvos, 2002; Bulet et al., 2005), whereas the knowledge about AMPs from other invertebrates
9 remains limited.

10
11 Annelids might be considered as a basal group which is closest to a putative common ancestor of other
12 coelomate invertebrates. Therefore their immunity is a fascinating object of study for comparative
13 immunologists. Besides this, annelids are important players in the benthic communities as they are
14 critical for biomass production in marine and freshwater ecosystems. Paradoxically, the existing
15 knowledge about mechanisms of their host defence is largely based on the studies of Oligochaetes
16 (Cooper et al., 2004; Kauschke et al., 2007; Bilej et al., 2010) or Hirudineans (Tasiemski & Salzet,
17 2010), which are relatively specialized branches of Annelids, whereas more basal Polychaetes still fall
18 out of scope of investigation.

19
20 More specifically, just a few species of annelids were investigated as possible sources of AMPs (rev.
21 in Tasiemski, 2008). Among polychaetes these are: *Perinereis aibuchitensis* (Pan et al., 2004), *Nereis*
22 *diversicolor* (Tasiemski et al., 2007) - two closely related genera - and *Arenicola marina*
23 (Ovchinnikova et al., 2004). There is no structural homology between AMPs from listed species as
24 well as with any other published AMP. Up to date, only hedistin from *N. diversicolor* was studied in
25 terms of its functional activity within the host. Hedistin was shown to be exclusively and constitutively
26 expressed in so called NK cells (one of the coelomocyte types) (Tasiemski et al., 2007).

27
28 Arenicins-1 and -2, identified by our group in *A. marina*, are 21 residue peptides. They differ by one
29 amino acid replacement and their β -hairpin structure is supported by one disulphide bond
30 (Ovchinnikova et al., 2004). Structural properties make them feasible for chemical and recombinant
31 synthesis (Kolodkin et al., 2006; Ovchinnikova et al., 2007). These peptides have been extensively
32 studied during the last decade (Lee et al., 2007; Andra et al, 2008; Park & Lee, 2009; Andra et al,
33 2009; Cho & Lee, 2011). Collectively, structural and functional properties make them promising
34 candidates for preclinical studies. However, from the comparative immunological perspective, it is
35 important to establish the functional role of these peptides for the host defence of the organism.

36
37 In the present study we addressed the question of the physiological functions of arenicin 1 and 2 in the
38 lugworm body as components of its immune system with a specific focus on the spatial distribution of
39 their localization in the tissues and the pattern of their expression upon infectious stimuli.

40 41 **Material and methods**

42 **1. Animals and tissues**

43 **1.1. Animals.** Adult lugworms (apprx. 2.5 cm length) *Arenicola marina* individuals were collected
44 from wild populations over the intertidal zone in the vicinity of the White Sea Marine biological
45 station of Saint Petersburg University (Russia). Animals were maintained in permanently aerating
46 static tanks with marine water for 5-7 days.

47
48 **1.2. Stimulation.** To induce immune responses worms were injected by 10^7 CFU suspension of heat
49 inactivated *Escherichia coli* ML35p, *Listeria monocytogenes* EGD, *Candida albicans* 820 (alone or in
50 combination) in 1 ml of sterile marine water (SMW), control animals received SMW; all microbial

1 strains were kindly provided by Prof. R.Lehrer, University of California, Los Angeles, USA. Control
2 animals were injected by equivalent volume of SMW. Coelomic fluid (CF) of 5 animals representing
3 each group was pooled at 24 and 48 h after immunization. CF was collected from the middle part of a
4 lungworm body with a sterile syringe (d = 0.6 mm) needle filled with an ice-cold SMW to prevent
5 coagulation: CF to SMW ratio remained 1 : 2. CF was sedimented at 400 g for 10 min at 4°C directly
6 after collection. Precipitated cells were suspended in cold SMW for smear preparation or used for
7 RNA extraction.

8
9 **1.3. Other tissues.** Before dissection animals were anesthetized in 5% MgCl₂ in SMW whereupon
10 pieces of body wall, salivary glands, foregut and midgut tissue were dissected, placed into sterile tubes
11 and fixed for RNA extraction.

12 13 **2. PCR**

14 **2.1. Template.** RNA was isolated from fixed and homogenized in PureZOL (Cat. No. #732-6890,
15 BioRad Inc, CA, USA) tissues, using the RNAeasy kit (Cat. No. 74104 Qiagen Inc, CA, USA). After
16 DNase I (Cat. No. #EN0521 Fermentas, Thermo Fisher Scientific Inc, MA, USA) treatment (30 min at
17 room temperature) and purification RNA quality was verified with NanoDrop® ND-1000 UV-Vis
18 Spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). 260/280 and 260/230 nm
19 absorbance ratios did not exceed 2.0 indicating appropriate RNA quality.

20
21 **2.2. Semiquantative PCR.** Arenicin-1, arenicin-2 and actin primers were custom made by Syntol C°
22 (Moscow, Russia). The sequences were as follows.

23 Arenicin-1: forward 5'-CTAATCCTGGCCATTTTCTGCG-3'; reverse 5'-
24 CCCTGAGCTGACTGGAAATAG-3'; product 338 bp.

25 Arenicin-2: forward 5'-GCGAGATCGGCTGGAGAG-3'; reverse 5'-CCCTGAGCTGACCGGAAG-
26 3'; product 254 bp.

27 Actin: forward 5'-CAAATCATGTTTCGAGACCTTC-3'; reverse 5'-GCTGATCCACATCTGTTGG3';
28 product 714 bp.

29
30 PCR was performed with SuperScript™ III One-Step RT-PCR System with Platinum® Taq DNA
31 Polymerase protocol (Cat. No. 12574-018, Invitrogen, Thermo Fisher Scientific Inc, MA, USA)
32 according to manufacturer's instructions. β -Actin gene was used as a control to normalize a template
33 loading. Amplification was carried out at C1000X cycler (BioRad Inc, CA, USA). The PCR products
34 were then sequenced in Resource Center for Molecular and Cellular Technologies of St. Petersburg
35 State University (St.Petersburg, Russia).

36
37 **2.3. qRT-PCR.** Real-time PCR was performed with iQ™ SYBR® Green Supermix (Cat. No. 170-
38 8880, BioRad, CA, USA). Five independent RNA isolations with following cDNA synthesis and RT
39 PCR were done for each experimental case. Reactions were conducted on CFX100 cycler (BioRad,
40 CA, USA). Actin was used as a housekeeping gene for normalization and results were analyzed by the
41 Δ CT method to calculate expression changes, P<0.05 was considered to be statistically significant.

42 43 **3. Antibody**

44 Recombinant arenicin-2 (Ovchinnikova et al., 2007) was conjugated with cargo-protein and used to
45 obtain polyclonal antiserum. The conjugate was compounded with complete Freund adjuvant (Cat. No.
46 F5881, Sigma-Aldrich, MO, USA) and injected into rabbit males at a dose 1 μ g per gram. The same
47 dose was used for secondary injection 2 weeks later. Serum samples were collected three times - 4, 6
48 and 8 weeks after first immunization. The titer of reactive antibody in the serum samples was
49 examined by immunoblotting against natural (in crude extract) and synthetic arenicin-1 (Kolodkin et
50 al., 2006). Active samples were pooled and stored at 4°C. Specific anti-arenicin antibody was purified

1 from the serum by affine chromatography using CNBr-activated Sepharose (Cat. No. GE17-0430-01
2 Sigma-Aldrich, MO, USA) conjugated with recombinant arenicin-2. Anti-arenicin AB was eluted by
3 NaCl gradient and desalted on Amicon Ultra centrifugal filters Ultracel 3K (Cat. No. Z677094-24EA,
4 Millipore, Merck KGaA, Germany). AB specificity was tested by immunoblotting against natural (in
5 the crude extract) and synthetic arenicin-1 (Kolodkin at al., 2006).

6 7 **4. Immunoblotting**

8 Crude extracts of coelomocytes were obtained by homogenization of coelomocytes (prepared as
9 described earlier) in 5% acetic acid. Particles were sedimented by centrifugation (10000 g, 40 min,
10 4°C) and supernatants were aspirated and accepted as crude extracts. Synthetic arenicin-1 was kindly
11 provided by Dr. Nickolai I. Kolodkin (Institute of Highly Pure Biopreparations, St-Petersburg, Russia).
12 Both crude extract (50 µg of total protein) and arenicin-1 (0.5 µg) were loaded into acidic urea 12.5%
13 PAAG (Panym & Chalkly, 1969) and transferred onto PVDF immune blot membrane (Cat. No. 162-
14 0177, BioRad Inc, CA, USA) in TetraCell Miniprotean Chamber (BioRad Inc, CA, USA) under
15 standard conditions. The membrane was equilibrated in PBST (3 times, 10 min), blocked by bovine
16 serum albumin (Cat. No. A2153, Sigma-Aldrich, MO, USA) overnight and stained initially by
17 polyclonal serum or purified anti-arenicin AB (1 hr) and further 1 hr by secondary sheep HRP-
18 conjugated anti-rabbit AB (Cat. No. #31463, Thermo Scientific, Thermo Fisher Scientific Inc, MA,
19 USA). DAB (Cat. No. D8001, Sigma-Aldrich, MO, USA) was used for visualization.

20 21 **5. Immunocytochemistry**

22 **5.1. Cross-sections.** Lugworms (apprx. 2.5 cm length) were anesthetized in 5% MgCl₂ in SMW, fixed
23 in 4% paraformaldehyde (Cat. No. 158127, Sigma-Aldrich, MO, USA), subjected to standard alcohol
24 dehydration and via xylol embedded into paraffin. Serial cross-sections (4 µm) were placed on
25 polylysine-coated slides (Cat. No. P0425, Sigma-Aldrich, MO, USA) and dried at 37°C.

26
27 **5.2. Coelomocytes smears.** Fresh coelomocytes in suspension were placed on polylysine-coated slides
28 (Cat. No. P0425, Sigma-Aldrich, MO, USA), left to dry out at room temperature and fixed with 4%
29 paraformaldehyde (Cat. No. 158127, Sigma-Aldrich, MO, USA). For *in vitro* phagocytosis, fresh
30 coelomocyte suspension was placed into the plastic plate for 10 min to adhere in the humid chamber at
31 4°C. The suspension of zymosan particles (Cat. No. Z4250, Sigma-Aldrich, MO, USA) was added
32 (MIO 10 : 1) and after incubation for 15 min the preparations were washed twice with SMW and fixed
33 in 4% paraformaldehyde.

34
35 **5.3. Antibody staining.** Prepared cross-sections or smears were incubated sequentially in PBST (3
36 times, 20 min), PBST with diluted sheep serum 1 : 5 (1 hr) and anti-arenicin AB (1 : 50) in sheep-
37 serum or rabbit IgG overnight. Preparations were washed by PBST (3 times, 20 min) and stained by
38 HRP-conjugated secondary AB (1 : 500) (Cat. No. #31463, Thermo Scientific, Thermo Fisher
39 Scientific Inc, MA, USA). AB was visualized with DAB (Cat. No. D8001, Sigma-Aldrich, MO, USA).
40 Some preparations were additionally stained by hematoxylin-eosin.

41 42 **Coelomic fluid plasma testing.**

43 CF was obtained as described earlier. Plasma was separated from cells by centrifugation (400 g, 10
44 min, 4°C). For component analysis of plasma it was loaded onto HPLC analytical column (C18, D 4.6
45 mm, length 20 cm, grain 5 µm, pore 300 Å, Vydec, The Nest Group Inc., MA, USA) and separated
46 using Agilent 1260 HPLC system (Agilent Technologies, CA, USA). Elution was performed with a
47 gradient of 0-60% acetonitrile in presence of 0.05% trifluoroacetic acid during 1 hr. Absorbance at 220
48 nm and 280 nm was used for detection.

1 For antimicrobial testing native plasma was concentrated (3 : 1) in VacSpeed System (Labconco, MO,
2 USA). Two variations of test were applied as described in Lehrer et al., 1989. In the overlay test the
3 sample was first separated in the acidic urea PAAG by electrophoresis and then PAAG was laid over
4 1% agarose gel (Cat. No. A0169, Type I, Sigma-Aldrich, MO, USA) with embedded microorganisms.
5 In the radial diffusion test the whole probe was put into the well of 1% agarose gel with immobilized
6 microorganisms as above (4×10^6 CFU per plate). After 2.5 hr of diffusion, growth medium (Tryptic
7 Soy Broth for bacteria, Cat. No. M011, or Sabouraud medium for fungi, Cat. No. M013, HiMedia
8 Laboratories, Mumbai, India) was added and plates were incubated overnight.

9 10 **Results.**

11 **1. Arenicins mRNA is ubiquitously expressed in the lugworm tissues.**

12 The presence of arenicins-1 and -2 transcripts were examined in coelomocytes, body wall, salivary
13 glands, foregut and midgut by semi-quantitative RT-PCR. Arenicins mRNA expression was detected
14 in all types of analysed samples, however the highest levels of both arenicin-1 and 2 mRNA were
15 observed in coelomocytes (**Fig.1**; signal of arenicin-2 was markedly attenuated in all samples, **S1**).
16 Detection of arenicin transcripts in several different compartments of the worm body suggests their
17 ubiquitous expression and potential involvement in different types of host defence - local or systemic.
18 The next step was to test if their expression was sensitive to microbial invasion.

19 20 **2. mRNA expression of arenicins does not depend on the microbial stimulation.**

21 To mimic the infectious process worms were injected with heat-inactivated laboratory strains of
22 microorganisms into the coelomic cavity. In order to investigate the possible role of different
23 PAMPs/PRRs interaction in mediating the response, worms were challenged with representatives of
24 the three major different lineages of microorganisms (gram-positive and gram-negative bacteria and
25 fungi) in different combinations. At 24 h post microbial challenge, samples of different tissues were
26 collected and arenicins mRNA expression was quantified by Real-Time PCR. There was no change in
27 the levels of arenicin-1 and -2 expression post-stimulation of coelomocytes compared to intact or
28 SMW-injected animals (**Fig.2**). Similarly, injection of heat-inactivated microbes into the coelomic
29 cavity did not induce changes in arenicins mRNA expression in the gut or body wall, compared to the
30 controls (**S2**). These results demonstrate that expression of arenicin-1 and arenicin-2 is constitutive and
31 does not depend either on the nature of the pathogen or on the presence/absence of pathogens in
32 coelomic cavity.

33 34 **3. Immunohistochemistry confirms ubiquitous expression of arenicins in the lugworm tissues.**

35 Anti-arenicins rabbit pAB were purified by affine chromatography against recombinant arenicin-2.
36 Their specificity was confirmed by Western-blot against crude coelomocyte extract, synthetic arenicin-
37 1 being a positive control (**Fig.3**). Western blotting demonstrated that pAB was specific for the mature
38 arenicins fraction of the extract, but lacked the ability to discriminate between two isoforms of
39 arenicin. After testing, the pAB were used for immunohistochemical analysis of lugworm tissues.

40
41 Paraffin-embedded tissue cross-sections were pAB stained to characterize mature peptide localization.
42 Several compartments including the outermost part of body wall and gut wall proved to be pAB-
43 immunoreactive (**Fig.4, A-B**), which correlated with the PCR result. The arenicin-positive part of the
44 body wall is continuous and represented by a thin cuticle and underlying epithelium lining.
45 Unexpectedly the arenicin signal came from chaetae complex (**Fig.4, C-F**). In addition there was
46 evident arenicin-positive staining, associated with ventral nervous chain (even some additional
47 background staining was also noted **Fig. 5**).

48
49 In the gut wall arenicins were expressed only in sparse individual cells (**Fig.6**). These cells appeared to
50 be intercalated into the epithelial layer and morphologically distinct from the others - they were larger

1 in size and rounded as opposed to cylindrical enterocytes. Presumably these cells could be considered
2 as a type of glandular cell (**Fig.6, E-G**). In addition, high levels of arenicin expression were detected in
3 tissues associated with ventral blood vessel in contrast to dorsal ones (**Fig.7, A-C**). Usually this
4 compartment is described as extravasal or chloragogue tissue (**Fig.7, D-F**) (Gardiner, 1992).
5 Coelomocytes were not visible on these sections.

6 7 **4. In *Arenicola marina* coelomocytes arenicins are predominantly localized in the vesicular** 8 **compartment.**

9 Next we examined the expression of arenicins in coelomocytes. AMPs translated from individual
10 transcripts (in contrast to proteolytic fragments) usually include signal peptide and undergo co-
11 translational transfer into the endoplasmic reticulum. This implicates their localization in the vesicular
12 apparatus of the cell. In our previous study (Ovchinnikova et al, 2004) it was demonstrated that
13 arenicins are synthesized as large precursors including pre-peptidic part. Here we wanted to test
14 whether arenicins are stored in granules or other cellular compartments.

15
16 Immunostaining of the preparations of the coelomic fluid smears demonstrated heterogeneity in the
17 morphology of the coelomocyte population which supports previously reported observations (rev. in
18 Vetvicka & Sima, 2009). Cells varied in morphology and the extent to which they spread on the
19 substrate correlated with the number and size of the granules in their cytoplasm. Generally two main
20 cell types were recognized on the smears. First were ones which possessed hyaline cytoplasm. They
21 spread extensively and contained minimal large granules near the nucleus (granulocytes I, **Fig.8, A-D**).
22 The second type was characterized by numerous medium-sized granules filling up almost all of the
23 cytoplasm (granulocytes II). Presumably, this well developed granular apparatus prevented these cells
24 from effective spreading, resulting in a more rounded form with just short and broad pseudopodia
25 (**Fig.8, E,F**). Additionally, putative juvenile cells (small cells with high nucleus / cytoplasm ratio
26 without any granules) were also regularly observed as a minor part of the population (**Fig. 8, B-D**).
27 Noteworthy, both types of mature granulocytes showed arenicins immunopositivity, whereas juvenile
28 cells did not, suggesting that arenicins expression might be associated with maturation of
29 coelomocytes. The arenicin signal was clearly associated with the granular apparatus of the cell,
30 regardless of its organisation.

31 32 **5. Coelomic fluid does not contain arenicins.**

33 AMPs stored in cytosolic granules can be liberated from cells via exocytosis. We investigated if
34 arenicins could be found in coelomic fluid plasma (CFP). CFP was fractionated by HPLC and major
35 fractions were analysed by electrophoresis. Neat, as well as concentrated (1 : 10) CFP appeared to be
36 free from protein containing fractions with only one macromolecular component (MW > 10 kDa) in
37 CFP detected. However, its detection was not regular. Notably, neither whole CFP nor any of its
38 fractions showed detectable antimicrobial activity. Injection of microbes into coelomic cavity had no
39 effect on the spectrum of CFP fractions as well as on the antimicrobial activity of CFP. Collectively,
40 these results suggest that CFP does not contain anti-microbial components either under normal
41 condition or after the microbial challenge.

42 43 **6. In the preparations of *in vitro* phagocytosis of lugworm coelomocytes arenicins are co-localised** 44 **with the phagocytized particle.**

45 One of the main functions of AMPs is their participation in the process of microbial killing inside the
46 phagolysosome. To test this possibility we immunostained the *in vitro* preparations of phagocytosis
47 with anti-arenicin pAb. Both types of coelomocytes readily phagocytized available zymosan particles.
48 In the individual arenicin-positive cells, engulfed particles were stained brown with a well visible
49 brown rim surrounding them (**Fig.8, H,I**), indicating the presence of arenicins within phagolysosomes.
50 This suggests involvement of these peptides in the killing of phagocytized microbes.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

Discussion

In the present study we characterized the tissue distribution of previously described antimicrobial peptide arenicin, represented by two isoforms which differed by a single amino-acid replacement in the mature peptide (Ovchinnikova et al., 2004). Also arenicins-1 and -2 differ in 4 amino acids in the pro- and in 2 amino acids in the pre-peptide parts, respectively. It is still not ascertained as to whether these two peptides are the products of the same gene or of two individual genes. An unambiguous interpretation of the noticeable lapse in the PCR signal strength from two isoforms is not possible. However, as we did not reveal any other principal difference in their expression pattern (except signal strength) and as both isoforms were effectively recognized by the pAB, we will discuss them together.

There were three major findings in this study.

Firstly, arenicins are expressed in a wide range of tissues of the lugworm body in addition to coelomocytes where they were initially identified. The list of the tissues expressing arenicins includes several epithelia, which implies involvement of arenicins into both systemic and epithelial branches of immunity.

Secondly, expression of arenicins (resembling that of hedistin from *N. diversicolor* (Tasiemski et al., 2007)) is not sensitive to the presence of different PAMPs in coelomic fluid.

Finally, although coelomocytes are capable of production and storage of arenicins, their secretion into coelomic fluid was negligible. However, our study did demonstrate that arenicins participate in the pathogen inactivation within the phagolysosome.

Lugworms are common dwellers in muddy habitats of subtidal and intertidal zones. Habitats such as these are abundant in microbes. Lugworms spend their lives engulfing and disgorging the surrounding sediments. Both epithelia - body wall and the gut - are under permanent threat of being attacked by microbial pathogens. Our findings supported this as we detected the expression of arenicins using both PCR and immunohistochemical methods in both critical organism boundaries - the body wall and gut. This characterizes arenicins as key players in the epithelial defense.

The outermost part of the body wall which was positive for arenicin consists of a thin cuticle and one layered epithelium. The principal cell type constituting the surface epithelium in polychaetes is represented by so called "supporting" cells. These cells bear apical microvilli which penetrate cuticle and are responsible for its synthesis (rev. in Gardiner, 1992). Generally cuticular components include proteins embedded into carbohydrate scaffolds (rev. in Gardiner, 1992). Immunopositivity for arenicins was detected in both the cuticle and the underlying cellular layer. It was also observed inside the chaetae, which are cuticular derivatives. These data suggest that arenicins are implicated in protection of the outer boundary of worm body, alongside with the other proteins which are synthesized and secreted into the cuticle by supporting cells. Likewise, an AMP from oligochaete *Pheretima tshiliensis* (soil living worm) - PP-1, was immunodetected in the mucus, covering the body (although it was not detected in epidermal cells) (Wang et al., 2003).

Midventrally situated neuronal cord, lying upon circular musculature of the body wall, also proved to be arenicin-positive. The structure of the *A. marina* nervous system is not comprehensively characterized yet. In general, in polychaetes the outermost part of both segmentary ganglia and intersegmentary connectives is represented by projections of glial supporting cells. These projections span outside the cord and into it to provide mechanical support against muscular compression. (Golding, 1992). In *A. marina* segmentary ganglia are not differentiated at all - neurons are evenly

1 spread laterally along the ventral nervous chain (Wells, 1950). This is very well observed in negative
2 control preparations (Fig. 5, A). Outer parts of the neural cord (an envelope and basolateral sectors)
3 showed strongest signal of arenicin presence. It is possible that glial cells or neurons or both express
4 arenicins inside the neural cord. The expression of two AMPs by glia and neurons was demonstrated in
5 the medicinal leech *Hirudo medicinalis*. The expression was up-regulated in response to injury and
6 microbial challenge implicating the involvement of AMPs in both defense and regeneration
7 (Schikorski et al., 2008). The complexity and the degree of differentiation of the polychaetes ventral
8 nerve cord never reaches the magnitude found in leeches (Golding, 1992). *A. marina* is accepted as an
9 organism, which is incapable of regenerating any part of its body (Bely, 2006). Additionally, there is
10 no reason to suspect the secretion of arenicins into coelomic cavity (see below). All these facts
11 preclude direct extrapolation of revealed AMPs neuronal function in leech to lugworm. The
12 functioning of arenicins in the nervous system still remains to be elucidated although their presence in
13 this compartment indirectly supports the multifunctional nature of AMPs *in vivo*.

14
15 The gut is the second principal compartment of the lugworm as it exists in permanent contact with the
16 surrounding mud. Interestingly, our study revealed that arenicins are also expressed in this area. The
17 polychaete midgut usually consist of enterocytes as a dominant population and one or several types of
18 secretory gland cells (rev. Saulnier-Michel, 1992). *A. marina* midgut contains both cell types
19 (Kaganovskaya, 1978). Arenicin-positive cells are scattered irregularly in the epithelial layer and
20 morphologically resemble gland cells. This implies that arenicins may be secreted into the gut lumen,
21 although this requires further investigation. Similarly, AMPs of another annelid worm - theromicin and
22 theromyzin from a leech *Theromyzon tessulatum* - were shown to be expressed both in intestinal and
23 epidermal cells (Tasiemski et al., 2004).

24
25 As in original study (Ovchinikova et al., 2004) it was reported that arenicins were isolated from
26 coelomocytes, hence we expected to detect their expression in these cells. We tested if the level of
27 arenicins mRNA expression is sensitive to different microbial stimuli by Real-Time PCR. To capture
28 the spectrum of potential PAMP/PRRs interactions which may be involved in the regulation of
29 arenicins expression we used three different heat-inactivated microbial strains (Gram-positive *L.*
30 *monocytogenes*, gram-negative *E.coli* and fungi *C.albicans*) representing the principal groups of
31 microbial pathogens. Notably, there was no difference in the levels of arenicin-1,2 expression upon
32 challenge with any of the above stimuli, including combined stimulation, compared to non-stimulated
33 controls. This suggests that arenicins expression is constitutive and could be explained by the nature of
34 lugworm biology, in that all tissues examined in this study are continually exposed to the surrounding
35 environment. As a consequence they have developed a certain degree of tolerance to various
36 microbiota. In such a scenario constitutive expression would be more beneficial in contrast to
37 inducible (as this was documented for some AMPs in epithelia of salivary glands and the ejaculatory
38 duct in insects (Ryu at al., 2004)).

39
40 We further addressed the functional role of arenicins as components of lugworm host defense. We
41 examined the possibility for arenicins to be released into the CF thus providing the first line of defense
42 acting as soluble factors. However, thorough analysis of CF by different biochemical methods did not
43 reveal any traces of arenicins. Furthermore, the analysis of CFP antimicrobial activity demonstrated
44 negative results, suggesting that in coelomic cavity, anti-microbial defense predominantly depends on
45 cell-mediated immunity. Next we proposed that the main function of arenicins would facilitate the
46 process of microbial killing within the phagolysosome. Cytoplasm of *A.marina* coelomocytes
47 demonstrated well developed vesicular apparatus strongly positive for arenicins indicating that
48 peptides are being stored in the vesicular compartment.

1 Generally there are several types of coelomocytes in polychaetes, such as granulocytes (syn.
2 amoebocytes), eleocytes and hemocytes. However, the last two are accepted to be absent in
3 Arenicolidae (rev. in Vetvicka & Sima, 2009). Granulocytes (amoebocytes) are further subdivided into
4 several subtypes, although relationships between those are not well understood. In *Arenicola* these
5 subtypes vary in their extent of development of granular apparatus and actin fiber networks (Chaga &
6 Persinina, 1998; Chaga et al., 1998). Chaga et al. separated the terms amoebocytes (few granules, well
7 developed actin fibers) and granulocytes (numerous granules, thin membrane associated actin
8 network), which corresponded to granulocytes of types I and II respectively in Dhainaut & Porchet-
9 Hennere nomenclature (Dhainaut & Porchet-Hennere, 1988). Both cell types originate from the same
10 source which are termed “juvenile cells” (Persinina & Chaga, 1994). Our data support this concept -
11 granulocytes of both types are clearly recognizable and even their morphology differs from described
12 previously (as they are actively spread on smears) as well as juvenile cells. Importantly, both
13 granulocyte types contain arenicins in their granules unlike juvenile cells, which have no developed
14 granular apparatus. This implies that arenicins expression in coelomocytes correlates with the
15 maturation of their granular apparatus.

16
17 Still there is unambiguous interpretation of an interrelation among their subpopulations. Moreover, an
18 evident demonstration of the common hematopoietic area is also lacking. Different derivatives of
19 coelomic peritoneum including extravasal tissue are most often proposed as candidates (rev. in
20 Gardiner, 1992; Vetvicka & Sima, 2009). Our data are in agreement with this suggestion. Extravasal
21 tissue (a derivative of coelothelium) was surmised for hematopoietic activity (rev. in Gardiner, 1992).
22 Its cells possess granular cytoplasm and participate in clearance of coelomic cavity via phagocytosis
23 (Braunbeck & Dales, 1984; Braunbeck & Dales, 1985). In the present study these cells along with
24 coelomocytes were detected as producers of arenicins. That finding may suggest potential lineal
25 relation between extravasal tissue and coelomocytes (as both populations were shown to possess
26 arenicin positive granules and be able to phagocytize). The fact that arenicins were absent in CF and
27 abundant in coelomocytes granules allows speculation about the role of arenicins in the killing of
28 phagocytized pathogens. Even though we have not demonstrated involvement of arenicins in
29 phagocytosis in extravasal tissue, we have shown this in the case of coelomocytes.

30
31 In granulocytes, immunopositivity was detected in granular apparatus, which was expected, as pre-pro-
32 arenicins include signal peptides. There are two main destinations for cytosolic granules - to be
33 exocytized or to fuse with the other vacuole. Immunohistochemistry of *in vitro* induced phagocytosis
34 of *A. marina* coelomocytes demonstrated the co-localization of arenicins with the internalized particles
35 suggesting involvement of arenicins in the process of pathogen inactivation (**Fig.7, H,I**). To our
36 knowledge this is the first report to demonstrate the role of AMP in phagocytosis within Annelids.

37
38 Coelomocytes were tested for induction of the arenicin gene upon infectious conditions and showed no
39 change in arenicin expression. This result of constitutive arenicins synthesis in coelomocytes is
40 concurrent with that of hedistin - an AMP constitutively expressed in coelomocytes of a benthic
41 polychaete *Nereis decersicolor* (Tasiemski et al., 2007). Unlike *A. marina* where coelomocytes are the
42 principal but not exclusive site of arenicins expression, in *N. diversicolor* the only one amoebocyte
43 population - NK cells or granulocytes type 3 - constitutively expresses hedistin. *Nereis* NK cells
44 possess minimal large granules in the cytoplasm where hedistin is likely stored as its transcript
45 encodes for a signal peptide. These cells are not phagocytic, but cytotoxic. They are able to actively
46 migrate to the site of infectious invasion and release their content of granules. There are no NK-like
47 cells in *Arenicola*. Despite both arenicin and hedistin being constitutively expressed in coelomocytes,
48 their functioning in the worm organism differs.

49

1 Active phagocytic cells and epithelia are the most common compartments of AMP expression in
2 metazoans (including mammals and insects). Interestingly, in the present study we revealed expression
3 of the same peptide in both compartments. Usually epithelia and phagocytes produce their own set of
4 AMPs (e.g. in humans - alpha-defensins 1-4 in neutrophils, LL-37 in monocytes, alpha-defensins 5-6
5 in intestine and different beta-defensins in skin and other borders) (De Smet & Contreras, 2005)).
6 Another remarkable fact is that there is no detectable antimicrobial activity in coelomic fluid, whereas
7 in other investigated annelids (*Thremyzon* or *Nereis*) AMPs are secreted by producing cells to
8 participate in host defense as humoral factors (Tasiemski et al., 2004; Tasiemski et al., 2007).
9 Signaling cascades, regulating AMP expression are thoroughly investigated in mammals and insects,
10 but still unexplored in annelids. It is an urgent problem to clarify whether or not variability in AMP
11 structure and functioning in annelids correlates with diversity or alternatively conservation of
12 regulatory mechanisms. In this study we did not address the molecular mechanism of regulation of
13 arenicin expression and this is a subject for future studies.

14

15 **Conclusion**

16 Our study demonstrated that arenicins are found in the tissues of the lugworm body (coelomocytes,
17 body wall, extravasal tissue and the gut) which provide the first line of defense against infections. This
18 supports the important role of arenicins as key components of both epithelial and systemic branches of
19 host defense. It was established that expression of arenicins is constitutive and does not depend on
20 stimulation of various infectious stimuli. In coelomocytes, arenicins function as killing agents inside
21 the phagolysosome, and may potentially carry this trait in extravasal tissue. In the gut and body wall
22 epithelia, arenicins are released from producing cells via secretion as they are found in the content of
23 the cuticle and inside the midgut gland cells. This study demonstrates that distribution and functioning
24 of arenicins in the lugworm share some features with AMPs of other annelids but overall their
25 characteristics are quite unique.

26

27 **Acknowledgement**

28 The project was supported by the research resource center 'Molecular and cell technologies' of St. Petersburg
29 State University, Russia. We thank Nikolai I.Kolokin and Tatiana V.Ovchinnikova for kindly providing
30 synthetic and recombinant arenicins, respectively. We also thank Marina A.Varfolomeeva (StPSU) and Dr.
31 Megan Jackson (QUB) for thorough review of the manuscript. Financial support for this project came from St-
32 Petersburg State University grant 1.0.140.2010 (PI Andrei I.Granovitch).

33

34 **References**

- 35 Andra, J., Hammer, M.U., Grotzinger, J., Jakovkin, I., Lindner, B., Vollmer, E., Fedders, H., Leippe, M.,
36 Gutschmann, T. (2009). Significance of the cyclic structure and of arginine residues for the antibacterial activity of
37 arenicin-1 and its interaction with phospholipid and lipopolysaccharide model membranes. *Biol. Chem.* 390(4),
38 337-349.
- 39 Andra, J., Jakovkin, I., Grotzinger, J., Hecht, O., Krasnosdembskaya, A., Goldmann, T., Gutschmann, T., Leippe,
40 M. (2008). Structure and mode of action of the antimicrobial peptide arenicin. *Biochem. J.* 410, 113-122.
- 41 Bely, A.E. (2006). Distribution of segment regeneration ability in the Annelida. *Integr. Comp. Biol.* 46(4), 508-
42 518.
- 43 Bilej, M., Procházková, P., Šilerová, M., & Josková, R. (2010). "Earthworm immunity", in *Invertebrate*
44 *Immunity*, ed. K. Söderhäll K. Springer US, 66-79.
- 45 Boman, H.G. (2003). Antibacterial peptides: basic facts and emerging concepts. *J. Internal Med.* 254(3), 197-
46 215.
- 47 Braunbeck, T., Dales, R.P. (1984). The role of the heart-body and of the extravasal tissue in disposal of foreign
48 cells in two polychaete annelids. *Tissue and Cell* 16(4), 557-563.
- 49 Braunbeck, T., Dales, R.P. (1985). The ultrastructure of the heart-body and extravasal tissue in the polychaete
50 annelids *Neoamphitrite figulus* and *Arenicola marina*. *J. Mar. Biol. Assoc. UK* 65(3), 653-662.
- 51 Bulet, P., Hetru, C., Dimarcq, J.L., Hoffmann, D. (1999). Antimicrobial peptides in insects; structure and
52 function. *Dev. Comp. Immunol.* 23(4), 329-344.

- 1 Bulet, P., Stocklin, R. (2005). Insect antimicrobial peptides: structures, properties and gene regulation. *Prot. Pept. Lett.* 12(1), 3-11.
- 2
- 3 Chaga, O.Y., Persinina, M.S., Dyakonova, M.Y., Podgornaya, O.I., Mikhailova, N.A. (1998). Studies on the
4 cytoskeleton coelomocytes in the polychaete *Arenicola marina*. I. Cytoskeleton rearrangement in the course of
5 coelomocyte differentiation. *Tsitologiya* 40, 209.
- 6 Cho, J., Lee, D.G. (2011). The antimicrobial peptide arenicin-1 promotes generation of reactive oxygen species
7 and induction of apoptosis. *Biochim. Biophys. Acta (BBA)-General Subjects* 1810(12), 1246-1251.
- 8 Cooper, E.L., Ru, B., Weng, N. (2004). "Earthworms: sources of antimicrobial and anticancer molecules", in
9 Complementary and Alternative Approaches to Biomedicine, ed. E.L. Cooper, and N. Yamaguchi. Springer US,
10 359-389.
- 11 Dhainaut, A., Porchet-Henneré, E. (1988). Haemocytes and coelomocytes. *The Ultrastructure of Polychaeta. Microfauna Marina* 4, 215-230.
- 12
- 13 Gardiner, S.L. (1992). "Polychaeta: general organization, integument, musculature, coelom, and vascular
14 system", in Microscopic anatomy of invertebrates, Annelida, ed. F.W. Harrison and S.L. Gardiner. Willey-Liss,
15 19-52.
- 16 Golding, D.W. (1992). "Polychaeta: nervous system", in Microscopic anatomy of invertebrates, Annelida, ed.
17 F.W. Harrison and S.L. Gardiner. Willey-Liss, 153-179.
- 18 Kaganovskaya, E.A. (1978). Electron-microscope study of ciliated and absorptive cells and their interrelations
19 in stomach epithelium of a polychaete *Arenicola marina*. *Tsitologiya* 20(8), 882-890.
- 20 Kauschke, E., Mohrig, W., Cooper, E.L. (2007). Coelomic fluid proteins as basic components of innate
21 immunity in earthworms. *Eur. J. Soil Biol.* 43, 110-115.
- 22 Kolodkin, N.I., Afonina, I.V., Krasnodembskaya, A.D., Smirnova, M.P., Shpen, V.M. (2006). Synthesis of
23 arenicin-1 - an antimicrobial peptide from coelomocytes of marine polychaete *Arenicola marina*. *J. Pept. Sci.*
24 12, 143-143.
- 25 Lee, J.U., Kang, D.I., Zhu, W.L., Shin, S.Y., Hahm, K.S., Kim, Y. (2007). Solution structures and biological
26 functions of the antimicrobial peptide, arenicin-1, and its linear derivative. *Pept. Sci.* 88(2), 208-216.
- 27 Lehrer, R.I., Barton, A., Daher, K.A., Harwig, S.S., Ganz, T., Selsted, M.E. (1989). Interaction of human
28 defensins with *Escherichia coli*. Mechanism of bactericidal activity. *J. Clinical Invest.* 84(2), 553.
- 29 Otero-González, A.J., Magalhães, B.S., Garcia-Villarino, M., López-Abarrategui, C., Sousa, D.A., Dias, S.C.,
30 Franco, O.L. (2010). Antimicrobial peptides from marine invertebrates as a new frontier for microbial infection
31 control. *FASEB J.* 24(5): 1320-1334.
- 32 Otvos, L. (2002). The short proline-rich antibacterial peptide family. *Cell. Mol. Life Sci. CMLS* 59(7): 1138-
33 1150.
- 34 Ovchinnikova, T.V., Aleshina, G.M., Balandin, S.V., Krasnodembskaya, A.D., Markelov, M.L., Frolova, E.I.,
35 Leonova, Y.F., Tagaev, A.A., Krasnodembsky, E.G., Kokryakov, V.N. (2004). Purification and primary
36 structure of two isoforms of arenicin, a novel antimicrobial peptide from marine polychaeta *Arenicola marina*.
37 *FEBS Lett.* 577(1), 209-214.
- 38 Ovchinnikova, T.V., Shenkarev, Z.O., Nadezhdin, K.D., Balandin, S.V., Zhmak, M.N., Kudelina, I.A., Finkina,
39 E.I., Kokryakov, V.N., Arseniev, A.S. (2007). Recombinant expression, synthesis, purification, and solution
40 structure of arenicin. *Biochem. Biophys. Res. Commun.* 360(1), 156-162.
- 41 Pan, W., Liu, X., Ge, F., Han, J., Zheng, T. (2004). Perinerin, a novel antimicrobial peptide purified from the
42 clamworm *Perinereis aibuhitensis* Grube and its partial characterization. *J. Biochem.* 135(3), 297-304.
- 43 Panyim, S., Chalkley, R. (1969). High resolution acrylamide gel electrophoresis of histones. *Archives Biochem.*
44 *Biophys.* 130(1), 337-46.
- 45 Park, C., Lee, D.G. (2009). Fungicidal effect of antimicrobial peptide arenicin-1. *Biochim. Biophys. Acta (BBA)-*
46 *Biomembranes* 1788(9), 1790-1796.
- 47 Pasupuleti, M., Schmidtchen, A., Malmsten, M. (2012). Antimicrobial peptides: key components of the innate
48 immune system. *Critical Rev. Biotechnol.* 32(2), 143-171.
- 49 Persinina, M.S., Chaga, O.Y. (1998). Renewal and differentiation of coelomic fluid cells in the polychaete,
50 *Arenicola marina*. IV. Ultrastructural study. *Tsitologiya* 40, 595-602.
- 51 Reddy, K.V.R., Yedery, R.D., Aranha, C. (2004). Antimicrobial peptides: premises and promises. *Int. J.*
52 *Antimicrobial Agents* 24(6), 536-547.
- 53 Ryu, J.H., Nam, K.B., Oh, C.T., Nam, H.J., Kim, S.H., Yoon, J.H., Seong, J.K., Yoo, M.A., Jang, I.H., Brey,
54 P.T., Lee, W.J. (2004). The homeobox gene Caudal regulates constitutive local expression of antimicrobial
55 peptide genes in *Drosophila* epithelia. *Mol. Cell. Biol.* 24(1), 172-185.

1 Saulnier-Michel, C. (1992). "Polychaeta: digestive system", in Microscopic anatomy of invertebrates. Annelida,
2 ed. F.W. Harrison and S.L. Gardiner. Willey-Liss, 53-69.
3 Schikorski, D., Cuvillier-Hot, V., Leippe, M., Boidin-Wichlacz, C., Slomianny, C., Macagno, E., Salz, M.,
4 Tasiemski, A. (2008). Microbial challenge promotes the regenerative process of the injured central nervous
5 system of the medicinal leech by inducing the synthesis of antimicrobial peptides in neurons and microglia. *J.*
6 *Immunol.* 181(2), 1083-1095.
7 De Smet, K., Contreras, R. (2005), Human antimicrobial peptides: defensins, cathelicidins and histatins.
8 *Biotechnol. Lett.* 27(18), 1337-1347.
9 Tasiemski, A., Vandenbulcke, F., Mitta, G., Lemoine, J., Lefebvre, C., Sautière, P.E., Salz, M. (2004).
10 Molecular characterization of two novel antibacterial peptides inducible upon bacterial challenge in an annelid,
11 the leech *Theromyzon tessulatatum*. *J. Biol. Chem.* 279(30), 30973-30982.
12 Tasiemski, A., Schikorski, D., Le Marrec-Croq, F., Pontoire-Van Camp, C., Boidin-Wichlacz, C., Sautière, P.E.
13 (2007). Hedistin: A novel antimicrobial peptide containing bromotryptophan constitutively expressed in the NK
14 cells-like of the marine annelid, *Nereis diversicolor*. *Dev. Comp. Immunol.* 31(8): 749-762.
15 Tasiemski, A. (2008). Antimicrobial peptides in annelids. *J. Invertebr. Sur i* 5, 75-82.
16 Tasiemski, A., Salz, M. (2010). "Leech immunity: from brain to peripheral responses", in Invertebrate
17 Immunity, ed. K. Söderhäll. Springer US, 80-104.
18 Vetvicka, V., Sima, P. (2009). Origins and functions of annelid immune cells: the concise survey. *Invert. Surv.*
19 *J.* 6, 138-143.
20 Wang, X., Wang, X., Zhang, Y., Qu, X., Yang, S. (2003). An antimicrobial peptide of the earthworm *Pheretima*
21 *tschiliensis*: cDNA cloning, expression and immunolocalization. *Biotechnol. Lett.* 25: 1317-1323.
22 Wells, G.P. (1950). The anatomy of the body wall and appendages in *Arenicola marina* L., *Arenicola claparedii*
23 Levinsen and *Arenicola ecaudata* Johnston. *J. Mar. Biol. Assoc. UK* 29(01), 1-44.
24 Yount, N.Y., Bayer, A.S., Xiong, Y.Q., Yeaman, M.R. (2006). Advances in antimicrobial peptide
25 immunobiology. *Pept. Sci.* 84(5), 435-458.
26
27
28

29 **Figure captions.**

30
31 **Fig.1. mRNA expression levels of arenicin-1 (A) relatively to actin (B) measured by PCR.** Expression
32 levels in coelomocytes (1), body wall (2), pharynx (3), intestine (4).
33

34 **Fig.2. mRNA expression levels of arenicin-1 measured by RT-PCR relatively to actin in coelomocytes**
35 **after immunization.** Microbial mixture 24h post infection (1), *C.albicans* 24h post infection (2), *E.coli* 48h (3)
36 and 24h post infection (4), *L.monocytogenes* 24h post infection (5), PBS 24h post infection (6), intact animals
37 (7). Data are expressed as mean±SD.
38

39 **Fig.3. Polyclonal antibody specificity testing by Western blot.** Acidic urea PAAG (A) and pAB staining
40 visualized by DAB (B). Left line - crude extract, right line - synthetic arenicin-1.
41

42 **Fig.4. Arenicin-positive immunoreactivity of lugworm body wall.** A-B: general view of dorsal body part; C-
43 D: body wall; E-F - chaetae complex; B,D,G,H - negative controls; c - cuticle, cc - coelomic cavity, ch -
44 chaetae, ep - body wall epithelium, in - intestine, m - metanephridia.
45

46 **Fig.5. Arenicin-positive immunoreactivity of lugworm nervous system.** A-B: sector of dorsal body part; A:
47 negative control; ep - body wall epithelium, lm - longitudinal musculature, vcn - ventral nervous chain.
48

49 **Fig.6. Arenicin-positive immunoreactivity of lugworm intestine.** A,C: general view of intestine; B,D-F:
50 sparse immunopositive cells; C,F - negative controls; cc - coelomic cavity, gc - presumable gland cells, in -
51 intestine.
52

53 **Fig.7. Arenicin-positive immunoreactivity of lugworm vascular system.** A,B: general view of dorsal and
54 ventral vessels respectively; E,F: extravasal tissue; C,D: negative controls; cc - coelomic cavity, dv - dorsal
55 vessel, in - intestine, et - extravasal tissue, vv - ventral vessel.

1
2 **Fig.8. Arenicin-positive immunoreactivity of lugworm coelomocytes.** A-F: coelomic fluid smears; A-B:
3 granulocytes I and juvenile cells and C-D: corresponding negative controls; E,F: granulocytes II; G-I: in vitro
4 phagocytosis preparations where H: negative control; jc - juvenile cells, pp - phagocytized particles.

Figure 1.JPEG




Figure 2.JPEG





Figure 3.JPEG

A



B



Figure 4.JPEG




Figure 5.TIF





Figure 6.JPEG









Figure 7.JPEG




Figure 8.JPEG

