Linking Genes to Microbial Biogeochemical Cycling Lessons from Arsenic

Zhu, Y-G., Xue, X-M., Kappler, A., Rosen, B. P., & Meharg, A. A. (2017). Linking Genes to Microbial Biogeochemical Cycling Lessons from Arsenic. *Environmental science & technology*, 51(13), 7326-7339. <https://doi.org/10.1021/acs.est.7b00689>

Published in:
Environmental science & technology

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
Copyright © 2017 American Chemical Society. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

1 Linking genes to microbial biogeochemical cycling: lessons from arsenic

2 Yong-Guan Zhu, ^{†, ‡, #, *} Xi-Mei Xue, ^{†, #} Andreas Kappler, [‡] Barry P Rosen, [§] Andrew A
3 Meharg [⊥]

4 [†] Key Laboratory of Urban Environment and Health, Institute of Urban Environment,
5 Chinese Academy of Sciences, Xiamen, China

6 [‡] State Key Laboratory of Urban and Regional Ecology, Research Center for
7 Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, China

8 [‡] Geomicrobiology, Center for Applied Geosciences, University of Tübingen,
9 Tübingen 72076, Germany

10 [§] Department of Cellular Biology and Pharmacology, Herbert Wertheim College of
11 Medicine, Florida International University, Miami, FL 33199, U.S.A.

12 [⊥] Institute for Global Food Security, Queen's University Belfast, Belfast BT9 5HN,
13 United Kingdom

14
15 [#]These authors contributed equally to the article.

16 ^{*}Address correspondence to Yong-Guan Zhu

17 Address: Institute of Urban Environment, Chinese Academy of Sciences, 1799 Jimei
18 Road, Xiamen 361021, China, Phone number: +86(0)592 6190997, Fax number:
19 +86(0)592 6190977, Email address: ygzhu@rcees.ac.cn

20

Abstract

The biotransformation of arsenic is highly relevant to the arsenic biogeochemical cycle. Identification of the molecular details of microbial pathways of arsenic biotransformation coupled with analyses of microbial communities by meta-omics can provide insights into detailed aspects of the complexities of this biocycle. Arsenic transformations couple to other biogeochemical cycles, and to the fate of both nutrients and other toxic environmental contaminants. Microbial redox metabolism of iron, carbon, sulfur and nitrogen affects the redox and bioavailability of arsenic species. In this critical review we illustrate the biogeochemical processes and genes involved in arsenic biotransformations. We discuss how current and future metagenomic-, metatranscriptomic-, metaproteomic-, and metabolomic-based methods will help to decipher individual microbial arsenic transformation processes, and their connections to other biogeochemical cycle. These insights will allow future use of microbial metabolic capabilities for new biotechnological solutions to environmental problems. To understand the complex nature of inorganic and organic arsenic species and the fate of environmental arsenic will require integrating systematic approaches with biogeochemical modeling. Finally, from the lessons learned from these studies of arsenic biogeochemistry, we will be able to predict how the environment changes arsenic, and, in response, how arsenic biotransformations change the environment.

1. Introduction

Biogeochemical cycles are interconnected through redox reactions and other biotransformations.¹ Biogeochemical processes such as the cycling of a particular element are likely to be mediated by multiple microbes and are often linked to other biogeochemical processes. For example, redox changes of arsenic are mediated by diverse arsenate (As(V))-reducing and arsenite (As(III))-oxidizing microbes. Arsenic biogeochemical cycling is often coupled to the cycling of iron (Fe),² carbon (C)³ and nitrogen (N),⁴ and to the dynamics of elements/ions associated with the arsenic redox cycle, such as sulfur (S).⁵ Coupling of biogeochemical cycles has recently received attention. The study of coupled biogeochemical cycles offers a scientific basis for major current environmental problems.⁶

The proteins catalyzing physiological processes in living organisms are influenced by geological, physical and chemical forces and therefore continuously evolve and redistribute chemical species involved in biogeochemical cycles. Genetic analysis is the key to understand the arsenic biogeochemical cycle. Once the genes associated with the reactions and the environmental signals that affect gene expression are understood, we will be able to predict how microbial metabolism influences arsenic biogeochemical cycling. In this review, we focus on the known genes involved in arsenic biotransformations and the effect of other elements on arsenic biogeochemistry. We highlight the effects of other elements on arsenic metabolism and the current state of meta-omics research in microbial arsenic metabolism. Finally, we discuss how integration of meta-omics information into biogeochemical models can allow us to predict the possible biotransformation of other elements.

2. Arsenic metabolism: from genes to biogeochemical processes

Organisms have evolved various strategies to transform arsenic for detoxification or energy metabolism.⁷ An overview about the enzymatic pathways for arsenic biotransformations is presented in Figure 1, and related microbial genes are summarized in Table 1.

2.1 The arsenic redox cycle

The earliest microorganisms evolved in an anoxic environment, where the predominant arsenic species was most probably reduced As(III), with little oxidized As(V). The physiological activities of the earliest microorganisms were, therefore, largely driven by anaerobic metabolic processes,⁸ and we propose that As(III) bioavailability was a driving force for the evolution or acquisition of genes encoding anaerobic respiratory pathways.³³ For example, the photosynthetic purple sulfur bacterium *Ectothiorhodospira* PHS-1 carries out anoxygenic photosynthesis using As(III) as an electron donor in the light^{34, 35, 36} and uses As(V) as an electron acceptor in the dark.³⁷ The chemolithoautotrophic As(III)-oxidizer *Alkalilimnicola ehrlichii* MLHE-1 utilizes As(III) as an electron donor and nitrate as an electron acceptor in energy-generating respiratory chains.³⁸ Microorganisms with similar metabolic versatility probably evolved quite early. These microbes could cope with extreme growth conditions, such as high concentrations of As(III) or low oxygen, similar to those that existed in an primordial anoxic biosphere.

Since As(III) was probably the primary bioavailable arsenic species on the early anoxic Earth, it was the inevitable choice for organisms to utilize As(III) as an electron donor to produce energy.³⁹ As(III) oxidation is catalyzed by the enzyme As(III) oxidase, which is composed of two different subunits, a large subunit (α) having molybdopterin and a [3Fe-4S] cluster (AioA) and a smaller subunit (β)

incorporating a Rieske-type [2Fe-2S] cluster (AioB).⁴⁰ The cluster of *aioA* and *aioB* genes (*aio* operon) usually consists of *aioS* and *aioR* genes, encoding for a two-component signal transduction pair, AioS (sensor histidine kinase)/AioR (transcriptional regulator), which regulates expression of *aio* genes via recognizing As(III).⁹ The operon sometimes has an *aioX* gene that encodes an As(III)-binding protein involved in As(III)-based signaling and regulation of As(III) oxidation,¹² a *cytC* gene encoding a cytochrome c that is required for efficient As(III) oxidation in *Ochrobactrum tritici* SCII24,⁴¹ or a *moeA* gene encoding MoeA protein that synthesizes the molybdenum cofactor of AioAB oxidase.⁹ Recently, a new type of As(III) oxidase, ArxA that exhibited both As(V) reductase and As(III) oxidase activities *in vitro*,¹⁰ was identified in *A. ehrlichii* MLHE-1.⁴² In *Ectothiorhodospira* sp. PHS-1 these genes code for As(III) oxidation coupled to photosynthesis.^{35, 11} In addition to *arxA*, the MLHE-1 and PHS-1 *arx* operons, contain four other genes *arxB2*, *arxB*, *arxC*, and *arxD*, that encode two proteins with [4Fe-4S] centers, a membrane anchoring and quinol oxidoreductase subunit and a TorD-like molybdoenzyme chaperone respectively.¹¹ An adjacent and divergent gene cluster, *arxXSR*, encodes putative regulatory proteins, a periplasmic substrate-binding protein specific for phosphate (ArxX), a two-component histidine kinase sensor (ArxS), and a response regulator (ArxR).¹¹ ArxA has higher sequence similarity to the ArrA subunit than to AioA, and fills the phylogenetic gap between As(III) oxidases and As(V) reductases.^{42, 11}

Note that As(III) oxidation by anaerobes would have produced As(V) in the absence of an oxygen-containing atmosphere, which opened a niche for As(V)-respiring microbes prior to the Great Oxidation Event (GOE).³³ Dissimilatory As(V)-respiring prokaryotes (DARPs) evolved pathways to take advantage of the

appearance of As(V) as a terminal electron acceptor. This new energy-generating respiratory chain utilized the respiratory As(V) reductase, ArrAB, that reduce the less toxic As(V) to the more toxic and potentially more mobile As(III).^{40, 43, 44} ArrAB is a heterodimer consisting of a large catalytic subunit (ArrA) and a small subunit (ArrB).^{15, 16} The *arr* operon also includes *arrC*, *arrD*, *arrS*, and *arrR*. Their gene products are ArrC, a membrane-bound As(V) reductase subunit, ArrD, a As(V) reductase chaperon, ArrS, a sensor histidine kinase and ArrR, a transcriptional regulator respectively.¹⁷ A phylogenetic analysis was conducted to search for molybdenum-bis (pyranopterin guanine dinucleotide)-containing catalytic subunits of representative enzymes. This complex iron sulfur molybdoenzyme family includes Arr, Aio, Arx, polysulfide reductase, and nitrate reductase. The results indicate that Arr clusters most likely evolved from polysulfide reductases.¹⁷

After the GOE, As(III) in oceans mostly oxidized to As(V), a new environmental toxin. As(V) enters the cells of most organisms adventitiously via phosphate uptake systems.⁴⁵ As a consequence, early life had to evolve novel strategies for coping with new (potentially toxic) arsenic species. As described in more detail below, nearly every extant microbe has ArsB or Acr3 efflux permeases for As(III) detoxification, so it is reasonable to assume that organisms that arose before the GOE already had an As(III) efflux system. When As(V) became the predominant soluble species, all cells had to do was to reduce As(V) to As(III), the substrate of ArsB or Acr3, and they would become resistant to As(V). A number of independently-evolved As(V) reductases arose in a variety of organisms using a small molecular mass protein As(V) reductases (one of several types of ArsC or Acr2 reductases). The ArsC system conferred by the *ars* operon is the most well studied mechanism of arsenic detoxification and resistance (for details see the previous review).¹⁸ Most recently, a

glutathione S-transferase B (GstB) was found to mediate an alternate pathway which conferred As(V) resistance to *E. coli* mutant cells lacking *arsC* by directly reducing As(V) to As(III).²¹ These enzymes all use small molecule proteins such as glutaredoxin (Grx) or reduced glutathione (GSH) coupled to thioredoxin (Trx) as electron donor. The Acr2 reductases evolved from proteins that incorporated the phosphate binding loop of phosphorprotein tyrosine phosphatases related to the cell phosphatase CDC25.⁴⁶ These phosphatase can be converted into As(V) reductases by just a few mutations,⁴⁷ indicating a facile evolutionary path.

2.2 The arsenic methylation cycle

In addition to oxidation and reduction of inorganic arsenic species, pathways for biotransformation of arsenic, including methylation and demethylation, organoarsenical degradation, evolved in early organisms. Interest in arsenic biomethylation began in 1800's with the observation that inorganic arsenic compounds used as wallpaper pigments were converted into Gosio gas (trimethylarsine) by fungi. More recent reports of methylated arsenical showed that arsenic methylation was widespread in the environment and detected in bacteria,⁴⁸ cyanobacteria,⁴⁹ algae,²³ protozoa.⁵⁰ Arsenic methylation is a common stratagem to detoxify arsenic. The highly toxic trivalent products are rapidly oxidized nonenzymatically in air to the less toxic pentavalent methylated arsenic species. Also, gaseous end-products such as trimethylarsine will emit to air, thus removing the product. Methylation is catalyzed by the enzyme As(III) S-adenosylmethionine (SAM) methyltransferase (EC 2.1.1.137), designated as AS3MT in animals and as ArsM in microorganisms. Expression of typical prokaryotic and archaeal *arsM* genes are regulated by the As(III)-responsive transcriptional repressor ArsR,²⁰ consistent with arsenic methylation being a detoxification pathway in the microbes. Expression of

arsM in some cyanobacteria appears to be constitutive,⁵¹ indicating that alternate detoxification pathways are used by microorganisms in which the expression of *arsM* is not regulated.⁵²

The degradation of environmental organoarsenicals has been documented for some time,^{53, 54} while few molecular mechanisms for these reactions have been demonstrated. Recently, a two-step pathway of MSMA reduction and demethylation was elucidated.⁵⁵ Although no reductases of pentavalent organoarsenicals have been identified as yet, the enzyme, ArsI, which catalyzes demethylation of trivalent organoarsenicals, was identified and characterized from the environmental isolate bacterium *Bacillus* sp. MD1²⁴ and from the cyanobacterium *Nostoc* sp. 7120.⁵⁶ ArsI, a non-heme iron-dependent dioxygenase with C-As lyase activity, cleaves the C-As bond in MAs(III), trivalent roxarsone, and other trivalent aromatic arsenicals. Putative ArsI orthologs were found only in bacterial species, suggesting that alternate pathways of organoarsenical demethylation might exist in other organisms.²⁴

2.3 The organoarsenical cycle

The arsenic concentration in seawater is around 1 to 2 µg per liter, mainly inorganic arsenic that is usually transformed into complex organoarsenical compounds by marine organisms.⁵⁷ Arsenosugars, first identified in 1981,⁵⁸ are commonly detected water-soluble arsenic species present in marine algae; arsenobetaine is the most abundant arsenic species in the majority of marine animals.⁵⁷ More complex organoarsenicals have been identified with the improvement of analytical techniques in recent years. Since the structure of an arsenosugar phospholipid (AsPL) from a brown alga *Undaria pinnatifida* was first identified,⁵⁹ AsPL has been found in algae⁶⁰ and cyanobacteria.⁶¹ Arsenic-containing fatty acids (AsFA) that were first identified in cod liver oil⁶² have now been found in algae⁶³ and various fish species.

^{64, 65} AsHC that were first reported in capelin ⁶⁶ have been detected in fish ^{64, 65} and algae. ⁶⁰ A new class of arsenolipids, trimethylarsenio fatty alcohols (TMA₃FOH), was reported in Capelin oil. ⁶⁷ Two new groups of arsenolipids, arsenic-containing phosphatidylcholines (AsPC) and arsenic-containing phosphatidylethanolamine (AsPE) from herring caviar, were characterized. ⁶⁸ In total, more than 20 arsenosugars and 70 arsenolipids have been identified in marine organisms that live in low phosphate and high salt environments. These organoarsenicals are not toxic for the marine plants and animals, but their function in these marine organisms is not known. Phytoplankton can substitute sulfur- and nitrogen-containing membrane lipids for membrane phospholipids, ⁶⁹ arsenolipids might be used in membranes in place of phospholipids due to the more similarity of As(V), than sulfate and nitrate, to inorganic phosphate. Thus As(V) could be used as a phosphate-sparing substitute in phosphate-limiting environments. A recent study on *Ectocarpus siliculosus* that was found to produce more arsenosugar phospholipids under low-phosphate than under normal phosphate conditions ⁷⁰ supports this hypothesis.

These organoarsenicals may be toxic to organisms that cannot biosynthesize them. *In vitro* toxicological characterization of three arsenic-containing hydrocarbons showed that cytotoxicity of the arsenic-containing hydrocarbons was comparable to that of As(III) for cultured human bladder and liver cells, ⁷¹ and arsenolipids were metabolized by humans to dimethylated arsenical species (DMAs) and other small molecular arsenic compounds, then excreted in the urine. ^{72, 73}

Although several potential pathways have been proposed for the synthesis of complex organoarsenicals, ⁸ few genes involved in these biotransformation have been identified. Even less is known about the degradation of these organoarsenicals, and more studies are needed on this front.

2.4 The arsenic thiolation cycle

Thioarsenates ($\text{H}_3\text{AsS}_n\text{O}_{4-n}$) are the dominant arsenic species in alkaline, arsenic-rich, sulfidic environments. These play a significant role in the arsenic biochemical cycle in sulfidic geothermal environments.^{74, 75, 76} Thioarsenates are transformed to As(V) and/or As(III) via exposing to oxidizing agents or increased pH,⁷⁷ by biological conversion by sulfur-oxidizing bacteria,^{75, 78} or by abiotic decomposition (desulfidation) with subsequent biological oxidation.^{77, 79} Thioarsenates contain reduced S^{2-} and oxidized As(V). They can serve both as electron donors and electron acceptors. For example, monothioarsenate can be used as an electron donor by *Thermocrinis ruber* OC 14/7/2,⁸⁰ and for anoxygenic photosynthesis by phototrophic purple sulfur bacteria growing in an alkaline environment.⁸¹ Recently, the haloalkaliphilic bacterium MLMS-1 can grow chemolithotrophically by oxidizing the S^{2-} of monothioarsenate to S^0 or SO_4^{2-} , while concurrently reducing As(V) to As(III).⁷⁸ In summary, various microbes have evolved to utilize thioarsenates that are widespread in sulfidic environments.

2.5 Arsenic efflux pathways

The best way to deal with toxic arsenicals in cells is acquisition of an efficient efflux system. As(III) efflux systems have been intensively studied in both microbes and higher organisms.^{82, 83} As(III) efflux in most bacteria is mediated by ArsB in an energy-dependent process, driven in *Staphylococcus aureus* by the membrane potential⁸⁴ and in *E. coli* by ATP hydrolysis that ArsA binds to ArsB to an ATP-driven arsenic-specific pump.²⁶ In the legume symbiont *S. meliloti*, an aquaglyceroporin (AqpS), instead of ArsB, has been identified to extrude As(III) from cells.²⁷ Acr3 has been shown to be an As(III)-efflux transporter in both bacteria and yeast, and provides a pathway for As(III) extrusion from cells.¹⁹ In fact, genes for

Acr3 are more wide-spread in bacteria and archaea than are *arsB* genes. The cytosolic As(III)/glutathione complex sequestered into vacuoles by an ABC-type transporter, Ycf1p (yeast cadmium factor protein), is the second pathway for As(III) detoxification in yeast *S. cerevisiae*.²⁸

Moreover, a novel mechanism for As(V) resistance was identified in a variety of microbes including *Pseudomonas aeruginosa*.²⁹ In these bacteria there are two genes that always go together, one encoding a typical glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and the second one, called *arsJ*, that encodes an organoarsenical efflux permease (ArsJ). GAPDH uses As(V) and glyceraldehyde 3-phosphate to form the extremely unstable organoarsenical 1-arsenol-3-phosphoglycerate, which is extruded from cells by ArsJ and immediately breaks down into As(V) and 3-phosphoglycerate. The net reaction is effectively As(V) extrusion, and the coupled reaction confers As(V) resistance to these microbes, the only known efflux pathway for As(V). Meanwhile the bacterial permease, ArsP, from *Campylobacter jejuni*, was demonstrated to be an efflux system specific for trivalent organoarsenicals.³⁰ It is more selective for the ancient organoarsenical MAs(III) than for the recently anthropogenically-developed antimicrobial aromatic arsenical growth promoters such as trivalent roxarsone. More and more arsenic reductases and trivalent arsenic-specific transporters identified show that arsenic reduction and efflux play an important role in arsenic biogeochemical cycling.

3. Coupling of arsenic biogeochemical cycling to other elements

Any biogeochemical process, such as the cycling of a particular element, is likely to be mediated by more than one organism, and often linked to other fundamental biogeochemical processes. Arsenic biotransformations are often coupled to the

cycling of C, Fe, S and N.^{85, 2, 86} The effect of turnover of these elements on microbes involved in arsenic biogeochemical cycling is summarized in Figure 2 which contains data based on previous studies^{8, 87, 88, 89} and as described below.

3.1 The effects of iron on arsenic biogeochemical cycling

The chemical speciation of arsenic and arsenic mobility in natural environments are strongly dependent on redox potential and pH. Under oxic conditions, As(V) is the predominated arsenic species, present mainly as H_2AsO_4^- at acid pH or HAsO_4^{2-} at alkaline pH. In anoxic environments, arsenic occurs primarily as reduced As(III) ($\text{As}(\text{OH})_3$ at neutral pH or H_2AsO_3^- at alkaline pH), and more mobile than As(V).⁹⁰ Moreover, pH will impact arsenic interactions with Fe, the sorption of As(V) onto amorphous iron oxide and goethite is higher than that of As(III) below pH 5-6, As(V) and As(III) sorption onto iron oxide are both relatively high at neutral pH, As(III) is more easily adsorbed to iron oxide than As(V) above pH 7-8.⁹¹

Transformation of arsenic-bearing Fe mineral phases strongly affects the bioavailability of arsenic within soils and aquifers due to direct and indirect interactions between the arsenic and Fe cycles including mineral formation, transformation, dissolution and redox reactions.^{89, 92, 93, 94} Previous studies from our laboratory showed that Fe plaque consisting of Fe(III) (oxyhydr)oxides, which was induced artificially through adding ferrous iron in solution to paddy soils, has high affinity for As(V), and reduced arsenic uptake by rice.^{95, 96, 97}

Fe(III)-reducing bacteria modulate arsenic mobility in the rhizosphere.^{98, 99, 100} Dissimilatory reduction of Fe(III) (oxyhydr)oxides to Fe(II) by dissimilatory iron-reducing bacteria (DIRB) can result either in the release of As(V) from poorly crystalline or more crystalline ferric minerals as well as from sorption sites within sediments,¹⁰¹ or in the binding of arsenic to the formed Fe(II) minerals.^{102, 103, 104, 105,}

¹⁰⁶ DIRB are commonly present in rice paddy soil, and mediate dissimilatory reduction of Fe(III) on the rice root-plaque. ¹⁰⁷ A study on the role of DIRB in arsenic release under a range of biogeochemical regimes indicated that Fe(III) reduction was stimulated by addition of acetate as a potential electron donor that resulted in a marked increase in the number of DIRB, reduction of As(V) to As(III), and arsenic release after Fe(III) reduction. ¹⁰⁰ If DARPs were used as Fe(III)-reducers, *Shewanella* sp. ANA-3 ¹⁰⁸ or *Sulfurospirillum barnesii* ¹⁰⁹ could release both As(III) and Fe(II) from ferrihydrite containing As(V) by reducing solid-phase As(V) and Fe(III). Eventually, most of the ferrihydrite matrix was liberated as Fe(II) and As(III) if sufficient organic electron donor was present. ¹¹⁰ There is more aluminum in the crust than iron. However, *S. barnesii* does not reductively dissolve the As(V)-aluminum hydroxide precipitate, ¹¹¹ so we did not include a detailed description of the effect of aluminum on arsenic biogeochemical cycling.

In addition to Fe(III) reduction that has the potential to mobilize or immobilize arsenic depending on geochemical conditions that lead to the formation of either dissolved Fe²⁺ or Fe(II) minerals, the formation of Fe(III) minerals under Fe(II)-oxidizing conditions has the potential to significantly immobilize arsenic and thus to lower its bioavailability. ^{112, 113} In particular for nitrate-reducing Fe(II)-oxidizing bacteria it has been shown that they form poorly soluble Fe(III) minerals and efficiently co-precipitate arsenic (Figure 2). ^{112, 113, 114} Besides nitrate-dependent Fe(II) oxidation, also microaerophilic Fe(II) oxidation has the potential to influence arsenic mobility and could even be used in biotechnical applications for arsenic removal, e.g. in drinking water filters ¹¹⁵ although it has been shown that in commercial drinking water filters the formation of iron biominerals by Fe(II)-oxidizing bacteria lowers arsenic removal from the water. ¹¹⁶ Moreover, a

thermo-acidophilic iron-oxidizing archaeon *Acidianus brierleyi*, has been used to immobilize As(III) in the copper refinery process by producing thermodynamically stable crystalline scorodite ($\text{FeAsO}_4 \cdot 2\text{H}_2\text{O}$).¹¹⁷

3.2 The effects of sulfur on arsenic biogeochemical cycling

More than 200 different arsenic-containing minerals have been found in the Earth's crust, and 20% are sulfides and sulfosalts.¹¹⁸ The behavior of arsenic is affected by abiotic or biological redox of sulfur, which can either release or immobilize arsenic.^{114, 119} Abiotic sulfide, a strong reductant under sulfate-reduced conditions, plays a critical role in arsenic solubility by forming pyrite (FeS_2), realgar (AsS), orpiment (As_2S_3),¹²⁰ arsenopyrite (FeAsS),¹²¹ or by reducing As(V).¹²²

In addition to abiotic factors, sulfate-reducing bacteria (SRB) also cause dramatic changes in Fe, sulfide and arsenic species by generating hydrogen sulfide¹²³ or elemental sulfur from sulfate,⁵ or by localized reprecipitation of released arsenic as As_2S_3 ¹²⁴ or FeAsS , which have low solubility.¹²⁵ Sulfur-oxidizing bacteria have the potential to immobilize arsenic by using free or arsenic-bound sulfur as an electron donor to directly or indirectly transform As(III) and thioarsenates to As(V),⁷⁵ or reduce As(V).^{126, 127} In brief, transformations involving sulfur have the potential to significantly impact the fate of environmental arsenic.

3.3 The effects of N on arsenic biogeochemical cycling

Nitrate is an ecologically significant oxidant that can support microbial oxidation of As(III) in the absence of oxygen. The first evidence that microbes are capable of linking anoxic As(III) oxidation to denitrification came from a field study in anoxic lake water columns.⁴ The absence or presence of nitrate affected the redox state of arsenic. As(III) was present where nitrate was depleted, but As(V) was the dominant species during anoxic nitrate-rich periods. Subsequently, a nitrate-dependent As(III)

oxidation bacterium *A. ehrlichii* strain MLHE-1 was found to be capable of coupling As(III) oxidation with partial denitrification of nitrate to nitrite.^{38, 128} Two other anoxic chemolithoautotrophic strains, *Azoarcus* strain DAO1 and *Sinorhizobium* strain DAO10, were able to oxidize As(III) and fix CO₂ via complete denitrification of nitrate to dinitrogen gas.¹²⁹ Biological nitrate-dependent As(III) oxidation is widespread in the environment, and potentially plays a significant role in arsenic biogeochemical cycling.¹³⁰

As mentioned above, nitrate also influences the bioavailability and mobility of arsenic indirectly by linking nitrate reduction to Fe(II) oxidation. Previous studies showed tight coupling between N, Fe, and arsenic in paddy soil.⁸⁵ Addition of nitrate reduced arsenic uptake by rice probably because i) the nitrate inhibited/reduced Fe(III) reduction leading to less arsenic mobilization and ii) nitrate-dependent Fe(II)-oxidizing bacteria stimulated Fe(II) oxidation, which led to arsenic co-precipitation with Fe(III) minerals in soil. Nitrate strongly affects arsenic cycling under anoxic conditions in nitrate-rich Upper Mystic Lake by microbially catalyzing As(III) to more particle-reactive As(V) and oxidizing Fe(II) to arsenic-sorbing particulate ferric oxides.⁴ Microbial nitrate-dependent Fe(II) oxidation in groundwater,¹³¹ freshwater sediments¹³² and marine sediments¹³³ has the potential to contribute to the reduction of arsenic mobility in various ecosystems.

3.4 The effects of organic matter on arsenic biogeochemical cycling

Natural organic matter (NOM) is widely distributed in the environment. NOM consists of heterogeneous mixtures of organic compounds with various structural and functional properties,¹³⁴ that influence the fate of arsenic by competitive adsorption and redox reactions,^{135, 136, 137} and by formation of arsenic-bearing organic-metal-complexes and mineral colloids.^{138, 139} NOM molecules have

combinations of carboxylic, amino, sulfhydryl, hydroxyl, esteric, phenolic, nitroso, and other functional groups.¹⁴⁰ They are considered to be an efficient geochemical trap for arsenic both under oxic and reducing conditions. Whereas As(V) is immobilized by binding to protonated amino groups of NOM³ or a nucleophile substitution reaction between As(V) and phenolic OH groups of NOM,¹⁴¹ As(III) is associated with NOM via phenolic OH or carboxyl groups of NOM by H-bonding, hydrophobic As(III)-NOM interactions,¹⁴¹ or via ternary As(III)-Fe(III)-NOM complexes that form bridges between Fe(III), arsenic oxyanions and the functional entities of NOM.¹⁴² In contrast, under sulfate-reducing conditions, the formation of a trigonal-pyramidal complex between As(III) and sulfhydryl groups of NOM could potentially be a sequestration mechanism for arsenic.¹⁴³

Organic matter could also change the fate of arsenic by influencing microbial communities or activities.^{144, 145, 146} NOM is thought to drive the reductive dissolution of Fe(III) (oxyhydr)oxides, thereby causing redox transformations of As(V) to As(III) and facilitate arsenic release.¹⁴⁷ In addition to the reductive dissolution of arsenic-bearing Fe(III) (oxyhydr)oxide phases, studies have revealed that addition of organic matter to paddy soil significantly increased arsenic methylation and volatilization.¹⁴⁸ Application of rice straw into soil increased arsenic accumulation in rice by influencing microbial processes involved in arsenic redox.¹⁴⁹

Humic acids, which are forms of NOM, can reduce As(V).^{150, 151, 152} Small organic molecules, including lactate, pyruvate, fumarate, malate, succinate, butyrate, citrate, acetate, glycerol, ethanol, and formate, can be used as an electron donor by DARPs for As(V) reduction.^{153, 154, 155, 156} *Desulfosporosinus* sp. Y5 has been found to couple even complex aromatic substrates such as phenol, syringic acid, benzoate, ferulic acid, and toluene, to As(V) reduction.¹⁵⁷ In fact, some DARPs can also respire

sulfate, selenite, Fe(III), thiosulfate, nitrate, or nitrite.^{153, 154} This diversity of electron donors and acceptors may be of benefit to microbes grown in environments where Fe, N, S, or C coexist with arsenic.

Apart from iron oxides and organic materials, silicon significantly decreased As(III) concentration, but increased the concentration of DMAs in both the vegetative and reproductive tissues of rice.¹⁵⁸ In brief, in addition to biological transformations, other inorganic elements, compounds/minerals and physico-chemical properties interact with arsenic in the environment. Coupling of arsenic with other elements makes it necessary to consider genes involved in cycling of other elements, such as Fe, N, S, and C when studying the arsenic biogeochemical cycling.^{2, 87, 88}

Known genes involved in arsenic biotransformation are readily determined in pure cultures. However, in the field, these genes cannot easily be quantified in bacterial communities with a limited number of primers, even though As(V)-reducing and As(III)-oxidizing bacteria are widely distributed in the environment.^{159, 160, 161} In addition, the speciation, fate and biogeochemical transformation processes of arsenic in the environment are much more complex than under laboratory conditions.^{162, 163} It is therefore necessary to apply more systematic and more comprehensive approaches such as metagenomics, metatranscriptomics, metaproteomics, and metabolomics to understand interactions between environmental microbes. These approaches will take into account local geochemical surroundings and neighboring organisms by analyzing DNA, RNA, proteins, and metabolites extracted directly from environmental samples.

4 Understanding biogeochemical arsenic cycling by application of “omics” methods and integrated modeling

4.1 Metagenomics

Metagenomics provides an inestimable window into the microbial world by characterizing microorganisms involved in difficult-to-elucidate but important biochemical pathways, as the overwhelming majority of microbes in the environment cannot be cultured in the laboratory. From a metagenomic library two arsenic-resistant bacteria and one novel As(V) resistance gene (*arsN*), which encodes a protein similar to acetyltransferases, were identified.²² Xiao et al applied metagenomic techniques to analyze genes associated with arsenic transformation. They analyzed five low-arsenic paddy soils using high-throughput sequencing and constructed a protein database of arsenic metabolizing genes. Their analysis shows that arsenic metabolism genes are ubiquitous and abundant, even in low-arsenic environments.¹⁶⁴ Metagenomics was also used to unravel the correlations between the microbes and arsenic transformation in different niches.^{165, 166, 167, 168, 169} Although metagenomics provides taxonomic and functional profiles of a microbial community, it does not demonstrate the levels of expression of the genes nor their physiological activity.¹⁷⁰ Therefore, metatranscriptomics is needed to delineate the active functional genes and communities.

4.2 Metatranscriptomics

Metatranscriptomics offers novel insights into the expression of functional genes and microbial activities of complex microbial communities at a specific moment or under specific environmental conditions by sequencing the total mRNAs extracted from natural microbial communities. Functional metatranscriptomics has potential for isolation and characterization of novel genes involved in heavy metal transformation.¹⁷¹ Metatranscriptomics enhances our understanding of microbial responses to their environment¹⁷² and the functional profile of a microbial community.¹⁷³ Recently, a transcriptomics meta-analysis was used to unravel the effect of As(III) on the

symbiotic interaction between the model legume *Medicago truncatula* and its symbiont *Ensifer* (syn. *Sinorhizobium*) *medicae* MA11.¹⁷⁴ This study identified the adaptive responses of the bacterial symbiont to arsenic exposure. This metatranscriptomic approach will be useful to study how microbes regulate their genes to adapt to the changes in environmental conditions, particularly arsenic concentrations.¹⁷⁵ The correlation between mRNA and protein inventories in environmental microbial communities is low when environmental conditions change rapidly. mRNA inventories respond rapidly and sensitively to the shift, while changes in protein inventories are slow.¹⁷⁶ As a consequence, microbial metaproteomics and metabolomics had to be used to identify the repertoire of proteins and small molecular metabolites that microbes use to adapt to complex and dynamic environments. In this way, the metabolic activities of a microbial community in specific environments at the moment of sampling could be elucidated.¹⁷⁰

4.3 Metaproteomics and metabolomics

Metaproteomics and metabolomics are the comprehensive methods by which proteins produced by microbes and metabolites released by microorganisms into the environment are characterized and quantified using a combination of liquid or gas chromatography-based separations and mass spectrometry-based identification techniques.^{177, 178} The study of microbial proteome and metabolome can provide valuable information about the function of microbial communities and the interactions of the microbial communities with the environment.^{179, 180} When the diversity of arsenic-adapted prokaryotic communities in mildly arsenic-contaminated sediments was analyzed using meta-proteomic and 16S rRNA amplification, the results indicated that the data sizes provided by metaproteomics are less than those provided by metagenomics and metatranscriptomics.¹⁸¹ So far, metabolomics has been used

mainly to analyze low molecular mass metabolites within a tissue, biofluid, a cell or cell compartment of an organism including plants, animals, bacteria, and humans exposed to an environmental stressor.¹⁸⁰ High-throughput metabolomics has been applied to the analysis of metabolites in the liver of mice when co-exposed to high fat and cholesterol diets and arsenic-contaminated drinking water.¹⁸² However, application of metaproteomic and metabolomic techniques to real environment is limited due to difficulties with amplification and the low quantities of extractable proteins and metabolites because of the interferences with many components present in complex environmental systems, such as soil.^{180, 183}

4.4 Integrating meta-omic techniques

From the above, meta-omics are key techniques in elucidating the dynamic and complex interactions between microbial communities and the environment. Integrating multiple meta-omic datasets will provide a complete exhibition from genes to biogeochemical cycles. Metatranscriptomic and metagenomic techniques were combined to detect large numbers of novel genes from complex marine microbial communities.¹⁷³ Datasets of meta-genomics were integrated with metabolomics to reveal how a microbial community interacts with the environment and responds to environmental parameters.¹⁸⁴ Taken together, meta-omic technologies offer an unprecedented opportunity to elucidate the functions of microbes that are not readily cultured under normal laboratory conditions in biogeochemical cycles.

Recently, "Arsenomics" was termed as an approach to focusing on the analysis of alterations in transcriptome, proteome and metabolome occurring in microbes exposed to arsenic.^{185, 186} With the application of meta-omics to environmental

science, we believe that Arsenomics will evolve to include the analysis of metagenomic, metatranscriptomic, metaproteomic, and metabolomic changes in microbial communities from the real environment where they are exposed to arsenic.

4.5 Integrating environmental meta-omics into biogeochemical models for arsenic

Microbes are ubiquitous in diverse environmental niches including soil, oceans, sediments, freshwater environments, and inside the body of animals or plants, and exert great influence on biogeochemical cycles in these habitats.¹⁸⁷ For example, microbes involved in arsenic transformation are ubiquitously distributed in paddy soils, resulting in various concentrations and percentage of inorganic and methylated arsenic species among different rice plants.^{188, 189, 190, 191} Since higher plants appear not to methylate arsenic,¹⁹² microbial methylation is probably the primary source of methylated arsenic in plants, which occurs in soil prior to plant uptake.

In situ measurements or prediction of arsenic transformations contribute to analysis of the dynamics of arsenic and prediction of arsenic bioavailability. *In situ* measurements of As(V) reduction in Mono Lake, California (dissolved inorganic arsenic ~ 200 μ M), made with radiotracers (⁷³As and ³⁵S) of mass balance considerations, revealed that As(V) reduction occurred in the monimolimnion waters with the highest rates between 18 and 19 meters (rate, ~ 5.9 μ M/day) and sulfate reduction rates increased with depth at depths of 21 meters and below with the highest rates at 28 meters (rate, ~ 2.3 μ M/day).¹⁹³ The radioisotope method was further employed to examine the As(V) and sulfate reduction processes in sediments of two arsenic-rich soda lakes, Mono Lake (moderately salt, ~ 90 g/L) and Searles Lake (saturated salt, ~ 340 g/L).¹⁹⁴ The rate constant [k] of As(V) reduction was 0.103 to 0.04 h⁻¹ in Mono Lake and 0.012 to 0.002 h⁻¹ in Searles Lake, and sulfate reduction

was only detected in Mono Lake ($k=7.6\times 10^4$ to 3.2×10^{-6} h⁻¹). Denatured gradient gel electrophoresis (DGGE) of 16S rRNA genes amplified from Mono Lake and Searles Lake sediment DNA indicated that microbial communities from two sediments were distinct from each other. More *arrA* gene signal was found in Mono Lake than in Searles Lake, where higher As(V) reduction activity was observed, due to PCR biases, the presence of novel *arrA* genes, or higher expression of low-abundance *arrA* genes.

¹⁹⁴

Characterization of genes involved in arsenic biotransformation and application of multiple meta-omics in environment analysis will lead to insights into the microbial world, as limited information on functional genes cannot predict the status of arsenic in the environment. The information of field/*in situ* characterization of functional genes/functional microbial communities and biogeochemical fluxes should be integrated into biogeochemical models to complete the transition from lab to the field, from biochemistry to biogeochemistry, and from genes-genomics to microbial communities. This integration will help to predict the dynamics of arsenic in the environment, and to improve the effectiveness of mitigation technologies. Many strategies have been developed to model low complexity environments. For example, recent work integrating environmental genomics and qPCR in biogeochemical models explored the nexus between microbial community and geochemistry in the Arabian Sea oxygen minimum zone.¹⁹⁵ Metabolic processes coupling C, N, and S transformations in the Saanich Inlet oxygen-starved zone were integrated using a biogeochemical model that integrates multi-omic information and geochemistry.¹⁹⁶ These studies indicated that such integrated modeling approaches can provide a novel insight into microbial metabolic networks in water bodies, and pave a road for prediction of elemental cycling.

540

541 **5 Perspectives**

542 In summary, as arsenic biotransformations are catalyzed by a suite of enzymes from
543 diverse environmental organisms, and these are coupled to biogeochemical cycles of
544 other elements such as Fe, S, and N. As more and more genomes are being sequenced,
545 more genes directly or indirectly involved in arsenic metabolism will be discovered
546 and characterized. With the development of new technologies, we anticipate rapid
547 advances in analytical chemistry, microbiology and genomics that will improve our
548 understanding of how microbial metabolic pathways contribute to and govern
549 complex environmental processes. In the future, integrating meta-omic datasets into
550 biogeochemical models will improve the ability of prediction and offer a deeper
551 insight into arsenic biogeochemical processes in diverse niches.

552 In future studies, it will be necessary to analyze the interaction between
553 organisms and the environment using additional meta-omics approaches at different
554 spatio-temporal scales. Geochemical analyses in combination with genetic analyses
555 will provide insights into the specific roles of the complex biochemical pathways in
556 the global arsenic biogeochemical cycle. More importantly, integrating modeling
557 approaches linking arsenic biogeochemical cycle with meta-omics data should be
558 developed to predict the dynamic of arsenic species in water, sediments and soils and
559 provide our society and authorities with the tools necessary for limiting arsenic
560 pollution, improving remediation and providing safe drinking water and food.

561

562 **Acknowledgments**

Our research was supported by National Natural Science Foundation of China (41571130063), Program of Chinese Academy of Sciences (XDB15020402), National Natural Science Foundation of China (21507125), and International Science & technology Cooperation Program of China (2011DFB91710). BPR was supported by NIH grants GM55425 and ES023779.

Reference

- (1) Bashkin V N. *Modern biogeochemistry*; Springer Science & Business Media, 2002.
- (2) Oremland, R. S.; Stolz, J. F. The ecology of arsenic. *Science* (80). **2003**, 300, 939–944 DOI: 10.1126/science.1081903.
- (3) Thanabalasingam, P.; Pickering, W. F. Arsenic sorption by humic acids. *Environ. Pollut. (Series B)* **1986**, 12 (3), 233–246 DOI: 10.1016/0143-148X(86)90012-1.
- (4) Senn, D. B.; Hemond, H. F. Nitrate controls on iron and arsenic in an urban lake. *Science* (80). **2002**, 296, 2373–2376 DOI: 10.1126/science.1072402.
- (5) Saalfeld, S.; Bostick, B. Changes in iron, sulfur, and arsenic speciation associated with bacterial sulfate reduction in ferrihydrite-rich systems. *Environ. Sci. Technol.* **2009**, 43, 8787–8793 DOI: 10.1021/es901651k.
- (6) Finzi, A. C.; Cole, J. J.; Doney, S. C.; Holland, E. A.; Jackson, R. B. Research frontiers in the analysis of coupled biogeochemical cycles. *Front Ecol Env.* **2011**, 9 (1), 74–80 DOI: 10.1890/100137.
- (7) Mukhopadhyay, R.; Rosen, B. P.; Phung, L. T.; Silver, S. Microbial arsenic: from geocycles to genes and enzymes. *FEMS Microbiol Rev* **2002**, 26, 311–325 DOI: 10.1111/j.1574-6976.2002.tb00617.x.
- (8) Zhu, Y.-G.; Yoshinaga, M.; Zhao, F.-J.; Rosen, B. P. Earth Abides Arsenic Biotransformations. *Annu. Rev. Earth Planet. Sci.* **2014**, 42 (1), 443–467 DOI: 10.1146/annurev-earth-060313-054942.
- (9) Sardiwal, S.; Santini, J. M.; Osborne, T. H.; Djordjevic, S. Characterization of a two-component signal transduction system that controls arsenite oxidation in the chemolithoautotroph NT-26. *FEMS Microbiol. Lett.* **2010**, 313 (1), 20–28 DOI: 10.1111/j.1574-6968.2010.02121.x.
- (10) Richey, C.; Chovanec, P.; Hoeft, S. E.; Oremland, R. S.; Basu, P.; Stolz, J. F. Respiratory arsenate reductase as a bidirectional enzyme. *Biochem. Biophys. Res. Commun.* **2009**, 382 (2), 298–302 DOI: 10.1016/j.bbrc.2009.03.045.
- (11) Zargar, K.; Conrad, A.; Bernick, D. L.; Lowe, T. M.; Stolz, V.; Hoeft, S.; Oremland, R. S.; Stolz, J.; Saltikov, C. W. ArxA, a new clade of arsenite oxidase within the DMSO reductase family of molybdenum oxidoreductases. *Environ. Microbiol.* **2012**, 14 (7), 1635–1645 DOI:

- 10.1111/j.1462-2920.2012.02722.x.
- (12) Liu, G.; Liu, M.; Kim, E. H.; Maaty, W. S.; Bothner, B.; Lei, B.; Rensing, C.; Wang, G.; McDermott, T. R. A periplasmic arsenite-binding protein involved in regulating arsenite oxidation. *Environ. Microbiol.* **2012**, *14* (7), 1624–1634 DOI: 10.1111/j.1462-2920.2011.02672.x.
- (13) Chen, J.; Bhattacharjee, H.; Rosen, B. P. ArsH is an organoarsenical oxidase that confers resistance to trivalent forms of the herbicide monosodium methylarsenate and the poultry growth promoter roxarsone. *Mol. Microbiol.* **2015**, *96* (5), 1042–1052 DOI: 10.1111/mmi.12988.
- (14) Xue, X. M.; Yan, Y.; Xu, H. J.; Wang, N.; Zhang, X.; Ye, J. ArsH from *Synechocystis* sp. PCC 6803 reduces chromate and ferric iron. *FEMS Microbiol. Lett.* **2014**, *356* (1), 105–112 DOI: 10.1111/1574-6968.12481.
- (15) Afkar, E.; Lisak, J.; Saltikov, C.; Basu, P.; Oremland, R. S.; Stolz, J. F. The respiratory arsenate reductase from *Bacillus selenitireducens* strain MLS10. *FEMS Microbiol. Lett.* **2003**, *226* (1), 107–112 DOI: 10.1016/S0378-1097(03)00609-8.
- (16) Krafft, T.; Macy, J. M. Purification and characterization of the respiratory arsenate reductase of *Chrysiogenes arsenatis*. *Eur. J. Biochem.* **1998**, *255* (3), 647–653 DOI: 10.1046/j.1432-1327.1998.2550647.x.
- (17) Van Lis, R.; Nitschke, W.; Duval, S.; Schoepp-Cothenet, B. Arsenics as bioenergetic substrates. *Biochim. Biophys. Acta - Bioenerg.* **2013**, *1827* (2), 176–188 DOI: 10.1016/j.bbabi.2012.08.007.
- (18) Mukhopadhyay, R.; Rosen, B. P. Arsenate reductases in prokaryotes and eukaryotes. *Environ. Health Perspect.* **2002**, *110*, 745–748 DOI: 10.1289/ehp.02110s5745.
- (19) Wysocki, R.; Goffeau, A. Isolation of three contiguous genes, ACR1, ACR2 and ACR3, involved in resistance to arsenic compounds in the yeast *Saccharomyces cerevisiae*. *Yeast* **1997**, *13*, 819–828 DOI: 10.1002/(SICI)1097-0061(199707)13.
- (20) David J. Thomas; Barry P. Rosen. Arsenic Methyltransferase. In *Encyclopedia of Metalloproteins*; Uversky, V. N., Kretsinger, R. H., Permyakov, E. E. a., Eds.; Science+Business Media: New York, 2013; pp 138–143 DOI: 10.1007/978-1-4614-1533-6.
- (21) Chrysostomou, C.; Quandt, E. M.; Marshall, N. M.; Stone, E.; Georgiou, G. An alternate pathway of arsenate resistance in *E. coli* mediated by the glutathione S-transferase GstB. *ACS Chem. Biol.* **2015**, *10* (3), 875–882 DOI: 10.1021/cb500755j.
- (22) Chauhan, N. S.; Ranjan, R.; Purohit, H. J.; Kalia, V. C.; Sharma, R. Identification of genes conferring arsenic resistance to *Escherichia coli* from an effluent treatment plant sludge metagenomic library. *FEMS Microbiol Ecol* **2009**, *67*, 130–139 DOI: 10.1111/j.1574-6941.2008.00613.x.
- (23) Qin, J.; Rosen, B. P.; Zhang, Y.; Wang, G.; Franke, S.; Rensing, C. Arsenic detoxification and evolution of trimethylarsine gas by a microbial arsenite S-adenosylmethionine methyltransferase. *Proc. Natl. Acad. Sci.* **2006**, *103* (7), 2075–2080 DOI: 10.1073/pnas.0506836103.

- 639 (24) Yoshinaga, M.; Rosen, B. P. A C · As lyase for degradation of environmental organoarsenical
640 herbicides and animal husbandry growth promoters. *Proc. Natl Acad. Sci.* **2014**, *111* (21),
641 7701–7706 DOI: 10.1073/pnas.1403057111.
- 642 (25) Louis S. Tisa; Rose, B. P. Molecular characterization of an anion pump-The ArsB protein is the
643 membrane anchor for the ArsA protein. *J. Biol. Chem.* **1990**, *265*, 190–194.
- 644 (26) Lin, Y.; Walmsley, A. R.; Rosen, B. P. An arsenic metallochaperone for an arsenic
645 detoxification pump. *Proc. Natl Acad. Sci.* **2006**, *103* (42), 15617–15622 DOI:
646 10.1073/pnas.0603974103.
- 647 (27) Yang, H.; Cheng, J.; Finan, T. M.; Rosen, B. P. Novel pathway for arsenic detoxification in the
648 legume symbiont *Sinorhizobium meliloti*. *J. Bacteriol.* **2005**, *187* (20), 6991–6997 DOI:
649 10.1128/JB.187.20.6991.
- 650 (28) Ghosh, M.; Shen, J.; Rosen, B. P. Pathways of As (III) detoxification in *Saccharomyces*
651 *cerevisiae*. *Proc. Natl Acad. Sci.* **1999**, *96*, 5001–5006 DOI: 10.1073/pnas.96.9.5001.
- 652 (29) Chen, J.; Yoshinaga, M.; Garbinski, L. D.; Rosen, B. P. Synergistic interaction of
653 glyceraldehydes-3-phosphate dehydrogenase and ArsJ, a novel organoarsenical efflux
654 permease, confers arsenate resistance. *Mol. Microbiol.* **2016**, *100* (6), 945–953 DOI:
655 10.1111/mmi.13371.
- 656 (30) Chen, J.; Madegowda, M.; Bhattacharjee, H.; Rosen, B. P. ArsP: A methylarsenite efflux
657 permease. *Mol. Microbiol.* **2015**, *98* (4), 625–635 DOI: 10.1111/mmi.13145.
- 658 (31) Grondin, K.; Haimeur, A.; Mukhopadhyay, R.; Rosen, B. P.; Ouellette, M. Co-amplification of
659 the gamma-glutamylcysteine synthetase gene *gsh1* and of the ABC transporter gene *pgpA* in
660 arsenite-resistant *Leishmania tarentolae*. *EMBO J.* **1997**, *16* (11), 3057–3065 DOI:
661 10.1093/emboj/16.11.3057.
- 662 (32) Wang, L.; Chen, S.; Xiao, X.; Huang, X.; You, D.; Zhou, X.; Deng, Z. arsRBOCT arsenic
663 resistance system encoded by linear plasmid pHZ227 in *Streptomyces* sp. strain FR-008. *Appl*
664 *Env. Microbiol* **2006**, *72* (5), 3738–3742 DOI: 10.1128/AEM.72.5.3738.
- 665 (33) Oremland, R. S.; Saltikov, C. W.; Wolfe-Simon, F.; Stolz, J. F. Arsenic in the Evolution of
666 Earth and Extraterrestrial Ecosystems. *Geomicrobiol. J.* **2009**, *26* (V), 522–536 DOI:
667 10.1080/01490450903102525.
- 668 (34) McCann, S. H.; Boren, A.; Hernandez-maldonado, J.; Stoneburner, B.; Saltikov, C. W.; Stolz, J.
669 F.; Oremland, R. S. Arsenite as an Electron Donor for Anoxygenic Photosynthesis: Description
670 of Three Strains of *Ectothiorhodospira* from Mono Lake, California and Big Soda Lake ,
671 Nevada. *Life* **2016**, *7* (1), 1–14 DOI: 10.3390/life7010001.
- 672 (35) Hernandez-maldonado, J.; Sanchez-sedillo, B.; Stoneburner, B.; Boren, A.; Miller, L.; Mccann,
673 S.; Rosen, M.; Oremland, R. S.; Saltikov, C. W. The genetic basis of anoxygenic
674 photosynthetic arsenite oxidation. *Env. Microbiol* **2017**, *19*, 130–141 DOI:
675 10.1111/1462-2920.13509.
- 676 (36) Kulp, T. R.; Hoeft, S. E.; Asao, M.; Madigan, M. T.; Hollibaugh, J. T.; Fisher, J. C.; Stolz, J. F.;
677 Culbertson, C. W.; Miller, L. G.; Oremland, R. S. Arsenic(III) fuels anoxygenic photosynthesis
678 in hot spring biofilms from Mono Lake, California. *Science* **2008**, *321* (5891), 967–970 DOI:

10.1126/science.1160799.

(37) Hoeft, S. E.; Kulp, T. R.; Han, S.; Lanoil, B.; Oremland, R. S. Coupled arsenotrophy in a hot spring photosynthetic biofilm at Mono Lake, California. *Appl. Environ. Microbiol.* **2010**, *76* (14), 4633–4639 DOI: 10.1128/AEM.00545-10.

(38) Oremland, R. S.; Hoeft, S. E.; Santini, J. M.; Bano, N.; Hollibaugh, R. a; Hollibaugh, J. T. Anaerobic Oxidation of Arsenite in Mono Lake Water and by a Facultative, Arsenite-Oxidizing Chemoautotroph, Strain MLHE-1. *Appl. Environ. Microbiol.* **2002**, *68* (10), 4795–4802 DOI: 10.1128/AEM.68.10.4795.

(39) Kulp, T. R. Arsenic and primordial life. *Nat. Geosci.* **2014**, *7* (11), 785–786 DOI: 10.1038/ngeo2275.

(40) Stolz, J. F.; Basu, P.; Oremland, R. S. Microbial Arsenic Metabolism: New Twists on an Old Poison. *Microbe* **2010**, *5* (2), 53–59 DOI: 10.1128/microbe.5.53.1.

(41) Branco, R.; Francisco, R.; Chung, A. P.; Morais, P. V. Identification of an aox system that requires cytochrome c in the highly arsenic-resistant bacterium *Ochrobactrum tritici* SCH24. *Appl. Environ. Microbiol.* **2009**, *75* (15), 5141–5147 DOI: 10.1128/AEM.02798-08.

(42) Zargar, K.; Hoeft, S.; Oremland, R.; Saltikov, C. W. Identification of a novel arsenite oxidase gene, *arxA*, in the haloalkaliphilic, arsenite-oxidizing bacterium *Alkalilimnicola ehrlichii* strain MLHE-1. *J. Bacteriol.* **2010**, *192* (14), 3755–3762 DOI: 10.1128/JB.00244-10.

(43) Hery, M.; Gault, A. G.; Rowland, H. A. L.; Lear, G.; Polya, D. A.; Lloyd, J. R. Molecular and cultivation-dependent analysis of metal-reducing bacteria implicated in arsenic mobilisation in south-east asian aquifers. *Appl. Geochemistry* **2008**, *23* (11), 3215–3223 DOI: 10.1016/j.apgeochem.2008.07.003.

(44) Pederick, R. L.; Gault, A. G.; Charnock, J. M.; Polya, D. A.; Lloyd, J. R. Probing the biogeochemistry of arsenic: Response of two contrasting aquifer sediments from Cambodia to stimulation by arsenate and ferric iron. *J. Environ. Sci. Heal. Part A* **2007**, *42*, 1763–1774 DOI: 10.1080/10934520701564269.

(45) Yan, Y.; Ding, K.; Yu, X.; Ye, J.; Xue, X. Ability of Periplasmic Phosphate Binding Proteins from *Synechocystis* sp. PCC 6803 to Discriminate Phosphate Against Arsenate. *Water. Air. Soil Pollut.* **2017**, *228*, 148 DOI: 10.1007/s11270-017-3334-4.

(46) Bhattacharjee, H.; Sheng, J.; Ajees, A. A.; Mukhopadhyay, R.; Rosen, B. P. Adventitious arsenate reductase activity of the catalytic domain of the human Cdc25B and Cdc25C phosphatases. *Biochemistry* **2010**, *49* (4), 802–809 DOI: 10.1021/bi9019127.

(47) Zhou, R.; Royyuru, A.; Athma, P.; Suits, F.; Silverman, B. D. Magnitude and spatial orientation of the hydrophobic moments of multi-domain proteins. *Int. J. Bioinform. Res. Appl.* **2006**, *2* (2), 161–176 DOI: 10.1016/j.molbiopara.2006.03.009.

(48) Wang, P.; Sun, G.; Jia, Y.; Meharg, A. A.; Zhu, Y. A review on completing arsenic biogeochemical cycle: Microbial volatilization of arsines in environment. *J. Environ. Sci.* **2014**, *26*, 371–381 DOI: 10.1016/S1001-0742(13)60432-5.

(49) Yin, X.-X.; Chen, J.; Qin, J.; Sun, G.-X.; Rosen, B. P.; Zhu, Y.-G. Biotransformation and volatilization of arsenic by three photosynthetic cyanobacteria. *Plant Physiol.* **2011**, *156* (3),

- 1631–1638 DOI: 10.1104/pp.111.178947.
- (50) Yin, X. X.; Zhang, Y. Y.; Yang, J.; Zhu, Y. G. Rapid biotransformation of arsenic by a model protozoan *Tetrahymena thermophila*. *Environ. Pollut.* **2011**, *159* (4), 837–840 DOI: 10.1016/j.envpol.2010.12.033.
- (51) Ye, J.; Rensing, C.; Rosen, B. P.; Zhu, Y. G. Arsenic biomethylation by photosynthetic organisms. *Trends Plant Sci.* **2012**, *17* (3), 155–162 DOI: 10.1016/j.tplants.2011.12.003.
- (52) Xue, X.-M.; Ye, J.; Raber, G.; Francesconi, K. A.; Li, G.; Gao, H.; Yan, Y.; Rensing, C.; Zhu, Y.-G. Arsenic Methyltransferase is Involved in Arsenosugar Biosynthesis by Providing DMA. *Environ. Sci. Technol.* **2017**, *51* (3), 1224–1230 DOI: 10.1021/acs.est.6b04952.
- (53) Stolz, J. F.; Perera, E.; Kilonzo, B.; Kail, B.; Crable, B.; Fisher, E.; Ranganathan, M.; Wormer, L.; Basu, P. Biotransformation of 3-nitro-4-hydroxybenzene arsonic acid (Roxarsone) and release of inorganic arsenic by clostridium species. *Environ. Sci. Technol.* **2007**, *41* (3), 818–823 DOI: 10.1021/es061802i.
- (54) Lehr, C. R.; Polishchuk, E.; Radoja, U.; Cullen, W. R. Demethylation of methylarsenic species by *Mycobacterium neoaurum*. *Appl. Organomet. Chem.* **2003**, *17* (11), 831–834 DOI: 10.1002/aoc.544.
- (55) Yoshinaga, M.; Cai, Y.; Rosen, B. P. Demethylation of methylarsonic acid by a microbial community. *Environ. Microbiol.* **2011**, *13* (5), 1205–1215 DOI: 10.1111/j.1462-2920.2010.02420.x.
- (56) Yan, Y.; Ye, J.; Xue, X.-M.; Zhu, Y.-G. Arsenic Demethylation by a C · As Lyase in Cyanobacterium *Nostoc* sp. PCC 7120. *Environ. Sci. Technol.* **2015**, *49*, 14350–14358 DOI: 10.1021/acs.est.5b03357.
- (57) Edmonds, J. S.; Francesconi, K. A. Organoarsenic compounds in the marine environment. In *Organometallic compounds in the environment*; Craig, P. J., Ed.; Wiley: Chichester, 2013; pp 195–222 DOI: 10.1017/CBO9781107415324.004
- (58) Edmonds, J. S.; Francesconi, K. A. Arseno-sugars from brown kelp (*Ecklonia radiata*) as intermediates in cycling of arsenic in a marine ecosystem. *Nature* **1981**, *289*, 602–604 DOI: 10.1038/289602a0
- (59) Morita, M.; Shibata, Y. Isolation and identification of arseno-lipid from a brown alga, *Undaria pinnatifida* (Wakame). *Chemosphere* **1988**, *17* (6), 1147–1152 DOI: 10.1016/0045-6535(88)90180-4.
- (60) Garcia-Salgado, S.; Raber, G.; Raml, R.; Magnes, C.; Francesconi, K. A. Arsenosugar phospholipids and arsenic hydrocarbons in two species of brown macroalgae. *Environ. Chem.* **2012**, *9*, 63–66 DOI: 10.1071/EN11164.
- (61) Xue, X.-M.; Raber, G.; Foster, S.; Chen, S.-C.; Francesconi, K. A.; Zhu, Y.-G. Biosynthesis of arsenolipids by the cyanobacterium *Synechocystis* sp. PCC 6803. *Environ. Chem.* **2014**, *11* (5), 506–513 DOI: 10.1071/EN14069.
- (62) Rumpler, A.; Edmonds, J. S.; Katsu, M.; Jensen, K. B.; Goessler, W.; Raber, G.; Gunnlaugsdottir, H.; Francesconi, K. A. Arsenic-containing long-chain fatty acids in cod-liver oil: A result of biosynthetic infidelity? *Angew. Chemie - Int. Ed.* **2008**, *47* (14), 2665–2667

- DOI: 10.1002/anie.200705405.
- (63) Raab, A.; Newcombe, C.; Pitton, D.; Ebel, R.; Feldmann, J. Comprehensive analysis of lipophilic arsenic species in a brown alga (*Saccharina latissima*). *Anal. Chem.* **2013**, 85 (5), 2817–2824 DOI: 10.1021/ac303340t.
- (64) Taleshi, M. S.; Raber, G.; Edmonds, J. S.; Jensen, K. B.; Francesconi, K. A. Arsenolipids in oil from blue whiting *Micromesistius poutassou*—evidence for arsenic-containing esters. *Sci. Rep.* **2014**, 4, 7492 DOI: 10.1038/srep07492.
- (65) Taleshi, M. S.; Edmonds, J. S.; Goessler, W.; Ruiz-Chancho, M. J.; Raber, G.; Jensen, K. B.; Francesconi, K. A. Arsenic-containing lipids are natural constituents of sashimi tuna. *Environ. Sci. Technol.* **2010**, 44 (4), 1478–1483 DOI: 10.1021/es9030358.
- (66) Taleshi, M. S.; Jensen, K. B.; Raber, G.; Edmonds, J. S.; Gunnlaugsdottir, H.; Francesconi, K. A. Arsenic-containing hydrocarbons: natural compounds in oil from the fish capelin, *Mallotus villosus*. *Chem. Commun.* **2008**, 39 (39), 4706–4707 DOI: 10.1039/b808049f.
- (67) Amayo, K. O.; Raab, A.; Krupp, E. M.; Gunnlaugsdottir, H.; Feldmann, J. Novel identification of arsenolipids using chemical derivatizations in conjunction with RP-HPLC-ICPMS/ESMS. *Anal. Chem.* **2013**, 85 (19), 9321–9327 DOI: 10.1021/ac4020935.
- (68) Viczek, S. A.; Jensen, K. B.; Francesconi, K. A. Arsenic-containing Phosphatidylcholines: a New Group of Arsenolipids Discovered in Herring Caviar. *Angew. Chemie Int. Ed.* **2016**, 55, 5259–5262 DOI: 10.1002/anie.201512031.
- (69) Van Mooy, B. a S.; Fredricks, H. F.; Pedler, B. E.; Dyhrman, S. T.; Karl, D. M.; Koblížek, M.; Lomas, M. W.; Mincer, T. J.; Moore, L. R.; Moutin, T.; et al. Phytoplankton in the ocean use non-phosphorus lipids in response to phosphorus scarcity. *Nature* **2009**, 458, 69–72 DOI: 10.1038/nature07659.
- (70) Petursdottir, A. H.; Fletcher, K.; Gunnlaugsdottir, H.; Krupp, E.; Kupper, F. C.; Feldmann, J. Environmental effects on arsenosugars and arsenolipids in *Ectocarpus* (Phaeophyta). *Environ. Chem.* **2016**, 13 (1), 21–33 DOI: 10.1071/EN14229.
- (71) Meyer, S.; Matissek, M.; Müller, S. M.; Taleshi, M. S.; Ebert, F.; Francesconi, K. A.; Schwerdtle, T. *In vitro* toxicological characterisation of three arsenic-containing hydrocarbons. *Metallomics* **2014**, 6, 1023–1033 DOI: 10.1039/c4mt00061g.
- (72) Schmeisser, E.; Goessler, W.; Francesconi, K. A. Human metabolism of arsenolipids present in cod liver. *Anal. Bioanal. Chem.* **2006**, 385 (2), 367–376 DOI: 10.1007/s00216-006-0401-x.
- (73) Taylor, V.; Goodale, B.; Raab, A.; Schwerdtle, T.; Reimer, K.; Conklin, S.; Karagas, M. R.; Francesconi, K. A. Human exposure to organic arsenic species from seafood. *Sci. Total Environ.* **2017**, 580, 266–282 DOI: 10.1016/j.scitotenv.2016.12.113.
- (74) Planer-friedrich, B.; Suess, E.; Scheinost, A. C.; Wallschla, D. Arsenic speciation in sulfidic waters: Reconciling contradictory spectroscopic and chromatographic evidence. *Anal. Chem.* **2010**, 82 (24), 10228–10235 DOI: 10.1021/ac1024717.
- (75) Fisher, J. C.; Wallschlager, D.; Planer-Friendrich B., H. J. T. A new role for sulfur in arsenic cycling. *Environ. Sci. Technol.* **2008**, 42 (1), 81–85 DOI: 10.1021/es0713936.
- (76) Planer-Friedrich, B.; London, J.; Mccleskey, R. B.; Nordstrom, D. K.; Wallschlager, D.

- Thioarsenates in geothermal waters of yellowstone National Park: Determination, preservation, and geochemical importance. *Environ. Sci. Technol.* **2007**, *41* (15), 5245–5251 DOI: 10.1021/es070273v.
- (77) Taylor, P.; Planer-friedrich, B.; Fisher, J. C.; Hollibaugh, J. T.; Süß, E.; Planer-friedrich, B.; Fisher, J. C.; Hollibaugh, J. T.; Elke, S.; Wallschl, D. Oxidative Transformation of Trithioarsenate Along Alkaline Geothermal Drainages — Abiotic versus Microbially Mediated Processes. *Geomicrobiol J* **2009**, *26*, 339–350 DOI: 10.1080/01490450902755364.
- (78) Planer-friedrich, B.; Hartig, C.; Lohmayer, R.; Suess, E.; Mccann, S. H.; Oremland, R. Anaerobic chemolithotrophic growth of the haloalkaliphilic bacterium strain MLMS-1 by disproportionation of monothioarsenate. *Env. Sci Technol* **2015**, *49*, 6554–6563 DOI: 10.1021/acs.est.5b01165.
- (79) Härtig, C.; Planer-friedrich, B. Thioarsenate transformation by filamentous microbial mats thriving in an alkaline, sulfidic hot Spring. *Environ. Sci. Technol.* **2012**, *46* (8), 4348–4356 DOI: 10.1021/es204277j.
- (80) Hartig, C.; Lohmayer, R.; Kolb, S.; Horn, M. A.; Inskeep, W. P.; Planer-friedrich, B. Chemolithotrophic growth of the aerobic hyperthermophilic bacterium *Thermocrinis ruber* OC 14/7/2 on monothioarsenate and arsenite. *FEMS Microbiol Ecol* **2014**, *90*, 747–760 DOI: 10.1111/1574-6941.12431.
- (81) Edwardson, C. F.; Planer-friedrich, B.; Hollibaugh, J. T. Transformation of monothioarsenate by haloalkaliphilic, anoxygenic photosynthetic purple sulfur bacteria. *FEMS Microbiol Ecol* **2014**, *90*, 858–868 DOI: 10.1111/1574-6941.12440.
- (82) Tamaki, S.; Frankenberger, W. T. J. Environmental biochemistry of arsenic. In *Reviews of environmental contamination and toxicology*; Ware, G. W., Ed.; Springer New York: United States, 1992; Vol. 124, pp 79–110 DOI: 10.1007/978-1-4612-2864-6_4
- (83) Ali, W.; Isayenkov, S. V.; Zhao, F. J.; Maathuis, F. J. M. Arsenite transport in plants. *Cell. Mol. Life Sci.* **2009**, *66* (14), 2329–2339 DOI: 10.1007/s00018-009-0021-7.
- (84) Broer, S.; Ji, G.; Broer, A.; Silver, S. Arsenic efflux governed by the arsenic resistance determinant of *Staphylococcus aureus* plasmid pI258. *J. Bacteriol.* **1993**, *175* (11), 3480–3485 DOI: 10.1128/jb.175.11.3480-3485.1993.
- (85) Chen, X.-P.; Zhu, Y.-G.; Hong, M.-N.; Kappler, A.; Xu, Y.-X. Effects of different forms of nitrogen fertilizers on arsenic uptake by rice plants. *Environ. Toxicol. Chem.* **2008**, *27* (4), 881–887 DOI: 10.1897/07-368.1.
- (86) Handley, K. M.; Mcbeth, J. M.; Charnock, J. M.; Vaughan, D. J.; Wincott, P. L.; Polya, D. A.; Lloyd, J. R. Effect of iron redox transformations on arsenic solid-phase associations in an arsenic-rich, ferruginous hydrothermal sediment. *Geochim. Cosmochim. Acta* **2013**, *102*, 124–142 DOI: 10.1016/j.gca.2012.10.024.
- (87) Carlson, H. K.; Clark, I. C.; Blazewicz, S. J.; Iavarone, A. T.; Coates, J. D. Fe(II) oxidation is an innate capability of nitrate-reducing bacteria that involves abiotic and biotic reactions. *J. Bacteriol.* **2013**, *195* (14), 3260–3268 DOI: 10.1128/JB.00058-13.
- (88) Bertin, P. N.; Heinrich-Salmeron, A.; Pelletier, E.; Goulhen-Chollet, F.; Arsène-Ploetze, F.;

839 Gallien, S.; Lauga, B.; Casiot, C.; Calteau, A.; Vallenet, D.; et al. Metabolic diversity among
840 main microorganisms inside an arsenic-rich ecosystem revealed by meta- and proteo-genomics.
841 *ISME J.* **2011**, 5 (11), 1735–1747 DOI: 10.1038/ismej.2011.51.

842 (89) Melton, E. D.; Swanner, E. D.; Behrens, S.; Schmidt, C.; Kappler, A. The interplay of
843 microbially mediated and abiotic reactions in the biogeochemical Fe cycle. *Nat. Rev. Microbiol.*
844 **2014**, 12 (12), 797–809 DOI: 10.1038/nrmicro3347.

845 (90) Masscheleyn, P. H.; Delaune, R. D.; Patrick, Jr., W. H. Effect of redox potential and pH on
846 arsenic speciation and solubility in a contaminated soil. *Environ. Sci. Technol.* **1991**, 25 (8),
847 1414–1419 DOI: 10.1021/es00020a008.

848 (91) Dixit, S.; Hering, J. G. Comparison of arsenic(V) and arsenic(III) sorption onto iron oxide
849 minerals: Implications for arsenic mobility. *Environ. Sci. Technol.* **2003**, 37 (18), 4182–4189
850 DOI: 10.1021/es030309t.

851 (92) Amstaetter, K.; Borch, T.; Larese-Casanova, P.; Kappler, A. Redox transformation of arsenic
852 by Fe(II)-activated goethite (-FeOOH). *Environ. Sci. Technol.* **2010**, 44 (1), 102–108 DOI:
853 10.1021/es901274s.

854 (93) Borch, T.; Kretzschmar, R.; Kappler, A.; Van Cappellen, P.; Ginder-Vogel, M.; Voegelin, A.;
855 Campbell, K. Biogeochemical redox processes and their impact on contaminant dynamics.
856 *Environ. Sci. Technol.* **2010**, 44 (1), 15–23 DOI: 10.1021/es9026248.

857 (94) Liu, X.; Fu, W.; Da, E.; Shi, X.; Cao, Y.; Rathinasabapathi, B.; Chen, Y.; Ma, L. Q. Microbial
858 siderophores and root exudates enhanced goethite dissolution and Fe/As uptake by
859 As-hyperaccumulator *Pteris vittata*. *Environ. Pollut.* **2017**, 223, 230–237 DOI:
860 10.1016/j.envpol.2017.01.016.

861 (95) Liu, W. J.; Zhu, Y. G.; Smith, F. A.; Smith, S. E. Do iron plaque and genotypes affect arsenate
862 uptake and translocation by rice seedlings (*Oryza sativa* L.) grown in solution culture? *J. Exp.*
863 *Bot.* **2004**, 55 (403), 1707–1713 DOI: 10.1093/jxb/erh205.

864 (96) Liu, W. J.; Zhu, Y. G.; Hu, Y.; Williams, P. N.; Gault, A. G.; Meharg, A. A.; Charnock, J. M.;
865 Smith, F. A. Arsenic sequestration in iron plaque, its accumulation and speciation in mature
866 rice plants (*Oryza sativa* L.). *Environ. Sci. Technol.* **2006**, 40 (18), 5730–5736 DOI:
867 10.1021/es060800v.

868 (97) Chen, Z.; Zhu, Y. G.; Liu, W. J.; Meharg, A. A. Direct evidence showing the effect of root
869 surface iron plaque on arsenite and arsenate uptake in rice (*Oryza sativa*) roots. *New Phytol.*
870 **2005**, 165 (1), 91–97 DOI: 10.1111/j.1469-8137.2004.01241.x.

871 (98) Reyes, C.; Lloyd, J. R.; Saltikov, C. W. Geomicrobiology of Iron and Arsenic in Anoxic
872 Sediments. In *Arsenic Contamination of Groundwater*; Ahuja, S., Ed.; John Wiley & Sons, Inc.:
873 Hoboken, NJ, USA, 2008; pp 123–146 DOI: 10.1002/9780470371046.ch6

874 (99) Lloyd, J. R.; Oremland, R. Microbial Transformations of Arsenic in the Environment: From
875 Soda Lakes to Aquifers. *Elements* **2006**, 2, 85–90 DOI: 10.2113/gselements.2.2.85.

876 (100) Islam, F. S.; Gault, A. G.; Boothman, C.; Polya, D. A.; Charnock, J. M.; Chatterjee, D.; Lloyd,
877 J. R. Role of metal-reducing bacteria in arsenic release from Bengal delta sediments. *Nature*
878 **2004**, 430 (6995), 68–71 DOI: 10.1038/nature02638.

- 879 (101) Cummings, D. E.; Caccavo, F.; Fendorf, S.; Rosenzweig, R. F. Arsenic mobilization by the
880 dissimilatory Fe(III)-reducing bacterium *Shewanella alga* BrY. *Environ. Sci. Technol.* **1999**, *33*
881 (5), 723–729 DOI: 10.1021/es980541c.
- 882 (102) Wang, X.-J.; Chen, X.-P.; Kappler, A.; Sun, G.-X.; Zhu, Y.-G. Arsenic binding to iron (II)
883 minerals produced by an iron (III)-reducing aeromonas strain isolated from paddy soil. *Environ.*
884 *Toxicol. Chem.* **2009**, *28* (11), 2255–2262 DOI: 10.1897/09-085.1.
- 885 (103) Muehe, E. M.; Scheer, L.; Daus, B.; Kappler, A. Fate of arsenic during microbial reduction of
886 biogenic versus abiogenic As-Fe(III)-mineral coprecipitates. *Environ. Sci. Technol.* **2013**, *47*
887 (15), 8297–8307 DOI: 10.1021/es400801z.
- 888 (104) Muehe, E. M.; Morin, G.; Scheer, L.; Le Pape, P.; Esteve, I.; Daus, B.; Kappler, A. Arsenic(V)
889 Incorporation in Vivianite during Microbial Reduction of Arsenic(V)-Bearing Biogenic Fe(III)
890 (Oxyhydr)oxides. *Environ. Sci. Technol.* **2016**, *50* (5), 2281–2291 DOI:
891 10.1021/acs.est.5b04625.
- 892 (105) Islam, F. S.; Pederick, R. L.; Gault, A. G.; Adams, L. K.; Polya, D. A.; Charnock, J. M.; Lloyd,
893 J. R. Interactions between the Fe(III) -Reducing Bacterium *Geobacter sulfurreducens* and
894 Arsenate, and Capture of the Metalloid by Biogenic Fe(II). *Appl Env. Microbiol* **2005**, *71* (12),
895 8642–8648 DOI: 10.1128/AEM.71.12.8642.
- 896 (106) Islam, F. S.; Boothman, C.; Gault, A. G.; Polya, D. A.; Lloyd, J. R. Potential role of the
897 Fe(III)-reducing bacteria *Geobacter* and *Geothrix* in controlling arsenic solubility in Bengal
898 delta sediments. *Mineral. Mag.* **2005**, *69* (5), 865–875 DOI: 10.1180/0026461056950294.
- 899 (107) Wang, X.; Chen, X.; Yang, J.; Wang, Z.; Sun, G. Effect of microbial mediated iron plaque
900 reduction on arsenic mobility in paddy soil. *J. Environ. Sci.* **2009**, *21* (11), 1562–1568 DOI:
901 10.1016/S1001-0742(08)62456-0.
- 902 (108) Campbell, K. M.; Malasarn, D. Simultaneous Microbial Reduction of Iron(III) and Arsenic(V)
903 in Suspensions of Hydrous Ferric Oxide. *Env. Sci Technol* **2006**, *40* (19), 5950–5955 DOI:
904 10.1021/es0600476.
- 905 (109) Kocar, B. D.; Herbel, M. J.; Tufano, K. J.; Fendorf, S. Contrasting effects of dissimilatory
906 iron(III) and arsenic(V) reduction on arsenic retention and transport. *Environ. Sci. Technol.*
907 **2006**, *40* (21), 6715–6721 DOI: 10.1021/es061540k.
- 908 (110) Oremland, R. S.; Stolz, J. F. Arsenic, microbes and contaminated aquifers. *Trends Microbiol.*
909 **2005**, *13* (2), 45–49 DOI: 10.1155/2011/863623.
- 910 (111) Zobrist, J.; Dowdle, P. R.; Davis, J. A.; Oremland, R. S. Mobilization of arsenite by
911 dissimilatory reduction of adsorbed arsenate. *Environ. Sci. Technol.* **2000**, *34* (22), 4747–4753
912 DOI: 10.1021/es001068h.
- 913 (112) Hohmann, C.; Winkler, E.; Morin, G.; Kappler, A. Anaerobic Fe(II)-oxidizing bacteria show as
914 resistance and immobilize as during Fe(III) mineral precipitation. *Environ. Sci. Technol.* **2010**,
915 *44* (1), 94–101 DOI: 10.1021/es900708s.
- 916 (113) Hohmann, C.; Morin, G.; Ona-Nguema, G.; Guigner, J. M.; Brown, G. E.; Kappler, A.
917 Molecular-level modes of As binding to Fe(III) (oxyhydr)oxides precipitated by the anaerobic
918 nitrate-reducing Fe(II)-oxidizing *Acidovorax* sp. strain BoFeN1. *Geochim. Cosmochim. Acta*

- 919 **2011**, 75 (17), 4699–4712 DOI: 10.1016/j.gca.2011.02.044.
- 920 (114) Omoregie, E. O.; Couture, R.; Cappellen, P. Van; Corkhill, C. L.; Charnock, J. M.; Polya, D.
 921 A.; Vaughan, D.; Vanbroekhoven, K.; Lloyd, R. Arsenic Bioremediation by Biogenic Iron
 922 Oxides and Sulfides. *Appl. Environ. Microbiol.* **2013**, 79 (14), 4325–4335 DOI:
 923 10.1128/AEM.00683-13.
- 924 (115) Nitzsche, K. S.; Weigold, P.; Lösekann-Behrens, T.; Kappler, A.; Behrens, S. Microbial
 925 community composition of a household sand filter used for arsenic, iron, and manganese
 926 removal from groundwater in Vietnam. *Chemosphere* **2015**, 138, 47–59 DOI:
 927 10.1016/j.chemosphere.2015.05.032.
- 928 (116) Kleinert, S.; Muehe, E. M.; Posth, N. R.; Dippon, U.; Daus, B.; Kappler, A. Biogenic Fe(III)
 929 minerals lower the efficiency of iron-mineral-based commercial filter systems for arsenic
 930 removal. *Environ. Sci. Technol.* **2011**, 45 (17), 7533–7541 DOI: 10.1021/es201522n.
- 931 (117) Okibe, N.; Koga, M.; Morishita, S.; Tanaka, M.; Heguri, S.; Asano, S.; Sasaki, K.; Hirajima, T.
 932 Microbial formation of crystalline scorodite for treatment of As(III) -bearing copper refinery
 933 process solution using *Acidianus brierleyi*. *Hydrometallurgy* **2014**, 143, 34–41 DOI:
 934 10.1016/j.hydromet.2014.01.008.
- 935 (118) Mandal, B. K.; Suzuki, K. T. Arsenic round the world : a review. *Talanta* **2002**, 58, 201–235
 936 DOI: 10.1016/S0039-9140(02)00268-0.
- 937 (119) Corkhill, C. L.; Wincott, P. L.; Lloyd, J. R.; Vaughan, D. J. The oxidative dissolution of
 938 arsenopyrite (FeAsS) and enargite (Cu₃AsS₄) by *Leptospirillum ferrooxidans*. *Geochim.*
 939 *Cosmochim. Acta* **2008**, 72 (23), 5616–5633 DOI: 10.1016/j.gca.2008.09.008.
- 940 (120) O'Day, P. a; Vlassopoulos, D.; Root, R.; Rivera, N. The influence of sulfur and iron on
 941 dissolved arsenic concentrations in the shallow subsurface under changing redox conditions.
 942 *Proc. Natl. Acad. Sci. U. S. A.* **2004**, 101 (38), 13703–13708 DOI: 10.1073/pnas.0402775101.
- 943 (121) Kim, M.; Nriagu, J.; Haack, S. Arsenic species and chemistry in groundwater of southeast
 944 Michigan. *Environ. Pollut.* **2002**, 120, 379–390 DOI: 10.1016/S0269-7491(02)00114-8.
- 945 (122) Rochette, E. A.; Bostick, B. C.; Li, G.; Fendorf, S. Kinetics of arsenate reduction by dissolved
 946 sulfide. *Environ. Sci. Technol.* **2000**, 34 (22), 4714–4720 DOI: 10.1021/es000963y.
- 947 (123) Kuhn, A.; Sigg, L.; Jul, N. Arsenic Cycling in Eutrophic Lake Greifen, Switzerland: Influence
 948 of Seasonal Redox Processes. *Limnol. Ocean.* **1993**, 38 (5), 1052–1059 DOI:
 949 10.4319/lo.1993.38.5.1052.
- 950 (124) Newman, D. K.; Beveridge, T. J. Precipitation of Arsenic Trisulfide by *Desulfotomaculum*
 951 *auripigmentum*. *Appl. Environ. Microbiol.* **1997**, 63 (5), 2022–2028.
- 952 (125) Rittle, K. A.; Drever, J. I.; Colberg, P. J. S. Precipitation of Arsenic during Bacterial Sulfate
 953 Reduction. *Geomicrobiol. J.* **1995**, 13 (1), 1–11 DOI: 10.1080/01490459509378000.
- 954 (126) Hoefft, S. E.; Kulp, T. R.; Stolz, J. F.; Hollibaugh, J. T.; Oremland, R. S. Dissimilatory Arsenate
 955 Reduction with Sulfide as Electron Donor: Experiments with Mono Lake Water and Isolation
 956 of Strain MLMS-1, a Chemoautotrophic Arsenate Respirer. *Appl. Environ. Microbiol.* **2004**, 70
 957 (5), 2741–2747 DOI: 10.1128/AEM.70.5.2741.
- 958 (127) Hollibaugh, J. T.; Budinoff, C.; Hollibaugh, R. A.; Ransom, B.; Bano, N. Sulfide Oxidation

- 959 Coupled to Arsenate Reduction by a Diverse Microbial Community in a Soda Lake. *Appl Env.*
960 *Microbiol* **2006**, 72 (3), 2043–2049 DOI: 10.1128/AEM.72.3.2043.
- 961 (128) Hoeft, S. E.; Blum, J. S.; Stolz, J. F.; Tabita, F. R.; Witte, B.; King, G. M.; Santini, J. M.;
962 Oremland, R. S. *Alkalilimnicola ehrlichii* sp. nov., a novel, arsenite-oxidizing haloalkaliphilic
963 gammaproteobacterium capable of chemoautotrophic or heterotrophic growth with nitrate or
964 oxygen as the electron acceptor. *Int. J. Syst. Evol. Microbiol.* **2007**, 57 (3), 504–512 DOI:
965 10.1099/ij.s.0.64576-0.
- 966 (129) Rhine, E. D.; Phelps, C. D.; Young, L. Y. Anaerobic arsenite oxidation by novel denitrifying
967 isolates. *Environ. Microbiol.* **2006**, 8 (5), 899–908 DOI: 10.1111/j.1462-2920.2005.00977.x.
- 968 (130) Sun, W.; Sierra, R.; Fernandez, N.; Sanz, J. L.; Amils, R.; Legatzki, A.; Maier, R. M.; Amils,
969 R.; Field, J. A. Molecular Characterization and *In Situ* Quantification of Anoxic Arsenite
970 Oxidizing Denitrifying Enrichment cultures. *FEMS Microbiol Ecol.* **2009**, 68 (1), 72–85 DOI:
971 10.1007/s12671-013-0269-8.Moving.
- 972 (131) Senko, J. M.; Dewers, T. A.; Krumholz, L. R. Effect of Oxidation Rate and Fe(II) State on
973 Microbial Nitrate-Dependent Fe(III) Mineral Formation. *Appl. Environ. Microbiol.* **2005**, 71
974 (11), 7172–7177 DOI: 10.1128/AEM.71.11.7172.
- 975 (132) Weber, K. A.; Urrutia, M. M.; Churchill, P. F.; Kukkadapu, R. K.; Roden, E. E. Anaerobic
976 redox cycling of iron by freshwater sediment microorganisms. *Environ. Microbiol.* **2006**, 8,
977 100–113 DOI: 10.1111/j.1462-2920.2005.00873.x.
- 978 (133) Schippers, A.; Jorgensen, B. B. Biogeochemistry of pyrite and iron sulfide oxidation in marine
979 sediments. *Geochim. Cosmochim. Acta* **2002**, 66 (1), 85–92 DOI:
980 10.1016/S0016-7037(01)00745-1.
- 981 (134) Stevenson, F. J. *Humus chemistry: genesis, composition, reactions*, 2 nd.; John Wiley & Sons,
982 1994.
- 983 (135) Jiang, J.; Bauer, I.; Paul, A.; Kappler, A. Arsenic redox changes by microbially and chemically
984 formed semiquinone radicals and hydroquinones in a humic substance model quinone. *Environ.*
985 *Sci. Technol.* **2009**, 43 (10), 3639–3645 DOI: 10.1021/es803112a.
- 986 (136) Sharma, P.; Rolle, M.; Kocar, B. D.; Fendorf, S.; Kappler, A. Influence of Natural Organic
987 Matter on As Transport and Retention. *Environ. Sci. Technol.* **2010**, 45 (2), 546–553 DOI:
988 10.2134/jeq2003.1393.
- 989 (137) Sharma, P.; Kappler, A. Desorption of arsenic from clay and humic acid-coated clay by
990 dissolved phosphate and silicate. *J. Contam. Hydrol.* **2011**, 126 (3–4), 216–225 DOI:
991 10.1016/j.jconhyd.2011.08.005.
- 992 (138) Bauer, M.; Blodau, C. Arsenic distribution in the dissolved, colloidal and particulate size
993 fraction of experimental solutions rich in dissolved organic matter and ferric iron. *Geochim.*
994 *Cosmochim. Acta* **2009**, 73 (3), 529–542 DOI: 10.1016/j.gca.2008.10.030.
- 995 (139) Thomasarrigo, L. K.; Mikutta, C.; Byrne, J.; Barnettler, K.; Kappler, A.; Kretzschmar, R. Iron
996 and Arsenic Speciation and Distribution in Organic Flocs from Streambeds of an
997 Arsenic-Enriched Peatland. *Environ. Sci. Technol.* **2014**, 48, 13218–13228 DOI:
998 10.1021/es503550g.

- 999 (140) Macalady, D. L.; Ranville, J. F. The chemistry and geochemistry of natural organic matter
1000 (NOM). In *Perspectives in environmental chemistry*; Macalady, D. L., Ed.; Oxford University
1001 Press: New York, 1988; pp 94–137.
- 1002 (141) Buschmann, J.; Kappeler, A.; Lindauer, U.; Kistler, D.; Berg, M.; Sigg, L. Arsenite and
1003 arsenate binding to dissolved humic acids: Influence of pH, type of humic acid, and aluminum.
1004 *Environ. Sci. Technol.* **2006**, *40* (19), 6015–6020 DOI: 10.1021/es061057+.
- 1005 (142) Sharma, P.; Ofner, J.; Kappler, A. Formation of binary and ternary colloids and dissolved
1006 complexes of organic matter, Fe and As. *Environ. Sci. Technol.* **2010**, *44* (12), 4479–4485 DOI:
1007 10.1021/es100066s.
- 1008 (143) Langner, P.; Mikutta, C.; Kretzschmar, R. Arsenic sequestration by organic sulphur in peat. *Nat.*
1009 *Geosci.* **2011**, *5* (1), 66–73 DOI: 10.1038/ngeo1329.
- 1010 (144) Rowland, H. A. L.; Polya, D. A.; Lloyd, J. R.; Pancost, R. D. Characterisation of organic matter
1011 in a shallow, reducing, arsenic-rich aquifer, West Bengal. *Org. Geochem.* **2006**, *37*, 1101–1114
1012 DOI: 10.1016/j.orggeochem.2006.04.011.
- 1013 (145) Rowland, H. A. L.; Pederick, R. L.; Polya, D. A.; Pancost, R. D.; Dongen, B. E. Van; Gault, A.
1014 G.; Vaughan, D. J.; Bryant, C.; Anderson, B.; Lloyd, J. R. The control of organic matter on
1015 microbially mediated iron reduction and arsenic release in shallow alluvial aquifers, Cambodia.
1016 *Geobiology* **2007**, *5*, 281–292 DOI: 10.1111/j.1472-4669.2007.00100.x.
- 1017 (146) Rowland, H. A. L.; Boothman, C.; Pancost, R.; Gault, A. G.; Polya, D. A.; Lloyd, J. R. The
1018 Role of Indigenous Microorganisms in the Biodegradation of Naturally Occurring Petroleum,
1019 the Reduction of Iron, and the Mobilization of Arsenite from West Bengal. *J. Environ. Qual.*
1020 **2009**, *38*, 1598–1607 DOI: 10.2134/jeq2008.0223.
- 1021 (147) Bauer, M.; Blodau, C. Mobilization of arsenic by dissolved organic matter from iron oxides,
1022 soils and sediments. *Sci. Total Environ.* **2006**, *354* (2–3), 179–190 DOI:
1023 10.1016/j.scitotenv.2005.01.027.
- 1024 (148) Huang, H.; Jia, Y.; Sun, G.-X.; Zhu, Y.-G. Arsenic speciation and volatilization from flooded
1025 paddy soils amended with different organic matters-supporting information. *Environ. Sci.*
1026 *Technol.* **2012**, *46* (4), 2163–2168 DOI: 10.1021/es203635s.
- 1027 (149) Jia, Y.; Huang, H.; Chen, Z.; Zhu, Y. G. Arsenic uptake by rice is influenced by
1028 microbe-mediated arsenic redox changes in the rhizosphere. *Environ. Sci. Technol.* **2014**, *48* (2),
1029 1001–1007 DOI: 10.1021/es403877s.
- 1030 (150) Palmer, N. E.; Freudenthal, J. H.; Wandruszka, R. Von. Reduction of Arsenates by Humic
1031 Materials. *Environ. Chem.* **2006**, *3*, 131–136 DOI: 10.1071/EN05081.
- 1032 (151) Palmer, N. E.; Wandruszka, R. Von. Humic acids as reducing agents: the involvement of
1033 quinoid moieties in arsenate reduction. *Environ. Sci. Pollut. Res.* **2010**, *17*, 1362–1370 DOI:
1034 10.1007/s11356-010-0322-2.
- 1035 (152) Redman, A. D.; Macalady, D.; Ahmann, D. Natural organic matter affects arsenic speciation
1036 and sorption onto hematite. *Environ. Sci. Technol.* **2002**, *36* (13), 2889–2896 DOI:
1037 10.1021/es0112801.
- 1038 (153) Niggemyer, A.; Spring, S.; Stackebrandt, E. Isolation and Characterization of a Novel As (V)

- 1039 -Reducing Bacterium: Implications for Arsenic Mobilization and the Genus *Desulfotobacterium*.
 1040 *Appl. Environ. Microbiol.* **2001**, 67 (12), 5568–5580 DOI: 10.1128/AEM.67.12.5568.
- 1041 (154) Stolz, J. F.; Oremland, R. S. Bacterial respiration of arsenic and selenium. *FEMS Microbiol*
 1042 *Rev* **1999**, 23, 615–627 DOI: 10.1111/j.1574-6976.1999.tb00416.x.
- 1043 (155) Handley, K. M.; Hery, M.; Lloyd, J. R. Redox cycling of arsenic by the hydrothermal marine
 1044 bacterium *Marinobacter santoriniensis*. *Env. Microbiol* **2009**, 11 (6), 1601–1611 DOI:
 1045 10.1111/j.1462-2920.2009.01890.x.
- 1046 (156) Lear, G.; Song, B.; Gault, A. G.; Polya, D. A.; Lloyd, J. R. Molecular analysis of
 1047 arsenate-reducing bacteria within Cambodian sediments following amendment with acetate.
 1048 *Appl. Environ. Microbiol.* **2007**, 73 (4), 1041–1048 DOI: 10.1128/AEM.01654-06.
- 1049 (157) Liu, A.; Garcia-dominguez, E.; Rhine, E. D.; Young, L. Y. A novel arsenate respiring isolate
 1050 that can utilize aromatic substrates. *FEMS Microbiol Ecol* **2004**, 48, 323–332 DOI:
 1051 10.1016/j.femsec.2004.02.008.
- 1052 (158) Liu, W. J.; McGrath, S. P.; Zhao, F. J. Silicon has opposite effects on the accumulation of
 1053 inorganic and methylated arsenic species in rice. *Plant Soil* **2014**, 376 (1–2), 423–431 DOI:
 1054 10.1007/s11104-013-1991-7.
- 1055 (159) Escudero, L. V.; Casamayor, E. O.; Chong, G.; Pedro, C.; Demergasso, C. Distribution of
 1056 Microbial Arsenic Reduction, Oxidation and Extrusion Genes along a Wide Range of
 1057 Environmental Arsenic Concentrations. *PLoS One* **2013**, 8 (10), e78890 DOI:
 1058 10.1371/journal.pone.0078890.
- 1059 (160) Han, Y.; Fu, J.; Xiang, P.; Cao, Y.; Rathinasabapathi, B.; Chen, Y.; Ma, L. Q. Arsenic and
 1060 phosphate rock impacted the abundance and diversity of bacterial arsenic oxidase and reductase
 1061 genes in rhizosphere of As-hyperaccumulator *Pteris vittata*. *J. Hazard. Mater.* **2017**, 321,
 1062 146–153 DOI: 10.1016/j.jhazmat.2016.08.079.
- 1063 (161) Wang, X.; Rathinasabapathi, B.; Oliveira, L. M. de; Guilherme, L. R. G.; Ma, L. Q.
 1064 Bacteria-Mediated Arsenic Oxidation and Reduction in the Growth Media of Arsenic
 1065 Hyperaccumulator *Pteris vittata*. *Env. Sci Technol* **2012**, 46, 11259–11266 DOI:
 1066 10.1021/es300454b.
- 1067 (162) Zhang, S. Y.; Zhao, F. J.; Sun, G. X.; Su, J. Q.; Yang, X. R.; Li, H.; Zhu, Y. G. Diversity and
 1068 abundance of arsenic biotransformation genes in paddy soils from southern china. *Environ. Sci.*
 1069 *Technol.* **2015**, 49 (7), 4138–4146 DOI: 10.1021/acs.est.5b00028.
- 1070 (163) Jia, Y.; Huang, H.; Sun, G. X.; Zhao, F. J.; Zhu, Y. G. Pathways and relative contributions to
 1071 arsenic volatilization from rice plants and paddy soil. *Environ. Sci. Technol.* **2012**, 46 (15),
 1072 8090–8096 DOI: 10.1021/es300499a.
- 1073 (164) Xiao, K. Q.; Li, L. G.; Ma, L. P.; Zhang, S. Y.; Bao, P.; Zhang, T.; Zhu, Y. G. Metagenomic
 1074 analysis revealed highly diverse microbial arsenic metabolism genes in paddy soils with
 1075 low-arsenic contents. *Environ. Pollut.* **2016**, 211, 1–8 DOI: 10.1016/j.envpol.2015.12.023.
- 1076 (165) Lu, K.; Mahbub, R.; Cable, P. H.; Ru, H.; Parry, N. M. A.; Bodnar, W. M.; Wishnok, J. S.;
 1077 Styblo, M.; Swenberg, J. A.; Fox, J. G.; et al. Gut Microbiome Phenotypes Driven by Host
 1078 Genetics Affect Arsenic Metabolism. *Chem. Res. Toxicol.* **2014**, 27, 172–174 DOI:

- 1079 10.1021/tx400454z.
- 1080 (166) Cai, L.; Yu, K.; Yang, Y.; Chen, B. W.; Li, X. D.; Zhang, T. Metagenomic exploration reveals
1081 high levels of microbial arsenic metabolism genes in activated sludge and coastal sediments.
1082 *Appl. Microbiol. Biotechnol.* **2013**, 97 (21), 9579–9588 DOI: 10.1007/s00253-012-4678-8.
- 1083 (167) Rascovan, N.; Javier, M.; Martín P, V.; María, E. F. Metagenomic study of red biofilms from
1084 Diamante Lake reveals ancient arsenic bioenergetics in haloarchaea Metagenomic study of red
1085 biofilms from Diamante Lake reveals ancient arsenic bioenergetics in haloarchaea. *ISME J.*
1086 **2016**, 10, 299–309 DOI: 10.1038/ismej.2015.109.
- 1087 (168) Luo, J.; Bai, Y.; Liang, J.; Qu, J. Metagenomic Approach Reveals Variation of Microbes with
1088 Arsenic and Antimony Metabolism Genes from Highly Contaminated Soil. *PLoS One* **2014**, 9
1089 (10), e108185 DOI: 10.1371/journal.pone.0108185.
- 1090 (169) Sangwan, N.; Lambert, C.; Sharma, A.; Gupta, V.; Khurana, P.; Khurana, J. P.; Sockett, R. E.;
1091 Gilbert, J. A.; Lal, R. Arsenic rich Himalayan hot spring metagenomics reveal genetically
1092 novel predator – prey genotypes. *Environ. Microbiol. Rep.* **2015**, 7, 812–823 DOI:
1093 10.1111/1758-2229.12297.
- 1094 (170) Aguiar-pulido, V.; Huang, W.; Suarez-ulloa, V.; Cickovski, T.; Mathee, K.; Narasimhan, G.
1095 Metagenomics, metatranscriptomics, and metabolomics approaches for microbiome analysis.
1096 *Evol. Bioinforma.* **2016**, 12 (S1), 5–16 DOI: 10.4137/EBO.S36436.
- 1097 (171) Kour, M. Isolation and Characterization of Metal Resistance Genes by using
1098 Metatranscriptomic Approach. Master Dissertation, Thapar University, Patiala, Punjab-147004,
1099 2014.
- 1100 (172) Carvalhais, L. C.; Dennis, P. G.; Tyson, G. W.; Schenk, P. M. Application of
1101 metatranscriptomics to soil environments. *J. Microbiol. Methods* **2012**, 91 (2), 246–251 DOI:
1102 10.1016/j.mimet.2012.08.011.
- 1103 (173) Gilbert, J. A.; Field, D.; Huang, Y.; Edwards, R.; Li, W.; Gilna, P.; Joint, I. Detection of Large
1104 Numbers of Novel Sequences in the Metatranscriptomes of Complex Marine Microbial
1105 Communities. *PLoS One* **2008**, 3 (8), e3042 DOI: 10.1371/journal.pone.0003042.
- 1106 (174) Lafuente, A.; Patricia, P.; Molina-s, D.; Caviades, M. A.; Rodr, I. D. Unraveling the effect of
1107 arsenic on the model *Medicago – Ensifer* interaction: a transcriptomic meta-analysis. *ISME J.*
1108 **2015**, 205, 255–272 DOI: 10.1111/nph.13009.
- 1109 (175) Zhang, Y.; Chen, S.; Hao, X.; Su, J. Q.; Xue, X.; Yan, Y.; Zhu, Y. G.; Ye, J. Transcriptomic
1110 analysis reveals adaptive responses of an enterobacteriaceae strain LSJC7 to arsenic exposure.
1111 *Front. Microbiol.* **2016**, 7, 1–12 DOI: 10.3389/fmicb.2016.00636.
- 1112 (176) Moran, M. A.; Satinsky, B.; Gifford, S. M.; Luo, H.; Rivers, A.; Chan, L.; Meng, J.; Durham, B.
1113 P.; Shen, C.; Varaljay, V. A.; et al. Sizing up metatranscriptomics. **2012**, 7 (2), 237–243 DOI:
1114 10.1038/ismej.2012.94.
- 1115 (177) VerBerkmoes, N. C.; Denef, V. J.; Hettich, R. L.; Banfield, J. F. Functional analysis of natural
1116 microbial consortia using community proteomics. *Nat Rev Microbiol* **2009**, 7 (3), 196–205 DOI:
1117 10.1038/nrmicro2080.
- 1118 (178) Fiehn, O. Metabolomics - The link between genotypes and phenotypes. *Plant Mol. Biol.* **2002**,

- 1119 48 (1–2), 155–171 DOI: 10.1023/A:1013713905833.
- 1120 (179) Schneider, T.; Riedel, K. Environmental proteomics: Analysis of structure and function of
1121 microbial communities. *Proteomics* **2010**, *10* (4), 785–798 DOI: 10.1002/pmic.200900450.
- 1122 (180) Lankadurai, B. P.; Nagato, E. G.; Simpson, M. J. Environmental metabolomics: an emerging
1123 approach to study organism responses to environmental stressors. *Environ. Rev.* **2013**, *21*,
1124 180–205 DOI: 10.1139/er-2013-0011.
- 1125 (181) Halter, D.; Cordi, A.; Gribaldo, S.; Gallien, S.; Goulhen-Chollet, F.; Heinrich-Salmeron, A.;
1126 Carapito, C.; Pagnout, C.; Montaut, D.; Seby, F.; et al. Taxonomic and functional prokaryote
1127 diversity in mildly arsenic-contaminated sediments. *Res. Microbiol.* **2011**, *162* (9), 878–887
1128 DOI: 10.1016/j.resmic.2011.06.001.
- 1129 (182) Shi, X.; Wei, X.; Koo, I.; Schmidt, R. H.; Yin, X.; Kim, S. H.; Vaughn, A.; McClain, C. J.;
1130 Arteel, G. E.; Zhang, X.; et al. Metabolomic Analysis of the Effects of Chronic Arsenic
1131 Exposure in a Mouse Model of Diet-Induced Fatty Liver Disease. *J. Proteome Res.* **2014**, *13*,
1132 547–554 DOI: 10.1021/pr400719u.
- 1133 (183) Bastida, F.; Moreno, J. L.; Nicolás, C.; Hernández, T.; García, C. Soil metaproteomics: a
1134 review of an emerging environmental science. Significance, methodology and perspectives.
1135 *Eur. J. Soil Sci.* **2009**, *60* (6), 845–859 DOI: 10.1111/j.1365-2389.2009.01184.x.
- 1136 (184) Lu, K.; Abo, R. P.; Schlieper, K. A.; Michelle, E.; Levine, S.; Wishnok, J. S.; Swenberg, J. A.;
1137 Tannenbaum, S. R.; Fox, J. G. Arsenic Exposure Perturbs the Gut Microbiome and Its
1138 Metabolic Profile in Mice: An Integrated Metagenomics and Metabolomics Analysis. *Environ.*
1139 *Health Perspect.* **2014**, *122*, 284–291 DOI: 10.1289/ehp.1307429.
- 1140 (185) Tripathi, R. D.; Tripathi, P.; Dwivedi, S.; Dubey, S.; Chatterjee, S.; Chakrabarty, D.; Trivedi, P.
1141 K. Arsenomics: Omics of arsenic metabolism in plants. *Front. Physiol.* **2012**, *3*, 1–14 DOI:
1142 10.3389/fphys.2012.00275.
- 1143 (186) Sacheti, P.; Bhonsle, H.; Patil, R.; Kulkarni, M. J.; Srikanth, R.; Gade, W. Arsenomics of
1144 *Exiguobacterium* sp. PS (NCIM 5463). *RSC Adv.* **2013**, *3*, 9705–9713 DOI:
1145 10.1039/c3ra40897c.
- 1146 (187) Zhang, S.; Williams, P. N.; Luo, J.; Zhu, Y. Microbial mediated arsenic biotransformation in
1147 wetlands. *Front. Environ. Sci. eng.* **2017**, *11* (1), 1–11 DOI: 10.1007/s11783-017-0893-y.
- 1148 (188) Meharg, A. A.; Williams, P. N.; Adomako, E.; Lawgali, Y. Y.; Deacon, C.; Villada, A.;
1149 Cambell, R. C. J.; Sun, G.; Zhu, Y. G.; Feldmann, J.; et al. Geographical variation in total and
1150 inorganic arsenic content of polished (white) rice. *Environ. Sci. Technol.* **2009**, *43* (5),
1151 1612–1617 DOI: 10.1021/es802612a.
- 1152 (189) Sun, G. X.; Williams, P. N.; Carey, A. M.; Zhu, Y. G.; Deacon, C.; Raab, A.; Feldmann, J.;
1153 Islam, R. M.; Meharg, A. A. Inorganic arsenic in rice bran and its products are an order of
1154 magnitude higher than in bulk grain. *Environ. Sci. Technol.* **2008**, *42* (19), 7542–7546 DOI:
1155 10.1021/es801238p.
- 1156 (190) Sun, G. X.; Williams, P. N.; Zhu, Y. G.; Deacon, C.; Carey, A. M.; Raab, A.; Feldmann, J.;
1157 Meharg, A. A. Survey of arsenic and its speciation in rice products such as breakfast cereals,
1158 rice crackers and Japanese rice condiments. *Environ. Int.* **2009**, *35* (3), 473–475 DOI:

- 10.1016/j.envint.2008.07.020.
- (191) Norton, G. J.; Pinson, S. R. M.; Alexander, J.; McKay, S.; Hansen, H.; Duan, G.; Islam, M. R.; Islam, S.; Stroud, J. L.; Zhao, F.; et al. Variation in grain arsenic assessed in a diverse panel of rice (*Oryza sativa*) grown in multiple sites. *New Phytol.* **2011**, *193*, 650–664 DOI: 10.1111/j.1469-8137.2011.03983.x.
- (192) Lomax, C.; Liu, W. J.; Wu, L.; Xue, K.; Xiong, J.; Zhou, J.; McGrath, S. P.; Meharg, A. A.; Miller, A. J.; Zhao, F. J. Methylated arsenic species in plants originate from soil microorganisms. *New Phytol.* **2012**, *193* (3), 665–672 DOI: 10.1111/j.1469-8137.2011.03956.x.
- (193) Oremland, R. S.; Dowdle, P. R.; Hoeft, S.; Sharp, J. O.; Schaefer, J. K.; Miller, L. G.; Blum, J. S.; Smith, R. L.; Bloom, N. S.; Wallschlager, D. Bacterial dissimilatory reduction of arsenate and sulfate in meromictic Mono Lake, California. *Geochim. Cosmochim. Acta* **2000**, *64* (18), 3073–3084 DOI: 10.1016/S0016-7037(00)00422-1.
- (194) Kulp, T. R.; Hoeft, S. E.; Miller, L. G.; Saltikov, C.; Murphy, J. N.; Han, S.; Lanoil, B.; Oremland, R. S. Dissimilatory Arsenate and Sulfate Reduction in Sediments of Two Hypersaline, Arsenic-Rich Soda Lakes: Mono and Searles Lakes, California. *Appl. Environ. Microbiol.* **2006**, *72* (10), 6514–6526 DOI: 10.1128/AEM.01066-06.
- (195) Reed, D. C.; Algar, C. K.; Huber, J. A.; Dick, G. J. Gene-centric approach to integrating environmental genomics and biogeochemical models. *Proc. Natl Acad. Sci.* **2014**, *111* (5), 1879–1884 DOI: 10.1073/pnas.1313713111.
- (196) Louca, S.; Hawley, A. K.; Katsev, S.; Torres-Beltran, M.; Bhatia, M. P.; Kheirandish, S.; Michiels, C. C.; Capelle, D.; Lavik, G.; Doebeli, M.; et al. Integrating biogeochemistry with multiomic sequence information in a model oxygen minimum zone. *Proc. Natl. Acad. Sci.* **2016**, *113* (40), E5925–E5933 DOI: 10.1073/pnas.1602897113.

Legends

Figure 1. Proposed pathways for arsenic redox reactions and synthesis of novel organoarsenicals.

Figure 2. The model of effect of Fe, N, S and natural organic matter (NOM) on microbes involved in arsenic biogeochemical cycling highlights proteins associated with elemental metabolisms. Green ovals denote arsenic transporters, yellow ovals denote transmembrane enzymes. Red words are enzymes, blue words are related arsenic compounds. The full name of enzymes that were not mentioned in the text was provided in the follow, NarG, transmembrane nitrate reductase that drives the nitrate

1195 reduction to nitrite; Nas, cytoplasmic-assimilatory nitrate reductase that drives the
1196 nitrate reduction to nitrite; NapA, periplasmic-dissimilatory nitrate reductase; Nir /Nrf
1197 (associated with NapA), nitrite reductase that drives the nitrite reduction to nitric
1198 oxide; NorB/C, nitric oxide reductase that drives the nitric oxide reduction to nitrous
1199 oxide; NosZ, nitrous oxide reductase that drives the nitrous oxide reduction to
1200 nitrogen; Nif, nitrogenase that catalyzes the nitrogen fixation to ammonia; Hs,
1201 hydrazine synthase that catalyzes the production of nitrogen from nitrous oxide and
1202 ammonia; cyt, cytochrome.

1204 Table 1. The genes involved in arsenic metabolisms

Gene	Protein	Protein abbreviation	function	Reference
<i>As(III) oxidation</i>				
<i>aioA</i> and <i>aioB</i>	As(III) oxidase	AioAB	Oxidate As(III)	9
<i>arxA</i>	As(III) oxidase	ArxAB	Oxidate As(III)	10
<i>arxB/arxB2</i>	[4Fe-4S] containing protein	ArxB/ ArxB2	Unknown function	11
<i>aioX/arxX</i>	As(III)-binding protein	AioX/ArxX	Involved in As(III)-based signaling and regulation of As(III) oxidation	12, 11
<i>aioS/arxS</i>	sensor histidine kinase	AioS /ArxS	One part of two-component signal transduction system	9, 11
<i>aioR/arxR</i>	transcriptional regulator	AioR/ArxR	Regulate the expression of <i>aio/arx</i> operon	9, 11
<i>moeA</i>	molybdenum cofactor biosynthesis protein	MoeA	Synthesize the molybdenum cofactor of AioAB oxidase	9
<i>arxC</i>	membrane anchoring and quinol oxidoreductase subunit	ArxC	Involved in As(III) oxidation	11
<i>arxD</i>	TorD-like molybdoenzyme chaperone	ArxD	Involved in As(III) oxidation	11
<i>arsH</i>	organoarsenical oxidase	ArsH	Oxidate trivalent methylated and aromatic arsenicals, reduce chromium and iron	13, 14
<i>As(V) reduction</i>				
<i>arrA</i> and <i>arrB</i>	As(V) respiratory reductases	ArrAB	Reduce As(V)	15, 16
<i>arrC</i>	As(V) reductase membranous subunit	ArrC	Involved in As(V) reduction	17
<i>arrD</i>	As(V) reductase chaperon	ArrD	Involved in As(V) reduction	17
<i>arrS</i>	sensor histidine kinase	ArrS	Regulate the expression of <i>arr</i> operon	17
<i>arsC</i>	As(V) reductase	ArsC	Reduce As(V)	18
<i>ACR2</i>	As(V) reductase	ACR2	Reduce As(V)	19

<i>ACR1</i>	transcriptional regulatory protein	ACR1	Regulate the expression of <i>ACR</i> genes	19
<i>arrR/arsR</i>	arsenic-responsive repressor	ArrR /ArsR	Regulate the expression of <i>arr/ars</i> operon	17, 20
<i>GstB</i>	glutathione S-transferase B	GstB	Reduce As(V) to As(III) with reduced GSH	21
<i>arsN</i>	acetyltransferase	ArsN	Putative As(V) reductase	22
<i>Arsenic methylation and demethylation</i>				
<i>arsM</i>	As(III) S-adenosylmethionine (SAM) methyltransferases	ArsM	Methylate arsenic	23
<i>arsI</i>	A C · As lyase	ArsI	Catalyze demethylation of trivalent organoarsenicals	24
<i>Arsenic transport</i>				
<i>arsB</i>	As(III)-pump protein	ArsB	Extrude As(III) from the cell	25
<i>arsA</i>	As(III)-pump ATPase	ArsA	The catalytic subunit of an oxyanion-translocating ATPase	26
<i>arsD</i>	arsenical metallochaperone	ArsD	Transfer trivalent metalloids to ArsA	26
<i>apqS</i>	aquaglyceroporin	AqpS	Extrude As(III) from the cell	27
<i>ACR3</i>	As(III) permease	ACR3	Extrude As(III) from the cell	19
<i>Ycf1p</i>	yeast cadmium factor protein cytosolic	Ycf1p	Sequester cytosolic As(III)/glutathione complex into vacuoles	28
<i>arsJ</i>	organoarsenical efflux permease	ArsJ	Extrude organoarsenicals from the cell	29
<i>arsP</i>	efflux system specific for trivalent organoarsenicals	ArsP	Extrude trivalent organoarsenicals from the cell	30
<i>pgpA</i>	P-glycoprotein-related protein	PgpA	Recognize and transport thiol-metal conjugates	31
<i>Unknown functions</i>				
<i>arsO</i>	putative flavin-binding monooxygenase	ArsO	Unknown function	32
<i>arsT</i>	putative thioredoxin reductase	ArsT	Unknown function	32