**QUEEN'S
UNIVERSITY
BELFAST**

Seeking the sun in deep, dark places: mesopelagic sightings of ocean sunfishes (Molidae)

Phillips, N. D., Harrod, C., Gates, A. R., Thys, T. M., & Houghton, J. D. R. (2015). Seeking the sun in deep, dark places: mesopelagic sightings of ocean sunfishes (Molidae). *Journal of the Marine Biological Association of the United Kingdom*, 87, 1118-1126. <https://doi.org/10.1111/jfb.12769>

Published in:

Journal of the Marine Biological Association of the United Kingdom

Document Version:

Peer reviewed version

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2015 Cambridge University Press.

This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

1 **Seeking the Sun in Deep, Dark Places: Mesopelagic Sightings of Ocean**

2 **Sunfishes (Family Molidae)**

3 N. D. Phillips^{*1}, C. Harrod², A. R. Gates³, T. M. Thys⁴, and J. D. R. Houghton^{1,5}

4 ¹*School of Biological Sciences, Queen's University Belfast, MBC Building, 97 Lisburn Road,*
5 *Belfast BT9 7BL; ²Instituto de Ciencias Naturales Alexander von Humboldt, Universidad de*
6 *Antofagasta, Avenida Angamos 601, Antofagasta, Chile; ³National Oceanography Centre,*
7 *University of Southampton, European Way, Southampton, SO14 3ZH, UK ; ⁴California*
8 *Academy of Science, 55 Music Concourse Drive, Golden Gate Park, San Francisco, CA 94118*
9 *USA; ⁵Institute of Global Food Security, Queen's University Belfast, 123 Stranmillis Road,*
10 *Belfast, County Antrim, BT9 5AG; *Author to whom correspondence should be addressed.*
11 *Email: nphillips01@qub.ac.uk, Tel: +4428 9097 2620.*

12
13 Evidence is presented from publicly available remotely operated vehicle (ROV) footage
14 that suggests deep water ranging in ocean sunfishes (family Molidae) is more common than
15 typically thought, including a new maximum depth recorded for the southern sunfish (*Mola*
16 *ramsayi*).

17 Keywords: Diving Behaviour, Foraging, *Mola mola*, Remotely Operated Vehicle, SERPENT
18 Project.

19 Running Head: Mesopelagic Sunfish Sightings

The family Molidae, or the ocean sunfishes, is currently believed to comprise four widely distributed species: the ocean sunfish *Mola mola*, (Linnaeus, 1758), sharptail sunfish *Masturus lanceolatus* (Liénard, 1840), southern sunfish *Mola ramsayi* (Giglioli, 1883) and slender sunfish *Ranzania laevis* (Pennant, 1776), although the taxonomy remains uncertain, with the possible existence of several currently undescribed species (Bass *et al.*, 2005; for review see Pope *et al.*, 2010). Often seen basking at the surface, there was a long held perception that these species were rare, inactive drifters feeding solely on gelatinous zooplankton (see Pope *et al.*, 2010 for review). However, an increasing number of recent studies have shown these notions to be far from accurate, with evidence of long distance migrants travelling ca. 27 km day⁻¹ (Cartamil & Lowe, 2004), displaying deep diving behaviours (Sims *et al.*, 2009; Hays *et al.*, 2009) and consuming a mixed diet (Syväranta *et al.*, 2012; Nakamura & Sato, 2014). More specifically, there is now strong evidence of cryptic benthivory in these typically pelagic fishes (Harrod *et al.*, 2013) and records of deep water forays to feed on colonial gelata such as siphonophores (Nakamura & Sato, 2014; Nakamura *et al.*, 2015). Taken together, a picture of an active migrant feeding broadly within marine food webs has emerged, but observational data to validate behavioural studies remains difficult to come by (Houghton *et al.*, 2000). Serendipitous sightings of three sunfish species are presented here, sourced from industrial and research submersibles, providing further evidence of the deep ranging capabilities of the Molidae. More broadly, these records taken together with previous ROV sightings of a hammerhead shark (*Sphyrna lewini* (Griffith & Smith, 1834); Moore, S. & Gates, A.,R. *pers. comms.*), add weight to the idea that mesopelagic environments may be of greater importance than thought previously for taxa often considered as epipelagic.

Sightings data were sourced primarily from the SEPRENT project (Jones *et al.*, 2009), which constitutes a scientific partnership using existing industrial technology. As direct observations of marine life in the deep sea are often prohibitively expensive, this collaboration enables scientists to browse video footage shot by industrial ROVs, providing rare access to life in the open ocean and sightings of much understudied species *e.g.* oarfish *Regalecus glesne* (Ascanius, 1772) (Benfield *et al.*, 2013) and deep sea squid *Grimalditeuthis bonplandi* (Verany, 1839) (Hoving *et al.*, 2013). Other data were sourced from ROV footage available online or through personal communications (see Fig. 1 and Table I).

In total, 13 anecdotal sightings of Molidae were obtained at depths of up to 550 m (Table I). Eleven animals were recorded from depths of > 160 m, with eight of these individuals exceeding 200 m. Aside from the general pattern of mid-water ranging the Australian sighting of a *M. ramsayi* at 483 m probably constitutes the deepest record for this species to date. The only previous record of depth use for this species came from the Sea of Oman (Sea & Bejgan, 2014) where an individual was caught in a trawl at 85 m. For context, the maximum depths previously recorded from sunfishes were to 844 m by *M. mola* (Potter & Howell, 2011) and 670 m by *M. lanceolatus* (Harbison & Janssen, 1987), which suggests that 483 m is unlikely to represent a maxima for *M. ramsayi* given the gross morphological similarities between these species.

Previous studies have intimated that such movements of Molidae into deeper water may be a means of locating deep or vertically migrating zooplankton prey (Cartamil & Lowe, 2004; Sims *et al.*, 2009; Dewar *et al.*, 2010; Nakamura *et al.*, 2015). Typically, such forays to depth constitute daytime bouts of V-shaped dives punctuated by periods at the surface (Cartamil & Lowe, 2004; Dewar *et al.*, 2010). Recently, Nakamura *et al.* (2015) verified that the main

function of surfacing is the recovery of body temperature, with the fish increasing heat gain from the warm surface water by physiological regulation [see Hays, (2015) for review]. The data presented here may be aligned with such behaviour given that the greatest depths of sunfish recorded from the ROV footage (Table I) occurred during the day (06:00 – 18:00) but with some diving activity at night (with max. depth of 306 m recorded at 01:30). Such daytime excursions to mesopelagic depths (albeit rarely) have also been documented in leatherback turtles, *Dermochelys coriacea* (Vandelli, 1761), with the overall assertion that individuals were speculating for vertically ascending gelatinous zooplankton, rather than feeding at depth (Houghton *et al.*, 2008). From the ROV footage (Fig. I), no comment can be made on whether these observations constituted brief (i.e. speculative), or prolonged (i.e. feeding) excursions to depth, or the behaviour prior to or following such events. However, these data can provide visual observations over a broad geographical range to support the notion that ocean sunfishes routinely dive below the epipelagic zone.

With respect to prey acquisition at depth, low levels of mesopelagic light suggest that the Molidae possess adequate visual acuity at depth to hunt mobile prey, potentially in combination with other senses such as olfaction, (for review see Hara, 1994). Visual acuity, determined from immature *M. mola*, was calculated at 3.5-4.3 cycles per degree (Kino *et al.*, 2009), similar to values recorded from some adult sharks (e.g. 3.8: *Galeus melastomus* (Rafinesque, 1810), 2.8: *Etmopterus spinax* (Linnaeus, 1758); Bozzano and Collin, 2000) and higher than those from adult cetaceans (e.g. 2.7: *Lagenorhynchus obliquidens* (Gill, 1865), 2.6: *Delphinapterus leucas* (Pallas, 1776); Murayama & Somiya, 1998). As visual acuity typically increases over the lifetime of an individual (Fritsches & Marshall, 2003), this value is likely to increase for mature sunfishes. Eye size usually reflects the importance of vision to

a species (Walls, 1942) and the large eyes of the Molidae support the suggestion by Hays *et al.* (2003) that ambient light levels at prey field depths may be important in determining foraging success in marine predators. Ocean sunfish eyeball diameter data has only been published from juveniles (TL < 1 m), with maximum eye diameter 38 mm (Cleland, 1862). This is comparatively large when considering other pelagic predators (*Caracharodon carcharias* (Linnaeus, 1758); 37.1 mm, *S. lewini*; 25.5 mm: Lisney & Collin, 2007) with the overall inference that Molidae are well adapted to foraging at depth. Moreover, as gelatinous zooplankton may be difficult to locate while looking down through the water column, diving to depth may allow such species to be silhouetted against down-welling light during re-ascent.

Alternatively, prey densities at depth may simply be sufficient to warrant exploration in low-light conditions, although the mechanisms for detection are not yet fully understood. Davenport (1988) and Davenport & Balazs (1991) suggested that the Molidae and leatherback turtles may use luminescence to help in their search for deep prey (*e.g.* pyrosomes): however as such behaviour was not observed here, it is not possible to comment further. The potential for the ROV lights themselves to attract sunfishes must also be acknowledged, given that BRUVs (Baited Remote Underwater Video Systems) routinely employ illumination to entice animals closer to the camera (*e.g.* Fitzpatrick *et al.*, 2013, De Vos *et al.*, 2014, Harvey *et al.*, 2012). This matter aside, gelata are certainly present at depth in open water and can constitute 50-80% of the integrated standing crop by volume (Angel & Pugh, 2000; Houghton *et al.*, 2008). Qualitatively, there is evidence of putative prey in the Gulf of Mexico from the SERPENT archives (corresponding geographically to the 2012 *M. mola* sighting; Table I) including the scyphomedusa *Stygiomedusa gigantean* (Browne, 1910)

113 (Benfield & Graham, 2010), *Aurelia aurita*, (Linnaeus, 1758) *narcomedusae* (*Solmissus* spp.),
114 salps and ctenophores, suggesting deep water feeding behaviour warrants further
115 investigation.

116 Within this collection of mesopelagic encounters, there was one sighting of specific
117 interest where a *M. lanceolatus* was filmed interacting with the physical structure of the rig
118 itself. Given that the Molidae can be highly parasitised (Fraser-Brunner, 1951) and have
119 been observed undertaking many behaviours suggested to reduce parasite loads (e.g.
120 attracting sea birds; Abe *et al.*, 2012, aggregating at reef cleaning stations and breaching;
121 Konow *et al.*, 2006), the rig structure may offer a hard surface to rub against. In the
122 relatively featureless open ocean, such opportunities are lacking which may explain the
123 behaviour of the 2009 *M. lanceolatus* sighting at 167 m (see Table I: Subsea 2009a). The
124 video footage here clearly shows the *M. lanceolatus* using the platform's framework to
125 scratch itself, suggesting that such structures might provide a service for sunfishes
126 throughout the water column.

127 Another sighting of specific interest was recorded in 2004 to the West of Shetland (Table
128 I). This record was not collected from great depth, but in surface waters, where a *M. mola*
129 was observed at 60°N, close to the reported northerly limit of the *M. mola* range, where
130 they are sighted rarely. This northerly boundary may be temperature constrained as *M.*
131 *mola* in the N.E. Atlantic have been recorded spending 99% of the time in water
132 temperatures between 10-19°C (Sims *et al.*, 2009). Despite its northerly location, this
133 individual was sighted in September when the average sea surface temperature off Shetland
134 is 12.7°C (World Sea Temperatures, 2015), well within the stated *M. mola* temperature

range. Consequently, it is unlikely that the scarcity of *M. mola* sightings in this area reflects thermal tolerances alone.

In conclusion, this paper provides repeated evidence of mid-water ranging in the Molidae using anecdotal sightings from publically available ROV footage. These data suggest that sunfishes are more common in the mesopelagic zone than previously thought, in line with deep water forays of other typically epipelagic species. A new depth record is also presented for the *M. ramsayi* sighted at 483 m and a rare observation of a *M. mola* above 60°N. These data show the importance of collaborations between commercial enterprises and academics, providing access to anecdotal sightings that have the potential to significantly increase the understanding of rarely encountered fish species. Such evidence also serves to highlight that the increasing realisation that the Molidae are far more complex in their ecology than previously thought.

Acknowledgements

The authors would like to thank Efetobore Egbore and Holly Escott from Subsea 7, along with Paul Tyler from the National Oceanography Centre Southampton for their contribution of sunfish sightings. Thanks also to Lawrence Eagling for proof-reading the manuscript on more than one occasion. We would like to thank the three reviewers and Associate Editor for their comments which improved this manuscript.

References

154 Abe, T., Sekiguchi, K., Onishi, H., Muramatsu, K. & Kamito, T. (2012). Observations on a
 155 school of ocean sunfish and evidence for a symbiotic cleaning association with
 156 albatrosses. *Marine Biology* **159**, 1173–1176.

157 Angel, M.V. & Pugh, P.R. (2000). Quantification of diel vertical migration by macroplankton
 158 and micronektonic taxa in the northeast Atlantic. *Hydrobiologia* **440**, 161–179.

159 Bass, A.L., Dewar, H., Thys, T., Streelman, J.T. & Karl, S.A. (2005). Evolutionary divergence
 160 among lineages of ocean sunfish family, Molidae. *Marine Biology* **414**, 148–405.

161 Benfield, M.C. & Graham, W.M. (2010). *In situ* observations of *Stygiomedusa gigantea* in the
 162 Gulf of Mexico with a review of its global distribution and habitat. *Journal of the Marine*
 163 *Biological Association of the United Kingdom* **90**, 1079–1093.

164 Benfield, M.C., Cook, S., Sharuga, S. & Valentine, M. M. (2013). Five *in situ* observations of
 165 live oarfish *Regalecus glesne* by remotely operated vehicles in the oceanic waters of
 166 the northern Gulf of Mexico. *Journal of Fish Biology* **83**, 28–38.

167 Bozzano, A.A. & Collin, S.P (2000). Retinal ganglion cell topography in elasmobranchs. *Brain*
 168 *Behavioural Evolution* **55**, 191–208.

169 Cartamil, D.P. & Lowe, C.G. (2004). Diel movement patterns of ocean sunfish *Mola mola* off
 170 southern California. *Marine Ecology Progress Series* **266**, 245–253.

171 Cleland, J. (1862). On the anatomy of the short sun-fish (*Orthorogoriscus mola*). *Natural*
 172 *History Review* **2**, 170–185.

173 Davenport, J. (1988). Do diving leatherbacks pursue glowing jelly? *British Herpetological*
174 *Society Bulletin* **24**, 20–21.

175 Davenport, J. & Balazs, G.H. (1991). “Fiery Bodies” -are pyrosomes an important component
176 of the diet of leatherback turtles? *Analysis*, 33–38.

177 De Vos, L., Gotz, A., Winker, H. & Attwood, C.G. (2014). Optimal BRUVs (baited remote
178 underwater video system) survey design for reef fish monitoring in the Stilbaai Marine
179 Protected Area. *African Journal of Marine Science* **36**, 1-10.

180 Dewar, H., Thys, T., Teo, S.L.H., Farwell, C., O’Sullivan, J., Tobayama, T., Soichi, M.,
181 Nakatsubo, T., Kondo, Y., Okada, Y., Lindsay, D.J., Hays, G.C., Walli, A., Weng, K.,
182 Streelman, J.T. & Karl, S.A. (2010). Satellite tracking the world's largest jelly predator,
183 the ocean sunfish, *Mola mola*, in the Western Pacific. *Journal of Experimental Biology*
184 *Marine Biology and Ecology* **393**, 32-42.

185 Fitzpatrick, C., McLean, D. & Harvey, E.S. (2013). Using artificial illumination to survey
186 nocturnal reef fish. *Fisheries Research* **146**, 41-50.

187 Fraser-Brunner, A. (1951). The ocean sunfishes (Family Molidae). *Bulletin of the British*
188 *Museum (Natural History) Zoology* **1**, 87–121.

189 Fritsches, K.A., & Marshall, N.J. (2003). Retinal specializations in the blue marlin: eyes
190 designed for sensitivity to low light levels. *Marine and Freshwater Research* **4**, 333 –
191 341.

192 Hara, T.J. (1994). The diversity of chemical stimulation in fish olfaction and gustation.
 193 *Reviews in Fish Biology and Fisheries* **4**, 1-35.

194 Harbison, G. R. & Janssen, J. (1987). Encounters with a swordfish (*Xiphias gladius*) and
 195 sharptail mola (*Masturus lanceolatus*) at depths greater than 600 meters. *Copeia* **2**,
 196 511–513.

197 Harrod, C., Syväranta, J., Kubicek, L., Cappanera, V. & Houghton, J.D.R. (2013). Reply to
 198 Logan & Dodge: “Stable isotopes challenge the perception of ocean sunfish *Mola mola*
 199 as obligate jellyfish predators”. *Journal of Fish Biology* **82**, 10–6.

200 Harvey, E.S., Butler, J.J., McLean, D.L. & Shand, J. (2012). Contrasting habitat use of diurnal
 201 and nocturnal fish assemblages in temperate Western Australia. *Journal of*
 202 *Experimental Marine Biology and Ecology* **426**, 78-86.

203 Hays, G.C. (2015). New insights: animal-bourne cameras and accelerometers reveal the
 204 secrets of cyrtic species. *Journal of Animal Ecology* **84**, 587-589.

205 Hays, G.C., Broderick, A.C., Godley, B.J., Luschi, P. & Nichols, W.J. (2003). Satellite telemetry
 206 suggests high levels of fishing-induced mortality in marine turtles. *Marine Ecology*
 207 *Progress Series* **262**, 305–309.

208 Hays, G.C., Farquhar, M.R., Luschi, P., Teo, S.L.H. & Thys, T.M. (2009). Vertical niche overlap
 209 by two ocean giants with similar diets: Ocean sunfish and leatherback turtles. *Journal*
 210 *of Experimental Marine Biology and Ecology* **370**, 134–143.

211 Houghton, J.D.R., Doyle, T.K., Davenport, J., Wilson, R.P. & Hays, G.C. (2008). The role of
 212 infrequent and extraordinary deep dives in leatherback turtles (*Dermochelys coriacea*).
 213 *Journal of Experimental Biology* **211**, 2566-2575.

214 Houghton, J.D.R., Woolmer, A. & Hays, G.C. (2000). Sea turtle diving and foraging behaviour
 215 around the Greek Island of Kefalonia. *Journal of the Marine Biological Association of*
 216 *the UK* **80**, 761–762.

217 Hoving, H.J.T., Zeidberg, L.D., Benfield, M.C., Bush, S.L., Robison, B.H. & Vecchione, M.
 218 (2013). First in situ observations of the deep-sea squid *Grimalditeuthis bonplandi* reveal
 219 unique use of tentacles. *Proceedings of the Royal Society B: Biological Sciences* **280**,
 220 1769.

221 Kino, M., Miayzaki, T., Iwami, T. & Kohbara, J. (2009). Retinal topography of ganglion cells in
 222 immature ocean sunfish, *Mola mola*. *Environmental Biology of Fishes* **85**, 33–38.

223 Konow, N., Fitzpatrick, R. & Barnett, A. (2006). Adult emperor angelfish (*Pomacanthus*
 224 *imperator*) clean giant sunfishes (*Mola mola*) at Nusa Lembongan, Indonesia. *Coral*
 225 *Reefs* **25**, 208.

226 Lisney, T.J. & Collin, S.P. (2007). Relative eye size in elasmobranchs. *Brain, Behavior and*
 227 *Evolution* **69**, 266-279.

228 Murayama, T. & Somiya, H. (1998). Distribution of ganglion cells and object localizing ability
 229 in the retina of three cetaceans. *Fisheries Science* **64**, 27–30.

230 Nakamura, I., Goto, Y. & Sato, K. (2015). Ocean sunfish rewarm at the surface after deep
 231 excursions to forage for siphonophores. *Journal of Animal Ecology* **83**, 1365–2656.

232 Nakamura, I. & Sato, K. (2014). Ontogenetic shift in foraging habit of ocean sunfish *Mola*
 233 *mola* from dietary and behavioral studies. *Marine Biology* **161**, 1263–1273.

234 Pope, E.C., Hays, G.C., Thys, T.M., Doyle, T.K., Sims, D.W., Queiroz, N., Hobson, V., Kubicek,
 235 L. & Houghton, J. (2010). The biology and ecology of the ocean sunfish *Mola mola*: a
 236 review of current knowledge and future research perspectives. *Reviews in Fish Biology*
 237 *and Fisheries* **20**, 471–487.

238 Potter, I.F. & Howell, W.H. (2011). Vertical movement and behavior of the ocean sunfish,
 239 *Mola mola*, in the northwest Atlantic. *Journal of Experimental Marine Biology and*
 240 *Ecology* **396**, 138–146.

241 Sea, N.O. & Bejgan, N. (2014). The first record of Southern Ocean sunfish, *Mola ramsayi*
 242 from Northern Oman Sea, Iran. *Iranian Journal of Fisheries Sciences* **13**, 242–246.

243 Sims, D.W., Queiroz, N., Doyle, T.K., Houghton, J.D.R. & Hays, G.C. (2009). Satellite tracking
 244 of the World’s largest bony fish, the ocean sunfish (*Mola mola*) in the North East
 245 Atlantic. *Journal of Experimental Marine Biology and Ecology* **370**, 127–133.

246 Subsea (2009a) Subsea 7 sunfish sighting ROV footage. Available at
 247 <https://www.youtube.com/watch?v=2iwnvXCXeeY> (last accessed 10 March 2015).

248 Subsea (2009b) Subsea 7 sunfish sighting ROV footage. Available at
 249 <https://www.youtube.com/watch?v=WNNDem-1heg> (last accessed 10 March 2015).

250 Syväranta, J., Harrod, C., Kubicek, L., Cappanera, V. & Houghton, J.D.R. (2012). Stable
251 isotopes challenge the perception of ocean sunfish *Mola mola* as obligate jellyfish
252 predators. *Journal of Fish Biology* **80**, 225–31.

253 Walls, G.L. (1942). The vertebrate eye and its adaptive radiation. *Optometry and Vision*
254 *Science* **20**, 30–32.

255

256 **Electronic References**

257 CDFW (2008). The California Department of Fish and Wildlife ROV footage. Available at
258 https://www.youtube.com/watch?v=ETbF_UTzKqs (last accessed 10 March 2015).

259

260 Cordes, E. (2009). 2002 Seward Johnson II Fall Cruise Report. Available at
261 <http://deepseanews.com/2009/06/reflections-on-the-johnson-sea-link-dr-cordes/> (last
262 accessed 10 March 2015).

263

264 Jones, D.O.B., Gates, A.R., Curry, R.A., Thomson, M., Pile, A., & Benfield, M. (Eds) (2009).
265 SERPENT project media database archive. Available
266 at <http://archive.serpentproject.com/> (last accessed 12 Mar 2015).

267

268 World Sea Temperatures: Shetland average September Sea Temperatures. Available at
269 <http://www.seatemperature.org/europe/united-kingdom/shetland-september.htm>
270 (last accessed 10 March 2015).