



**QUEEN'S
UNIVERSITY
BELFAST**

Mountain hare transcriptome and diagnostic markers as resources to monitor hybridization with European hares

Marques, J., Ferreira, M., Farelo, L., Callahan, C., Hackländer, K., Jenny, H., Montgomery, W., Reid, N., Good, J., Alves, P. C., & Melo-Ferreira, J. (2017). Mountain hare transcriptome and diagnostic markers as resources to monitor hybridization with European hares. *Scientific Data*, 4, Article 170178 .
<https://doi.org/10.1038/sdata.2017.178>

Published in:
Scientific Data

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2017 the authors.

This is an open access article published under a Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

SCIENTIFIC DATA

OPEN Data Descriptor: Mountain hare transcriptome and diagnostic markers as resources to monitor hybridization with European hares

Received: 15 February 2017

Accepted: 13 October 2017

Published: 5 December 2017

João P. Marques^{1,2}, Mafalda S. Ferreira^{1,2}, Liliana Farelo¹, Colin M. Callahan³, Klaus Hackländer⁴, Hannes Jenny⁵, W. Ian Montgomery^{6,7}, Neil Reid^{6,7}, Jeffrey M. Good³, Paulo C. Alves^{1,2} & José Melo-Ferreira^{1,2}

We report the first mountain hare (*Lepus timidus*) transcriptome, produced by *de novo* assembly of RNA-sequencing reads. Data were obtained from eight specimens sampled in two localities, Alps and Ireland. The mountain hare tends to be replaced by the invading European hare (*Lepus europaeus*) in their numerous contact zones where the species hybridize, which affects their gene pool to a yet unquantified degree. We characterize and annotate the mountain hare transcriptome, detect polymorphism in the two analysed populations and use previously published data on the European hare (three specimens, representing the European lineage of the species) to identify 4 672 putative diagnostic sites between the species. A subset of 85 random independent SNPs was successfully validated using PCR and Sanger sequencing. These valuable genomic resources can be used to design tools to assess population status and monitor hybridization between species.

Design Type(s)	individual genetic characteristics comparison design • strain comparison design • population dynamics analysis objective
Measurement Type(s)	transcription profiling assay
Technology Type(s)	RNA sequencing
Factor Type(s)	geographic location
Sample Characteristic(s)	<i>Lepus timidus</i> • liver • Republic of Ireland • Switzerland

¹CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Universidade do Porto, Vairão 4485-661, Portugal. ²Departamento de Biologia, Faculdade de Ciências do Porto, Porto 4169-007, Portugal. ³Division of Biological Sciences, University of Montana, 32 Campus Drive, Missoula, MT 59812, USA. ⁴Institute of Wildlife Biology and Game Management, BOKU-University of Natural Resources and Life Sciences, Vienna 1180, Austria. ⁵Amt für Jagd und Fischerei Graubünden, Chur 7001, Switzerland. ⁶Institute of Global Food Security, School of Biological Sciences, Queen's University Belfast, Belfast BT9 5BN, UK. ⁷School of Biological Sciences, Queen's University Belfast, 97 Lisburn Road, Belfast BT9 7BL, UK. Correspondence and requests for materials should be addressed to J.M.-F. (email: jmeloferreira@cibio.up.pt).

Background & Summary

The mountain hare (*Lepus timidus*) is an Arcto-alpine species that was the most common and widely distributed hare species across Europe during the last glacial periods¹. Nowadays, the mountain hare is distributed from Fennoscandia to Eastern Siberia, but also occurs in isolated/refuge populations (e.g., Ireland, Scotland, the Alps, Poland, the Baltics and Japan), and in places where it has been introduced (Iceland, England, Faroe Islands and New Zealand) (see Fig. 1). Even though they are a popular game species and abundant within its range, mountain hares have sharply declined in some regions, particularly in areas of contact with the European hare (*Lepus europaeus*), where the latter tends to invade and replace the range of the former^{1–4}. Mountain and European hares share extensive natural and human-induced contact zones in Western Europe, from the British Isles to Scandinavia and Central Europe (Fig. 1). Climate change is predicted to affect lagomorphs extensively^{5,6} and, in particular, to accelerate the replacement of mountain hares by European hares in the contact zones, such as the Alps, Sweden or Ireland^{7,8}. The two species may hybridize when in contact, resulting in some genetic introgression^{9–13}, with potential effects on local adaptation¹⁴.

Even though the mountain hare and other hare species have been the subject of several population genetics studies, these have been mostly based on a few markers^{10,15–17}. Therefore, permanent genomic resources provide fundamental information to develop monitoring tools to evaluate population status and implement protective policies. In this work, we use high-throughput RNA sequencing to: i) generate genomic resources for the mountain hare; and, ii) use published data on the European hare¹⁸ to pinpoint candidate fixed differences between the species that can be used to build genotyping tools to monitor gene exchange in the contact zones. We here present the first mountain hare transcriptome, and the most complete among the currently available European *Lepus* transcriptomes.

Methods

A summary of the methodological workflow is shown in the flowchart of Fig. 2.

Sampling procedure and locations

Specimens from the Alps (see Fig. 1) were sampled during regular permit hunting in Grisons, Switzerland. Specimens from Ireland (see Fig. 1) were captured from the wild in Borris-in-Ossory, by the Irish Coursing Club (ICC) for scientific research purposes under National Parks & Wildlife (NPWS) licence No. C 337/2012 issued by the Department of Arts, Heritage and the Gaeltacht (dated 31/10/2012).




Figure 1. Approximate mountain and European hare distribution. Approximate distributions of the mountain hare, *Lepus timidus*, and the European hare, *L. europaeus*, in Eurasia with indication of the areas of contact and of broad geographic overlap between the species (distribution ranges were adapted from IUCN Spatial Data Resources; IUCN 2016⁵¹). Circles indicate the mountain hare sampling locations for this work (open circle—Ireland; closed circle—Alps).

Irish hares were dispatched humanely and in accordance with the licence conditions by means of lethal injection administered by Mr William Fitzgerald, Veterinary Laboratory Service Follow (MVB MVM CertCSM), from the Department of Agriculture, Food and the Marine, Regional Veterinary Laboratory, Hebron Road, Kilkenny, R95 TX39. Total RNA was isolated from 8 individuals.

RNA extraction

Liver tissue was freshly collected, immediately preserved in RNAlater and then stored at -80°C until RNA extraction. Prior to extraction, frozen samples were ground in liquid nitrogen with a ceramic mortar and pestle. Mortar and pestle were washed prior to extraction using a 6-step wash that includes the following washing reagents in order: 70% ethanol, tap water, 10% bleach, milli-Q water, RNase away (Thermo Fisher Scientific) and finishing with molecular grade H_2O . RNA extraction was performed using RNeasy Mini Kit according to manufacturer instructions.

RNA sequencing library preparation

The SureSelect Strand-Specific RNA Library Prep for Illumina Multiplexed Sequencing (Agilent Technologies) kit was used to prepare cDNA libraries for all samples. Library sizes were estimated using a Bioanalyzer 2,100 and quantified using KAPA Library quantification kit (KAPA BIOSYSTEMS). Equal molar concentrations of each library were pooled together for sequencing.

Sequence data processing and *de novo* transcriptome assembly

A detailed description of tools and commands used in the data analysis is shown in Table 1 (available online only). A first quality evaluation of obtained sequence reads (Data Citation 1) was performed with FastQC v0.11.5¹⁹. After read quality inspection, adapters were removed and quality trimming performed using TRIMMOMATIC v0.36²⁰, with instructions to remove the first ten bases, Illumina adapters, reads below 25 bp long and bases in the ends of reads with quality below 10, and to perform a 4-base sliding window trimming and cutting fragments with an average quality below 10. Trimmed-read quality was rechecked with FastQC (Data Citation 2). A *de novo* transcriptome assembly was then performed using all properly paired reads from the eight individuals in the dataset using TRINITY v2.2.0²¹, establishing RF as read orientation for a strand-specific assembly. In addition, as a complementary resource, *de novo* transcriptome assemblies for each of the two sampling localities were also performed. Transrate v1.0.3²² was used to evaluate assembly quality and completeness and to remove possible chimeras and poorly supported contigs. Cleaned reads were mapped back to the produced assembly and only the well-supported contigs were retained (Transrate optimal cut-off >0.024). In order to remove redundancy produced by using multi-sample data to perform the assembly, all contigs were clustered using CD-HIT-EST v4.6.4²³ with a 95% similarity threshold. Open reading frames were predicted with TransDecoder v3.0.0²⁴ to remove possible contaminants such as non-coding RNA and DNA contamination. The final filtered transcriptome comprised contigs with predicted open reading frame and/or rabbit (*Oryctolagus cuniculus*) or pfam annotation. Filtered transcriptome as well as raw assemblies are available in *Figshare* (Data Citation 2).




Figure 2. Methodological workflow. Flowchart of the RNA-sequencing setup and data analysis steps. Commands used in the analytical steps shown in bold are detailed in Table 1 (available online only).

Transcriptome annotation

Transcriptome annotation was performed adapting the protocol of Trinotate v3.0.1²⁴, using i) Conditional Reciprocal Best BLAST (crb-blast) v0.6.6²⁵ against the rabbit transcriptome reference (release 86) and Swiss-Prot database²⁶; ii) protein domain identification by HMMER v3.1b2²⁷ onto the PFAM database²⁸; iii) protein signal peptide through signal v 4.1²⁹; iv) transmembrane domain prediction using tmHMM v2.0³⁰; and v) eggNOG³¹, GO³² and Kegg³³ databases annotation. Annotation information was incorporated into an xlsx database (Data Citation 2).

SNP inference

SNP calling was performed separately for mountain hares (Data Citation 1) and European hares (Data Citation 3, from Amoutzias *et al.*¹⁸). The three European hare specimens represent the European lineage of the species¹⁸. First, reads from all the individuals were mapped to the filtered mountain hare *de novo* transcriptome with bwa-mem v0.7.15³⁴ with default parameters and read group information added to each sequencing lane-sample pair. The resulting alignments were converted to a binary file (bam format), sorted and submitted to fixmate step using SAMtools v1.3.1³⁵. Duplicate reads were removed using Picard v1.140 (<http://broadinstitute.github.io/picard>) with the option MarkDuplicates. Realignment and recalibration was performed with Genome Analysis Toolkit v3.6-0³⁶. Finally, SNP call was carried out using Reads2snp v2.0.64³⁷ using a threshold of 20 for site and mapping qualities, the paralog filter, a minimum coverage of 10X and a genotype probability >0.95. The resulting VCF file was deposited in Figshare (Data Citation 2). Only SNPs represented in all sampled specimens were retained.

Differentiation, admixture and Gene Ontology enrichment analysis

A set of random 5 502 SNPs, selected from independent contigs in order to reduce the linkage probability, was identified with VCFtools v0.1.14³⁸. PGDSpyder v2.1.1.0³⁹ was used to convert this file to the required file formats. Partitions of genetic diversity in the dataset were investigated with a Principal Components Analysis, using PLINK v1.90b3.45⁴⁰ and ggplot2 R package⁴¹ to plot the results. Additionally, the data were analysed using the admixture model implemented in STRUCTURE 2.3.4⁴², with three replicate runs with 1 million steps after a burn-in period of 200 000, and K = 2. Results were plotted using CLUMPACK⁴³. Gene Ontology enrichment analyses were performed for the collection of contigs/genes with fixed differences between mountain and European hare samples, and between mountain hare sampling localities. The analysis was based on the rabbit proteome annotations and performed with g:Profiler³⁴, applying the g:SCS multiple test correction and the 'best per parent group' hierarchical filter. The background set of genes was reduced to contigs with SNP information.

Independent SNP genotyping

A random set of 110 SNPs, inferred as potentially diagnostic between *L. timidus* and *L. europaeus*, was selected for independent validation using Sanger sequencing. DNA was extracted from two of the previously analysed mountain hare samples (one Alpine, Sample_3112, and one Irish, Sample_3103) and two other European hare specimens (sampled in Clermont-Ferrand—Sample—1569—Font-Romeu, Pyrenees—Sample—1550—in France during the regular hunting season). DNA extraction was performed using JETQUICK Tissue DNA Purification kit (Genomed). PCR primers were designed to be anchored in a single exon (taking into account intron-exon boundaries from the European rabbit reference genome) and to amplify a portion of 110 independent contigs containing at least one putative diagnostic SNP. The primer sets were designed using the Scrimmer pipeline⁴⁴, which depends on Primer3⁴⁵ to design and set the primer conditions. A third internal sequencing primer was designed. PCRs were performed using QIAGEN Multiplex PCR Master Mix (Qiagen) and the following thermal cycling profile: initial denaturation at 95 °C for 15', 35 cycles of denaturation at 95 °C for 30", annealing at 60–67 °C for 20" and elongation at 72 °C for 30", and a final extension step at 72 °C for 5'. PCR products were visually inspected under UV-light after electrophoresis in agarose gels stained with GelRed (Biotium), purified with Exonuclease I (New England Biolabs) and FastAP Thermosensitive Alkaline Phosphatase (Thermo Scientific), and sequenced using internal or, in a few cases, PCR primers in a ABI 3130xl genetic analyzer.

Code availability

Analyses in this work were performed with freely available open access tools mainly using command line versions (Table 1 (available online only)). Parameters are described in the methods section and software versions and commands used are detailed in Table 1 (available online only).

Data Records

Forty-eight raw FASTQ files were submitted to *NCBI Sequence Read Archive*, with accession number SRP095715 (Data Citation 1 and Tables 2 and 3). FASTQ files were divided in two sets, corresponding to the sampling localities (Ltim_Ireland and Ltim_Alps), and by biosample-specimen (SAMN06186748-3101, SAMN06186761-3102, SAMN06186762-3103 and SAMN06186763-3105; SAMN06186727-3112, SAMN06186728-3113, SAMN06186729-3114 and SAMN06186738-3116). In each biosample, six files were submitted, corresponding to three different Illumina HiSeq sequencing lanes and two read directions. Pre/post-cleaning FASTQC base quality pdf report (FASTQC.pdf) can be accessed in Figshare (Data Citation 2). This dataset is the core of this work and has not been released or analysed previously.

Trinity raw assemblies (Ltimidus_Trinity.fasta, LtimidusIreland_Trinity.fasta and LtimidusAlps_Trinity.fasta) were deposited on *Figshare* (Data Citation 2 and Table 4). The curated transcriptome assembly fasta files (LtimidusTranscriptome.cds.fasta and LtimidusTranscriptome.pep.fasta) and the annotated database file (LtimidusTranscriptome.xlsx) can also be found in *Figshare* (Data Citation 2).

The European hare data used here (Data Citation 3) was previously published by Amoutzias *et al.*¹⁸ (NCBI Sequence Read Archive, accession number SRP055741, samples SRR1823098, SRR1863103 and SRR1863605).

Mapping statistics (Table 5), SNP call VCF file (LtimVsLeur.vcf) and population/species diagnostic SNPs tables (Supplementary Tables 1) were deposited in *Figshare* (Data Citation 2).

Technical Validation

RNA integrity

The quality and quantity of each RNA sample was assessed using the 260/280 and 260/230 absorbance ratios estimated by an IMPLEN P330 NanoPhotometer and RNA Integrity Number (RIN) and concentration ($\mu\text{g}\mu\text{l}^{-1}$) with a Bioanalyzer 2,100 (Agilent Technologies). All samples had RIN values above 8.

RNA-Seq data quality

The Illumina HiSeq run produced a total raw output of 103 941 215 100 bp paired-end reads (207 882 430 total reads). Adapter removal and quality trimming decreased this number to 201 569 448 reads (97%) (Table 4). Final analysed reads passed the minimum quality parameters as established by FastQC.

Transcriptome assembly curation, annotation and quality

Cleaned reads were assembled into 272 183 contigs with a mean length of 594 bp and a N50 length of 839 bp (Table 4). After assembly curation with Transrate optimal cut-off >0.024 , clustering with a 95% similarity threshold and open reading frame prediction, were retained 25 868 transcripts with a mean length of 842 bp and a N50 length of 1 182 (Table 4).

Annotation using a conditional reciprocal best blast hit approach results in 16 772 (65%) annotated transcripts, of which 13 641 were annotated to the rabbit transcriptome and 15 955 to the Swiss-Prot database (Fig. 3). In order to reduce the number of non-annotated transcripts, the less stringent unidirectional blast hit was added to the database. Hits were recovered for 25 549 transcripts (99%) (Fig. 3).

The mountain hare transcriptome produced in this study represents an important improvement compared to the currently available transcriptomic resources for European *Lepus*—*L. granatensis*⁴⁶ and *L. europaeus*¹⁸ transcriptomes—as it performs better on several assembly statistics, such as reference coverage (42 versus 32% in *L. granatensis* and 40% in *L. europaeus*; using the rabbit transcriptome as reference).

Genetic variation, differentiation and gene ontology enrichment

In total, 218 057 526 reads (63%) were mapped to the filtered transcriptome—136 511 846 mountain hare reads (68%) and 81 545 680 European hare reads (57%) (see statistics in Table 5). After filtering, 159 629 high-quality SNPs were inferred, of which 41 182 (26%) were sequenced in all eleven specimens. A summary of polymorphic, shared and fixed SNPs is shown in Fig. 3. 4 672 putative species-diagnostic SNPs (considered when species presented alternative fixed alleles) were inferred (Data Citation 2, Supplementary Tables 1, also deposited in *Figshare*). The diagnostic power of our SNP set could be strongly reduced if any of the sequenced specimens was admixed (namely from the Alps, where the species overlap). We therefore conducted a Principal Component Analysis and a Bayesian Assignment analysis to assess our ability to separate the species. The results suggest that the analysed mountain and European hares are well differentiated with our SNP set, and only possible limited levels of admixture were found for Sample—3116 (Fig. 4). An extra table of putative species-diagnostic SNPs excluding that individual was therefore produced (Data Citation 2, Supplementary Table 4, also deposited in *Figshare*). 25 269 SNPs were inferred in the mountain hare, of which 12 548 and 18 591 were polymorphic in the

Sample ID	Species (population)	Tissue	Method	NCBI BioSample ID
Sample_3101	<i>Lepus timidus</i> (Ireland)	liver	RNA-seq	SAMN06186748
Sample_3102	<i>Lepus timidus</i> (Ireland)	liver	RNA-seq	SAMN06186761
Sample_3103	<i>Lepus timidus</i> (Ireland)	liver	RNA-seq	SAMN06186762
Sample_3105	<i>Lepus timidus</i> (Ireland)	liver	RNA-seq	SAMN06186763
Sample_3112	<i>Lepus timidus</i> (Alps)	liver	RNA-seq	SAMN06186727
Sample_3113	<i>Lepus timidus</i> (Alps)	liver	RNA-seq	SAMN06186728
Sample_3114	<i>Lepus timidus</i> (Alps)	liver	RNA-seq	SAMN06186729
Sample_3116	<i>Lepus timidus</i> (Alps)	liver	RNA-seq	SAMN06186738

Table 2. Summary of sample data information deposited in the NCBI database.

Sample ID	NCBI SRA runs accession	Raw reads	Mbytes
Sample_3101	SRR5133282	26,598,712	2,525
Sample_3102	SRR5133280	26,128,525	2,532
Sample_3103	SRR5133285	24,469,456	2,414
Sample_3105	SRR5133283	26,662,182	2,582
Sample_3112	SRR5133287	22,444,667	2,263
Sample_3113	SRR5133281	20,825,930	2,100
Sample_3114	SRR5133286	32,749,011	3,294
Sample_3116	SRR5133284	21,690,965	2,189

Table 3. Illumina RNA-seq data deposited in the NCBI database.

<i>Lepus timidus</i> transcriptome	Value
Raw Reads	207,882,430
Clean Reads	201,569,448
Mapped Reads	136,511,846
Raw <i>de novo</i> assembly (Trinity)	
Number of contigs	272,183
Largest (bp)	14,048
Smallest (bp)	201
N50 (bp)	839
Mean (bp)	594
Post assembly curation (TransRate)	
Number of contigs	113,694
Largest (bp)	14,048
Smallest (bp)	201
N50 (bp)	801
Mean (bp)	567
Post redundancy removal (CD-HIT-EST)	
Number of contigs	109,239
Largest (bp)	14,048
Smallest (bp)	201
N50 (bp)	765
Mean (bp)	554
Post open reading frame prediction (TransDecoder)	
Number of contigs	25,868
Largest (bp)	13,728
Smallest (bp)	297
N50 (bp)	1,182
Mean (bp)	842
Reference Coverage (%)	42

Table 4. Mountain hare transcriptome assembly and curation statistics.

Irish and Alpine samples respectively, and 126 were fixed between sampling localities (Data Citation 2, Supplementary Tables 5, deposited in *Figshare*). The ‘membrane part’ gene ontology term was found enriched in the collection of genes with fixed differences between the Irish and Alpine mountain hare samples, while terms ‘lipid metabolic process’, ‘small molecule catabolic process’, ‘extracellular space and acyl-CoA dehydrogenase activity’ were found enriched in genes with fixed differences between samples of the two species. Note however that even though the background gene set was controlled for, RNA-sequencing data does not provide an unbiased sample of information across different genes and these results may represent tissue-related functions.




Figure 4. Characterization of inferred SNPs in the sampled populations and species. (a) Relative proportion of the 41 182 SNPs mapped to the mountain hare transcriptome, summarized as polymorphic within each species and fixed or shared between *L. timidus* (mountain hare) and *L. europaeus* (European hare). The proportion is shown considering the complete *L. timidus* dataset (i) and only the Irish (ii) and Alpine (iii) populations. (b) STRUCTURE analysis to evaluate cluster membership and admixture proportions. Individuals are sorted by population and species. Mountain hare populations are shown in blue and European hare individuals in orange. (c) Principal Component Analysis (PCA) plot using one SNP per contig. The first principal component (PC1) splits species and the second (PC2) the sampled populations.

Usage Notes

These genomic resources (which greatly extend previously available marker sets; e.g.⁵⁰) will be useful for a variety of studies, particularly in the characterization of genetic diversity in mountain hare populations and on the development of hybridization monitoring tools. Note that SNPs were here inferred from an uneven and small species sample, and therefore any diagnostic genotyping assay built from this data

should be first tested with adequate sample sizes from pure parental populations of the species, before being applied to hybrid zones.

References

1. Thulin, C. G. The distribution of mountain hares *Lepus timidus* in Europe: A challenge from brown hares *L. europaeus*? *Mamm. Rev.* **33**, 29–42 (2003).
2. Reid, N. & Montgomery, W. I. Is naturalisation of the brown hare in Ireland a threat to the endemic Irish hare? *Biol. Environ.* **107**, 129–138 (2007).
3. Reid, N. European hare (*Lepus europaeus*) invasion ecology: Implication for the conservation of the endemic Irish hare (*Lepus timidus hibernicus*). *Biol. Invasions* **13**, 559–569 (2011).
4. Caravaggi, A., Montgomery, W. I. & Reid, N. Range expansion and comparative habitat use of insular, congeneric lagomorphs: invasive European hares *Lepus europaeus* and endemic Irish hares *Lepus timidus hibernicus*. *Biol. Invasions* **17**, 687–698 (2015).
5. Leach, K., Kelly, R., Cameron, A., Montgomery, W. I. & Reid, N. Expertly validated models and phylogenetically-controlled analysis suggests responses to climate change are related to species traits in the order Lagomorpha. *PLoS ONE* **10**, e0122267 (2015).
6. Leach, K., Montgomery, W. I. & Reid, N. Biogeography, macroecology and species' traits mediate competitive interactions in the order Lagomorpha. *Mammal Review* **45**, 88–102 (2015).
7. Acevedo, P., Jiménez-Valverde, A., Melo-Ferreira, J., Real, R. & Alves, P. C. Parapatric species and the implications for climate change studies: A case study on hares in Europe. *Glob. Chang. Biol.* **18**, 1509–1519 (2012).
8. Caravaggi, A. *et al.* Niche overlap of mountain hare subspecies and the vulnerability of their ranges to invasion by the European hare; the (bad) luck of the Irish. *Biol. Invasions* 1–20, doi:10.1007/s10530-016-1330-z (2016).
9. Thulin, C. G., Jaarola, M. & Tegelstrom, H. The occurrence of mountain hare mitochondrial DNA in wild brown hares. *Mol. Ecol.* **6**, 463–467 (1997).
10. Thulin, C. G., Fang, M. & Averianov, A. O. Introgression from *Lepus europaeus* to *L. timidus* in Russia revealed by mitochondrial single nucleotide polymorphisms and nuclear microsatellites. *Hereditas* **143**, 68–76 (2006).
11. Suchentrunk, F. *et al.* Introgressive hybridization in wild living mountain hares (*L. timidus varronis*) and brown hares (*L. europaeus*) and morphological consequences. *Mamm. Biol.* **70**, 39–40 (2005).
12. Melo-Ferreira, J., Alves, P. C., Freitas, H., Ferrand, N. & Boursot, P. The genomic legacy from the extinct *Lepus timidus* to the three hare species of Iberia: Contrast between mtDNA, sex chromosomes and autosomes. *Mol. Ecol.* **18**, 2643–2658 (2009).
13. Zachos, F. E., Ben Slimen, H., Hackländer, K., Giacometti, M. & Suchentrunk, F. Regional genetic in situ differentiation despite phylogenetic heterogeneity in Alpine mountain hares. *J. Zool.* **282**, 47–53 (2010).
14. Hughes, M., Reid, N., Montgomery, I. & Prodoehl, P. Verification of hybridisation between introduced European and native Irish hares. *North. Irel. Environ. Agency Res. Dev. Ser. Irel. Environ. Agency Res. Dev. Ser.* 11–11 (2011).
15. Hamill, R. M., Doyle, D. & Duke, E. J. Spatial patterns of genetic diversity across European subspecies of the mountain hare, *Lepus timidus* L. *Heredity* **97**, 355–365 (2006).
16. Melo-Ferreira, J. *et al.* Recurrent introgression of mitochondrial DNA among hares (*Lepus* spp.) revealed by species-tree inference and coalescent simulations. *Syst. Biol.* **61**, 367–381 (2012).
17. Melo-Ferreira, J. *et al.* Home-loving boreal hare mitochondria survived several invasions in Iberia: the relative roles of recurrent hybridisation and allele surfing. *Heredity* **112**, 265–273 (2014).
18. Amoutzias, G. D. *et al.* SNP identification through transcriptome analysis of the European brown hare (*Lepus europaeus*): Cellular energetics and mother's curse. *PLoS ONE* **11**, e0159939 (2016).
19. Andrews S. FastQC: a quality control tool for high throughput sequence data. Available online at: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>. 2010.
20. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible read trimming tool for Illumina NGS data. *Bioinformatics* **30**, 2114–2120 (2014).
21. Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).
22. Smith-Unna, R., Boursnell, C., Patro, R., Hibberd, J. M. & Kelly, S. TransRate: Reference-free quality assessment of de novo transcriptome assemblies. *Genome Res.* **26**, 1134–1144 (2016).
23. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: Accelerated for clustering the next-generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).
24. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494–1512 (2013).
25. Aubry, S., Kelly, S., Kumpers, B. M. C., Smith-Unna, R. D. & Hibberd, J. M. Deep Evolutionary Comparison of Gene Expression Identifies Parallel Recruitment of Trans-Factors in Two Independent Origins of C4 Photosynthesis. *PLoS Genet.* **10** (6) e1004365 (2014).
26. Boutet, E. *et al.* UniProtKB/Swiss-Prot, the Manually Annotated Section of the UniProt KnowledgeBase: How to Use the Entry View. *Methods Mol. Biol.* **1374**, 23–54 (2016).
27. Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: Interactive sequence similarity searching. *Nucleic Acids Res.* **39**, W29–W37 (2011).
28. Finn, R. D. *et al.* Pfam: The protein families database. *Nucleic Acids Research* **42**, D1, D222–D230, (2014).
29. Petersen, T. N., Brunak, S., von Heijne, G. & Nielsen, H. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat. Methods* **8**, 785–786 (2011).
30. Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E. L. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* **305**, 567–580 (2001).
31. Powell, S. *et al.* eggNOG v3.0: Orthologous groups covering 1133 organisms at 41 different taxonomic ranges. *Nucleic Acids Res.* **40** (2012).
32. Gene Ontology Consortium. The Gene Ontology Consortium. Gene ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25–29 (2000).
33. Kanehisa, M., Goto, S., Sato, Y., Furumichi, M. & Tanabe, M. KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Res.* **40**, D109–D114 (2012).
34. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**, 589–595 (2010).
35. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).
36. McKenna, A. H. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).
37. Gayral, P. *et al.* Reference-Free Population Genomics from Next-Generation Transcriptome Data and the Vertebrate-Invertebrate Gap. *PLoS Genet.* **9**, e1003457 (2013).
38. Danecek, P. *et al.* The variant call format and VCF tools. *Bioinformatics* **27**, 2156–2158 (2011).

39. Lischer, H. E. L. & Excoffier, L. PGDSpider: An automated data conversion tool for connecting population genetics and genomics programs. *Bioinformatics* **28**, 298–299 (2012).
40. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015).
41. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York, 2009.
42. Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using multilocus genotype data. *Genetics* **155**, 945–959 (2000).
43. Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A. & Mayrose, I. Clumpak: A program for identifying clustering modes and packaging population structure inferences across K. *Mol. Ecol. Resour.* **15**, 1179–1191 (2015).
44. Mořkovský, L., Pačes, J., Řídl, J. & Reifová, R. Scrimer: Designing primers from transcriptome data. *Mol. Ecol. Resour.* **15**, 1415–1420 (2015).
45. Untergasser, A. *et al.* Primer3-new capabilities and interfaces. *Nucleic Acids Res.* **40**, e115 (2012).
46. Marques, J. P. *et al.* Range expansion underlies historical introgressive hybridization in the Iberian hare. *Sci. Rep.* **7**, 40788 (2017).
47. Luo, X. *et al.* Comparative transcriptomics uncovers alternative splicing and molecular marker development in radish (*Raphanus sativus* L.). *BMC Genomics* **18**, 505 (2017).
48. Yu, Y. *et al.* SNP discovery in the transcriptome of white pacific shrimp *Litopenaeus vannamei* by next generation sequencing. *PLoS ONE* **9**, e87218 (2014).
49. Cánovas, A., Rincon, G., Islas-Trejo, A., Wickramasinghe, S. & Medrano, J. F. SNP discovery in the bovine milk transcriptome using RNA-Seq technology. *Mamm. Genome* **21**, 592–598 (2010).
50. Beugin, M.-P. *et al.* A single multiplex of twelve microsatellite markers for the simultaneous study of the brown hare (*Lepus europaeus*) and the mountain hare (*Lepus timidus*). *Ecol. Evol.* **7**, 3931–3939 (2017).
51. IUCN. IUCN Red List of Threatened Species. Version 2016-1 www.iucnredlist.org (2016).

Data Citations

1. NCBI Sequence Read Archive SRP095715 (2017).
2. Marques, J. P. *et al.* Figshare <http://dx.doi.org/10.6084/m9.figshare.c.3682042> (2017).
3. NCBI Sequence Read Archive SRP055741 (2016).

Acknowledgements

This work was funded by project HybridAdapt with reference FCT-ANR/BIA-EVF/0250/2012, supported by Portuguese National Funds through the Fundação para a Ciência e a Tecnologia, FCT. Additional support was obtained from POPH-QREN funds, from the European Social Fund and FCT (IF/00033/2014/CP1256/CT0005 FCT Investigator grant to JM-F, and SFRH/BD/115089/2016 and PD/BD/108131/2015 PhD grants to JPM and MSF respectively). Instrumentation, laboratory and computational support was provided by CIBIO NEW-GEN sequencing platform, supported by European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement no. 286431. Laboratory work at the University of Montana was performed in the UM Genomics Core, supported with a grant from the M.J. Murdock Charitable Trust. Transcriptome sequencing was performed through the Vincent J. Coates Genomics Sequencing Laboratory at University of California Berkeley, supported by NIH S10 Instrumentation Grants S10RR029668 and S10RR027303. Support was additionally obtained from FLAD (Luso-American Foundation) travel grant to MSF. We are grateful to René Gadiant and local hunters in Grisons for their tremendous effort in collecting samples under Alpine conditions. Laura Claffey issued the licence to take animals for scientific purposes in Ireland (dated 31/10/2012) from the Wildlife Licensing Unit, National Parks & Wildlife Service (NPWS), Department of Arts, Heritage & the Gaeltacht, and Mr Jimi Conroy, the local NPWS Conservation Officer consented to sampling within his jurisdiction. We thank the Irish Coursing Club (ICC) for providing hares, and in particular Mr D.J. Histon, Chief Executive and Secretary, for supporting the research. Thanks also to Mr William Fitzgerald, Department of Agriculture, Food and the Marine for dispatching animals in a humane manner.

Author Contributions

J.P.M. and J.M.-F. conceived the study. J.M.-F., P.C.A., N.R., W.I.M., K.H. and H.J. organized and performed the sampling. C.C. and M.S.F. performed laboratory work at the University of Montana under the supervision of J.M.G. L.F. performed laboratory work at CIBIO under the supervision of J.M.-F. J.P.M. analysed the data. J.P.M. and J.M.-F. wrote the paper with contributions from M.S.F. and L.F. All authors read, revised and approved the manuscript.

Additional Information

Supplementary Information accompanies this paper at <http://www.nature.com/sdata>

Competing interests: The authors declare no competing financial interests.

How to cite this article: Marques, J. P. *et al.* Mountain hare transcriptome and diagnostic markers as resources to monitor hybridization with European hares. *Sci. Data* **4**:170178 doi: 10.1038/sdata.2017.178 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party

material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

The Creative Commons Public Domain Dedication waiver <http://creativecommons.org/publicdomain/zero/1.0/> applies to the metadata files made available in this article.

© The Author(s) 2017