**QUEEN'S
UNIVERSITY
BELFAST**

The structures of 1-(2-halo-6-fluorophenylmethyl)-1-methylimidazolium bromide salts

Lorenzini, F., Marr, A. C., Saunders, G. C., & Thomas, H. (2018). The structures of 1-(2-halo-6-fluorophenylmethyl)-1-methylimidazolium bromide salts. *Journal of Fluorine Chemistry*.
<https://doi.org/10.1016/j.jfluchem.2018.03.006>

Published in:
Journal of Fluorine Chemistry

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2018 Elsevier.

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

**The structures of 1-(2-halo-6-fluorophenylmethyl)-1-methylimidazolium bromide
salts**

Fabio Lorenzini^a, Andrew C. Marr^a, Graham C. Saunders^{*,b} and Hayden Thomas^b

^a *School of Chemistry and Chemical Engineering, Queen's University Belfast, BT9
5AG, United Kingdom*

^b *School of Science, University of Waikato, Hamilton, 3240, New Zealand*

* To whom correspondence should be addressed. E-mail: g.saunders@waikato.ac.nz

Abstract -

The crystal structures of 1-(2-halo-6-fluorophenylmethyl)-1-methylimidazolium bromide salts and 1-(2-trifluoromethyl-6-fluorophenylmethyl)-1-methylimidazolium bromide [MeNC₃H₃NCH₂C₆H₃X-2-F-6]⁺.Br⁻ [X = F (**1**), Cl (**2**), Br (**3**), I (**4**), CF₃ (**5**)] have been determined. The crystal structure of the hydrate of salt **1** contains π - π stacked imidazolium $\cdots$ difluorophenyl $\cdots$ difluorophenyl $\cdots$ imidazolium units. Those of salts **2**, **4** and **5** possess bromide $\cdots$ halofluorophenyl $\cdots$ halofluorophenyl $\cdots$ bromide motifs comprising anion- π and π - π stacking interactions. That of salt **4** also contains bromide $\cdots$ iodide halogen bonding. The crystal structure of the sesquihydrate of salt **3** possesses bromine $\cdots$ halofluorophenyl $\cdots$ halofluorophenyl $\cdots$ bromine motifs.

Keywords

Imidazolium salt; hydrogen bonding; anion- π interaction; π - π stacking; halogen bonding; crystal structure; DFT calculations

1. Introduction

The crystal structures of fluoroaryl-substituted imidazolium salts are a valuable source of information on a range of non-covalent interactions, in particular, π - π stacking [1-5] and anion- π interactions [2,3,5-8], arising from the presence of fluorine atoms, along with charge-assisted hydrogen bonding [3,9]. These interactions have importance in crystal engineering and through the judicious choice of fluoroaryl substituents polar crystal structures can be realized [1-5]. Furthermore, the incorporation of other halogen atoms into the polyfluoroaryl group provides the opportunity for halogen bonding when the anion is a halide [5,6].

Control of interionic and intermolecular interactions is fundamental to crystal engineering [10]. It is expected that non-covalent interactions can be turned on or off for imidazolium halide salts by variation of the substituent aryl group. As an example, although the structures of the related cations of 1-pentafluorophenylmethyl-3-methylimidazolium bromide (ROTMEU) [7] and 1-(2-bromophenylmethyl)-3-methylimidazolium bromide (EVACIO) [11] adopt similar conformations, because of the difference in the electronic nature of the phenyl ring the crystal structure of the former possesses an anion- π interaction [2,3,5-8,12], whereas that of the latter does not (Figure 1). In order to further probe these structural features and non-covalent interactions we are investigating fluoroaryl-substituted imidazolium salts. Here we report the results of our structural study, supported by density functional theory calculations, into 1-phenylmethyl-3-methylimidazolium bromide salts with a fluorine atom and a halogen atom or a trifluoromethyl group occupying the *ortho* positions of the phenyl ring.

Figure 1. Overlay of the structures of the cations of 1-(pentafluorophenylmethyl)-3-methylimidazolium bromide (red) and 1-(2-bromophenylmethyl)-3-methylimidazolium bromide (cyan) with the bromide anion closest to the face of the phenyl ring.

2. Results and discussion

The 2,6-disubstituted phenylmethylimidazolium bromide salts $[\text{MeNC}_3\text{H}_3\text{NCH}_2\text{C}_6\text{H}_3\text{X-2-F-6}]^+.\text{Br}^-$ [$\text{X} = \text{F}$ (**1**), Cl (**2**), Br (**3**), I (**4**), CF_3 (**5**)] were

prepared by treatment of 1-methylimidazole with the appropriate benzyl bromide. The salts were characterized by mass spectrometry and ^1H , ^{13}C and ^{19}F NMR spectroscopy. The salts crystallized with one ion pair in the asymmetric unit, with that of **1** also containing one molecule of water, and that of **3** also containing one and a half molecules of water (the oxygen atom of the half water molecule lies on a crystallographic two-fold rotation axis). Crystal data are given in Table 1 and selected distances and angles are given in Table 2. The structures of the cations and positions of the nearest bromide anions and water molecules are shown in Figures 2–6. The bond distances and angles are similar to those calculated using the $\omega\text{B97X-D}$ functional [13] with the 6-311++G(2d,2p) basis set for the isolated cations in the gas phase.

Figure 2. The structure of the cation of 1-(2,6-difluorophenylmethyl)-3-methylimidazolium bromide (**1**) indicating the positions of the bromide anions and water molecules close to C(1), C(2) and C(3). Thermal ellipsoids are at the 50% level. Hydrogen atoms are omitted for clarity.

Figure 3. The structure of the cation of 1-(2-chloro-6-fluorophenylmethyl)-3-methylimidazolium bromide (**2**) indicating the positions of the bromide anions close to C(1), C(2) and C(3). Thermal ellipsoids are at the 50% level. Hydrogen atoms are omitted for clarity.

Figure 4. The structure of the cation of 1-(2-bromo-6-fluorophenylmethyl)-3-methylimidazolium bromide (**3**) indicating the positions of the bromide anions and water molecules close to C(1), C(2) and C(3), and a $[(\text{Br}^-)_2 \cdot 2\text{H}_2\text{O}]$ rhomboid. Thermal ellipsoids are at the 50% level. Hydrogen atoms are omitted for clarity.

Figure 5. The structure of the cation of 1-(2-iodo-6-fluorophenylmethyl)-3-methylimidazolium bromide (**4**) indicating the positions of the bromide anions close to C(1), C(2), C(3) and I. Thermal ellipsoids are at the 50% level. Hydrogen atoms are omitted for clarity.

planes of the of the imidazolium rings of **2**, **4** and **5**, which adopt almost identical conformations (Figure 7b), are parallel to the C(5)—C(6) bond with the N₂C—H bond *syn* to the phenyl ring. This conformation is similar to that of 1-pentafluorophenylmethyl-3-methylimidazolium bromide [7], with the exception that in this salt the N₂C—H bond is *anti* to the phenyl ring.

Figure 7. Overlay of the structures of the cations of (a) 1-(2,6-difluorophenylmethyl)-3-methylimidazolium bromide (**1**) (red) and of 1-(2-bromo-6-fluorophenylmethyl)-3-methylimidazolium bromide (**3**) (pink) indicating the positions of the closest bromide anions and the oxygen atoms of the water molecules (dark blue for **1**, light blue for **3**), and (b) 1-(2,6-chlorofluorophenylmethyl)-3-methylimidazolium bromide (**2**) (yellow), 1-(2,6-iodofluorophenylmethyl)-3-methylimidazolium bromide (**4**) (purple) and of 1-(2,6-difluorophenylmethyl)-3-methylimidazolium bromide (**1**) (red).

1-(2-trifluoromethyl-6-fluorophenylmethyl)-3-methylimidazolium bromide (**5**) (green) indicating the positions of the bromide anions closest to C(1).

Consistent with other imidazolium bromide salts [3,9] all five crystal structures possess bromide anions lying close to both the plane of the imidazolium ring and C(1) (Table 3). The C(1)⋯Br⁻ distances are considerably less than the sum of the van der Waals' radius of carbon (1.70 Å [14]) and the corrected value of the van der Waals radius for bromide (2.35 Å [15]), and the N–C(1)⋯Br⁻ angles are between 100 and 150° suggestive of charge-assisted hydrogen-bonding [16,17] between H(1) and the bromide anion. The interactions between the cations and this bromide anion were calculated to be attractive by 338 to 372 kJ mol⁻¹ (Table 3), which is consistent with other imidazolium bromide salts [3,9]. The magnitudes of the interactions for **2**, **4** and **5** in which the bromide anion lies above one face of the fluorophenyl ring, suggestive of an anion– π interaction [18,19], are more than 10 kJ mol⁻¹ higher than those of **1** and **3**. Consistent with this observation, the interactions between bromide anion and 2-chloro-6-fluorotoluene, 2-iodo-6-fluorotoluene and 2-trifluoromethyl-6-fluorotoluene in the same relative positions as the respective imidazolium salts were calculated to be attractive by 11, 21 and 12 kJ mol⁻¹ respectively. The anion– π interaction between bromide and hexafluorobenzene has been calculated to be -38 kJ mol⁻¹ when the bromide anion is 3.37 Å along the normal to the ring's centroid [19]. The lower values for **2**, **4** and **5** are presumably a consequence of the less electron poor character of the arene and the non-optimal position of the bromide anion (Table 2); it is further from the

ring's plane and situated on the normal to the ring carbon attached to the most electron-withdrawing substituent, C(7) for **2** and **4**, and C(11) for **5**.

All the crystal structures also possess bromide anions lying close to the plane of the imidazolium ring and close to C(2) with geometric parameters (Table 3) that also suggest charge-assisted hydrogen-bonding [16,17] between H(2) and the bromide anion. Although the C(2)⋯Br⁻ distances are similar for all the salts, the bromide anion is displaced away from N(1) in **1** and **3** because of an interaction with a water molecule which is close to C(3) (Figures 1 and 3). The N(1)—C(1)⋯Br⁻ angles are ca. 40° greater and the N(2)—C(1)⋯Br⁻ angles are ca. 50° smaller respectively for **1** and **3** compared to **2**, **4** and **5**. The interactions between the cations and this bromide anion were calculated to be attractive by ca. 300 kJ mol⁻¹ (Table 3), which is consistent with other imidazolium bromide salts [3,5,9]. The magnitude of the interactions is smaller for **1** and **3**, which is presumed to be a consequence of the positions of the bromide anions being less favourable for hydrogen bonding. In both these structures this bromide anion lies approximately on the normal to F(7), ca. 4.5 Å from the plane the halophenyl ring, and over 5 Å from the centroid, and consequently there is no significant anion- π interaction (Table 3).

The crystal structures of **2**, **4** and **5** also possesses bromide anions close to C(3) with geometric parameters (Table 3) that also suggest charge-assisted hydrogen-bonding [16,17]. The interactions between the cations and the anion in this position were calculated to be attractive by ca. 300 kJ mol (Table 3), which is ca. 10% lower than the values calculated for 1-(4-chloro-2,3,5,6-tetrafluorophenylmethyl)-3-benzylimidazolium bromide [5]. In contrast, there are two molecules of water close to C(3) in the structures of **1** and **3**. The

C(3)⋯O distances, which are close to the sum of the van der Waals radii of carbon and oxygen (3.22 Å [14]), and the N(2)—C(3)⋯O and C(2)—C(3)⋯O angles suggest hydrogen bonding between H(3) and the water molecules. The interactions were calculated to be attractive by ca. 15 kJ mol⁻¹ (Table 3). Pairs of water molecules are linked by hydrogen bonding to two bromide anions forming [(Br⁻)₂.(H₂O)₂] rhomboids, which are a common motif in bromide hydrates [3,20]. For each structure the centre of the rhomboid lies on a crystallographic inversion centre. In the structure of **3** the rhombi are linked into chains parallel to the *a* axis by hydrogen bonding to other water molecules, the oxygen atoms of which lie on crystallographic two-fold rotation axes (O⋯O 2.815(4) Å, O⋯O⋯O 97.2(1)°).

The structure of **4** possesses a bromide anion close to the iodine atom. The I⋯Br distance, which is ca. 1 Å less than the sum of van der Waals' radius of iodine (1.98 Å [14]) and the corrected value of the van der Waals radius for bromide (2.35 Å [15]), and almost linear C—I⋯Br angle (Table 3) are indicative of halogen bonding [21]. The I⋯Br distance is consistent with those between bromide anions and polyfluoroiodobenzenes (3.02 to 3.64 Å) [22]. The energy of this interaction is similar to that for 1-(4-iodo-2,3,5,6-tetrafluorobenzyl)-3-benzylimidazolium bromide [5]. This is most likely a coincidence, since the electrostatic interaction between the ions is likely to be stronger for **4**, for which the distance between the anion and the centre of positive charge, the midpoint of N(1) and N(2), is 7.317 Å compared to 10.399 Å for 1-(4-iodo-2,3,5,6-tetrafluorobenzyl)-3-benzylimidazolium bromide, whilst the interaction between the bromide anion and iodoarene is expected to be weaker for the less electron poor arene of **4**.

In all the crystal structures there are pairs of adjacent cations related by a centre of inversion and with their halofluorophenyl rings parallel and in close proximity. For **1** each difluorophenyl ring lies close to and almost parallel to an imidazolium ring of another adjacent cation forming motifs of apparently π - π stacked imidazolium $\cdots$ difluorophenyl $\cdots$ difluorophenyl $\cdots$ imidazolium units (Figure 8). To investigate the interaction between two the difluorophenyl groups a study DFT of a pair of 1-(2,6-difluorophenylmethyl)imidazole molecules and a pair of 2,6-difluorotoluene molecules using the experimentally determined positions of the relevant atoms of **1** were undertaken. The interactions between the two 1-(2,6-difluorophenylmethyl)imidazole molecules and the two 2,6-difluorotoluene molecules were calculated to be attractive by 55 and 19 kJ mol⁻¹ respectively. The difference indicates a significant interaction between the imidazole and the phenyl rings. For **2**, **4** and **5** there are bromide $\cdots$ halofluorophenyl $\cdots$ halofluorophenyl $\cdots$ bromide motifs comprising anion- π and π - π stacking interactions (that of **2** is shown in Figure 9). Similar arrangements involving metal complexes have been reported previously [23]. The interactions between the two 1-(2-halo-6-fluorophenylmethyl)imidazole molecules and the two 2-halo-6-fluorotoluene molecules using experimentally determined positions of the relevant atoms were calculated to be attractive by ca. 35 kJ mol⁻¹ (Table 3). There is little difference between the values calculated for the two different molecules because, unlike for **1**, the imidazole rings are orientated away from the other molecule (Figure 9). The larger magnitude of the interactions between the 2-halo-6-fluorotoluene molecules compared to that of **1** is consistent with the shorter distance between the planes and centroids of the aryl rings. For **3** there are bromine $\cdots$ halofluorophenyl $\cdots$ halofluorophenyl $\cdots$ bromine motifs with the covalently

bonded bromine atoms directed at, and close to, the centroids of the bromofluorophenyl rings of the central cations, suggesting lone pair- π interactions (Figure 10). The π - π stacking interactions between 1-(2-bromo-6-fluorophenylmethyl)imidazole molecules and 2-bromo-6-fluorotoluene molecules with parallel aryl rings using experimentally determined positions of the relevant atoms were calculated to be attractive by ca. 40 kJ mol⁻¹ (Table 3). The larger magnitude is consistent with the aryl rings of **3** being closer than those of the other salts. The Br $\cdots$ C₆(plane) distance and C-Br $\cdots$ C₆(centroid) angle (Table 3) are consistent with a lone pair- π interaction in which the bromine atom acts as the acceptor [24]. This interaction between two 2-bromo-6-fluorotoluene molecules using experimentally determined positions of the relevant atoms was calculated to be attractive by 12 kJ mol⁻¹.

Figure 8. The stacking of the cations of 1-(2,6-difluorophenylmethyl)-3-methylimidazolium bromide (**1**). Thermal ellipsoids are at the 50% level. Hydrogen atoms are omitted for clarity.

Figure 9. The stacking of the cations of 1-(2-chloro-6-fluorophenylmethyl)-3-methylimidazolium bromide (**2**). Thermal ellipsoids are at the 50% level. Hydrogen atoms are omitted for clarity.

Figure **10**. The stacking of the cations of 1-(2-bromo-6-fluorophenylmethyl)-3-methylimidazolium bromide (**3**). Thermal ellipsoids are at the 50% level. Hydrogen atoms are omitted for clarity.

3. Conclusions

The crystal structures of 1-(2-halo-6-fluorophenylmethyl)-1-methylimidazolium bromide salts possess common features: charge-assisted hydrogen bonding between bromide anions and H(1) and H(2), and π - π stacking between phenyl rings, which are related by a centre of inversion. The presence of water molecules in the crystal structures of **1** and **3** does not affect these similarities, but it does lead to significant differences between the conformations of the cations, the absence of anion- π interactions, and hydrogen bonding between H(3) and water molecules instead of bromide anions. Only the crystal structure of **4** possesses halogen bonding, which is presumably a consequence of the interaction between the iodine atom and the bromide anion being stronger than that between the other halogen atoms and a bromide anion. The relative strengths of the interactions are consistent with those observed previously for imidazolium salts [3,5,9]: $\text{H}(1)\cdots\text{Br}^- > \text{H}(2)\cdots\text{Br}^- \approx \text{H}(3)\cdots\text{Br}^- > \text{I}\cdots\text{Br}^- > \pi\cdots\text{Br}^- > \pi\cdots\pi$.

4. Experimental

4.1 Instrumentation

The ^1H , $^{13}\text{C}\{^1\text{H}\}$ and ^{19}F NMR spectra were recorded in CDCl_3 using Bruker DRX300 or DPX400 spectrometers. ^1H (300.13 or 400.14 MHz) were referenced internally using the residual protio solvent resonance relative to SiMe_4 (δ 0), ^{13}C (75.48

or 100.61 MHz) externally to SiMe₄ (δ 0), and ¹⁹F (282.40 or 376.47 MHz) externally to CFCl₃ (δ 0). All chemical shifts are quoted in δ (ppm), using the high frequency positive convention, and coupling constants in Hz. Electrospray mass spectra were recorded on a Bruker Daltonics micrOTOF spectrometer.

4.2 Materials

1-Methylimidazole (Aldrich), and 2-halo-6-fluorobenzyl bromides and 2-trifluoromethyl-6-fluorobenzyl bromide (Apollo Scientific) were used as supplied.

4.3 Preparation of 1-(2,6-difluorophenylmethyl)-3-methylimidazolium bromide (1)

1-Methylimidazole (0.57 g, 6.93 mmol) was added to 2,6-difluorophenylmethyl bromide (1.45 g, 6.99 mmol) in dichloromethane (30 cm³). The mixture was left at ambient temperature for 96 h. The solvent was removed by rotary evaporation affording the product as a white powder. Yield 1.89 g (94%). MS: C₁₁H₁₁F₂N₂ requires 209.0890; found [M - Br]⁺ 209.0913. ¹H (CDCl₃/(CD₃)₂SO): δ = 10.07 (1H, s, N₂CH), 7.43 (1H, t, ³J_{HH} = 1.8 Hz, HCCH), 7.34 (1H, m, H_{para}), 7.18 (1H, t, ³J_{HH} = 1.8 Hz, HCCH), 6.29 (2H, m, H_{meta}), 5.50 (2H, s, CH₂), 3.98 (3H, s, CH₃). ¹³C{¹H} NMR (CDCl₃/(CD₃)₂SO): δ = 162.8 (dd, ¹J_{CF} = 252 Hz, ³J_{CF} = 6 Hz, CF), 137.7 (s, N₂CH), 132.5 (t, ²J_{CF} = 10 Hz, C_{para}), 124.1 (s, CHCH), 121.6 (s, CHCH), 112.3 (dm, ²J_{CF} = 24 Hz, C_{meta}), 109.2 (t, ²J_{CF} = 18 Hz, C_{ipso}), 41.0 (t, ³J_{CF}, NCH₂), 36.7 (NCH₃). ¹⁹F (CDCl₃/(CD₃)₂SO): δ = -113.80 (2F, m).

4.4 Preparation of 1-(2-chloro-6-fluorophenylmethyl)-3-methylimidazolium bromide (2)

1-Methylimidazole (0.58 g, 7.00 mmol) and 2-chloro-6-fluorophenylmethyl bromide (1.54 g, 6.87 mmol) were treated as in 4.3. Yield 1.81 g (86%). MS: $C_{11}H_{11}^{35}ClFN_2$ and $C_{11}H_{11}^{37}ClFN_2$ require 225.0595 and 227.0595; found $[M - Br]^+$ 225.0630 and 227.0691. 1H ($(CD_3)_2SO$): δ = 9.18 (1H, s, N_2CH), 7.72 (1H, s, $HCCH$), 7.71 (1H, s, $HCCH$), 7.58 (1H, td, $^3J_{HH} = 8.2$, $^3J_{HF} = 6.3$ Hz, C_6H_3), 7.49 (1H, dt, $J_{HH} = 8.2$ Hz, $J_{HH} = 1.1$ Hz, C_6H_3), 7.40 (1H, ddd, $J_{HF} = 9.5$ Hz, $J_{HH} = 8.2$, $J_{HH} = 1.3$ Hz, C_6H_3), 5.56 (2H, d, $J = 1.6$ Hz, NCH_2), 3.84 (3H, s, NCH_3). $^{13}C\{^1H\}$ NMR ($(CD_3)_2SO$): δ = 163.4 (d, $^1J_{CF} = 251$ Hz, CF), 137.3 (N_2CH), 135.4 (d, $J_{CF} = 5$ Hz, C_6), 133.1 (d, $J_{CF} = 10$ Hz, C_6), 126.6 (d, $J_{CF} = 3$ Hz, C_6), 124.4 (s, $HCCH$), 122.9 (s, $HCCH$), 120.3 (d, $J_{CF} = 17$ Hz, C_6), 115.9 (d, $^2J_{CF} = 22$ Hz, C_{ipso}), 44.2 (d, $^3J_{CF} = 4$ Hz, NCH_2), 36.4 (s, NCH_3). ^{19}F (d_6 -DMSO): δ = -111.89 (1F, m).

4.5 Preparation of 1-(2-bromo-6-fluorophenylmethyl)-3-methylimidazolium bromide (3)

1-Methylimidazole (0.155g, 1.88 mmol) and 2-bromo-6-fluorophenylmethyl bromide (0.505 g, 1.88 mmol) were treated as in 4.3. Yield 0.635 g (96%). MS: $C_{11}H_{11}^{79}BrFN_2$ and $C_{11}H_{11}^{81}BrFN_2$ require 269.0090 and 271.0186; found $[M - Br]^+$ 269.0087 and 271.0068. 1H ($(CD_3)_2SO$): δ = 9.22 (1H, s, N_2CHN), 7.76 (1H, s, $HCCH$), 7.73 (1H, s, $HCCH$), 7.64 (1H, d, $^3J_{HH} = 7.9$ Hz, C_6H_3), 7.51 (1H, td, $^3J_{HH} = 8.4$ Hz, $J_{HF} = 6.2$ Hz, C_6H_3), 7.46 (1H, ddd, $^3J_{HF} = 9.6$ Hz, $J_{HH} = 8.4$ Hz, $J_{HH} = 1.2$ Hz, C_6H_3), 5.56 (2H, d, $^3J_{HF} = 1.5$ Hz, NCH_2), 3.86 (3H, s, NCH_3). $^{13}C\{^1H\}$ NMR ($(CD_3)_2SO$): δ = 163.4 (d, $^1J_{CF} = 251$ Hz, CF), 137.2 (N_2CH), 133.5 (d, $J_{CF} = 10$ Hz, C_6), 129.9 (d, $J_{CF} = 3$ Hz, C_6), 125.6 (d, $J_{CF} = 4$ Hz, C_6), 124.5 (s, $HCCH$), 122.9 (s, $HCCH$), 121.6 (d, $J_{CF} = 17$ Hz, C_6), 116.4 (d, $^2J_{CF} = 22$ Hz, C_{ipso}), 46.7 (d, $^3J_{CF} = 3$ Hz, NCH_2), 36.4 (s, NCH_3). ^{19}F ($(CD_3)_2SO$): δ = -110.71 (m).

4.6 Preparation of 1-(2-iodo-6-fluorophenylmethyl)-3-methylimidazolium bromide (4)

1-Methylimidazole (0.135 g, 1.65 mmol) and 2-iodo-6-fluorophenylmethyl bromide (0.505 g, 1.60 mmol) were treated as in 4.3. Yield 0.595 g (93%). MS: $C_{11}H_{11}FIN_2$ requires 316.9951; found $[M - Br]^+$ 316.9960. 1H ($(CD_3)_2SO$): δ = 9.16 (1H, s, N_2CH), 7.86 (1H, d, J_{HH} = 7.7 Hz, C_6H_3), 7.76 (1H, m, $HCCH$), 7.72 (1H, m, $HCCH$), 7.41 (1H, ddd, J_{HF} = 9.6 Hz, J_{HH} = 8.4, 1.1 Hz, C_6H_3), 7.32 (1H, td, J_{HH} = 8.0 Hz, J_{HF} = 6.1 Hz, C_6H_3), 5.52 (2H, d, J_{HF} = 1.8 Hz, NCH_2), 3.86 (3H, s, NCH_3). $^{13}C\{^1H\}$ NMR ($(CD_3)_2SO$): δ = 162.7 (d, $^1J_{CF}$ = 252 Hz, CF), 137.1 (s, N_2CH), 136.4 (d, J_{CF} = 3 Hz, C_6), 134.0 (d, J_{CF} = 9 Hz, C_6), 124.5 (s, $HCCH$), 124.3 (d, J_{CF} = 16, C_6), 122.9 (s, $HCCH$), 116.9 (d, $^2J_{CF}$ = 22 Hz, C_{ipso}), 103.9 (s, C_6), 51.2 (d, $^3J_{CF}$ = 4, NCH_2), 36.4 (s, NCH_3). ^{19}F ($(CD_3)_2SO$): δ = -109.88 (m)

4.7 Preparation of 1-(2-trifluoromethyl-6-fluorophenylmethyl)-3-methylimidazolium bromide (5)

1-Methylimidazole (ca. 0.1 g, 1.3 mmol) and 2-bromo-6-fluorophenylmethyl bromide (0.478 g, 1.86 mmol) were treated as in 4.3. Yield 0.36 g (82%). MS: $C_{12}H_{11}F_4N_2$ requires 259.0858; found $[M - Br]^+$ 259.1499. 1H ($CDCl_3/(CD_3)_2SO$): 9.52 (s, 1H, N_2CH), 7.64 (s, 1H, $HCCH$), 7.63 (1H, m, C_6H_3), 7.59 m (1H, m, C_6H_3), 7.43 (1H, dd, J = 9.3 Hz, J = 9.3 Hz, C_6H_3), 7.31 (s, 1H, $HCCH$), 5.85 (2H, s, CH_2), 3.94 (3H, s, CH_3). $^{13}C\{^1H\}$ ($CDCl_3/(CD_3)_2SO$): δ = 162.2 (d, $^1J_{CF}$ = 250 Hz, CF), 137.4 (s, N_2CH), 132.8 (d, J_{CF} = 10 Hz, C_6), 130.8 (q, $^2J_{CF}$ = 31 Hz, CCF_3), 124.4 (s, $HCCH$), 123.2 (qd, $^1J_{CF}$ = 275 Hz, $^4J_{CF}$ = 4 Hz, CF_3), 122.8 (dq, J_{CF} = 6 Hz, J_{CF} = 5 Hz, C_6), 122.0 (s, $HCCH$),

120.8 (d, $J_{\text{CF}} = 22$ Hz, C_6), 118.2 (d, $^2J_{\text{CF}} = 17$ Hz, C_{ipso}), 43.6 (s, CH_2), 36.7 (s, CH_3).

^{19}F ($\text{CDCl}_3/(\text{CD}_3)_2\text{SO}$): $\delta = -57.84$ (3F, s, CF_3), -111.16 (1F, m, C_6F).

4.8 X-ray crystallography

Crystals of **1** and **5** were grown from dichloromethane. Crystals of **2** were grown from diffusion of pentane into a solution in dichloromethane. Crystals of **3** and **4** were grown from a mixture of dichloromethane, tetrahydrofuran and methanol. Crystal data are listed in Table 1. Diffraction data were collected on an Agilent SuperNova, single source at offset, Atlas diffractometer with graphite-monochromated $\text{Cu}-\text{K}\alpha$ radiation. The structures of **1** – **5** were solved using Olex2 [25] structure solution programme using Charge Flipping and refined with the olex2.refine [26] refinement package using Gauss-Newton minimization. The non-hydrogen atoms were refined with anisotropic thermal parameters. Hydrogen atom positions were added in idealized positions and a riding model with fixed thermal parameters ($U_{ij} = 1.2U_{eq}$ for the atom to which they are bonded (1.5 for CH_3)) was used for subsequent refinements. The function minimized was $\Sigma[w(|F_o|^2 - |F_c|^2)]$ with reflection weights $w^{-1} = [\sigma^2 |F_o|^2 + (g1P)^2 + (g2P)]$ where $P = [\max |F_o|^2 + 2|F_c|^2]/3$. CCDC 1583317 - 1583321 (**1** – **5** respectively) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

4.9 DFT calculations

DFT calculations were performed using Gaussian09 [27] with the long range corrected functional ω B97X-D [13] method with the 6-311++G(2d,2p) basis set, except for iodine for which the 6-311G(d) basis set was used. The energies of interaction were calculated as the difference between the difference between the energy of the species and the sum of those of its components. The C—H and O—H bond distances of the experimental were changed to 1.083 and 0.960 Å before calculation of their energies [28].

5. Acknowledgement

We thank the University of Waikato for support and Dr J. Burrows for technical assistance.

References

- [1] J.M. Serrano-Becerra, S. Hernández-Ortega, D. Morales-Morales, J. Valdés-Martínez, Bottom-up design and construction of a non-centrosymmetric network through π – π stacking interactions, *CrystEngComm* 11 (2009) 226–228.
- [2] G.C. Saunders, On the importance of π – π stacking and cation–anion interactions in the construction of non-centrosymmetric networks of bromide salts of imidazolium cations bearing arene and polyfluoroarene rings, *CrystEngComm* 13 (2011) 1801–1803.
- [3] H.I. Althagbi, A.J. Edwards, B.K. Nicholson, D.A. Reason, G.C. Saunders, S.A. Sim, D.A. van der Heijden, Arene-Perfluoroarene-Anion Stacking and Hydrogen

Bonding Interactions in Imidazolium Salts for the Crystal Engineering of Polarity, *Cryst. Grow. Des.* 16 (2016) 174–188.

- [4] M.A. Oosterwijk, G.C. Saunders, W. Zou, Polyfluoroarene–arene π – π stacking in three directions to provide crystal polarity, *J. Fluorine Chem.* 188 (2016) 80–84.
- [5] H.I. Althagbi, D.R. Bernstein, W.C. Crombie, J.R. Lane, D.K. McQuiston, M.A. Oosterwijk, G.C. Saunders, W. Zou, The crystal structures of 1-(4-halo-2,3,5,6-tetrafluorophenyl)-3-benzylimidazolium bromides: the relative importance of anion– π , lone pair– π , π – π stacking and halogen bonding interactions, *J. Fluorine Chem.* 206 (2018) 61–71.
- [6] V.L. Arcus, D.R. Bernstein, W.C. Crombie, G.C. Saunders, Infinite stacking of alternating polyfluoroaryl rings and bromide anions, *CrystEngComm* 15 (2013) 9841–9843.
- [7] M. Giese, M. Albrecht, A. Valkonen, K. Rissanen, The pentafluorophenyl group as π -acceptor for anions: a case study, *Chem. Sci.* 6 (2015) 354–359.
- [8] G.C. Saunders, H.P. Thomas, Crystal structure and theoretical calculations of 1-(4-trifluoromethyl-2,3,5,6-tetrafluorophenyl)-3-benzylimidazolium bromide, *J. Struct. Chem.* 58 (2017) 211–215.
- [9] S. Tsuzuki, H. Tokuda, M. Mikami, Theoretical analysis of the hydrogen bond of imidazolium C₂–H with anions, *Phys. Chem. Chem. Phys.* 9 (2007) 4780–4784.
- [10] (a) D. Braga, G.R. Desiraju, J.S. Miller, A.G. Orpen, S.L. Price, Innovation in crystal engineering, *CrystEngComm* 4 (2002) 500–509.
(b) D. Braga, Crystal engineering, Where from? Where to?, *Chem. Commun.* (2003) 2751–2754.

- (c) D. Braga, L. Brammer, N.R. Champness, New trends in crystal engineering, *CrystEngComm*, 7 (2005) 1–19.
- (d) L. Brammer, Developments in inorganic crystal engineering, *Chem. Soc. Rev.* 33 (2004) 476–489.
- (e) S.K. Seth, D. Sarkar, A.D. Jana, T. Kar, On the Possibility of Tuning Molecular Edges To Direct Supramolecular Self-Assembly in Coumarin Derivatives through Cooperative Weak Forces: Crystallographic and Hirshfeld Surface Analyses, *Cryst Grow. Des.* 11 (2011) 4837–4849.
- [11] J.-Y. Dong, T.-P. You, 3-(2-Bromobenzyl)-1-methyl-1*H*-imidazol-3-ium bromide, *Acta Crystallogr., Sect. E: Struct. Rep. Online* 67 (2011) o1568.
- [12] (a) P. Manna, S.K. Seth, A. Bauzá, M. Mitra, S.R. Choudhury, A. Frontera, S. Mukhopadhyay, pH Dependent Formation of Unprecedented Water–Bromide Cluster in the Bromide Salts of PTP Assisted by Anion– π Interactions: Synthesis, Structure, and DFT Study, *Cryst. Grow. Des.* 14 (2014) 747–755.
- (b) P. Manna, S.K. Seth, M. Mitra, S.R. Choudhury, A. Bauzá, A. Frontera, S. Mukhopadhyay, Experimental and Computational Study of Counterintuitive $\text{ClO}_4^- \cdots \text{ClO}_4^-$ Interactions and the Interplay between $\pi^+ - \pi$ and Anion $\cdots \pi^+$ Interactions, *Cryst. Grow. Des.* 14 (2014) 5812–5821.
- [13] J.-D. Chai, M. Head-Gordon, Long-range corrected hybrid density functionals with damped atom–atom dispersion corrections, *Phys. Chem. Chem. Phys.* 10 (2008) 6615–6620.
- [14] (a) A. Bondi, van der Waals Volumes and Radii, *J. Phys. Chem.* 68 (1964) 441–451.

- (b) R.S. Rowland, R. Taylor, Intermolecular Nonbonded Contact Distances in Organic Crystal Structures: Comparison with Distances Expected from van der Waals Radii, *J. Phys. Chem.* 100 (1996) 7384–7391.
- [15] C. Estarellas, A. Bauzá, A. Frontera, D. Quiñonero, P.M. Deyà, On the directionality of anion– π interactions, *Phys. Chem. Chem. Phys.* 13 (2011) 5696–5702.
- [16] (a) M. Meot-Ner, The Ionic Hydrogen Bond, *Chem. Rev.* 105 (2005) 213–284.
 (b) S.A. Katsyuba, M.V. Vener, E.E. Zvereva, Z. Fei, R. Scopelliti, J.G. Brandenburg, S. Siankevich, P.J. Dyson, Quantification of Conventional and Nonconventional Charge-Assisted Hydrogen Bonds in the Condensed and Gas Phases, *J. Phys. Chem. Lett.* 6 (2015) 4431–4436.
- [17] (a) E. Arunan, G.R. Desiraju, R.A. Klein, J. Sadlej, S. Scheiners, I. Alkorta, D.C. Clary, R.H. Crabtree, J.J. Dannenberg, P. Hobza, H.G. Kjaergaard, A.C. Legon, B. Mennunucci, D.J. Nesbitt, Definition of the hydrogen bond (IUPAC Recommendations 2011), *Pure Appl. Chem.* 83 (2011) 1637–1641.
 (b) A. Elaiwi, P.B. Hitchcock, K.R. Seddon, N. Srinivasan, Y.-M. Tan, T.C. Welton, J.A. Zora, Hydrogen Bonding in Imidazolium Salts and its Implications for Ambient-temperature Halogenoaluminate(III) Ionic Liquids, *J. Chem. Soc., Dalton Trans.* (1995) 3467–3472.
 (c) C.B. Aakeröy, K.R. Seddon, The Hydrogen Bond and Crystal Engineering, *Chem. Soc. Rev.* (1993) 397–407.
 (d) G.R. Desiraju, A Bond by Any Other Name, *Angew. Chem. Int. Ed.* 50 (2001) 52–59.

- [18] (a) D. Quiñonero, C. Garau, C. Rotger, A. Frontera, P. Ballester, A. Costa, P.M. Deyà, Anion- π Interactions: Do They Exist?, *Angew. Chem. Int. Ed.* 41 (2002) 3389–3392.
- (b) M. Giese, M. Albrecht, K. Rissanen, Anion- π interactions with fluoroarenes, *Chem. Rev.* 115 (2015) 8867–8895.
- [19] I. Alkorta, I. Rozas, J. Elguero, Interaction of Anions with Perfluoro Aromatic Compounds, *J. Am. Chem. Soc.* 124 (2002) 8593–8598.
- [20] For example: (a) F.R. Ahmed, E.J. Gabe, The crystal and molecular structure of 6,6-diphenyl-3,3-diethyl-3-azabicyclo[3.1.0]hexane bromide monohydrate, *Acta Crystallogr.* 17 (1964) 603–608.
- (b) J.Z. Gougoutas, J. Johnson, Structure and solid-state chemistry of 3-carboxy-2-naphthalenediazonium bromide, *J. Am. Chem. Soc.* 100 (1978) 5816–5820
- (c) T.C.W. Mak, A centrosymmetric hydrogen-bonded cyclic (H₂O.Br)₂ system. Crystal structure of 1-methyl-1,3,5,7-tetraazaadamantan-1-ium bromide hydrate, *Inorg. Chem.* 23 (1984) 620–622.
- (d) T. Taga, K. Machida, N. Kimura, S. Hayashi, J. Umemura, T. Takenaka, Structure of dodecyldimethylpropylammonium bromide monohydrate, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* 42 (1986) 608–610.
- (e) P.J. Barnard, M.V. Baker, S.J. Berners-Price, B.W. Skelton, A.H. White, Dinuclear gold(I) complexes of bridging bidentate carbene ligands: synthesis, structure and spectroscopic characterisation, *Dalton Trans.* (2004) 1038–1047.
- (f) L. Dobrzanska, *cis*-(Carbonato- κ^2O,O')(*rac*-5,5,7,12,12,14-hexamethyl-1,4,8,11-tetraazacyclotetradecane- κ^4N)chromium(III) bromide monohydrate, *Acta Crystallogr., Sect. E: Struct. Rep. Online* 61 (2005) m1625–m1627.

- (g) V. Vitske, C. König, O. Hubner, E. Kaifer, H.-J. Himmel, Syntheses of the First Coordination Compounds of the New Strong Molecular Electron Donor and Double Proton Sponge 1,4,5,8-Tetrakis(tetramethylguanidino)naphthalene, *Eur. J. Inorg. Chem.* (2010) 115–126.
- (h) A. Das, A.D. Jana, S.K. Seth, B. Dey, S.R. Choudhury, T. Kar, S. Mukhopadhyay, N.J. Singh, I.-C. Hwang, K.S. Kim, Intriguing $\pi^+ - \pi$ Interaction in Crystal Packing, *J. Phys. Chem. B* 114 (2010) 4166–4170.
- (i) S. Patil, A. Deally, B. Gleeson, H. Müller-Bunz, F. Paradisi, M. Tacke, Novel benzyl-substituted *N*-heterocyclic carbene–silver acetate complexes: synthesis, cytotoxicity and antibacterial studies, *Metallomics* 3 (2011) 74–88.
- (j) A. Basu, G. Das, Encapsulation of a discrete cyclic halide water tetramer $[X_2(H_2O)_2]^{2-}$, $X = Cl^-/Br^-$ within a dimeric capsular assembly of a tripodal amide receptor, *Chem. Commun.* 49 (2013) 3997–3999.
- (k) J. Beckmann, J. Bolsinger, A. Duthie, P. Finke, Diarylhalotelluronium(IV) cations $[(8-Me_2NC_{10}H_6)_2TeX]^+$ ($X = Cl, Br, I$) stabilized by intramolecularly coordinating N-donor substituents, *Dalton Trans.* 42 (2013) 12193–12202.
- (l) N. Lah, K. Lapanje, Bromidobis[2-(pyridin-2-yl)methanol- κ^2N,O]copper(II) bromide monohydrate, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* 69 (2013) 624–626.
- (m) K.P. Zhuravlev, A.V. Vologzhanina, V.A. Kudryashova, Z.S. Klemenkova, V.I. Tsaryuk, Influence of outer-sphere anions in europium tetrahydrofuran-2-carboxylates on the Eu^{3+} luminescence center, analyzed by methods of X-ray diffraction and optical spectroscopy, *Polyhedron* 56 (2013) 109–115.

- (n) A. Caballero, S. Bennett, C.J. Serpell, P.D. Beer, Iodo-imidazolium salts: halogen bonding in crystals and anion-templated pseudorotaxanes, *CrystEngComm* 15 (2013) 3076–3081.
- [21] (a) P. Metrangolo, H. Neukirch, T. Pilati, G. Resnati, Halogen bonding based recognition processes: a world parallel to hydrogen bonding, *Acc. Chem. Res.* 38 (2005) 386–395.
- (b) R. Metrangolo, G. Resnati, T. Pilati, S. Biella, Halogen Bonding in Crystal Engineering, *Struct. Bond.* 126 (2008) 105–136.
- [22] (a) J. Grebe, G. Geisler, K. Harms, K. Dehnicke, Donor-acceptor complexes of halide ions with 1,4-diiodotetrafluorobenzene, *Z. Naturforsch., B. Chem. Sci.* 54 (1999) 77–86.
- (b) H.M. Yamamoto, R. Kato, Design, Preparation, and Characterization of Novel ET Salts with Supramolecular Assembly. Sheet, Chain, and Pore Structures Based on Difluorotetraiodobenzene, *Chem. Lett.* (2000) 970–971.
- (c) S. Triguero, R. Llusar, V. Polo, M. Fourmigué, Halogen Bonding Interactions of *sym*-Triiodotrifluorobenzene with Halide Anions: A Combined Structural and Theoretical Study, *Cryst. Growth Des.* 8 (2008) 2241–2247.
- (d) G. Cavallo, S. Biella, J. Lu, P. Metrangolo, T. Pilati, G. Resnati, G. Terraneo, Halide anion-templated assembly of di- and triiodoperfluorobenzenes into 2D and 3D supramolecular networks, *J. Fluorine Chem.* 131 (2010) 1165–1172.
- (e) J. Viger-Gravel, S. Leclerc, I. Korobkov, D.L. Bryce, Correlation between ^{13}C chemical shifts and the halogen bonding environment in a series of solid paradiiidotetrafluorobenzene complexes, *CrystEngComm* 15 (2013) 3168–3177.

- [23] (a) A. Bhattacharyya, P.K. Bhaumik, A. Bauzá, P.P. Jana, A. Frontera, M.G.B. Drew, S. Chattopadhyay, A combined experimental and computational study of supramolecular assemblies in ternary copper(II) complexes with a tetradentate N4 donor Schiff base and halides, *RSC Adv.* 4 (2014) 58643–58651.
- (b) S.R. Choudhury, P. Gamez, A. Robertazzi, C.-Y. Chen, H.M. Lee, S. Mukhopadhyay, Experimental Observation of Supramolecular Carbonyl- $\pi/\pi-\pi/\pi$ -carbonyl and Carbonyl- $\pi/\pi-\pi/\pi$ -anion Assemblies Supported by Theoretical Studies, *Cryst. Grow. Des.* 8 (2008) 3773–3784.
- (c) M. Roy, A. Bauzá, A. Frontera, S. Banerjee, S. Halder, A. Saha, An experimental and computational investigations of supramolecular anion- $\pi/\pi-\pi/\pi$ -anion assemblies in mononuclear Zn(II) complexes with a versatile tetradentate N-donor Schiff base ligand, *Polyhedron* 102 (2015) 764–772.
- [24] (a) F.J. Strieter, D.H. Templeton, Crystal structure of the carbon tetrabromide p-xylene complex, *J. Chem. Phys.*, 37 (1962) 161–164.
- (b) S.V. Rosokha, I.S. Neretin, T.Y. Rosokha, J. Hecht, J.K. Kochi, Charge-transfer character of halogen bonding: Molecular structures and electronic spectroscopy of carbon tetrabromide and bromoform complexes with organic σ - and π -donors, *Heteroatom Chem.* 17 (2006) 449–459.
- (c) S.M. Hubig, S.V. Lindeman, J.K. Kochi, Charge-transfer bonding in metal–arene coordination, *Coord. Chem. Rev.* 200-202 (2000) 831–873.
- (d) D.S. Reddy, D.C. Craig, G.R. Desiraju, Supramolecular Synthons in Crystal Engineering. 4. Structure Simplification and Synthon Interchangeability in Some Organic Diamondoid Solids, *J. Am. Chem. Soc.* 118 (1996) 4090–4093.

- [25] O.V. Dolomanov, L.J. Bourhis, R.J. Gildea, J.A.K. Howard, H. Puschmann, *OLEX2: a complete structure solution, refinement and analysis program*, J. Appl. Cryst. 42 (2009) 339–341.
- [26] L.J. Bourhis, O.V. Dolomanov, R.J. Gildea, J.A.K. Howard, H. Puschmann, The anatomy of a comprehensive constrained, restrained refinement program for the modern computing environment – *Olex2* dissected, Acta Cryst. A 71 (2015) 59–75.
- [27] M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G.A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H.P. Hratchian, A.F. Izmaylov, J. Bloino, G. Zheng, J.L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J.A. Montgomery Jr., J.E. Peralta, F. Ogliaro, M. Bearpark, J.J. Heyd, E. Brothers, K.N. Kudin, V.N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J.C. Burant, S.S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J.M. Millam, M. Klene, J.E. Knox, J.B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R.E. Stratmann, O. Yazyev, A.J. Austin, R. Cammi, C. Pomelli, J.W. Ochterski, R.L. Martin, K. Morokuma, V.G. Zakrzewski, G.A. Voth, P. Salvador, J.J. Dannenberg, S. Dapprich, A.D. Daniels, Ö. Farkas, J.B. Foresman, J.V. Ortiz, J. Cioslowski, D.J. Fox, Gaussian 09, Revision A. 2, Gaussian, Inc, Wallingford, CT, 2009.
- [28] (a) F.H. Allen, A Systematic Pairwise Comparison of Geometric Parameters Obtained by X-ray and Neutron Diffraction, Acta Crystallogr., Sect. B: Struct. Sci. 42 (1986) 515–522.

(b) F.H. Allen, I.J. Bruno, Bond lengths in organic and metal-organic compounds revisited: $X-H$ bond lengths from neutron diffraction data, *Acta Crystallogr., Sect. B: Struct. Sci., Cryst. Eng. Mat.* 69 (2013) 624–626.

Table 1

Crystallographic data for 1-(2,6-difluorophenylmethyl)-3-methylimidazolium bromide **1**, 1-(2-chloro-6-fluorophenylmethyl)-3-methylimidazolium bromide **2**, 1-(2-bromo-6-fluorophenylmethyl)-3-methylimidazolium bromide **3**, 1-(2-iodo-6-difluorophenylmethyl)-3-methylimidazolium bromide **4**, and 1-(2-trifluoromethyl-6-fluorophenylmethyl)-3-methylimidazolium bromide **5**.^a

	1	2	3	4	5
Formula	C ₁₁ H ₁₁ BrF ₂ N ₂ .H ₂ O	C ₁₁ H ₁₁ BrClFN ₂	C ₁₁ H ₁₁ Br ₂ FN ₂ .1½H ₂ O	C ₁₁ H ₁₁ BrFIN ₂	C ₁₂ H ₉ BrF ₄ N ₂
Formula weight	307.14	305.58	377.05	397.03	339.13
Crystal system	monoclinic	triclinic	monoclinic	triclinic	monoclinic
Space group	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> $\bar{1}$	<i>I</i> 2/ <i>c</i>	<i>P</i> $\bar{1}$	<i>P</i> 2 ₁ / <i>c</i>
<i>a</i> , Å	7.32808(12)	6.9865(11)	13.8188(4)	7.4492(4)	8.59143(17)
<i>b</i> , Å	12.4953(2)	8.8271(7)	8.7968(2)	8.9471(5)	10.82231(17)
<i>c</i> , Å	13.7069(2)	10.1670(13)	22.6679(8)	10.3194(5)	14.1621(3)

$\alpha, ^\circ$	—	76.806(9)	—	76.142(5)	—
$\beta, ^\circ$	97.1464(15)	75.514(13)	92.576(3)	81.048(4)	104.443(2)
$\gamma, ^\circ$	—	85.464(10)	—	67.455(5)	—
$V, \text{\AA}^3$	1245.34(4)	590.88(14)	2752.74(14)	615.17(6)	1275.17(4)
Z	4	2	8	2	4
$D_c (\text{g cm}^{-3})$	1.638	1.717	1.820	2.143	1.766
Crystal size (mm^3)	$0.033 \times 0.105 \times 0.146$	$0.075 \times 0.169 \times 0.440$	$0.141 \times 0.151 \times 0.216$	$0.057 \times 0.068 \times 0.306$	$0.232 \times 0.302 \times 0.540$
$\mu (\text{mm}^{-1})$	4.639	6.729	7.526	24.173	4.775
θ range ($^\circ$)	$4.81 \rightarrow 74.00$	$4.60 \rightarrow 74.51$	$3.90 \rightarrow 74.05$	$4.42 \rightarrow 73.97$	$5.21 \rightarrow 73.75$
Total reflections	7,147	5,733	7,667	6,540	12,304
Unique reflections (R_{int})	2,443 (0.0173)	2,312 (0.0645)	2,716 (0.212)	2,410 (0.0416)	2,530 (0.0506)

Observed reflections [$I > 2\sigma(I)$]	2,226	1927	2,610	2,316	2,104
Parameters	163	146	167	146	172
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0213$ $wR_2 = 0.0541$	$R_1 = 0.0723$ $wR_2 = 0.1732$	$R_1 = 0.0283$ $wR_2 = 0.0744$	$R_1 = 0.0309$ $wR_2 = 0.0792$	$R_1 = 0.0312$ $wR_2 = 0.0708$
R indices (all data)	$R_1 = 0.0255$ $wR_2 = 0.0557$	$R_1 = 0.0909$ $wR_2 = 0.1848$	$R_1 = 0.0311$ $wR_2 = 0.0753$	$R_1 = 0.0345$ $wR_2 = 0.0825$	$R_1 = 0.0428$ $wR_2 = 0.0775$
Weighting scheme	$w = 1/[\sigma^2(F_o)^2 + \{0.0270 (F_o^2 + 2F_c^2)/3\}^2 + 0.8870 (F_o^2 + 2F_c^2)/3]$	$w = 1/[\sigma^2(F_o)^2 + \{0.0330 (F_o^2 + 2F_c^2)/3\}^2 + 12.4940 (F_o^2 + 2F_c^2)/3]$	$w = 1/[\sigma^2(F_o)^2 + \{0.0450 (F_o^2 + 2F_c^2)/3\}^2 + 9.2540 (F_o^2 + 2F_c^2)/3]$	$w = 1/[\sigma^2(F_o)^2 + \{0.0550 (F_o^2 + 2F_c^2)/3\}^2 + 0.6490 (F_o^2 + 2F_c^2)/3]$	$w = 1/[\sigma^2(F_o)^2 + \{0.0350 (F_o^2 + 2F_c^2)/3\}^2 + 1.0700 (F_o^2 + 2F_c^2)/3]$
Max., min. $\Delta\rho$ (eÅ ⁻³)	0.402, -0.405	1.988, -1.918	0.700, -0.885	1.510, -1.272,	0.496, -0.636
Goodness of fit on F^2	1.052	1.048	1.021	1.041	1.054

^a Estimated standard deviations are given in parentheses. The data were collected at 100.0(1) K using Cu K α radiation ($\lambda = 1.54184$ Å).

Table 2

Selected bond distances (Å) and angles (°) for 1-(2,6-difluorophenylmethyl)-3-methylimidazolium bromide **1**,
1-(2-chloro-6-fluorophenylmethyl)-3-methylimidazolium bromide **2**, 1-(2-bromo-6-fluorophenylmethyl)-3-methylimidazolium bromide **3**,
1-(2-iodo-6-difluorophenylmethyl)-3-methylimidazolium bromide **4**, and 1-(2-trifluoromethyl-6-fluorophenylmethyl)-3-methylimidazolium bromide **5**.^a

	1		2		3		4		5	
	exptl	calc ^b	exptl	calc ^b	exptl	calc ^b	exptl	calc ^b	exptl	calc ^b
N(1)—C(1)	1.335(2)	1.328	1.333(10)	1.327	1.332(3)	1.327	1.342(5)	1.327	1.333(3)	1.328
N(2)—C(1)	1.329(2)	1.328	1.322(10)	1.328	1.332(3)	1.328	1.322(5)	1.328	1.326(3)	1.327
N(1)—C(2)	1.378(2)	1.374	1.379(10)	1.374	1.376(3)	1.373	1.388(5)	1.373	1.378(3)	1.375
N(2)—C(3)	1.380(2)	1.375	1.399(10)	1.376	1.382(3)	1.374	1.377(5)	1.374	1.381(4)	1.376
C(2)—C(3)	1.349(3)	1.354	1.349(11)	1.353	1.356(4)	1.354	1.355(6)	1.354	1.357(4)	1.353
N(1)—C(5)	1.474(2)	1.473	1.491(9)	1.472	1.473(3)	1.476	1.474(5)	1.477	1.482(3)	1.472

N(2)—C(4)	1.466(2)	1.460	1.474(9)	1.460	1.467(3)	1.460	1.469(5)	1.460	1.470(3)	1.460
C(5)—C(6)	1.501(3)	1.498	1.498(11)	1.501	1.508(3)	1.498	1.513(6)	1.499	1.512(4)	1.508
C(7)—F	1.356(2)	1.347	1.362(10)	1.347	1.368(3)	1.340	1.363(3)	1.341	1.358(3)	1.347
C(11)—X ^c	1.357(2)	1.342	1.748(9)	1.744	1.891(3)	1.898	2.114(4)	2.111	1.510(3)	1.510
N(1)—C(1)—N(2)	108.47(16)	108.8	108.7(6)	108.8	108.4(2)	108.8	108.2(3)	108.8	108.4(2)	108.9
C(1)—N(1)—C(2)	108.57(15)	108.7	109.0(6)	108.6	109.1(2)	108.7	108.6(3)	108.6	108.9(2)	108.6
C(1)—N(1)—C(5)	125.32(15)	125.5	123.1(7)	125.6	124.9(2)	126.4	127.9(3)	126.4	127.0(2)	125.5
C(1)—N(2)—C(3)	108.68(15)	108.5	108.4(6)	108.5	108.5(2)	108.5	109.5(3)	108.5	109.0(2)	108.4
C(1)—N(2)—C(4)	124.68(16)	125.8	125.4(6)	125.8	125.6(2)	125.7	124.5(3)	125.7	125.2(2)	126.0
N(1)—C(5)—C(6)	111.57(14)	111.9	112.0(6)	111.9	112.0(2)	112.4	113.0(3)	112.3	113.7(2)	111.7

C(1)—N(1)—C(5)—C(6)	-111.3(2)	94.7	1(1)	102.1	118.1(3)	21.0	-9.2(6)	17.1	36.8(4)	110.9
C(2)—N(1)—C(5)—C(6)	68.5(2)	-80.9	178.0(7)	-74.7	-62.4(3)	159.8	169.7(4)	-162.9	-148.1(2)	-67.1
N(1)—C(5)—C(6)—C(7)	-109.6(2)	97.1	-89.8(9)	-81.8	105.3(3)	-105.1	-74.6(5)	78.4	84.4(3)	-70.1
N(1)—C(5)—C(6)—C(11)	76.0(2)	-83.1	95(1)	98.7	-75.8(3)	76.0	109.7(5)	-102.1	-100.3(3)	110.6
$\angle$ C ₃ N ₂ C ₆ ^d	73.0(4)	71.1	85(2)	71.9	76.7(4)	84.9	76.2(7)	85.3	77.9(7)	79.4

^a Estimated standard deviations are given in parentheses.

^b The calculations were performed using the ω B97X-D functional and 6-311++G(2d,2p) basis set, except for the iodine atom, for which the 6-311G(d) basis set was used.

^c **1** X = F, **2** X = Cl, **3** X = Br, **4** X = I, **5** X = CF₃

^d The angle between the planes defined by the non-hydrogen atoms of the imidazolium and phenyl rings.

Table 3

Selected interionic bond distances (Å), angles (°) and energies of interaction (kJ mol⁻¹) for 1-(2,6-difluorophenylmethyl)-3-methylimidazolium bromide **1**, 1-(2-chloro-6-fluorophenylmethyl)-3-methylimidazolium bromide **2**, 1-(2-bromo-6-fluorophenylmethyl)-3-methylimidazolium bromide **3**, 1-(2-iodo-6-difluorophenylmethyl)-3-methylimidazolium bromide **4**, and 1-(2-trifluoromethyl-6-fluorophenylmethyl)-3-methylimidazolium bromide **5**.^a

	1	2	3	4	5
C(1)⋯Br ⁻	3.648(1)	3.568(3)	3.612(3)	3.562(5)	3.486(3)
N(1)—C(1)⋯Br ⁻	139.0(1)	145.0(5)	101.0(1)	102.2(3)	143.2(2)
N(2)—C(1)⋯Br ⁻	110.5(1)	102.6(5)	144.1(2)	145.5(3)	103.4(2)
C ₃ N ₂ ⋯Br ⁻ ^b	0.769(2)	0.922(9)	1.282(3)	0.942(6)	1.115(3)
C ₆ ⋯Br ⁻ ^c	—	4.190(9)	—	4.108(4)	4.334(3)
C ₆ [†] ⋯Br ⁻ ^d	—	4.304(9)	—	4.334(4)	4.511(3)

E [H(1)⋯Br] ^e	-338	-354	-343	-372	-357
E [C ₆ ⋯Br] ^e	—	-11	—	-21	-12
C(2)⋯Br ⁻	3.760(2)	3.660(7)	3.547(3)	3.709(4)	3.758(3)
N(1)—C(2)⋯Br ⁻	145.1(1)	104.2(5)	158.0(2)	105.5(3)	97.9(2)
C(3)—C(2)⋯Br ⁻	103.0(1)	148.3(6)	95.2(2)	147.8(3)	154.4(2)
C ₃ N ₂ ⋯Br ^b	1.144(2)	0.328(8)	0.120(3)	0.118(5)	0.443(3)
C ₆ ⋯Br ^{-c}	4.382(2)	—	4.563(3)	—	—
C ₆ [†] ⋯Br ^{-d}	5.487(2)	—	5.941(3)	—	—
E [H(2)⋯Br] ^e	-285	-310	-296	-323	-314
E [C ₆ ⋯Br] ^e	-2	—	-2	—	—

C(3)⋯Br ⁻	—	3.841(9)	—	3.769(5)	3.900(3)
N(2)—C(3)⋯Br ⁻	—	92.6(4)	—	129.9(3)	53.9(1)
C(2)—C(3)⋯Br ⁻	—	139.5(6)	—	102.5(3)	104.4(2)
C ₃ N ₂ ⋯Br ^{-b}	—	2.256(9)	—	2.385(5)	1.059(3)
E [H(3)⋯Br ⁻] ^e	—	-300	—	-290	-302
C(3)⋯O	3.213(3), 3.413(3)	—	3.474(4), 3.768(4)	—	—
N(2)—C(3)⋯O	104.6(1), 151.3(1)	—	135.8(2), 90.3(2)	—	—
C(2)—C(3)⋯O	139.6(1), 86.0(1)	—	117.0(2), 162.50(2)	—	—
C ₃ N ₂ ⋯O	1.317(3), 1.503(3)	—	0.013(4), 0.039(4)	—	—

E [H(3)⋯OH ₂] ^e	-14, -13	—	-18, -14	—	—
I⋯Br ⁻	—	—	—	3.3095(5)	—
C(11)—I⋯Br ⁻	—	—	—	174.9(1)	—
C ₆ F ₄ I⋯Br ⁻	—	—	—	0.344(4)	—
E [I⋯Br] ^e	—	—	—	-269	—
E [C ₆ ⋯Br] ^{e,f}	—	—	—	-54	—
O⋯Br ⁻	3.314(2), 3.342(2)	—	3.286(2), 3.314(3)	—	—
Br⋯O⋯Br ⁻	106.27(4)	—	106.89(7)	—	—
O⋯Br⋯O	73.73(4)	—	73.11(6)	—	—

$\text{C}_6 \cdots \text{C}_6^{\text{c}}$	3.726(4)	3.54(1)	3.410(4)	3.590(6)	3.751(4)
$\text{C}_6^{\dagger} \cdots \text{C}_6^{\dagger \text{d}}$	4.196(4)	3.99(1)	3.654(4)	4.087(6)	3.901(4)
$\text{E} [\text{C}_6 \cdots \text{C}_6]^{\text{e,g}}$	-55	-37	-42	-37	-37
$\text{E} [\text{C}_6 \cdots \text{C}_6]^{\text{e,h}}$	-19	-37	-37	-33	-32
$\text{C}_3\text{N}_2 \cdots \text{C}_6^{\dagger \text{b,d}}$	3.431(3)	—	—	—	—
$\angle \text{C}_3\text{N}_2 \text{C}_6^{\text{b,c}}$	6.4(1)	—	—	—	—
$\text{Br} \cdots \text{C}_6^{\text{c}}$	—	—	3.345(3)	—	—
$\text{Br} \cdots \text{C}_6^{\dagger \text{d}}$	—	—	3.403(3)	—	—
$\text{C}(11) - \text{Br} \cdots \text{C}_6^{\dagger \text{d}}$	—	—	174.4(1)	—	—

E [Br...C ₆] ^{e,i}	—	—	-12	—	—

^a Estimated standard deviations are given in parentheses.

^b C₃N₂ represents the plane defined by the non-hydrogen atoms of the imidazolium ring.

^c C₆ represents the plane defined by the six carbon atoms of the halotetrafluorophenyl ring.

^d C₆[†] represents the centroid of the six carbon atoms of the halotetrafluorophenyl ring.

^e The calculations were performed using the ωB97X-D functional and 6-311++G(2d,2p) basis set, except for the iodine atom, for which the 6-311G(d) basis set was used.

^f The anion–π interaction between bromide anion and 2-iodo-6-fluorotoluene based on the atom positions of **4**.

^g The π–π stacking interaction between adjacent imidazole molecules based on the atom positions of the salts.

^h The π–π stacking interaction between adjacent 2-halo-6-fluorotoluene molecules based on the atom positions of the salts.

ⁱ The lone pair–π interaction between two molecules of 2-bromo-6-fluorotoluene based on the atom positions of **3**.