**QUEEN'S
UNIVERSITY
BELFAST**

Whole grain intakes in Irish adults findings from the National Adults Nutrition Survey (NANS)

O'Donovan, C. B., Devlin, N. F., Buffini, M., Walton, J., Flynn, A., Gibney, M. J., Nugent, A. P., & McNulty, B. A. (2018). Whole grain intakes in Irish adults findings from the National Adults Nutrition Survey (NANS). *European Journal of Nutrition*. <https://doi.org/10.1007/s00394-018-1615-3>

Published in:
European Journal of Nutrition

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© Springer-Verlag GmbH Germany, part of Springer Nature 2018. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

1 **Whole grain intakes in Irish Adults: findings from the National Adults Nutrition Survey**
2 **(NANS)**

3 Clare B. O'Donovan¹, Niamh F. Devlin¹, Maria Buffini¹, Janette Walton^{2,3}, Albert Flynn², Michael
4 J. Gibney¹, Anne P. Nugent^{1,4} and Breige A. McNulty¹

6 **Author affiliations**

7 ¹UCD Institute of Food and Health, School of Agriculture and Food Science, University College
8 Dublin, Dublin 4, Ireland

9 ²School of Food and Nutritional Sciences, University College Cork, Cork, Ireland.

10 ³Department of Biological Sciences, Cork Institute of Technology, Cork, Republic of Ireland.

11 ⁴Department of Biological Sciences, Institute for Global Food Security, Queens University Belfast,
12 Northern Ireland.

14 **Corresponding author**

15 Dr. Anne Nugent, UCD Institute of Food and Health, School of Agriculture and Food Science,
16 University College Dublin, Dublin 4, Ireland. Email – anne.nugent@ucd.ie Telephone – (087)
17 9864415

19 **Acknowledgement**

20 This work was funded by the Irish Government, Department of Agriculture, Food and the Marine
21 under the 'Food for Health Research Initiative' 2007-2012 and Project 13 F 542 – National Nutritional
22 Databases for Public Health and New Product Development. AF and MJG were original grant holders
23 on the National Adult Nutrition Survey. AF, JW contributed to the design and execution of the study
24 and provided expert advice throughout. NFD and COD carried out the data analysis and wrote the
25 first draft. BAMcN and APN contributed to the design of the study, data analysis, paper editing and
26 review. MB contributed to data analysis and review. All authors critically reviewed the manuscript
27 and approved the final version submitted for publication.

28

29 **Abstract**

30 *Purpose:* Observational studies link high whole grain intakes to reduced risk of many chronic
31 diseases. This study quantified whole grain intakes in the Irish adult population and examined the
32 major contributing sources. It also investigated potential dietary strategies to improve whole grain
33 intakes.

34 *Methods:* Whole grain intakes were calculated in a nationally representative sample of 1500 Irish
35 adults using data from the most recent national food survey, the National Adult Nutrition Survey
36 (NANS). Food consumption was assessed, at brand level where possible, using a 4-day semi-
37 weighed food diary with whole grain content estimated from labels on a dry matter basis.

38 *Results:* Mean daily whole grain intakes were 27.8 ± 29.4 g/d, with only 19% of the population
39 meeting the quantity specific recommendation of 48g per day. Wheat was the highest contributor to
40 whole grain intake at 66%, followed by oats at 26%. High whole grain intakes were associated with
41 higher dietary intakes of fibre, magnesium, potassium, phosphorus and a higher alternative
42 Mediterranean Diet Score. Whole grain foods were most frequently eaten at breakfast time.
43 Regression analysis revealed that consumption of an additional 10g of whole grain containing
44 'ready to eat breakfast cereals', 'rice or pastas', or 'breads' each day would increase intake of whole
45 grains by an extra 5g, 3.5g and 2.7g respectively.

46 *Conclusions:* This study reveals low intakes of whole grains in Irish adults. Recommending cereals,
47 breads and grains with higher whole grain content as part of public health campaigns could improve
48 whole grain intakes.

49

50 **Keywords**

51 Whole grain intakes; adults; Ireland; food surveys; intake patterns

52

Introduction

The health implications associated with whole grain consumption have been widely documented, with previous research demonstrating that increasing whole grain intakes in the diet can result in positive health outcomes [1]. At a nutrient level, high whole grain consumption has been associated with diets rich in vitamins A, C, E, B6, fibre, folate, magnesium, potassium, phosphorus, calcium and iron [2, 3]. In terms of health, a number of meta-analyses have concluded that high whole grain intakes are associated with a reduced risk of many non-communicable diseases such as cardiovascular disease [4, 5], type 2 diabetes [6] and some cancers [7, 8]. Although, findings from intervention trials are not so convincing [1], it has been postulated that differences in study design, and methods of whole grain calculation make it difficult to compare findings [9]. Specific mechanisms behind any beneficial effect of whole grain intake remain unclear, with proposed explanations including the functional components of whole grains e.g. carotenoids, beta-glucan, inulin, phenolic acid, dietary fibre [10] and/or their nutrient profile [11]. It is also postulated that whole grains exert their benefits through actions such as reducing inflammation, improving blood lipid profile, reducing body weight, lowering blood pressure and exerting positive metabolic and hormonal effects due to phytochemicals present [5, 12, 13, 14].

With growing evidence supporting the inclusion of whole grain foods in the diet, intake patterns across and within populations worldwide are more routinely assessed and dietary recommendations increasingly mention whole grains. In certain countries such as Denmark and the US, specific quantitative intake recommendations of 75g and 48g per day have been established, whilst in other countries (Ireland, the UK and France), recommendations remain vague, and simply encourage whole grain varieties to be chosen over refined sources [15]. This may be due to the difficulties in establishing a uniform global definition of whole grain, due to issues such as the extent of processing permitted and which grains should be included in the definition [1]. Ferruzzi and colleagues proposed that the lowest amount a whole grain food should contain is 8g per 30g product (27g/100g) and based on this, the food product could be labelled as a 'whole grain food' [16], with similar thresholds endorsed recently by the Healthgrain forum (≥ 30 g whole grain ingredients) [17].

Whole grain intakes in Irish children and teenagers have previously been reported using data from the National Children's Food Survey and the National Teens' Food Survey [18], with no information on the intake patterns of older age groups. The aim of the current paper is to investigate the whole grain intakes in the Irish adult population, assess the major contributing sources and to investigate potential dietary strategies to improve whole grain intakes in the Irish population.

86 **Methods**

87 *National Adult Nutrition Survey (NANS)*

88 The National Adult Nutrition Survey (NANS) is a cross-sectional food survey, completed between
89 2008 and 2010, and conducted by the Irish Universities Nutrition Alliance (IUNA, www.iuna.net).
90 Ethical approval was obtained from University College Cork Clinical Research Ethics Committee of
91 the Cork Teaching Hospitals and the Human Ethics Research Committee of University College
92 Dublin (ECM 3 (p) 4 September 2008). Written consent was obtained from all participants in
93 accordance with the Declaration of Helsinki. A more detailed description of the survey methodology
94 has been reported elsewhere [19]. Because the Republic of Ireland does not have a national
95 identification system for adults, a database of names and addresses held by Data Ireland (National
96 Postal Service) was used to randomly select persons in 20 geographical clusters across the country,
97 selected to provide proportional representation across the urban-rural continuum. In total, 1500 adults
98 aged 18-90 years (740 men and 760 women) took part, with a response rate of 60%. A sample of
99 1500 free-living adults to represent a population of >4 million people participated in the dietary
100 survey. The sample size was chosen to deliver at least 100 individuals in the least-populated age and
101 sex subgroups. There were few exclusion criteria, other than pregnancy/lactation and inability to
102 complete the survey because of disability. The sample was representative of the Irish population with
103 respect to gender, age, location and social class as per the 2006 Irish census (Central Statistics Office
104 2007).

106 Food and beverage intake was determined using a 4-day semi-weighed food diary and assessed using
107 WISP V3.0 (Tinuviel Software, Anglesey, UK), based on data from the McCance and Widdowson's
108 The Composition of Foods (5th and 6th editions) supplemented with Irish food codes, to calculate
109 nutrient intakes. Each food/ beverage consumed was recorded and allocated an individual food code
110 and brand code with the participant recording the time of eating, the eating location and self-defining
111 each eating occasion as either a meal or a snack. The alternate Mediterranean Diet Index (aMED) was
112 calculated per individual as a marker of diet quality [20]. Under-reporters were determined as those
113 individuals who had a reported energy intake less than their basal metabolic rate (BMR) multiplied
114 by PAL 1.1 [21] and removed from subsequent analysis. Supplement use was also recorded.

116 From this food and nutrient database, all whole grain foods were identified (at brand level) and a new
117 whole grain database formed, the method for which has previously been described [18]. A whole
118 grain food was defined as any food containing a whole grain ingredient as listed in the Healthgrain
119 forum whole grain definition e.g. whole grain wheat, oats, rice, maize/corn, barley and rye [22]. Foods

were included in this study regardless of the whole grain content of the foods. Whole grain foods were then grouped into 1 of 7 groups for further analysis (Table 1). The total whole grain content per 100g of each food, and the source of whole grain in the food e.g. wheat, rye were also recorded. The amount of whole grain ‘actually consumed’ was then calculated by multiplying the amount of whole grain per 100g by the weight of food consumed. Following this, the dry weights of all whole grains were calculated to correct for water content as recommended by Ross and colleagues [8]. The dry weights of each whole grain ingredient were calculated by multiplying the % dry matter by the total weight of the food in 100g per day [23]. The % dry matter were obtained from McCance and Widdowson’s Cereal and cereal products supplement [24]. All data are presented on a % dry matter basis.

Statistical Analysis

Statistical analyses were carried out using SPSS® for Windows™ statistical software package version 20.0 (SPSS Inc., Chicago, IL, USA). The mean and standard deviations were calculated for the daily intake of total whole grain according to sex, age group, social class and eating location for the total population and for consumers of whole grain. As the distribution of the data approximated normality [17], statistical differences within the groups were detected either using independent t-tests (gender) or by one-way analysis of variance, adjusting for age group, social class and eating location as appropriate. Bonferroni post hoc tests were conducted and where appropriate corrected at a family level for type 1 error a rate of 5%.

The percentage contribution of each whole grain food group made to the diet was analysed, and the main whole grain sources were identified. Simple meal pattern analysis was carried out to assess the most common meals contributing to whole grain intakes whereby the proportion of total whole grain consumed was calculated for each meal type as defined by the participants in the food diaries (i.e. % of daily whole grain consumed at breakfast, lunch, dinner, snacks). Differences in mean daily intakes of macro- and micro-nutrients for consumers versus non-consumers of whole grain were evaluated using independent samples t-test.

Further analysis was carried out to determine the percentage of Irish adults not meeting US minimum recommendations for whole grain intakes (48g/day) [25].

Multiple linear regression analysis was conducted to explore the relationship of the seven whole grain food groups that contributed the most to total whole grain intakes. This was conducted on energy adjusted mean daily whole grain intakes using a backward process which was considered the best fit for the model.

154 **Results**

155 Mean and median daily intakes of whole grain (g/day) are presented in Table 2 for Irish adults (aged
156 18-90 years) and are presented for gender, age group, social class and eating location, for both the
157 total population and consumers only. Over 90% of Irish adults are consumers of whole grain foods,
158 with mean daily intakes of 27.8 ± 29.4 g (total population), increasing to 30.6 ± 29.5 g for consumers
159 only. Males consumed more whole grain ($P < 0.001$) in comparison to females, with mean daily intakes
160 of $31.6 \text{g} \pm 35.7$ versus $24.1 \text{g} \pm 20.9$ for the total population and $35.0 \text{g} \pm 35.9$ versus $26.3 \text{g} \pm 20.5$ for
161 consumers only. No intake differences were found across the age groups, however, there were
162 significant differences observed across social class and eating locations. The highest mean intake of
163 whole grains (31.1 ± 30.0 g/day) was identified in those individuals classed as professional/managerial
164 which was significantly ($P < 0.05$) higher than those individuals with skilled or manual jobs ($22.7 \pm$
165 23.5 g/day).

167 ***Whole grain recommendation***

168 The percentage of Irish adults meeting the US minimum whole grain recommendation of 48g/day
169 [25] was 19%, with little variation in compliance according to age group (Table 3).

171 ***Grain sources and food sources of whole grain***

172 Wheat was the major grain source consumed by the Irish adult population, and accounted for 66% of
173 all whole grain intakes. Oats were the next greatest source of whole grain (26%), while barley, rice,
174 rye and maize made up the lesser amounts ($< 8\%$) (Figure 1). Table 4 presents the food sources of
175 whole grain, by food group and across the tertiles of intake i.e. where the lowest consumers of whole
176 grain are presented in tertile 1 and the highest consumers in tertile 3. The major food group
177 contributing to whole grain intake in Irish adults was 'bread and rolls' (47% contribution) followed
178 by 'ready to eat breakfast cereals (RTEBC)' (26%) and 'other breakfast cereals' e.g. porridge (7%).
179 Across the tertiles, those in the highest tertiles had significantly ($P < 0.05$) greater intakes of all food
180 groups except 'rice, pasta and their dishes' and 'meat products, tofu & yoghurts'.

183 ***Food sources of whole grain at mealtimes***

184 Table 5 presents the food sources of whole grain consumed at different mealtimes and as snacks, for
185 the Irish adult population. Breakfast was the greatest contributor of whole grain intakes, providing
186 over 49% of the daily whole grain consumption. Of this, RTEBC and breads & rolls contributed most
187 significantly, providing nearly 40% and 35% of whole grain intake at breakfast respectively. At lunch,

188 dinner and snacks, whole grain bread and roll products contributed most to total whole grain intakes
189 at 80%, 58% and 44% respectively. Of note, whole grain containing sweet biscuits and related
190 products, and savoury biscuits and related products were more likely to be consumed as snacks
191 providing 30% and 15% of whole grain consumed at this eating occasion.

192

193 ***Consumers versus non-consumers of whole grain***

194 Comparing energy-adjusted nutrient intakes (% energy, mg/10MJ or µg/10MJ) across consumers
195 and non-consumers of whole grain, Table 6 shows non-consumers to have significantly lower intakes
196 of carbohydrate, fibre, potassium, phosphorus, magnesium and higher total fat intakes in comparison
197 with high whole-grain consumers. The aMED was significantly higher in tertile 3 in comparison with
198 non/lower/medium consumers of whole grain.

199

200 ***Proposed strategies to increase intakes of whole grains in the Irish population***

201 Table 7 presents the impact of increasing whole grain intakes across food groups. This analysis shows
202 that by increasing consumption of whole-grain food groups has a varied impact on overall intakes.
203 For example, if an individual was to increase their intake of RTEBC by 10g per day, their daily whole
204 grain intake would increase by nearly 5.4g. In contrast, a 10g daily increase in intakes of whole grain
205 containing bread and rolls or other breakfast cereals would increase daily whole grain intakes by
206 2.75g and 0.14g respectively. An additional 10g per day of savoury biscuits, crackers and popcorn
207 would increase daily whole grain intake by 7.5g.

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223 Discussion

224 The current paper describes whole grain intakes in Irish adults using data from the NANS study. For
225 the total population, the mean intake of whole grains was 27.8 g per day (median 20g/day), with over
226 91% of Irish adults identified as consumers of whole grains. Whole grain intakes differed in terms of
227 gender and social class, with males and those from higher social classes having higher intakes. In
228 comparison with the US quantity specific whole grain intake recommendation of 48g/day [25], only
229 19% of the Irish adult population met this minimum guideline. Approximately 50% of all whole
230 grain was consumed by Irish adults at breakfast, double the amount consumed at lunch, the second
231 most frequent whole grain eating occasion, with a reliance on wheat sources. Regression analysis
232 demonstrated that increased intakes of whole grain containing foods such as RTEBCs, rice and pastas
233 and bread by as little as 10g per day, could potentially enhance actual whole grain intakes in Irish
234 adults.

236 Dietary whole grain intakes have previously been described for Irish children and teenagers with
237 mean daily intakes of 18.5 and 23.2 g/d respectively and with over 90% of children aged 5-12 years
238 and over 86% of teenagers aged 13-17 years consuming whole grains (intakes for consumers only of
239 20.5 and 16.9 g/day respectively) [18]. Ready-to-eat breakfast cereals made the greatest contribution
240 to whole grain intakes in these Irish children and teens (44-59% of whole grain), followed by bread
241 and rolls (14-26% of whole grain) [18]. The reverse order was observed in the present study of Irish
242 adults, perhaps suggesting lowered consumption of whole grain containing breakfast cereals in adults
243 and a greater reliance on breads and rolls for whole grain intake. In all studies, wheat was the major
244 contributing grain, followed by oats. Our intake patterns (median daily intake 20g per day) are in line
245 with those reported from other countries [3, 26, 28, 29]. Recent data from the UK also reported a
246 median daily intake of 20g for whole grains in adults, with whole-grain breads and ready to eat
247 breakfast cereals as primary and secondary sources and wheat the main grain [26]. However, unlike
248 the current study which reported absolute intakes, energy-adjusted whole grain intakes appeared to
249 increase with age in the UK population, with females having higher whole grain intakes [26].
250 Australian consumption is reported as 21g/day for adults [28], with comparable values estimated for
251 US adults aged 19-50y in 2010 as 0.72 oz equivalents (approximately <20g/d) [29]. In contrast, very
252 low consumption levels are reported for France (mean daily intake of 4.7g per day [27] and Italy
253 (mean daily intake of 3.7 g per day [30]). The highest daily intakes are reported for Denmark where
254 a successful health campaign increased whole grains by 72% from an average of 32g/d in 2000-2004
255 to 55g/d in 2011-2012 [31]. Similar efforts are needed to boost whole grain intakes by Irish adults.

257 Differences between consumers and non-consumers of whole grains were investigated in this Irish
258 population, with non-consumers having significantly lower energy-adjusted intakes of dietary fibre,
259 potassium, phosphorus, magnesium and iodine, and higher total fat and MUFA (% energy) compared
260 with consumers of whole grain. In the UK, higher intakes of whole grains were associated with
261 significantly higher intakes of dietary fibre, iron, calcium, vitamin E, potassium, phosphorus,
262 magnesium, thiamin, vitamin B12 and vitamin D [32]. These findings are in line with studies which
263 report a healthier eating pattern in those with higher intakes of whole grains [2, 27, 32, 33]. For
264 example, in a subsample of the Scandinavian cohort 'HELGA', the intake of whole grains was
265 directly associated with intakes of vegetables, fruits, dairy products, fish and shellfish and inversely
266 associated with intakes of white bread, red meat and cakes and biscuits [33]. Similarly, in Irish adults,
267 high whole grain intakes were associated with a higher aMed Score which indicates greater
268 compliance to a Mediterranean style diet and may be viewed a marker of higher dietary quality.

270 In the current study, we completed a simple investigation of strategies to understand patterns of whole
271 grain intake and to increase their consumption in the Irish population as low whole grain intakes were
272 observed. The overwhelming majority of whole grain containing foods were consumed at breakfast
273 suggesting opportunities for encouraging whole grain options at other eating occasions, particularly
274 lunch and dinner. Wheat was the major grain form consumed, implying potential for other grain
275 sources, particularly oats. Further, using regression analysis, it became apparent that simply
276 increasing the quantities of whole grain foods as currently consumed resulted in variable increases in
277 daily whole grain intake. For example, consuming an additional 10g of whole-grain containing
278 RTEBC per day yielded an additional 5.5g whole grain. While 10g extra of whole grain containing
279 'rice, pasta and their dishes' yielded an extra 3.5g whole grain, 10g more of 'bread and rolls' provided
280 an extra 2.75g. Hence simple substitution of food choices has the potential to enhance population
281 whole grain intakes albeit that any such substitution will also be impacted by portion size and amounts
282 usually consumed per serving. We note that the single greatest impact was by consuming 10g extra
283 of the group 'savory biscuits, crackers and popcorn' (7.5g additional whole grain), a food group
284 likely to contain levels of nutrients deemed less acceptable for public health e.g. fat, sugar or salt but
285 typically associated with smaller portion sizes. Overall an opportunity exists for the food industry to
286 develop more whole grain-rich foods with acceptable sensory qualities, yet without concomitant high
287 salt and sugar contents. Use of a universally agreed labelling system for whole grain containing foods,
288 as suggested by the Healthgrain forum [17] may encourage development of such foods while also
289 highlighting to public health advocates and consumers options which are more healthful.

290

291 Currently there are no specific whole grain recommendations set in Ireland, or in the UK, making it
292 is difficult to place in context current intake patterns and the impact of any strategies to increase
293 whole grain intakes. In the absence of such national recommendations, dietary whole grain intakes
294 were compared to the US recommendation of 48g/day [25], with only 19% of individuals in this Irish
295 adult cohort each achieving this guideline. In additional analysis, we also used an alternative approach
296 to assess mean population compliance [34]. Such an approach acknowledges the distribution of intake
297 of any nutrient or ingredient in a population and accepts that dietary guidelines are targeted at
298 populations, rather than individuals. Hence, it is not necessary for all individuals in a population to
299 each achieve this goal, rather that mean population intakes should satisfy the recommendation [34].
300 However, using this approach, there remained poor population compliance with the whole grain
301 recommendation, with less than half (47%) of the entire NANS study population having an overall
302 mean intake of 48g/day. Irrespective of method used, such low intakes, suggests a need for more
303 specific whole grain recommendations which reflect the importance of having a diet high in whole
304 grains for additional benefits above and beyond dietary fibre, as discussed by Seal and colleagues
305 [15]. Observational studies consistently link high whole grain intakes and a reduced risk of CVD [4],
306 type 2 diabetes [5,6], metabolic syndrome [5] and a number of different cancers [8]. It is likely that
307 more specific whole grain recommendations would aid successful public health campaigns to
308 improve whole grain intakes similar to that of Denmark [31]. In the US, a health claim pertaining to
309 whole grain intakes is permitted based on the strength of the evidence to support the benefits of a diet
310 high in whole grains, however, no such health claim is currently approved by EFSA [15]. It is likely
311 that a more specific whole grain recommendation would make it easier to transfer the health message
312 across to consumers due to clearer labelling regarding whole grains on food products. The results of
313 this current study and from previous Irish cohorts in Irish children and teens, highlights the need for
314 a more specific whole grain recommendation and targeted public health campaigns similar to
315 Denmark [17,31].

316

317 This study has some noteworthy strengths such as the large sample size which is representative of the
318 Irish population. Furthermore, dietary data was collected using a 4-day semi-weighed food diary
319 which accounts for potential intra-variation in dietary habits within participants. Furthermore, where
320 possible, food packaging was obtained to ensure a higher level of accuracy regarding the dietary data
321 and the ability to ascertain whole grain composition at a brand level where feasible. The current
322 analysis was conducted following the removal of under-reporters which ensures absence of mis-
323 reporting in the dataset. A limitation of the current study is the lack of a biomarker for whole grain
324 intake to validate the dietary data collected. Alkylresorcinols have been identified as valid biomarkers

325 of whole-grain wheat and rye intake and may be useful for future studies investigating whole grain
326 intake [35].

327

328 In conclusion, this study is the first to report whole grain intakes in Irish adults using data from the
329 national food survey, NANS. We reported low whole grain intakes in Irish adults similar to Irish
330 children and teens and other countries such as the UK and France. We also show how the selection
331 of whole-grain containing RTEC, breads and pastas/rice has the potential to enhance daily intakes.
332 These findings support the argument for the development of a more specific whole grain
333 recommendation in Ireland and universally agreed labelling in light of the proposed health benefits
334 for a diet high in whole grains. Public health campaigns are warranted to try and improve whole grain
335 intakes in the Irish population across all age groups. Future work should also investigate the proposed
336 metabolic health benefits of consuming a diet high in whole grains in this population.

337

338

339 **Ethical Standards**

340 This study was approved by the University College Cork Clinical Research Ethics Committee of the
341 Cork Teaching Hospitals and the Human Ethics Research Committee of University College Dublin
342 (ECM 3 (p) 4 September 2008) and performed in accordance with the ethical standards laid down in
343 the 1964 Declaration of Helsinki and its later amendments. All persons gave their informed consent.

344

345 **Conflict of Interest**

346 APN and MJG have previously received unrestricted research funding from Cereal Partners
347 Worldwide, with no influence in the current analysis or study. There are no other conflicts of interest
348 to declare.

349

350 **References**

- 351 1. Seal CJ, Brownlee IA (2015) Whole-grain foods and chronic disease: evidence from
352 epidemiological and intervention studies. *Proc Nutr Soc* 74: 313-319.
- 353 2. O'Neil CE, Nicklas TA, Zhanovec M, Cho S. (2010) Whole-grain consumption is associated with
354 diet quality and nutrient intake in adults: the National Health and Nutrition Examination Survey,
355 1999-2004. *J Am Diet Assoc* 110: 1461-1468.
- 356 3. Reicks M, Jonnalagadda S, Albertson AM, Joshi N. (2014) Total dietary fiber intakes in the US
357 population are related to whole grain consumption: results from the National Health and Nutrition
358 Examination Survey 2009 to 2010. *Nutr Res* 34: 226-234.

359 4. Mellen PB, Walsh TF, Herrington DM (2008) Whole grain intake and cardiovascular disease: a
360 meta-analysis. *Nutr Metab Cardiovasc Dis* 18: 283-290.

361 5. Ye EQ, Chacko SA, Chou EL, Kugizaki M, Liu S. (2012) Greater whole-grain intake is associated
362 with lower risk of type 2 diabetes, cardiovascular disease, and weight gain. *J Nutr* 142, 1304-1313.

363 6. Aune D, Norat T, Romundstad P, Vatten LJ. (2013) Whole grain and refined grain consumption
364 and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies.
365 *Eur J Epidemiol* 28: 845-858.

366 7. Aune D, Chan DS, Lau R, Vieira R, Greenwood DC, Kampman E, Norat T. (2011) Dietary fibre,
367 whole grains, and risk of colorectal cancer: systematic review and dose-response meta-analysis of
368 prospective studies. *BMJ* 343: d6617.

369 8. Jacobs DR, Jr., Marquart L, Slavin J, Kushi LH. (1998) Whole-grain intake and cancer: an
370 expanded review and meta-analysis. *Nutr Cancer* 30: 85-96.

371 9. Ross AB, Kristensen M, Seal CJ, Jacques P, McKeown N. (2015) Recommendations for reporting
372 whole-grain intake in observational and intervention studies. *Am J Clin Nutr* 101: 903-907.

373 10. Borneo R, Leon AE (2012) Whole grain cereals: functional components and health benefits. *Food*
374 *Funct* 3: 110-119.

375 11. Slavin J (2004) Whole grains and human health. *Nutr Res Rev* 17: 99-110.

376 12. Tighe P, Duthie G, Vaughan N, Brittenden J, Simpson WG, Duthie S, Mutch W, Wahle K, Horgan
377 G, Thies F. (2010) Effect of increased consumption of whole-grain foods on blood pressure and other
378 cardiovascular risk markers in healthy middle-aged persons: a randomized controlled trial. *Am J Clin*
379 *Nutr* 92: 733-740.

380 13. Kristensen M, Toubro S, Jensen MG, Ross AB, Riboldi G, Petronio M, Buge S, Tetens I, Astrup
381 A. (2012) Whole grain compared with refined wheat decreases the percentage of body fat following
382 a 12-week, energy-restricted dietary intervention in postmenopausal women. *J Nutr* 142: 710-716.

383 14. Lefevre M, Jonnalagadda S (2012) Effect of whole grains on markers of subclinical inflammation.
384 *Nutr Rev* 70: 387-396.

385 15. Seal CJ, Nugent AP, Tee ES, Thielecke F. (2016) Whole-grain dietary recommendations: the need
386 for a unified global approach. *Br J Nutr* 115, 2031-2038.

387 16. Ferruzzi MG, Jonnalagadda SS, Liu S, Marquart L, McKeown N, Reicks M, Riccardi G, Seal C,
388 Slavin J, Thielecke F, van der Kamp J, Webb D (2014) Developing a standard definition of whole-
389 grain foods for dietary recommendations: summary report of a multidisciplinary expert roundtable
390 discussion. *Adv Nutr* 5: 164-176.

391 17. Ross AB, Van der Kamp JW, King R, Le KA, Mejbourn H, Seal CJ, Thielecke F, Healthgrain
392 Forum. (2017) Perspective: a definition for whole-grain food products – recommendations from the
393 Healthgrain forum. *Adv Nutr* 8: 5252-31.

394 18. Devlin NF, McNulty BA, Gibney MJ, Thielecke F, Smith H, Nugent AP. (2013) Whole grain
395 intakes in the diets of Irish children and teenagers. *Br J Nutr* 110: 354-362.

396 19. Cashman KD, Muldowney S, McNulty B, Nugent A, FitzGerald AP, Kiely M, Walton J, Gibney
397 MJ, Flynn A. (2013) Vitamin D status of Irish adults: findings from the National Adult Nutrition
398 Survey. *Br J Nutr* 109: 1248-1256.

399 20. Fung TT, McCullough ML, Newby PK, Manson JE, Meigs JB, Rifani N, Willett WC, Hu FB.
400 (2005) Diet-quality scores and plasma concentrations of markers of inflammation and endothelial
401 dysfunction. *Am J Clin Nutr* 82: 163-173.

402 21. Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Coward AW, Prentice AM. (1991)
403 Critical evaluation of energy intake data using fundamental principles of energy physiology: 1.
404 Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr* 45: 569-581.

405 22. van der Kamp JW, Poutanen K, Seal CJ, Richardson DP (2014) The HEALTHGRAIN definition
406 of 'whole grain'. *Food Nutr Res* 58: doi: 10.3402/fnr.v58.22100.

407 23. Jones AR, Mann KD, Kuznesof SA, Richardson DP, Seal CJ (2017) The whole grain content of
408 foods consumed in the UK. *Food Chem* 214: 453-459.

409 24. Holland B, Unwin I, Buss D (1988) Cereals and cereal products: The third supplement to
410 McCance & Widdowson's the composition of foods. 4th ed. Nottingham.

411 25. US Department of Agriculture; US Department of Health and Human Services (2015) Nutrition
412 and your health: Dietary Guidelines for Americans, 2015-2020, 8th edn. Government Printing Office,
413 Washington DC.

414 26. Mann KD, Pearce MS, McKeivith B, Thielecke F, Seal CJ. (2015) Low whole grain intake in the
415 UK: results from the National Diet and Nutrition Survey rolling programme 2008-11. *Br J Nutr* 113:
416 1643-1651.

417 27. Bellisle F, Hebel P, Colin J *et al.* (2014) Consumption of whole grains in French children,
418 adolescents and adults. *Br J Nutr* 112: 1674-1684.

419 28. Galea LM, Beck EG, Probst YC, Cashman CJ. (2017) Whole Grain intake of Australians
420 estimated from a cross-sectional analysis of dietary intake data from the 2011 – 13 Australian Health
421 Survey. *Public Health Nutr* 20 (12): 2166-2172.

422 29. McGill CR, Fulgoni VL, Devareddy L (2015) Ten-year trends in fibre and whole grain intakes
423 and food sources for the United States Population: National Health and Nutrition Examination Survey
424 2001-2010. *Nutrients* 7: 1119-1130.

425 30. Sette S, D'Addezio L, Piccinelli Ri, Hopkins S, Le Donne S, Ferrari M, Mistura L, Turrini A
426 (2017) Intakes of whole grain in an Italian sample of children, adolescents and adults. *Eur J Nutr* 56
427 (2): 521-533.

428 31. Kyro C, Tjonneland A (2016) Whole grains and public health. *BMJ* 353, i3046.

- 429 32. Mann KD, Pearce MS, McKeivith B, Thielecke F, Seal CJ. (2015) Whole grain intake and its
430 association with intakes of other foods, nutrients and markers of health in the National Diet and
431 Nutrition Survey rolling programme 2008-11. *Br J Nutr* 113: 1595-1602.
- 432 33. Kyro C, Skeie G, Dragsted LO, Christensen J, Overvad K, Hallmans G, Johansson I, Lund E,
433 Slimani N, Johnsen NF, Halkjær J, Tjønneland A, Olsen A. (2011) Intake of whole grains in
434 Scandinavia is associated with healthy lifestyle, socio-economic and dietary factors. *Public Health*
435 *Nutr* 14: 1787-1795.
- 436 34. Wearne SJ, Day MJ (1999) Clues for the development of food-based dietary guidelines: how are
437 dietary targets being achieved by UK consumers? *Br J Nutr* 81 Suppl 2: S119-126.
- 438 35. Landberg R, Kamal-Eldin A, Andersson A, Vessby B, Aman P. (2008) Alkylresorcinols as
439 biomarkers of whole-grain wheat and rye intake: plasma concentration and intake estimated from
440 dietary records. *Am J Clin Nutr* 87: 832-838.

441

442 **List of Figures**

443

444 **Fig. 1** Contribution (%) of food group categories to mean daily whole grain intakes in whole
445 grain consumers

446

447

CONFIDENTIAL

Table 1 Description of the foods included in each of the 7 whole grain food groups

Food Group	Foods included
Rice, pasta and their dishes	Brown and whole grain rice, wholemeal pasta, risottos, pasta bakes, lasagne, pasta and rice salads
Breads and rolls ^a	Wholemeal, brown, brown soda and granary breads and rolls. Other whole grain breads including scones, pitta breads and bread mix.
RTEBC ^b	All ready to eat breakfast cereals
Other breakfast cereals	Porridge and cooked breakfast cereals
Sweet biscuits, cereal bars and desserts	Plain and chocolate digestive biscuits, oat based biscuits, chewy, crunchy and baked cereal bars, flapjacks, cheesecake and biscuit cake.
Savoury biscuits, crackers and popcorn	Oatcakes, crackers, shop bought, microwaveable, toffee, butter and sweet popcorn
Meat products, tofu and yoghurts	White and black pudding, tofu products, cereal based yoghurts

^aAll breads and rolls were cross-checked at brand level and with manufacturers to ensure ingredients were whole grain containing.

^bRTEBC, ready to eat breakfast cereals

448

449

Table 2 Descriptive analysis of whole grain intake (g/d) for Irish adults (total population and whole grain consumers only)

	Total Population (n = 1051)							Consumers Only (n = 957; 91.1%)						
	n	%	Mean	Median	SD	P97.5	P	n	%	Mean	Median	SD	P97.5	P
Total Population	1051	100	27.8	20.0	29.4	102.7		957	91.1	30.6	22.5	29.5	108.8	
Sex														
Male	523	49.8	31.6	18.5	35.7	130.4	***	473	49.4	35.0	23.8	35.9	142.1	***
Female	528	50.2	24.1	20.4	20.9	76.1		484	50.6	26.3	21.6	20.5	78.0	
Age														
Age group 1 [†]	377	35.9	29.5	21.0	33.2	118.7	NS	335	35.0	33.2	23.7	33.4	120.3	NS
Age group 2 [†]	308	29.3	27.5	20.6	27.7	96.6		280	29.3	30.3	23.8	27.6	97.2	
Age group 3 [†]	204	19.4	27.2	20.1	27.0	85.3		191	20.0	29.1	21.4	27.0	86.4	
Age group 4 [†]	162	15.4	25.4	15.8	26.0	94.3		151	15.8	27.2	19.7	26.0	96.3	
Social Class^v														
Professional /managerial	484	46.1	31.1 ^a	22.9	30.0	109.3	*	450	47.0	33.5 ^a	24.6	29.9	110.5	*
Non-manual	183	17.4	25.6 ^{a,b}	16.4	25.3	90.3		169	17.7	27.7 ^{a,b}	21.0	25.2	90.7	
Skilled manual	142	13.5	22.7 ^b	15.3	23.5	84.3		125	13.1	25.8 ^b	20.9	23.4	86.8	
Semi-skilled and unskilled	205	19.5	27.0 ^{a,b}	15.5	34.7	121.6		178	18.6	31.1 ^{a,b}	20.1	35.5	132.3	
Eating Location														
Home	1050	99.9	24.6 ^b	16.5	27.7	96.2	***	956	99.9	27.0 ^b	19.1	27.8	98.1	***
Other Home	306	29.1	1.7 ^a	0.0	4.9	16.2		285	29.8	1.8 ^a	0	5.1	16.6	
Outside Home	862	82.0	3.4 ^a	0.0	8.0	25.8		784	81.9	3.7 ^a	0	8.3	28.0	

^{abc}Values with unlike superscript letters denote significant differences between mean daily intakes of whole grain (ANOVA with Scheffé post hoc tests): *P<0.05, **P<0.01,

***P<0.001, NS: not significant (P≥0.05). [†]Age group 1: 18-35 years, Age group 2: 36-50 years, Age group 3: 51-64 years, Age group 4: 65-90 years. ^vExcluding data for 37 students (missing data)

Table 3. Percentage of Irish adults meeting US whole grain recommendations of 48g/day [25]

Total population <i>n</i> 1051	
	(%) ^a
All adults (18-90 yrs)	19.3
Adults (18-35 yrs)	21.0
Adults (36 – 50 yrs)	18.8
Adults (51 – 64 yrs)	16.7
Adults (65 – 90 yrs)	19.8

^a refers to the percentage of individuals in the Irish population who met the dietary recommendations for minimal whole grains of 48g/day [25].

Table 4 Intake (g/10MJ/day) and contribution (%) of food groups to mean daily whole grain intakes in an Irish adult population across tertiles of intake in whole grain consumers

	Tertiles of mean daily whole grain intakes (g/10MJ/d)																
	Whole grain consumers				Tertile 1				Tertile 2				Tertile 3				P
					n=319				n=319				n=319				
	n=957				n=319				n=319				n=319				
%Cons	Mean	SD	%Ctr	%Cons	Mean	SD	%Ctr	%Cons	Mean	SD	%Ctr	%Cons	Mean	SD	% Ctr		
Rice, pasta & their dishes	6.6	17.9	11.3	2.3	0.9	6.8	4.9	0.5	4.4	12.8	6.8	2.1	14.4	20.1	11.8	4.4	NS
Bread & rolls	76.1	21.1	20.8	46.6	54.9	6.9 ^a	5.0	40.9	83.7	17.5 ^b	12.3	52.7	89.7	33.1 ^c	25.8	46.4	***
Ready to eat breakfast cereals	46.4	22.8	18.4	25.6	16.9	6.6 ^a	4.3	11.6	49.2	16.4 ^b	12.1	28.4	73.0	30.9 ^c	19.7	36.8	***
Other breakfast cereals	32.5	5.2	12.7	7.4	31.7	2.1 ^a	1.3	14.2	37.0	2.9 ^{a,b}	5.2	3.8	28.8	11.7 ^b	21.3	4.3	***
Sweet biscuits, cereal bars & desserts	33.6	3.3	4.5	6.6	31.7	2.4	2.8	13.3	32.3	2.9	3.2	3.9	37.0	4.3	6.3	2.6	*
Savoury biscuits, crackers & popcorn	15.8	11.9	10.7	6.4	9.4	6.1 ^a	4.3	6.5	17.6	13.4 ^b	11.3	8.1	20.4	13.4 ^{b,c}	11.3	4.6	*
Meat products, tofu & yoghurts	15.9	2.7	2.2	5.0	21.3	2.9	2.3	13.1	11.0	2.4	1.7	1.2	15.4	2.7	2.4	0.9	NS

% Cons - % consumers of each whole grain food group; % Ctr -% contribution of food groups to whole grain intakes within each tertile

^{abc} Values with unlike superscript letters denote significant differences between the mean values of whole grain intake across the tertiles of whole grain intake (adjusted for gender)

(ANCOVA with Bonferroni correction): * P<0.05; ** P<0.01; *** P<0.001; NS, P≥0.05

Table 5 Patterns of intake (%) of all whole grain and of whole grain containing food groups as consumed at meals and snacks

	Breakfast	Lunch	Dinner	Snacks
All sources of whole grain	49.5	22.7	10.3	17.3
<i>Of which</i>				
Rice, pasta and their dishes	0.0	2.2	11.7	0.0
Bread and rolls	35.5	79.6	58.0	44.2
RTEBC	39.8	1.1	0.8	8.1
Other breakfast cereals	18.0	0.1	0.1	1.3
Sweet biscuits, cereal bars and desserts	2.0	7.3	16.2	29.5
Savoury biscuits, crackers and popcorn	1.2	5.4	6.0	15.3
Meat, tofu and yoghurts	3.3	4.5	7.1	1.7

Table 6 Mean energy and nutrient intakes for non-consumers and across the tertiles of whole grain intake in Irish adults

	Non-consumers (n=94)		Tertile 1 (n=319)		Tertile 2 (n=319)		Tertile 3 (n=319)		P value
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Energy (MJ)	8.9 ^a	2.3	9.2 ^a	2.7	9.0 ^a	2.4	10.1 ^b	2.7	***
Protein (% energy)	15.7	3.9	16.3	3.3	16.4	3.2	16.9	3.5	NS
Carbohydrate (% energy)	40.2 ^a	7.7	41.8 ^{ad}	6.8	42.6 ^{bd}	6.9	44.6 ^d	6.1	***
Total sugars (% energy)	15.4	6.4	17	6.2	17.2	5.8	17.4	4.9	NS
Fat (% energy)	35.7 ^a	7	35.0 ^a	6.1	34.0 ^{ab}	6.4	33.2 ^b	5.9	**
Saturated fat (% energy)	13.7	4.3	13.9	3.3	13.4	3.4	13	3.2	NS
MUFA (% energy)	13.5 ^a	3	12.8 ^a	2.6	12.2 ^b	2.7	12.0 ^b	2.5	***
PUFA (% energy)	6	2.3	6	2.3	6.2	2.5	6	2.1	NS
Dietary fibre (g/10MJ)	16.5 ^a	6.3	19.4 ^b	6.5	23.4 ^c	7.2	26.9 ^d	7.2	***
Sodium (mg/10MJ)	2892.9	616.2	2942.4	675.9	2914.5	653.3	2913.5	598.3	NS
Iron (mg/10MJ)	16.4	21.8	17.2	22.2	15.7	16.3	19.8	25.1	NS
Calcium (mg/10MJ)	1062.8	574	1071.4	406.5	1161	453.8	1163.6	402.7	NS
Magnesium (mg/10MJ)	295.8 ^a	106.3	316.9 ^a	115.6	349.2 ^b	83.8	382.5 ^c	82	***
Potassium (mg/10MJ)	3478.7 ^{ab}	1467.7	3444.7 ^b	741.1	3666.1 ^a	713.8	3732.3 ^a	817	**
Phosphorus (mg/10MJ)	1453.3 ^a	276.9	1548.3 ^b	308	1648.4 ^c	302.4	1723.6 ^d	295.1	***
Iodine (µg/10MJ)	160.5 ^a	96.9	165.7 ^a	83.5	178.8 ^{ab}	86.4	189.3 ^b	81.5	***
Vitamin D (µg/10MJ)	4.1	5.5	5.0	6.0	5.8	7.5	6.2	9.3	NS
Thiamin (mg/10MJ)	2.3	4.3	3.8	10.9	3.3	8.3	3.9	10.4	NS
Riboflavin (mg/10MJ)	2.6	4	3.7	10	3.8	8.9	4.1	9.7	NS
Niacin (mg/10MJ)	32.1	31.6	30.7	15.9	33.6	41.3	35	19.5	NS
Vitamin B6 (mg/10MJ)	3.7	4.2	4.5	8.5	4.8	9.7	5.1	10.1	NS
Vitamin B12 (µg /10MJ)	6.7	5.8	9.3	23.9	11.3	69.5	8.2	11.4	NS
Folate (µg/10MJ)	354.2 ^a	184.8	393.3 ^a	201.3	425.0 ^{ac}	206.9	461.9 ^{bc}	207.0	***
aMED	2.0 ^a	1.1	2.5 ^b	1.3	3.4 ^c	1.6	3.9 ^d	1.5	***

aMED, alternative Mediterranean diet score. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, NS, not significant ($P \geq 0.05$). ^{abc} Values with unlike superscript letters denote significant differences across the tertiles of whole grain intake (adjusted for gender) (ANCOVA with Bonferroni correction)

CONFIDENTIAL

Table 7 Associations (unstandardised coefficients) of whole grain intake (g/10MJ/d) to predict the importance of whole grain food groups to increase whole grain intakes

Whole grain food group	β	SE	P
Savoury biscuits, crackers & popcorn	0.755	0.058	<0.001
RTEBC	0.549	0.017	<0.001
Rice, pasta & their dishes	0.352	0.030	<0.001
Bread & rolls	0.275	0.007	<0.001
Sweet biscuits, cereal bars & desserts	0.166	0.032	<0.001
Other breakfast cereals	0.014	0.005	0.002

Note: The unstandardised coefficient for whole grain ready-to-eat breakfast cereals (RTEBC) is 0.549 meaning that for each 1g increase, there would be an increase in whole grain consumption of 0.549. Therefore, if an individual was to add 10g of a RTEBC to the daily diet, the potential impact would be an increase of 5.4g in daily whole grain intakes.