**QUEEN'S
UNIVERSITY
BELFAST**

Absence of CALR Mutations in Idiopathic Erythrocytosis Patients with Low Serum Erythropoietin Levels

Catherwood, M. A., Graham, A., Cuthbert, R. J. G., Garrec, C., Gardie, B., Girodon, F., Laird, S., Cross, N. C. P., & McMullin, M. F. (2018). Absence of CALR Mutations in Idiopathic Erythrocytosis Patients with Low Serum Erythropoietin Levels. *Acta haematologica*, 139(4), 217-219. <https://doi.org/10.1159/000489006>

Published in:
Acta haematologica

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Absence of *CALR* mutations in idiopathic erythrocytosis patients with low serum erythropoietin levels

MA Catherwood¹, A Graham¹, RJG Cuthbert¹, C Garrec^{2,3}, B Gardie^{2,3}, F Girodon^{4,5}, S Laird⁶, NCP Cross⁶, MF McMullin^{1,7}

¹Clinical Haematology, Belfast City Hospital, BHSCT Belfast, Northern Ireland.

²Ecole Pratique des Hautes Etudes (EPHE), PSL Research University, France.

³INSERM U892, CNRS 6299, Université de Nantes, France.

⁴Service d'Hématologie Biologique CHU Dijon, France.

⁵INSERM, UMR866, University of Burgundy Franche-Comté, Dijon, France.

⁶Wessex Regional Genetics Laboratory, Salisbury, UK

⁷Centre for Medical Education (CME), Queen's University Belfast, Northern Ireland.

Correspondence to:

Prof. Mary Frances McMullin, Clinical Haematology, Belfast City Hospital, BHSCT Belfast, BT9 7AB. Northern Ireland.

Tel 02895048008: FAX: 02895048007: e-mail:m.mcmullin@qub.ac.uk

Key words: Idiopathic erythrocytosis, JAK2, calreticulin, erythropoietin

Short Title: *CALR* mutations in idiopathic erythrocytosis

Idiopathic erythrocytosis (IE) is a heterogeneous collection of rare haematologic disorders which can be either sporadic or familial in origin (McMullin *et al*, 2005). An erythrocytosis occurs when there is an increased red cell mass and an elevated haematocrit. IE is distinct from polycythaemia vera (PV) as the red cell hyperplasia present in IE is not accompanied by elevations in the megakaryocytic or granulocytic lineages and PV is characterized by low serum erythropoietin (Epo) levels, whereas IE is associated with a wide range of Epo levels. IE can be divided into two main groups. The first group consists of those with a low EPO level suggesting an unidentified molecular defect possibly involving components of the Epo signal transduction pathway (de la Chapelle *et al*, 1993). The second group are those with inappropriately normal or raised EPO level which suggests a secondary unidentified cause possibly affecting the oxygen sensing pathway. These patients are rarely observed in clinical practice, and little is known regarding their clinical characteristics, natural history and best management.

Recently, novel frameshift mutations in exon 9 of the calreticulin (*CALR*) gene were found in patients with *JAK2* or *MPL* unmutated primary myelofibrosis and essential thrombocythemia (ET) (Klampfl *et al*, 2013; Nangalia *et al*, 2013). Even though mutations of *CALR* have been reported in significant numbers in MPN it was initially thought that *CALR* mutations did not occur in PV. However in 2014 two *JAK2* negative PV patients were described that harboured *CALR* mutations (52 bp deletion) (Broseus *et al*, 2014) with further cases described in other studies (Chauveau *et al*, 2017).

The aims of this study were to assess the prevalence of *CALR* mutations in (i) a well characterized cohort of IE patients with low serum erythropoietin levels and (ii) a broad cohort of query myeloproliferative neoplasm (MPN) cases referred for molecular genetic investigation.

Initially we studied 79 patients with IE. Thirty eight were enrolled from a UK cohort and 41 from a French cohort.

Ethical approval was granted by the local review committees and informed consent was collected according to the Helsinki Declaration.

All the patients had low to normal serum EPO level (Table 1) and none of them carried *JAK2*V617F or *JAK2* exon 12 mutations as determined by allele specific PCR. Clinical and laboratory details of the patients are shown in Table 1. The procedures followed were in accordance with the Declaration of Helsinki and all patients gave informed written consent. *CALR* exon 9 mutations were screened by fragment length analysis according to Klampfl *et al*, 2013.

The patients were mostly males in both the groups with a higher frequency in the French cohort. Patients in both cohorts had similar characteristics (Table 1). No *CALR* mutations were detected in any of the 79 IE patients included in this study with low to normal EPO levels.

To further explore the possibility of *CALR* mutations in PV we audited samples referred for MPN panel analysis at the Wessex Regional Genetics Laboratory (WRGL) in Salisbury (Jones *et al*, 2015). Since October 2016 all query MPN referrals to the WRGL have been tested using an amplicon-based next generation sequencing approach for *JAK2* V617F, *JAK2* exon 12, *MPL* exon 10 and *CALR* exon 9 mutations irrespective of their phenotype. Of 2306 referrals, 137 (5.9%) tested positive for *CALR* +1 frameshift mutations. None of the 659 referrals that specified an erythroid phenotype PV, ?PV, high haematocrit or high haemoglobin) tested positive for *CALR* mutations.

In previous studies 4 *CALR* mutations have been identified in so called *JAK2* negative PV in patients more accurately described as *JAK2* negative unexplained erythrocytosis. In the first study only 2 patients were studied (Broseus *et al*, 2014). Both cases were *JAK2* negative PV and the *CALR* mutation was a 52bp deletion, commonly termed type 1 mutation. Of note, both patients had thrombocytosis in addition to erythrocytosis. In the second study a large cohort of 578 patients were examined (Chauveau *et al*, 2017). Of the two patients with *CALR* mutations one was subsequently considered to have ET with slightly increased red cell mass and the other considered to have idiopathic erythrocytosis. Both cases harboured the 52bp deletion. Of interest, in the patient with idiopathic erythrocytosis the *CALR* mutation was detected with a low allele burden of approximately 5%.

The underlying defect in cases with unexplained erythrocytosis provides a clinical conundrum. Our findings indicate that *CALR* mutations are absent in a cohort of 79 patients with IE and also in a large, broad real-life cohort of patients with an erythroid phenotype referred for MPN panel testing. We therefore consider that the screening of *CALR* mutations in this situation is not warranted as a standalone investigation.

“Disclosure Statement”

The authors declare no conflicts of interest.

References

1. McMullin MF, Bareford D, Campbell P, Green AR, Harrison C, Hunt B, Oscier D, Polkey MI, Reilly JT, Rosenthal E, Ryan K, Pearson TC, Wilkins B; General Haematology Task Force of the British Committee for Standards in Haematology. Guidelines for the diagnosis, investigation and management of polycythaemia/erythrocytosis. *Br J Haematol*, 2005;**130**:174–195
2. de la Chapelle A, Träskelin A-L, Juvonen E (1993). Truncated erythropoietin receptor causes dominantly inherited human erythrocytosis. *Proc Natl Acad Sci*; 1993;**90**:4495-4499.
3. Klampfl T, Gisslinger H, Harutyunyan AS, Nivarthi H, Rumi E, Milosevic JD, Them NC, Berg T, Gisslinger B, Pietra D, Chen D, Vladimer GI, Bagienski K, Milanesi C, Casetti IC, Sant'Antonio E, Ferretti V, Elena C, Schischlik F, Cleary C, Six M, Schalling M, Schönegger A, Bock C, Malcovati L, Pascutto C, Superti-Furga G, Cazzola M, Kralovics R. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med* 2013;**369**:2379-2390.
4. Nangalia J, Massie CE, Baxter EJ, Nice FL, Gundem G, Wedge DC, Avezov E, Li J, Kollmann K, Kent DG, Aziz A, Godfrey AL, Hinton J, Martincorena I, Van Loo P, Jones AV, Guglielmelli P, Tarpey P, Harding HP, Fitzpatrick JD, Goudie CT, Ortmann CA, Loughran SJ, Raine K, Jones DR, Butler AP, Teague JW, O'Meara S, McLaren S, Bianchi M, Silber Y, Dimitropoulou D, Bloxham D, Mudie L, Maddison M, Robinson B, Keohane C, Maclean C, Hill K, Orchard K, Tauro S, Du MQ, Greaves M, Bowen D, Huntly BJ, Harrison CN, Cross NC, Ron D, Vannucchi AM, Papaemmanuil E, Campbell PJ, Green AR. Somatic CALR mutations in myeloproliferative neoplasms with nonmutated JAK2. *N Engl J Med* 2013;**369**:2391-2405.
5. Jones AV, Ward D, Lyon M, Leung W, Callaway A, Chase A, Dent CL, White HE, Drexler HG, Nangalia J, Mattocks C, Cross NC. Evaluation of methods to detect CALR mutations in myeloproliferative neoplasms. *Leuk Res*. 2015;**39**:82-87.
6. Broséus J, Park JH, Carillo S, Hermouet S, Girodon F. Presence of calreticulin mutations in JAK2 negative polycythemia vera. *Blood* 2014;**124**:3964-3966.
7. Chauveau A, Nibourel O, Tondeur S, Paz DL, Mansier O, Paul F, Wemeau M, Preudhomme C, Lippert E, Ugo V; French Intergroup of Myeloproliferative Neoplasms (2017). Absence of CALR mutations in JAK2-negative polycythemia. *Haematologica* 2017;**102**:e15-e16.
8. Percy MJ, Scott LM, Erber WN, Harrison CN, Reilly JT, Jones FG, Green AR, McMullin MF. *Haematologica* 2007;**92**:1607-1614.

Authorship

AG analysed the data, wrote the paper and performed the molecular analysis. MC, AG, SL and BG performed the molecular analysis. RJGC contributed clinical data and followed the patients. MFM, NCPC and FG designed the study, recruited patients and critically reviewed the manuscript. All authors have read and approved the final version.

Table 1. Patients characteristics.

UK

	Mean	Range/percentage
Sex, female:male	11:27	29% female
Age	54	27-75
Haemoglobin level, g/L	198	169-249
Hematocrit, %	57	49-79
Platelet count, 10 ⁹ /L	220	119-467
White blood cells, 10 ⁹ /L	8.3	4.2-15.6
Erythropoietin level, mIU/mL	5.8	1-9.9

FRENCH

	Mean/ratio	Range/percentage
Sex, female:male	3:38	5% female
Age	48	13-79
Haemoglobin level, g/L	186	161-230
Hematocrit, %	52.7	46-66
Platelet count, 10 ⁹ /L	215	139-350
White blood cells, 10 ⁹ /L	7.7	5.34-9.14
Erythropoietin level, mIU/mL	11.35	1.7-30

BOTH

	Mean/ratio	Range/percentage
Sex, female:male	14:65	18% female
Age	51	13-79
Haemoglobin level, g/L	192	161-249
Hematocrit, %	54	46-79
Platelet count, 10 ⁹ /L	218.5	119-467
White blood cells, 10 ⁹ /L	8	4.2-15.6
Erythropoietin level, mIU/mL	8.6	1.7-30