**QUEEN'S
UNIVERSITY
BELFAST**

Microclimate has a greater influence than macroclimate on the availability of infective *Haemonchus contortus* larvae on herbage in a warmed temperate environment

Wang, T., Vineer, H. R., Morrison, A., van Wyk, J. A., Bolajoko, M-B., Bartley, D. J., & Morgan, E. R. (2018). Microclimate has a greater influence than macroclimate on the availability of infective *Haemonchus contortus* larvae on herbage in a warmed temperate environment. *Agriculture, Ecosystems and Environment*, 265, 31-36. <https://doi.org/10.1016/j.agee.2018.05.029>

Published in:
Agriculture, Ecosystems and Environment

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

1 **Microclimate has a greater influence than macroclimate on the**
2 **availability of infective *Haemonchus contortus* larvae on herbage in a**
3 **warmed temperate environment**

4
5 Tong Wang^{1,2}, Hannah Rose Vineer¹, Alison Morrison³, Jan A. van Wyk⁴,

6 Muhammad-Bashir Bolajoko¹, David J. Bartley³, Eric R. Morgan^{1,5*}

7 ¹ University of Bristol, School of Biological Sciences, Bristol Life Sciences Building,
8 24 Tyndall Avenue, Bristol BS8 1TQ, United Kingdom.

9 ² Faculty of Veterinary Medicine, University of Calgary, 2500 University Drive NW,
10 Calgary, AB, T2N 1N4, Canada.

11 ³ Moredun Research Institute, Pentlands Science Park, Bush Loan, Penicuik EH26 0PZ,
12 United Kingdom.

13 ⁴ Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University
14 of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa.

15 ⁵ School of Biological Sciences, Queen's University Belfast, 97 Lisburn Road, Belfast,
16 BT9 7BL, United Kingdom.

17 *Corresponding author: eric.morgan@bristol.ac.uk

Abstract

Global climate change is altering epidemiological patterns of gastrointestinal nematode infections in grazing livestock, including through effects of temperature and moisture on the availability of infective third-stage larvae (L3). While considerable experimental effort has been devoted to the influences of climate on L3 development and survival in major nematode species, knowledge of effects on L3 migration out of faeces and onto herbage is more limited. In this study, we examined elements of this process for *Haemonchus contortus* in controlled and natural climates. The effect of temperature on migration rate from faeces was quantified and found to peak at 15 °C. In glasshouses, a 3 °C difference in mean temperature failed to produce a statistically significant difference in the number of L3 reaching herbage after a single rainfall event, and faecal moisture content (FMC) did not decline significantly more rapidly at the higher temperatures. Most larvae left the faeces and reached the grass within 3 hours after simulated rainfall. On natural pasture in temperate summer, FMC was strongly affected by microclimate, with shade and long grass both significantly slowing drying. Results suggest that microclimate is important in determining FMC and larval migration, and that its effects can be greater than those of macroclimate, e.g. moderate differences in average ambient temperature. More work is needed to develop a full predictive understanding of larval availability in natural settings, which is the product of interacting factors acting on overlapping parasite cohorts.

Key words: nematode, parasite, climate change, transmission, epidemiology, faecal moisture content, temperature, rainfall, migration

1. Introduction

The abomasal nematode parasite *Haemonchus contortus* causes major problems to the health and productivity of sheep and goats, especially in tropical and sub-tropical areas with adequate rainfall (Waller, 1997; Perry and Randolph, 1999; O'Connor et al., 2006; van Dijk et al., 2010; Besier et al., 2016). Improved conditions for transmission are further predicted in temperate areas under climate change scenarios (Rose et al., 2016), where resistance to anthelmintic drugs threatens to undermine control (Rose et al. 2015a). Temperature and moisture have long been known to be important for the development, survival and translation of infective third stage *H. contortus* larvae (L3) onto herbage (Veglia, 1915; Rossanigo and Gruner, 1995; O'Connor et al., 2007; 2008). Previous experiments have attempted to characterize the effects of individual climatic factors on components of the *H. contortus* life cycle under controlled conditions (e.g. Hsu & Levine, 1977; Coyne & Smith, 1992), and relationships between climate and infection pressure have been widely observed in the field (e.g. Rose 1963; Onyali et al., 1990; Troell et al., 2005; Silva et al., 2008).

How temperature and rainfall interact to influence L3 migration out of faeces has received less attention (Todd et al., 1976; Hsu & Levine, 1977, Pandey et al., 1989; Van Dijk and Morgan, 2011; Wang et al., 2014). Under climate change scenarios, variability in temperature and rainfall is predicted to increase (Mason et al., 1999; Pinault, 2012), so the effect of short-term climatic conditions on larval availability, e.g. following brief rainfall events, and the modifying influence of pasture microclimate, might significantly alter transmission patterns. Increased temperatures could hypothetically

accelerate larval movement out of faeces, while simultaneously leading to faster drying of faeces and thus a shorter period of migration. The net effect of warmer temperatures on migration is therefore hard to predict.

This study aims to determine how temperature and rainfall interact to influence migration of infective *H. contortus* larvae from faeces onto grass, and how this is modified by variation in microclimate. Hence, to assess the likely net impact of realistic gradual climate warming on L3 availability in temperate environments. The overall objective is to improve ability to predict infection patterns under climate change, in order to inform farm adaptation strategies (Morgan and Wall 2009), as part of wider attempts to understand ecological responses of parasites to global change (Cable et al. 2017).

2. Materials and methods

2.1 General methods

2.1.1 Mono-infected faecal samples

Faecal samples from mono-cultures of MHco3(ISE) isolate *H. contortus* in sheep (Rose Vineer et al., 2017) were provided by the Moredun Research Institute (Edinburgh, U.K.). Briefly, lambs were born and reared indoors to ensure that they were free of pre-existing nematode infections. Donor lambs were drenched orally by syringe with 5,000 L3 in a suspension of tap water. Faecal egg counts (FEC) were checked 21 days later, and individual sheep with sufficient egg output were harnessed and a bag attached to collect faeces over a 24-hour period. Faecal samples were placed in a sealed plastic bag

and sent by post to the laboratory in Bristol, to arrive within 24 hours. A modified McMaster method (MAFF, 1986) was used to verify estimated egg density on arrival and ensure that development of eggs was not discernible before starting the experiments.

2.1.2 Larval culture and recovery

For Experiments 1 and 2 below, L3 were prepared as follows. Faeces containing *H. contortus* eggs were placed in six-well plates (Sterilin, U.K.) with a loose lid and incubated at 20 °C for seven days (Versatile Environmental Test Chamber, Sanyo, Japan). Lids were removed briefly each day to ensure enough air exchange for development, without the danger of excessive drying. Faeces containing fully developed L3 were used for the experiments, which aimed to test larval emergence from faeces. Following the various treatments, L3 were harvested from cultured faeces using a modified Baermann technique (Gruner, 1986), and species identity confirmed using the identification keys described in Van Wyk & Mayhew (2013).

2.1.3 Larval extraction from herbage

The method used in Experiment 2 to recover L3 from herbage was derived from several different studies (Martin et al., 1990; Couvillion, 1993; Van Bezooijen, 2006). Grass was cut using scissors at the base of the plant, as close to the soil as possible without including soil or roots, and immediately wrapped in a square of woven cotton muslin cloth, forming a loose ball, and the muslin secured with a clip. A 250 ml inverse conical flask was filled with water until it reached about 1cm below the rim, a few drops

of detergent were added, and herbage samples were submerged in the water. After standing for 12 hours, samples were removed and set aside, and the water decanted, leaving about 10ml of water containing the sediment, including larvae and debris. This residue was agitated and poured into a soft-walled flexible plastic test tube and centrifuged for 2 minutes at 1500rpm (c. 400g). Supernatant was then carefully removed by siphoning from the top of the tube, leaving 1ml of sediment. Sucrose solution of specific gravity 1.17 was added to fill two thirds of the tube, which was then centrifuged again at 400g to float the larvae. In order to capture all the larvae, the test tube was clamped with haemostatic forceps about 5mm below the surface, and inverted to pour the larval suspension into a beaker. The top portion of the clamped tube was then rinsed with water to recover any remaining larvae, and the rinsing water combined with the larval suspension. The whole suspension was then transferred to a standard test tube, topped up to 10ml with water, and inverted several times to mix the larval suspension. A 1ml aliquot of the suspension was transferred into a Sedgewick Rafter nematode counting chamber together with one drop of Lugol's iodine, and the larvae counted under $\times 40$ total magnification. The number of larvae obtained was multiplied by 10 to estimate total number present in the herbage sample.

The efficiency of the above method was evaluated as follows. Herbage from ungrazed pasture was harvested as above, and 10 equal 5 g aliquots cut into 1 cm pieces and each wetted with 1ml larval suspension that contained 399 ± 25 (mean $\pm$ standard deviation) *H. contortus* L3, evenly distributed by pipette throughout the herbage. After settling for 30 minutes, larvae were recovered from herbage using the above method

and counted to estimate the recovery rate.

2.1.4 Rainfall simulation

A portable rainfall simulator was applied in Experiment 2 below to standardize rainfall delivery. This consisted of a pressurized nozzle with a rubber hose connected to a tap, from which water was released at uniform intensity. For the purpose of calibration, the simulator was run for 1 minute, 1 m above an array of cylindrical beakers, and the vertical height and volume of water recovered from each beaker was recorded. The average value showed that rainfall was delivered at a rate of 1.4 mm/min, and this was applied to estimates of rainfall requirements for larval migration.

2.2 Experimental design

This study aimed to quantify the rate of L3 migration from faeces at a range of constant temperatures without moisture limitation (experiment 1). Secondly, to quantify L3 migration following temporary rewetting of faeces in controlled environments (experiment 2) and, finally, the rate of drying of faeces on natural pasture (experiment 3).

2.2.1 Experiment 1: Effect of temperature on the migration of H. contortus L3 from faeces

Migration of L3 was assessed using a mesh sieve apparatus based on that in Wang et al. (2014). Five faecal pellets containing *H. contortus* L3 were placed on a sieve 10

cm in diameter and of aperture 0.9 mm, which was inserted into a plastic cup. In order to create ideal moisture for migration, a fine water mist was applied to the pellets using a pressurized garden sprayer (Spraymist 1.25l, Hozelock, UK) for 20 s from a distance of around 1 m. The experimental units were then transferred immediately to, and maintained in, covered plastic boxes at relative humidity (RH) of 95 %, which were kept stored at 7, 10, 15, 20, 25 and 33 °C for one hour, with four units in each box. Faecal pellets were then transferred to a series of new sieves, each of which was inserted into a new plastic cup, and the same spraying treatment was once more applied to keep the faeces moist, before again placing them into the constant temperature boxes. Larvae that had migrated from the faeces onto the first sieve during incubation were washed into the respective cups and allowed to sediment for 12 h, after which the supernatant was siphoned off with a pipette. The bottom 10 ml of water containing larvae was mixed and a 1 ml aliquot was transferred into a nematode counting chamber. The process of larval recovery was repeated eight times at intervals of one hour. At the end of the experiment, the number of larvae remaining in the pellets was estimated by Baermann extraction, enabling calculation of the proportion of total recoverable L3 present that had migrated out of the faeces at each time point, and at each temperature. Owing to the ideal moisture conditions and short experimental time, mortality of L3 was ignored.

For each temperature treatment, the mean cumulative proportion of L3 that had left the faeces was calculated for each time point (h) and then logistically transformed as follows:

$$y = \log_{10} \left(\frac{x}{1-x} \right) \quad (1)$$

where x is the proportion of L3. Linear regression of y against time (h) was applied to identify the intercept on the horizontal axis, which identifies the number of hours needed for 50 % of L3 to migrate out of faeces (M_{50}) (Azam et al., 2012).

The hourly migration rate, γ , for each temperature was then calculated as follows:

$$\gamma = -\ln(0.5)/M_{50} \quad (2)$$

Regression analysis was then used to estimate the relationship between migration rate and temperature.

2.2.2 Experiment 2: Migration of larvae onto grass in baseline and warmed temperate conditions

The experiment was conducted in two glasshouses, one of which was cooled by air conditioning by around 3°C (baseline = cool) relative to the unregulated glasshouse (warmed). Use of glasshouses allowed for comparison of larval dynamics between treatments similar in the range and pattern of temperature fluctuation, but consistently different in mean temperature level. Mean temperatures in the glasshouses during the experiment were $19.4 \pm 1.0^\circ\text{C}$ (mean $\pm$ standard deviation; range 17.8-22.5) and $22.5 \pm 2.3^\circ\text{C}$ (range 19.1-28.7). The temperature difference between glasshouses was constrained by the ability of the air conditioning unit to cool the large uninsulated air space under solar irradiation, and in any case approximate to those observed in temperate summers, and match medium term climate change projections (Rose et al. 2015a). Relative air humidity averaged 76% in the cool and 62% in the warm greenhouse. Grass turf squares measuring 20 x 20 cm were used to evaluate larval

194 migration onto herbage in the hours after a single rainfall event: these dimensions were
195 chosen because Skinner & Todd (1980) showed that over 90% of larvae on grass were
196 found within 10cm of the faeces from which they originated. Turf was approximately
197 2 cm thick and composed of natural grass turf purchased from a garden centre, with no
198 prior exposure to livestock faeces. Experiments were conducted in daylight, beginning
199 around 9am.

200 A pilot study was first conducted to determine the approximate duration of the
201 window of opportunity for L3 migration following a single rainfall event. Five moist
202 faecal pellets containing L3 (average 2824 per square, standard deviation 659) were
203 placed on each turf square and four replicates were included in each glasshouse. A
204 single simulated rainfall event of approximately 2 mm was applied with the calibrated
205 rainfall simulator immediately after faecal deposition, based on the threshold amount
206 needed for larval migration from hydrated pellets in Wang et al. (2014). Six and 24
207 hours after rain, faecal pellets were carefully transferred to a new turf square with
208 forceps and the old square was collected in a sealed plastic bag and brought to the
209 laboratory. Herbage was processed using the technique described in 2.1.3 above to
210 recover migrated L3 from the grass. L3 remaining in the faeces were recovered by the
211 modified Baermann technique at the end of the experiment. Recovery of larvae
212 sequestered in the soil-root layer was also attempted by the Baermann method, but none
213 were recovered and this compartment was ignored in the main experiment.

214 The main experiment followed the same method of the pilot and used the same
215 faecal material but included two extensions. Firstly, additional turf squares with faecal

pellets were placed in equal numbers beside the experimental replicates to determine FMC at intervals by weighing the pellets and drying in an oven at 90 °C for 12 hours before re-weighing. Secondly, L3 recovery was undertaken every three hours, for the first nine hours after rainfall. L3 remaining in the faeces were again recovered by the modified Baermann technique after 24 hours.

The number of L3 recovered at each time point (N_i) was expressed as the proportion of the total number of L3 estimated to be present by adding those recovered at all time points from the grass and from the faeces at the end of the experiment. The arithmetic mean proportion of L3 recovered was plotted against time for each glasshouse, and subjected to arcsine transformation prior to formal analysis. A general linear model (GLM) was used to test the effect of variables on the proportion of L3 present on the herbage, with Tukey's *post hoc* test applied to significant outcomes. The variables included were glasshouse condition (cool or warm) and collection time (3 h and 6 h). Recovery at 9 h was not analyzed because no additional L3 were recovered from herbage 9 h after rain. The effect of glasshouse condition (cool or warm) on FMC after rainfall was analyzed using the Mann-Whitney U test. Temperature data and L3 results were then applied together to determine the duration of the 'migration window'. Thus, the known number of starting L3 in each replicate, the proportion of L3 observed to leave faeces, and the rate of migration derived from the regression on temperature derived in section 2.2.1 above, were used to estimate the duration for which migration was possible. Observed and predicted numbers of L3 during that period were compared using Pearson's correlation. Model efficiency (EF; Mayer & Butler, 1993) was applied

to evaluate the model (Equation 3):

$$EF = 1 - \frac{\sum(y_i - \hat{y}_i)^2}{\sum(y_i - \bar{y})^2} \quad (3)$$

where y_i = observed number of migrated L3 and $\hat{y}$ = predicted number of migrated L3.

2.2.3 Experiment 3: Faecal moisture loss in the field

A field study was carried out to investigate the rate of moisture loss from faeces on pasture under natural conditions, on un-grazed pasture in a sheep farming area in south-west France. The pasture was mown every few weeks, but unevenly such that it included different heights of grass. During the experiment, the air temperature taken at surface level varied from 21 °C to 37 °C, with wide diurnal fluctuations. Faecal samples were collected from grazing lambs on a nearby farm. Faecal moisture loss was measured under four different conditions, with four replicates for each group: short grass (< 1 cm) with and without shade and long grass (10-25 cm) with and without shade. Shade was produced by fixing a square of cardboard 15 cm above the ground, allowing for circulation of air beneath the canopy. Faeces were placed on plastic mesh (1 mm grid size) to facilitate transfer for weighing and to reduce loss from dung-burying beetles. All replicates were placed in position at 12:30 pm on day 1 and weighed five times, at 1 pm, 5 pm, and 8 pm on day 1 and 10:30 am and 5 pm on day 2. FMC was plotted against time for each treatment, then arcsine transformed to stabilise the variance and analysed in a GLM with grass length and shade included as factors.

3. Results

3.1 Experiment 1: Effect of temperature on the migration of *H. contortus* L3 from faeces

Migration rate was lowest at low and high temperatures, so a polynomial regression model was fitted, which predicted maximum migration rate at 15 °C (Fig. 1).

3.2 Experiment 2: Migration of larvae onto grass in baseline and warmed conditions

The proportion of *H. contortus* L3 recovered from grass using the modified Baermann technique was 0.68 ± 0.14 (mean of 10 replicates $\pm$ standard deviation).

Results of the pilot study showed that almost all migration onto grass occurred between 0 h and 6 h after wetting by simulated rainfall (Fig. 2), with no significant difference between cool and warm glasshouses.

In the main experiment, FMC on the day of deposition was 64 % in both glasshouses and increased to 69 % (cool house) and 68% (warm house) after rainfall, then decreased gradually. Change in FMC in warm and cool glasshouses did not differ significantly over the course of the experiment (Mann-Whitney $U_{46} = 284.5$, $p = 0.947$) (Fig. 3).

There was no significant difference in L3 migration between warm and cool glasshouses ($F_{1, 18} = 0.8$, $p = 0.783$), but the proportion of L3 recovered differed with time of collection ($F_{2, 18} = 24.8$, $p < 0.001$). *Post hoc* Tukey's HSD tests showed that significantly more L3 were recovered from herbage in the first collection, i.e. more larvae migrated onto the grass 0–3 h (mean proportion = $0.57 \pm$ standard deviation 0.26) than 3–6 h (0.05 ± 0.05) after rain ($p < 0.001$) (Fig. 4). No further L3 were recovered 9 h after rain. On average, a proportion of 0.38 ± 0.21 of L3 remained in the faeces at the end of the 24-hour experiment.

In the first 3 h after rainfall, the cumulative number of L3 observed correlated with that predicted from temperature (section 3.1) and the duration of permissive FMC (Pearson $r = 0.912$, $p = 0.002$), notwithstanding some over- and under-prediction (Fig. 5). Model efficiency (EF) was calculated to be 0.64, a reasonably good fit to the available data.

3.3 Experiment 3: Faecal moisture loss in the field

FMC change in the field in the 52 h following deposition is shown in Fig. 6. There was a decreasing trend of FMC over time in all treatments, briefly interrupted by a light rain shower between 28 and 44 h post deposition. Faecal weight decreased most significantly in the ‘short sun’ group, with the lowest FMC recorded as 37 %, after 52 h. Faeces became desiccated more slowly in long grass or under artificial shade. Statistically, both sunlight ($F_{1,218} = 67.517$, $p < 0.001$) and grass length ($F_{1,218} = 12.759$, $p < 0.001$) significantly affected FMC after 52 h. Dung beetles actively transferred and buried faecal samples throughout the experiments, resulting in about 30 % of the samples being removed or destroyed in this way and thus excluded from the analysis.

4. Discussion

The results presented provide first direct estimates on rates of *H. contortus* larval migration from faeces and how this is affected by faecal drying, while also making preliminary comparisons of the relative influences of macroclimate and microclimate on larval migration. While there have been some previous studies on the effect of

temperature on the vertical migration of gastrointestinal nematode larvae onto herbage (e.g. Silangwa & Todd, 1964; Callinan & Westcott, 1986), none have specifically reported on the impact of temperature on horizontal migration out of faeces. In the absence of moisture limitation, our results showed a convex curve, with the fastest migration occurring at intermediate temperatures around 15 °C. There was very little difference in average migration rate between 7 and 20 °C, and rates decreased at 25-30 °C. This was unexpected given the general tendency of metabolism to accelerate with increasing temperature, and could be an adaptation to conserve limited energy resources (Fenton and Rands, 2004) or to avoid exposure to adverse conditions for survival such as lethal ultraviolet radiation high on herbage in sunny conditions (Van Dijk et al., 2009), by remaining in protected micro-environments. Although the experimental conditions and methods were different, Buckley (1940) and Callinan and Westcott (1986) also regarded 15 °C to be the optimal temperature for the migration of trichostrongylid L3, while Rees (1950) and Callinan and Westcott (1986) further found that extremes of very high and very low temperatures inhibited directional larval movement. Even at temperatures of 25-30 °C in the present study, however, more than 10 % of L3 were able to leave the faeces per hour. It is clear from the present and previous results (Wang et al., 2014) that, at least in temperate conditions, temperature is a less important factor than availability of moisture for the migration process, which occurs rapidly when conditions permit.

In the glasshouses, temperature fluctuated with outdoor temperature, but remained in a range favourable for larval migration. The average overall temperature difference

between the two glasshouses was 3.1 °C, which equates to moderate global climate warming scenarios for temperate areas (Rose et al. 2016). Our working hypothesis before the experiment was that hydrated faeces dry faster at high temperatures and the window of opportunity for migration following wetting by a rainfall event will therefore be shorter than at low temperatures. In fact, faecal drying rate was similar in both greenhouses, with only 9 % FMC loss in the first 9 hours, and ambient temperature consequently made little difference to L3 migration rate. In field conditions, different microclimatic conditions showed strong effects on faecal drying rates. Faeces deposited in sunshine on short grass dried much more quickly than those placed in long grass under shade, and within 24 hours FMC was below 60% and therefore too dry to permit L3 migration following light rainfall (Wang et al. 2014). Under artificial shade or that provided by long grass, FMC remained well above 60% for at least twice as long, prolonging the window during which larval migration would occur following a light rain shower.

By quantifying L3 migration rates under different conditions, the present study provides additional information to support mechanistic prediction of climatic impacts on infection patterns. Thus, most migration onto grass occurred within 3 h after rain, with migration declining thereafter. An average of 38 % of L3 remained in faecal pellets 6-24 h after rain, indicating the closure of the migration window within 6 h, presumably as a result of disappearance of the surface water film (Van Dijk et al. 2011; Wang et al. 2014). Under continuous moisture, L3 migration is generally complete within 24 hours (Silangwa and Todd, 1964; Callinan and Westcott, 1986). Further studies will be needed

to more thoroughly characterize L3 migration across a wider range of temperature, rainfall and other environmental conditions such as wind (Krecek et al., 1992) and irradiation (Van Dijk et al. 2009), in order to refine existing predictive models (e.g. Rose et al. 2015b; Walker et al. 2018). The present results show that factors influencing microclimate on pasture should be taken into account in such studies, and could be especially important in environments more extreme than considered here. The interface between faeces, soil and herbage on natural pasture also mediates larval migration onto grass, with complex L3 movement between these compartments (Rose et al., 2015b), while soil moisture is already known to strongly influence L3 development in faeces (Khadijah et al. 2013a,b) and possibly translation onto pasture (Khadijah et al. 2013c). Dung burial, e.g. by invertebrates, has been shown to affect L3 availability (Fincher, 1973; Gronvold, 1979; Waghorn et al. 2002; Sands and Wall 2018), with net effects probably depending on climate, soil type and biodiversity. Although larval recovery from soil was unsuccessful in the experiments reported on here, possibly due to the dense turf structure and shallow soil/root layer with little free soil present, sequestration in soil could be an important modifier of larval availability under natural field conditions. Previous results vary widely, sometimes finding many larvae in soil (Callinan & Westcott, 1986; Krecek, et al., 1995; Leathwick, et al., 2011) and sometimes not (Crofton, 1948; Rose & Small, 1985).

The experiments reported here used a laboratory-maintained strain of *H. contortus* whose origin is obscure and likely tropical, and this might have affected its response to climate (Besier et al. 2016). The temperature-development profile of this isolate was

found to be similar to field isolates from Europe (Rose Vineer et al. 2016). Nevertheless, use of field populations for future experiments on climatic drivers of transmission would be preferable, to take account of natural variation and potential adaptation to local environments. Work on related nematode species has further shown wide within-population variability in response to climate, which might constitute adaptation to unpredictable environmental conditions (Van Dijk and Morgan 2010).

This study shows the potential importance of microclimate, in contrast to macroclimate, in determining rate of faecal drying and hence the effects of subsequent rainfall on L3 release. In the field, pasture will contain faecal pellets at a wide range of times since deposition, and in a range of microclimates, as well as larvae in soil, dung beetles and the herbage mat; the effect of rainfall on larval migration onto herbage is therefore likely to be complex. More sophisticated models and supporting experiments will therefore be needed to adequately predict the effects of climate change on larval availability.

Acknowledgements

This study was funded by the UK Biotechnology and Biological Sciences Research Council (BBSRC) under CIDLID project (BB/H00940X/1 RISCNET) and EU FP7 STREP project GLOWORM (FP7-KBBE-2011-5-288975). EM is supported by BBSRC projects BB/M003949/1 and BB/R010250/1. We are grateful to Hannah Johnston and Sharon Shin for assistance with the laboratory and glasshouse experiments, and to the community of La Grand' Mothe, Brigueil-le-Chantre, 86290

France, for kind permission to use communal land for the field experiment. We appreciate the constructive comments of three anonymous reviewers.

References

- Andersen, F.L., Levine, N.D., 1968. Effect of desiccation on survival of the free-living stages of *Trichostrongylus colubriformis*. J. Parasitol. 54, 117-128.
- Azam, D., Onyinye, M.U., Ashraf, S., Gamal, A. Abd-Allah., Morgan, E.R., 2012. Temperature and the development and survival of infective *Toxocara canis* larvae. Parasitol. Res. 110, 649-656.
- Besier, R.B., Kahn, L.P., Sargison, N.D., Van Wyk, J.A., 2016. The pathophysiology, ecology and epidemiology of *Haemonchus contortus* infection in small ruminants. Adv. Parasitol. 93, 95-143.
- Bryan, R.P., Kerr, J.D., 1989. Factors affecting the survival and migration of the free-living stages of gastrointestinal nematode parasites of cattle in central Queensland. Vet. Parasitol. 30, 315-326.
- Byrd, D.W., Barker, K.R., Ferris, H., Nusbaum, C.J., Griffin, W.E., Small, R.H., Stone, C.A., 1976. Two semi-automatic elutriators for extracting nematodes and certain fungi from soil. J. Nematol. 8, 206-212.
- Buckley, J., 1940. Observations on the vertical migrations of infective larvae of certain bursate nematodes. J. Helminthol. 18, 173-182.
- Callinan, A. & Westcott, J.M., 1986. Vertical distribution of trichostrongylid larvae on herbage and in soil. Int. J. Parasitol. 16, 241-244.

414 Cable, J., Barber, I., Boag, B., Ellison, A., Morgan, E.R., Murray, K., Pascoe, E.L., Sait,
415 S.M., Wilson, A.J. & Booth, M., 2017. Global change, parasite transmission and disease
416 control: lessons from ecology. *Phil. Trans. R. Soc. B Biol. Sci.* 372, 1719, 20160088.

417 Couvillion, C.E., 1993. Estimation of the numbers of trichostrongylid larvae on pasture.
418 *Vet. Parasitol.* 46, 197-203.

419 Coyne, M.J. & Smith, G., 1992. The development and mortality of the free-living stages
420 of *Haemonchus contortus* in laboratory culture. *Int. J. Parasitol.* 22, 641-650.

421 Fenton, A and Rands, S.A, 2004. Optimal parasite infection strategies: a state-
422 dependent approach. *Int. J. Parasitol.* 34, 813–821.

423 Crofton , H. D., 1948. The ecology of immature phases of trichostrongyle neamtodes.
424 (1) The vertical distribution of infective larvae of *Trichostrongylus retortaeformis* in
425 relation to their habitats. *Parasitology* 39, 17-25.

426 Fincher, G.T., 1973. Dung beetles as biological control agents for gastrointestinal
427 parasites of livestock. *J. Parasitol.* 59, 396-399.

428 Grønvold, J., 1979. On the possible role of earthworms in the transmission of
429 *Ostertagia ostertagi* third-stage larvae from feces to soil. *J. Paratitol.* 65, 831-832.

430 Gruner, L., 1986. Strongyle larval recovery from ovine faeces sampled on pasture:
431 efficiency of the Baermannization and epidemiological interest of the technique. In:
432 IVth International Symposium of Veterinary Laboratory Diagnosticians Abstracts June
433 2-6. Amsterdam: The Netherlands, pp. 186-189.

434 Hsu, C.K. & Levine, N.D., 1977. Degree-day concept in development of infective
435 larvae of *Haemonchus contortus* and *Trichostrongylus colubriformis* under constant
436 and cyclic conditions. Am. J. Vet. Res. 38, 1115-1119.

437 Khadijah, S., Kahn, L.P., Walkden-Brown, S.W., Bailey, J.N., Bowers, S.F., 2013a.
438 Effect of simulated rainfall timing on faecal moisture and development of *Haemonchus*
439 *contortus* and *Trichostrongylus colubriformis* eggs to infective larvae. Vet. Parasitol.
440 192, 199-210.

441 Khadijah, S., Kahn, L.P., Walkden-Brown, S.W., Bailey, J.N., Bowers, S.F., 2013b.
442 Soil moisture modulates the effects of the timing and amount of rainfall on faecal
443 moisture and development of *Haemonchus contortus* and *Trichostrongylus*
444 *colubriformis* to infective third stage larvae. Vet. Parasitol. 196, 347-357.

445 Khadijah, S., Kahn, L.P., Walkden-Brown, S.W., Bailey, J.N., Bowers, S.F., 2013c.
446 Translation of *H. contortus* and *T. colubriformis* from egg to establishment in grazing
447 sheep is unaffected by rainfall timing, rainfall amount and herbage height under
448 conditions of high soil moisture in the Northern Tablelands of NSW. Vet. Parasitol. 197,
449 204-211.

450 Krecek, R.C., Groeneveld, H.T. & Maritz, J.I., 1992. A preliminary study of the effect
451 of microclimate on third-stage larvae of *Haemonchus contortus* and *Haemonchus*
452 *placei* on irrigated pasture. Int. J. Parasitol. 22, 747-752.

453 Krecek, R. C., Hartman, R., Groeneveld, H. T. & Thorne, A., 1995. Microclimatic effect
454 on vertical migration of *Haemonchus contortus* and *Haemonchus placei* third-stage
455 larvae on irrigated Kikuyu pasture. Onderstepoort J. Vet. Res. 62, 117-122.

456 Leathwick, D. M., Miller, C. M. & Waghorn, T. S., 2011. Development and spatial
457 distribution of the free-living stages of *Teladorsagia circumcincta* and *Trichostrongylus*
458 *colubriformis* on pasture: A pilot study. New Zealand Vet. J. 59, 272-278.

459 MAFF, 1986. Fisheries and Food, Reference Book. In: Manual of Veterinary
460 Parasitological Laboratory Techniques. London: Ministry of Agriculture, HMSO. pp. 5.

461 Martin, R.R., Beveridge, I., Pullman, A.L. & Brown, T.H., 1990. A modified technique
462 for the estimation of the number of infective nematode larvae present on pasture, and
463 its application in the field under South Australian conditions. Vet. Parasitol. 37, 133-
464 143.

465 Mason, S.J. et al., 1999. Changes in extreme rainfall events in South Africa. Climate
466 Change 41, 249-257.

467 Mayer, D.G. & Butler, D.G., 1993. Statistical validation. Ecol. Model. 68, 21-32.

468 Morgan, E.R. & Wall, R., 2009 Climate change and parasitic disease: farmer mitigation?
469 Trends Parasitol. 25, 308-313

470 Morgan, E.R. & Van Dijk, J., 2012. Climate and the epidemiology of gastrointestinal
471 nematode infections of sheep in Europe. Vet. Parasitol. 189, 8-14.

472 O'Connor, L.J., Walkden-Brown, S.W. & Kahn, L.P., 2006. Ecology of the free-living
473 stages of major trichostrongylid parasites of sheep. Vet. Parasitol. 142, 1-15.

474 O' Connor, L.J., Kahn, L.P. & Walkden-Brown, S.W., 2007. Moisture requirements for
475 the free-living development of *Haemonchus contortus*: Quantitative and temporal
476 effects under conditions of low evaporation. Vet. Parasitol. 150, 128-138.

477 O' Connor, L.J., Kahn, L.P. & Walkden-Brown, S.W., 2008. Interaction between the

478 effects of evaporation rate and amount of simulated rainfall on development of the free-
479 living stages of *Haemonchus contortus*. Vet. Parasitol. 155, 223-234

480 Onyali, I.O., Onwuliri, C. & Ajayi, J.A., 1990. Development and survival of
481 *Haemonchus contortus* larvae on pasture at Vom, Plateau State, Nigeria. Vet. Res. Com.
482 14, 211-216.

483 Pandey, V.S., Chaer, A. & Dakkak, A., 1989. Effect of temperature on development of
484 the free-living stages of *Ostertagia circumcincta*. Vet. Parasitol. 32, 193-197.

485 Pandey, V.S., Chaer, A. & Dakkak, A., 1993. Effect of temperature and relative
486 humidity on survival of eggs and infective larvae of *Ostertagia circumcincta*. Vet.
487 Parasitol. 49, 219-227.

488 Perry, B. D. & Randolph, T. F., 1999. Improving the assessment of the economic impact
489 of parasitic diseases and of their control in production animals. Vet. Parasitol. 84, 145-
490 168.

491 Pinault, J.L., 2012. Global warming and rainfall oscillation in the 5–10 yr band in
492 Western Europe and Eastern North America. Climate Change 114, 621-650.

493 Rossanigo, C.E. & Gruner, L., 1995. Moisture and temperature requirements in faeces
494 for the development of free-living stages of gastrointestinal nematodes of sheep, cattle
495 and deer. J. Helminthol. 69, 357-362.

496 Rees, G., 1950. Observations on the vertical migration of third stage larvae of
497 *Haemonchus contortus* (Rud) on experimental plots of *Lolium perenne* S24 in relation
498 to meteorological and micrometeorological factors. J. Parasitol. 40, 127-143.

499 Rose, J. H., 1963. Observations on the free-living stages of the stomach worm

500 *Haemonchus contortus*. Parasitology. 53, 469-481.

501 Rose, J. H. & Small, A. J., 1985. The distribution of the infective larvae of sheep gastro-
502 intestinal nematodes in soil and on herbage and the vertical migration of
503 *Trichostrongylus vitrinus* larvae through the soil. J. Helminthol. 59, 127-135.

504 Rose, H., Rinaldi, L., Bosco, A., Mavrot, F., De Waal, T., Skuce, P., Charlier, J.,
505 Torgerson, P.R., Hertzberg, H., Hendrickx, G., Vercruysse, J., Morgan, E.R., 2015a.
506 Widespread anthelmintic resistance in European farmed ruminants: a systematic review.
507 Vet. Rec. 176, 21, doi:10.1136/vr.102982.

508 Rose, H., Wang, T., Van Dijk, J., & Morgan, E.R., 2015b. GLOWORM-FL: A
509 simulation model of the effects of climate and climate change on the free-living stages
510 of gastro-intestinal nematode parasites of ruminants. Ecol. Model. 297, 232-245.

511 Rose, H., Caminade, C., Bolajoko, M.B., Phelan, P., van Dijk, J., Baylis, M., Williams,
512 D., Morgan, E.R., 2016. Climate-driven changes to the spatio-temporal distribution of
513 the parasitic nematode, *Haemonchus contortus*, in sheep in Europe, Global Change Biol.
514 22, 1271-1285.

515 Rose Vineer, H., Steiner, T., Knapp-Lawitzke, F., Bull, K., von Son-de Fernex, E.,
516 Bosco, A., Hertzberg, H., Demeler, J., Rinaldi, L., Morrison, A.A., Skuce, P., Bartley,
517 D.J., Morgan, E.R., 2016. Implications of between-isolate variation for climate change
518 impact modelling of *Haemonchus contortus* populations. Vet. Parasitol. 229, 144-149.

519 Sands, B., Wall, R., 2017. Dung beetles reduce livestock gastrointestinal parasite
520 availability on pasture. J. Appl. Ecol. 54, 1180-1189.

521 Silangwa, S.M. & Todd, A.C., 1964. Vertical migration of trichostrongylid larvae on

522 grasses. J. Parasitol. 50, 278-285.

523 Silva, B.F. et al., 2008. Vertical migration of *Haemonchus contortus* third stage larvae
524 on *Brachiaria decumbens* grass. Vet. Parasitol 158, 85-92.

525 Skinner, W.D. & Todd, K.S., 1980. Lateral migration of *Haemonchus contortus* larvae
526 on pasture. Am. J. Vet. Res. 41, 395-398.

527 Todd, K.S., Levine, N.D. & Boatman, P.A., 1976. Effect of temperature on survival of
528 free-living stages of *Haemonchus contortus*. Am. J. Vet. Res. 37, 991-992.

529 Troell, K., Waller, P.J & Hoglund, J., 2005. The development and overwintering
530 survival of free-living larvae of *Haemonchus contortus* in Sweden. J. Helminthol. 79,
531 373-379.

532 Van Bezooijen, J., 2006. Methods and Techniques for Nematology. Wageningen:
533 Wageningen University. p. 19.

534 Van Dijk, J., De Louw, M., Kalis, L. & Morgan, E.R., 2009. Ultraviolet light increases
535 mortality of nematode larvae and can explain patterns of larval availability at pasture.
536 Int. J. Parasitol. 39, 1151-1156.

537 Van Dijk, J., Morgan, E.R., 2010. Variation in the hatching behaviour of *Nematodirus*
538 *battus*: Polymorphic bet hedging? Int. J. Parasitol. 40, 675-681.

539 Van Dijk, J., Sargison, N.D., Kenyon, F., Skuce, P.J., 2010. Climate change and
540 infectious disease: helminthological challenges to farmed ruminants in temperate
541 regions. Animal. 4, 377-392.

542 Van Dijk, J., Morgan, E.R., 2011. The influence of water on the migration of infective
543 trichostrongyloid larvae onto grass. Parasitology. 138, 1-9.

544 Van Wyk, J.A., Mayhew, E., 2013. Morphological identification of parasitic nematode
 545 infective larvae of small ruminants and cattle: A practical lab guide. Onderstepoort J.
 546 Vet. Res. 80, 1-14.

547 Veglia, F., 1915. The anatomy and life history of *Haemonchus contortus* (Rud.), 3rd and
 548 4th Reports of the Director of Veterinary Research, 347-500.

549 Waghorn, T.S, Leathwick, D.M, Chen, L.Y, Gray, R.A.J, Skipp, R. A, 2002. Influence
 550 of nematophagous fungi, earthworms and dung burial on development of the free-living
 551 stages of *Ostertagia (Teladorsagia) circumcincta* in New Zealand. Vet. Parasitol. 201,
 552 119-129

553 Waghorn, T.S, Reynecke, D.P, Oliver, A.M, Miller, C.M, Vlassoff, A, Koolaard, J.P,
 554 Leathwick, D.M., 2011. Dynamics of the free-living stages of sheep intestinal parasites
 555 on pasture in the North Island of New Zealand. 1. Patterns of seasonal development. N.
 556 Z. Vet. J. 59, 279-286.

557 Walker, J.G., Evans, K.E., Rose Vineer, H., Van Wyk, J.A., Morgan, E.R., 2018.
 558 Prediction and attenuation of seasonal spillover of parasites between wild and domestic
 559 ungulates in an arid mixed-use system. J. Appl. Ecol. 2018;00:1–11.
 560 <https://doi.org/10.1111/1365-2664.13083>

561 Waller, P.J., 1997. Anthelmintic resistance. Vet. Parasitolo, 72, 391-412.

562 Wang, T., Van Wyk, J.A., Morrison, A., Morgan, E.R., 2014. Moisture requirements for
 563 the migration of *Haemonchus contortus* third stage larvae out of faeces. Vet. Parasitol.
 564 204, 258–264.

565

Figure legends

Fig. 1. Regression of hourly migration rate out of sheep faeces against temperature for *Haemonchus contortus* L3. $R^2 = 0.292$, $F_{2,21} = 4.341$, $p = 0.026$; regression equation $y = -0.0002T^2 + 0.0065T + 0.1417$.

Fig. 2. Recovery of *Haemonchus contortus* L3 from grass 6 and 24 hours following wetting of sheep faeces by simulated rainfall, and from the faeces by Baermann extraction after 24 hours. Warm and cooled treatments were applied in glasshouses at 22.5 and 19.4 °C respectively. Error bars represent standard error. There was no significant difference in FMC change between the two greenhouses.

Fig. 3. Mean faecal moisture content (FMC) change in sheep faecal samples placed on grass turf in warm (22.5 °C) and cooled (19.4 °C) glasshouses. Artificial rainfall was administered once at the starting point (9 am). Error bars represent standard error. There was no significant difference in FMC change between the two greenhouses.

Fig. 4. Proportion of *Haemonchus contortus* L3 recovered from herbage after different

time periods (0-3, 3-6 and 6-9 hours) following a single artificial rainfall event applied to sheep faeces containing mature L3, and the proportion of L3 remaining in faeces after 24 hours (Baermann). The same rainfall treatment was applied in warm (22.5 °C) and cooled (19.4 °C) greenhouses. Error bars indicate standard error.

Fig. 5. The number of L3 observed in the first 3 hours after artificial rainfall and the number predicted from the regression of migration rate against temperature (Pearson $r = 0.912$, $p = 0.002$). Data from warm and cooled greenhouses were pooled for analysis. The solid line represents unity (i.e. the line of perfect prediction).

Fig. 6. Change in faecal moisture content (FMC) of sheep faeces over time, in the field under four different treatments. Error bars represent standard error of four replicates. There was a rain event between 28 and 44 hours; the amount of rain was not measured. Short / long = short and long grass; sun / shade = unshaded or shaded by cardboard canopy. The dotted line represents the approximate lower FMC threshold that stops L3 migration from faeces (Wang et al. 2014).