**QUEEN'S
UNIVERSITY
BELFAST**

Face Masks Reduce the Release of *Pseudomonas aeruginosa* Cough Aerosols when Worn for Clinically-Relevant Time Periods

CF Cough Aerosol Group (2018). Face Masks Reduce the Release of *Pseudomonas aeruginosa* Cough Aerosols when Worn for Clinically-Relevant Time Periods. *American Journal of Respiratory and Critical Care Medicine*, 198(10), 1339-1342. <https://doi.org/10.1164/rccm.201805-0823LE>

Published in:
American Journal of Respiratory and Critical Care Medicine

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© 2018 by the American Thoracic Society. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Face masks reduce the release of *Pseudomonas aeruginosa* cough aerosols when worn for clinically-relevant time periods

Rebecca E Stockwell^{1,2}, Michelle E Wood^{1,2,3}, Congrong He⁴, Laura J Sherrard⁵, Emma L Ballard⁶, Timothy J Kidd^{1,7}, Graham R Johnson⁴, Luke D Knibbs⁸, Lidia Morawska⁴, Scott C Bell^{*1,2,3}; CF Cough Aerosol Group.

CF cough aerosol group members:

Maureen Peasey, Christine Duplancic, Kay A Ramsay, Nassib Jabbour, Peter O'Rourke, Claire E Wainwright, Peter D Sly

¹Lung Bacteria Group, QIMR Berghofer Medical Research Institute, 300 Herston Road, Herston QLD 4006, Australia

²Faculty of Medicine, The University of Queensland, Herston QLD 4006, Australia

³Adult Cystic Fibrosis Centre, The Prince Charles Hospital, 627 Rode Road, Chermside, QLD 4032, Australia

⁴International Laboratory for Air Quality and Health, Queensland University of Technology, Brisbane QLD 4000, Australia

⁵Queen's University Belfast, 97 Lisburn Road, Belfast BT9 7BL, United Kingdom

⁶Statistical Support Group, QIMR Berghofer Medical Research Institute, 300 Herston Road, Herston QLD 4006, Australia

⁷School of Chemistry and Molecular Biosciences, The University of Queensland, St Lucia QLD 4067, Australia

⁸School of Public Health, The University of Queensland, Herston QLD 4006, Australia

Email:

rebecca.stockwell@qimrberghofer.edu.au

michelle.wood2@health.qld.gov.au

c.he@qut.edu.au

l.sherrard@qub.ac.uk

emma.ballard@qimrberghofer.edu.au

t.m.kidd@uq.edu.au

g.johnson@qut.edu.au

l.knibbs@uq.edu.au

l.morawska@qut.edu.au

scott.bell@qimrberghofer.edu.au

***Corresponding author.** Lung Bacteria Group, QIMR Berghofer Medical Research Institute,
300 Herston Road, Herston, Brisbane QLD 4006, Australia.

Tel: +61 7 3139 4770; Email: scott.bell@qimrberghofer.edu.au

Author contributions: G.R.J., L.D.K., T.J.K., R.E.S., L.J.S., L.M. and S.C.B. conceived and designed the experiment. S.C.B., T.J.K. and L.M. led the funding applications with other members of the CF cough aerosol group (C.E.W and P.D.S.). M.E.W. and S.C.B. recruited the study participants. R.E.S. and C.H. conducted the cough studies. G.R.J. acquired the aerosol data. R.E.S. performed microbiological analysis. E.L.B. led the data analysis. R.E.S. and S.C.B. provide overall responsibility for the data and wrote the manuscript, with input from all co-authors. M.P., C.D., K.A.R., N.J., P.O., C.E.W and P.D.S. provided support to the study including analysis and/or microbiology expertise and/or clinical supervision.

Funding support: The project was funded by Cystic Fibrosis Foundation Therapeutics USA (BELL14AO). T.J.K. acknowledges National Health and Medical Research Council (NHMRC) Early Career (GNT10884488) and ERS-EU RESPIRE2 Marie Skłodowska-Curie Postdoctoral Research (#4571-2013) Fellowship support. L.D.K. acknowledges an NHMRC Early Career Fellowship (APP1036620). R.E.S acknowledges The Prince Charles Hospital Foundation and Advance Queensland PhD Scholarships.

List ONE descriptor number that best classifies the subject of your manuscript: 9.17
Cystic Fibrosis: Translational & Clinical Studies

Total word count for the body of the manuscript = 1328

Abstract

Introduction: The cystic fibrosis (CF) infection control guidelines recommend that people with CF wear face-masks when in communal areas of hospitals. Recently, we reported short-term wear of face-masks (~10-minutes) reduced the release of *Pseudomonas aeruginosa* aerosols during coughing. However, there is limited evidence to determine if face-masks continue to be effective at reducing the release of infectious cough aerosols after longer wear times. **Methods:** We recruited 25 people with CF and chronic *P. aeruginosa* infection and 10 healthy volunteers. All participants underwent up to 5 cough tests in a validated cough rig: 1) uncovered cough; 2) coughing with surgical mask worn for 10-minutes; 3) coughing with surgical mask worn for 20-minutes; 4) coughing with a surgical mask worn for 40-minutes; 5) coughing with an N95 respirator worn for 20-minutes (optional). The wear time of the mask included a 5-minute cough period in the aerosol collection rig. Sputum samples and cough aerosols were collected from participants with CF as previously described. All participants rated their level of comfort post-test. **Results:** Surgical masks and N95 respirators were effective at reducing the release of *P. aeruginosa* aerosols during coughing after 40-minutes total wear and 20-minutes total wear respectively. Both participants with CF and healthy volunteers rated the surgical masks as more comfortable compared to N95 respirators. **Conclusions:** Surgical masks were the preferred interface to wear as source control and were effective at reducing the release of *P. aeruginosa* aerosols during coughing after 40-minutes of total wear.

Introduction

Aerosol dissemination of respiratory pathogens may contribute to person-to-person transmission in people with cystic fibrosis (CF) [1]. This evolving knowledge of transmission modes has led to an update of the CF Foundation Infection and Prevention Control Guidelines recommending people with CF wear surgical masks in communal hospital areas to prevent the spread of CF respiratory pathogens [2]. These guidelines recommended the use of surgical masks as source control despite limited evidence for this application.

We recently demonstrated that short-term wear of face-masks (10-minutes total wear) significantly reduces the release of *Pseudomonas aeruginosa* aerosols during coughing in people with CF [3]. These findings are consistent with an earlier study of people with CF that reached the same conclusion after very short-term wear of surgical masks (worn for 21 coughs) [4]. These results and a recent editorial to our short-term mask wear study [5] support our current aim to investigate the effectiveness, tolerability and functionality of face-masks as source control after extended wear.

Methods

We recruited 25 people with CF and chronic *P. aeruginosa* infection [6] from the Adult Cystic Fibrosis Centre, The Prince Charles Hospital, Brisbane, Australia. Ten healthy volunteers were recruited from hospital and research staff to assess mask comfort and mask weight change. All participants performed up to five randomly ordered tests in a validated cough system [7]: 1) uncovered cough; 2) coughing with surgical mask worn for 10-minutes; 3) coughing with surgical mask worn for 20-minutes; 4) coughing with surgical mask worn for 40-minutes; 5) coughing with N95 mask worn for 20-minutes [3, 7]. The N95 test was an optional test based on the poor comfort ratings observed in our earlier mask study [3].

The duration of the mask wear tests were selected based on observation of patients moving around communal areas of the hospital described here. Two types of masks were tested: “surgical mask” [Catalogue # 47107; Halyard FLUIDSHIELD Level 3 Fog-Free Procedure Mask ($\Delta P < 2.5$), Georgia, USA] and “N95 mask” [Catalogue # 46827 (small) or 46727 (regular), Halyard FLUIDSHIELD N95 Particulate Filter Respirator and Surgical Mask, Georgia, USA]. New masks were used for each test. The total wear time of the masks included 1-minute positioning of the participant into the rig, 2-minutes of tidal breathing with HEPA-filtered air, a 5-minute cough period, followed by another 2-minutes of tidal breathing. Cough aerosol collection, sputum processing and *P. aeruginosa* genotyping were performed as previously described [1, 3, 8]. All participants rated their comfort levels after each test [3, 9]. All masks were weighed before and following each test.

SPSS version 25 was used for statistical analysis. Participants with CF were stratified by the amount of aerosol colony forming units (CFU) produced during the uncovered cough test: high producer (total CFU was ≥ 10) or no/low producer (total CFU was < 10) [3]. Categorical variables were examined using Pearson Chi-squared test or Fisher’s Exact test. Continuous variables were examined using a Student t-test or Mann-Whitney U test. CFU were log transformed and the paired t-test examined changes over time. The McNemar-Bowker test was used to examine comfort scores over time. The Wilcoxon Signed Rank Test was used to examine the change in mask weight over time.

Results

P. aeruginosa was cultured from the sputum of 25/25 participants with CF and was cultured in cough aerosols during the uncovered cough test of 20/25 participants (Table 1). *P. aeruginosa*

was cultured from cough aerosols of 9/20 participants during any of the surgical mask tests (10-minutes, 20-minutes and 40-minutes total wear time) and 4/20 participants during the N95 mask test of 20-minutes total wear. The CFU were significantly reduced for the surgical mask tests compared with the uncovered cough test ($p<0.001$). Between mask tests, the CFU count remained similar as the duration of surgical mask wear increased as well as between mask types (Table 1). The *P. aeruginosa* strain types found in the cough aerosols were genetically indistinguishable from the paired sputum sample of each participant.

Participants with CF rated surgical masks less comfortable than healthy volunteers for all test durations (surgical mask: 10-minutes, $p=0.001$; 20-minutes, $p=0.007$; 40-minutes, $p=0.023$; N95: 20-minutes, $p=0.018$) (Table 2). Participants with CF were more tolerant of surgical mask wear (good comfort) after 10- and 20-minutes total wear time if they had higher lung function, yet this difference was lost after 40-minutes of surgical mask wear. N95 masks were rated less comfortable in both participants with CF and healthy volunteers (Table 2).

The change in mask weight for each test ranged from no weight change to a maximum weight change of 0.02g and was comparable between participants with CF and healthy volunteers (Table 2). There was a minor increase in surgical mask weight (median change, 0.01g) after 40-minutes compared with 10-minutes wear ($p=0.031$) (Table 2). No statistical differences in mask weight change were seen in other time or mask type comparisons (Table 2).

Discussion

Our study demonstrates that face masks worn for clinically-relevant time periods are effective at reducing the release of potentially infectious aerosols during coughing in people with CF. These results extend upon our earlier observations that demonstrated surgical masks and N95

156 masks were both effective at reducing the release of infectious cough aerosols when the mask
157 wear was of shorter duration [3]. The outcomes of our studies demonstrate that surgical masks
158 are effective and tolerable as source control [3] and support the CF Foundation (USA)
159 recommendations for surgical mask wear to reduce the risk of CF pathogen transmission in the
160 hospital setting [2].

161
162 Surgical masks were the preferred mask type for source control in terms of comfort, which is
163 similar to our short-term wear mask study findings [3]. Healthy volunteers tolerated the surgical
164 masks better than those with CF and participants with CF who had higher lung function
165 tolerated surgical masks better. When the comfort of surgical masks was assessed after extended
166 wear in this cohort, a major finding was that the comfort ratings remained unchanged regardless
167 of wear time for both people with and without CF. Therefore, surgical masks are not only
168 effective but are also well tolerated by participants after 40-minutes total wear.

169
170 An accompanying editorial of our recent mask study [3] questioned if mask dampness may
171 affect the ability of the mask to function as source control after prolonged wear times [5]. The
172 CF infection control guidelines indicate that masks being used as source control should be
173 replaced when damp [2] and excessive moisture accumulation was a common reason for
174 surgical mask replacement in people with tuberculosis using surgical masks as source control
175 [10]. Our data indicates that although there was evidence of surgical mask moisture
176 accumulation after 40-minutes total wear (estimated by increased weight), the surgical mask
177 continued to function effectively as source control mitigating this concern.

178
179 There are several limitations to this study: 1) The infectious dose of *P. aeruginosa* is unknown
180 and therefore the infection risk cannot be determined; 2) Participants remained in view of staff

181 while wearing the masks and this may have modified the extent to which participants interfered
182 with the mask leading to an incorrect estimation on the masks protective effects; 3) Participants
183 were seated during the cough testing and this may have impacted on the participant's ability to
184 cough freely; 4) While some participants experienced episodes of spontaneous cough during
185 testing, we were unable to differentiate between spontaneous and voluntary cough. Therefore,
186 the protective effects of the masks may be overestimated; 5) The effectiveness and tolerability
187 of masks is reported in adults only and these characteristics need to be studied in children; 6)
188 Our study had a maximum wear time of 40-minutes and the effectiveness of masks worn for
189 longer periods is unknown; 7) We did not assess inward protection provided by masks but this
190 has been highlighted as an understudied field of research [11].

191
192 Our study confirms the effectiveness of surgical masks at reducing the release of *P. aeruginosa*
193 cough aerosols in people with CF and provides evidence of patient tolerability and functionality
194 of these masks as source control after 40-minutes of total wear.

195
196 Acknowledgements: We thank Dr Farhad Salimi for his aerosol support to the study. We thank
197 Greg Flohr and staff from the Central Pathology Laboratory (Royal Brisbane and Women's
198 Hospital), Pathology Queensland for microbiological support to the study. We thank the Adult
199 CF Centre team in supporting recruitment to the studies. We also thank all the participants in
200 the study for supporting the work.

202 Table 1: Demographic and clinical characteristics of the study participants

	Group			Production level in CF participants		
	Healthy (n = 10)	CF (n = 25)	p-value	No/low (<10 CFU) (n = 14)	High (≥ 10 CFU) (n = 11)	p-value
Participant characteristics						
Age, years, mean (SD)	37.3 (12.3)	33.3 (9.0)	0.29	36.7 (9.3)	28.9 (6.9)	0.029
Sex, male, n (%)	6 (60.0)	15 (60.0)	1.00	9 (64.3)	6 (54.5)	0.70
Body mass index (BMI), kg/m ² , mean (SD)	24.6 (3.5)	22.8 (3.2)	0.14	22.5 (3.7)	23.1 (2.6)	0.63
FEV ₁ % predicted, mean (SD)	92.6 (9.2)	53.8 (20.8)	<0.001	54.2 (23.2)	53.3 (18.2)	0.91
Mean <i>P. aeruginosa</i> sputum concentration, x 10 ⁷ CFU/mL (95% CI) ^a	n/a	5.2 (2.1 – 12.9)	-	1.9 (0.7 – 5.7)	18.3 (4.7 – 70.9)	0.008
Participants with <i>P. aeruginosa</i> detected in cough aerosols						
Uncovered cough test						
n (%)	n/a	20 (80.0)	-	9 (64.3)	11 (100.0)	
Mean CFU (95% CI) ^a		17 (7 - 43)		2 (1 – 4)	75 (34 – 165)	<0.001 ^d
Surgical mask tests						
10-minutes total wear						
n (%)	n/a	9 (36.0)	-	1 (7.1)	8 (72.7)	
Mean CFU (95% CI) ^a		4 (1 – 10)		1	5 (1 – 13)	
20-minutes total wear						
n (%)	n/a	9 (36.0)	-	1 (7.1)	8 (72.7)	
Mean CFU (95% CI) ^a		4 (1 – 10)		1	4 (1 – 11)	0.99 ^e
40-minutes total wear						
n (%)	n/a	9 (36.0)	-	1 (7.1)	8 (72.7)	
Mean CFU (95% CI) ^a		3 (1 – 7)		1	4 (1 – 9)	0.56 ^f , 0.64 ^g
N95 mask test [~]						
20-minutes total wear (n=23)						
n (%)	n/a	4 (17.4)	-	0 (0.0) ^b	4 (40.0) ^c	
Mean CFU (95% CI) ^a		2 (0 – 6)		n/a	2 (0 – 6)	0.19 ^h

203 *Definitions:* FEV₁, forced expiratory volume in 1 second; CFU, colony forming unit; CFU/mL, CFU per millilitre of sputum; SD, standard deviation;
204 CI, confidence interval, n/a, not applicable ~ Optional test
205
206 ^ageometric mean
207 ^bparticipant number (n) = 13
208 ^cparticipant number (n) = 10
209 ^dUncovered cough (geometric mean CFU) compared to each surgical mask test (geometric mean CFU)
210 ^eSurgical masks (geometric mean CFU): 10-minutes *versus* 20-minutes total wear
211 ^fSurgical masks (geometric mean CFU): 10-minutes *versus* 40-minutes total wear
212 ^gSurgical masks (geometric mean CFU): 20-minutes *versus* 40-minutes total wear
213 ^hMean CFU surgical mask 20-minutes total wear *versus* mean CFU N95 mask 20-minutes total wear

214 Table 2: Summary of mask properties

Mask properties	Mask comfort			Mask weight change	
	Healthy n (%)	CF n (%)	p-value	All participants Median (IQR)	p-value
Uncovered cough comfort level			1.00	n/a	n/a
Poor	0 (0.0 %)	0 (0.0 %)			
Sufficient	1 (10.0 %)	2 (8.0 %)			
Good	9 (90.0 %)	23 (92.0 %)			
Coughing wearing a surgical mask – 10-minutes total wear			0.001	0.01g (0.00g – 0.02g)	n/a
Poor	1 (10.0 %)	0 (0.0 %)			
Sufficient	0 (0.0 %)	15 (60.0 %)			
Good	9 (90.0 %)	10 (40.0 %)			
Coughing wearing a surgical mask – 20-minutes total wear			0.007	0.01g (0.00g – 0.02g)	0.73 ^a
Poor	1 (10.0 %)	1 (4.0 %)			
Sufficient	0 (0.0 %)	13 (52.0 %)			
Good	9 (90.0 %)	11 (44.0 %)			
Coughing wearing a surgical mask – 40-minutes total wear			0.023	0.02g (0.01g – 0.03g)	0.25 ^b , 0.031 ^c
Poor	1 (10.0 %)	2 (8.0 %)			
Sufficient	1 (10.0 %)	15 (60.0 %)			
Good	8 (80.0 %)	8 (32.0 %)			
Coughing wearing N95 mask -20-minutes total wear			0.018	0.02g (0.00g – 0.04g)	0.21 ^d
Poor	0 (0.0 %)	11 (47.8 %)			
Sufficient	7 (77.8 %)	8 (34.8 %)			
Good	2 (22.2 %)	4 (17.4 %)			

215

216 ^aSurgical mask weight change: 10-minutes *versus* 20-minutes total wear

217 ^bSurgical mask weight change: 20-minutes *versus* 40-minutes total wear

218 ^cSurgical mask weight change: 10-minutes *versus* 40-minutes total wear

219 ^dMask weight change after 20-minutes wear: surgical mask *versus* N95 mask

References:

1. Knibbs, L.D., Johnson, G.R., Kidd, T.J., Cheney, J., Grimwood, K., Kattenbelt, J.A., O'Rourke, P.K., Ramsay, K.A., Sly, P.D., Wainwright, C.E., Wood, M.E., Morawska, L., and Bell, S.C. Viability of *Pseudomonas aeruginosa* in cough aerosols generated by persons with cystic fibrosis. *Thorax*. 2014;69(8):740-5
2. Saiman, L., Siegel, J.D., LiPuma, J.J., Brown, R.F., Bryson, E.A., Chambers, M.J., Downer, V.S., Fliege, J., Hazle, L.A., Jain, M., Marshall, B.C., O'Malley, C., Pattee, S.R., Potter-Bynoe, G., Reid, S., Robinson, K.A., Sabadosa, K.A., Schmidt, H.J., Tullis, E., Webber, J., and Weber, D.J. Infection prevention and control guideline for cystic fibrosis: 2013 update. *Infect Control Hosp Epidemiol*. 2014;35 Suppl 1:S1-s67
3. Wood, M.E., Stockwell, R.E., Johnson, G.R., Ramsay, K.A., Sherrard, L.J., Jabbour, N., Ballard, E., O'Rourke, P., Kidd, T.J., Wainwright, C.E., Knibbs, L.D., Sly, P.D., Morawska, L., and Bell, S.C. Face Masks and Cough Etiquette Reduce the Cough Aerosol Concentration of *Pseudomonas aeruginosa* in People with Cystic Fibrosis. *Am J Respir Crit Care Med*. 2018;197(3):348-355
4. Driessche, K.V., Hens, N., Tilley, P., Quon, B.S., Chilvers, M.A., de Groot, R., Cotton, M.F., Marais, B.J., Speert, D.P., and Zlosnik, J.E. Surgical masks reduce airborne spread of *Pseudomonas aeruginosa* in colonized patients with cystic fibrosis. *Am J Respir Crit Care Med*. 2015;192(7):897-9
5. Simmonds, N.J. and Bush, A. The Man in the Paper Mask: One (Mask) for All and All for . . . Cystic Fibrosis? *Am J Respir Crit Care Med*. 2018;197(3):281-283
6. Ramsay, K.A., Sandhu, H., Geake, J.B., Ballard, E., O'Rourke, P., Wainwright, C.E., Reid, D.W., Kidd, T.J., and Bell, S.C. The changing prevalence of pulmonary infection in adults with cystic fibrosis: A longitudinal analysis. *J Cyst Fibros*. 2017;16(1):70-77

- 244 7. Johnson, G.R., Knibbs, L.D., Kidd, T.J., Wainwright, C.E., Wood, M.E., Ramsay, K.A.,
245 Bell, S.C., and Morawska, L. A Novel Method and Its Application to Measuring
246 Pathogen Decay in Bioaerosols from Patients with Respiratory Disease. PLoS One.
247 2016;11(7):e0158763
- 248 8. Syrmis, M.W., Kidd, T.J., Moser, R.J., Ramsay, K.A., Gibson, K.M., Anuj, S., Bell,
249 S.C., Wainwright, C.E., Grimwood, K., Nissen, M., Sloots, T.P., and Whiley, D.M. A
250 comparison of two informative SNP-based strategies for typing *Pseudomonas*
251 *aeruginosa* isolates from patients with cystic fibrosis. BMC Infect Dis. 2014;14:307
- 252 9. Gregoret, C., Confalonieri, M., Navalesi, P., Squadrone, V., Frigerio, P., Beltrame, F.,
253 Carbone, G., Conti, G., Gamna, F., Nava, S., Calderini, E., Skrobik, Y., and Antonelli,
254 M. Evaluation of patient skin breakdown and comfort with a new face mask for non-
255 invasive ventilation: a multi-center study. Intensive Care Med. 2002;28(3):278-84
- 256 10. Dharmadhikari, A.S., Mphahlele, M., Stoltz, A., Venter, K., Mathebula, R., Masotla, T.,
257 Lubbe, W., Pagano, M., First, M., Jensen, P.A., van der Walt, M., and Nardell, E.A.
258 Surgical face masks worn by patients with multidrug-resistant tuberculosis: impact on
259 infectivity of air on a hospital ward. Am J Respir Crit Care Med. 2012;185(10):1104-9
- 260 11. MacIntyre, C.R., and Chughtai, A.A. Facemasks for the prevention of infection in
261 healthcare and community settings. BMJ. 2015;350:h694