**QUEEN'S
UNIVERSITY
BELFAST**

Evolving southwest African response to abrupt deglacial North Atlantic climate change events

Chase, B. M., Boom, A., Carr, A. S., Carré, M., Chevalier, M., Meadows, M. E., Pedro, J. B., Stager, J. C., & Reimer, P. J. (2015). Evolving southwest African response to abrupt deglacial North Atlantic climate change events. *Quaternary Science Reviews*, 121, 132-136. <https://doi.org/10.1016/j.quascirev.2015.05.023>

Published in:
Quaternary Science Reviews

Document Version:
Early version, also known as pre-print

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© 2015 The authors

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Evolving southwest African response to abrupt deglacial North Atlantic climate change events

Brian M. Chase^{a*}, Arnoud Boom^b, Andrew S. Carr^b, Matthieu Carré^a, Manuel Chevalier^a, Michael E. Meadows^c, Joel B. Pedro^d, J. Curt Stager^e, and Paula J. Reimer^f

^aCentre National de Recherche Scientifique, UMR 5554, Institut des Sciences de l'Evolution de Montpellier, Département Environnements, Université Montpellier 2, Bat.22, CC061, Place Eugène Bataillon, 34095 Montpellier, cedex 5, France.

^bDepartment of Geography, University of Leicester, Leicester, LE1 7RH, UK.

^cDepartment of Environmental and Geographical Science, University of Cape Town, Private Bag X3, Rondebosch 7701, South Africa.

^dCentre for Ice and Climate, University of Copenhagen, DK-2100, Copenhagen, Denmark.

^eNatural Sciences, Paul Smith's College, Paul Smiths, NY 12970, USA.

^fSchool of Geography, Archaeology and Palaeoecology, Queen's University Belfast, Belfast, BT7 1NN, Northern Ireland, UK.

*Correspondence to: brian.chase@univ-montp.fr

1 **Highlights**

- 2 • 19.4 kyr multi-proxy records from a rock hyrax midden from SW Africa.
- 3 • Aridification events at 34°S concurrent with the Younger Dryas and 8.2 ka events.
- 4 • Influence of bipolar-seesaw replaced by atmospheric teleconnection after 14.6 ka.

Abstract

Climate change during the last deglaciation was strongly influenced by the ‘bipolar seesaw’, producing antiphase climate responses between the North and South Atlantic. However, mounting evidence demands refinements of this model, with the occurrence of abrupt events in southern low to mid latitudes occurring in-phase with North Atlantic climate. Improved constraints on the north-south phasing and spatial extent of these events are therefore critical to understanding the mechanisms that propagate abrupt events within the climate system. We present a 19,400 year multi-proxy record of climate change obtained from a rock hyrax midden in southernmost Africa. Arid anomalies in phase with the Younger Dryas and 8.2 ka events are apparent, indicating a clear shift in the influence of the bipolar seesaw, which diminished as the Earth warmed, and was succeeded after ~14.6 ka by the emergence of a dominant interhemispheric atmospheric teleconnection.

Keywords: southern Africa, palaeoclimate, hyrax middens, bipolar seesaw, Atlantic Overturning Meridional Circulation

Introduction

While some studies have reported interhemispheric synchrony and symmetry during extreme climate disturbances such as the Younger Dryas cold reversal (YD; 12.9-11.6 ka (Lowe et al., 2008)) (Denton and Hendy, 1994; Goede et al., 1996), abrupt changes in Northern Hemisphere climates (North Greenland Ice Core Project members, 2004) have also been associated with anti-phase responses in the Southern Hemisphere (Kaplan et al., 2010; Putnam et al., 2010). Such antiphase responses are hypothesised to be driven by the oceanic Atlantic Overturning Meridional Circulation (AMOC), which draws heat from the Southern Hemisphere into the North Atlantic, but which is sensitive to disruption by ice and freshwater discharges (Broecker, 1998; McManus et al., 2004). Reduction and intensification of ocean heat transport during northern stadial (cold) and interstadial (warm) intervals leads to the alternating build-up and extraction of Southern Hemisphere heat; the so-called bipolar seesaw (Broecker, 1998; Stocker, 1998; Stocker and Johnsen, 2003).

An increasing number of records suggest that the relative warmth of the Northern Hemisphere's Bølling-Allerød interstadial coincided with the Antarctic Cold Reversal (ACR; 14.7-13.0 ka) (Pedro et al., 2011; Putnam et al., 2010), and that the marked northern cooling of the YD was a period of rising temperatures and glacial retreat in the southern high (Pedro et al., 2011) to mid-latitudes (Kaplan et al., 2010). However, a lack of reliable evidence from the low southern latitudes has still prevented a full assessment of the bipolar seesaw hypothesis, including the location of its 'fulcrum'. Such information is vital to test simulations, which are currently showing no consensus on the spatial extent of past (or future) abrupt climate change events (Kageyama et al., 2010).

To address this problem, we explore the regional impact of key perturbations in the North Atlantic using a multi-proxy record from the arid SW Cape region of South Africa (Fig. 1). The region lies at the juncture between southern Africa's three dominant climate systems: the South Atlantic anticyclone, the tropical easterlies, and the austral westerlies (Tyson, 1986). Approximately 75% of the region's precipitation falls during winter, when the westerlies and their related cold fronts migrate northward, advecting moisture from the southern Atlantic to the mountains of the SW Cape (Reason et al., 2006). In the dry summer months, the westerlies and the South Atlantic Anticyclone shift southward, limiting frontal system influence and blocking tropical moisture-bearing systems from the Indian Ocean and tropical Atlantic (Reason et al., 2006).

Little is known about SW Africa's environmental history, mainly due to its aridity and marked rainfall seasonality, which allows for few wetland sediment records. Rock hyrax (*Procavia capensis*) middens have emerged in this setting as valuable archives of palaeoenvironmental information (Carr et al., 2010; Chase et al., 2013; Chase et al., 2015; Chase et al., 2009; Chase et al., 2011; Chase et al., 2012; Meadows et al., 2010; Quick et al., 2011; Scott and Bousman, 1990; Scott et al., 2005; Scott and Vogel, 2000; Scott and Woodborne, 2007). As hyraxes use discrete locations as latrines, deposits of sub-fossilised urine (hyraceum) accumulate in their shelters, much like stalagmites in a cave. These finely laminated amber-like deposits preserve a wide range of proxies, including pollen, charcoal, and stable isotopes, all of which can provide insight into past environmental conditions (Chase et al., 2011; Valsecchi et al., 2013) (see Supplementary Information).

Results

The records presented here were obtained from two sections of a 53 cm thick midden collected from De Rif, in the Driehoek Valley of the Cederberg Mountains (32°26'45"S, 19°13'15"E, 1151 m amsl.) (Chase et al., 2011; Valsecchi et al., 2013). Chronologies spanning the past 19,400 years were established using 29 ^{14}C AMS dates (see Supplementary Information). Together, the De Rif midden records reveal coherent patterns of marked environmental variability since the Last Glacial Maximum (LGM; Fig. 2). Highlighted here are aspects of the records that primarily reflect changes in hydroclimate. In a region dominated by C_3 plants, the hyraceum $\delta^{13}\text{C}$ record primarily reflects variations in water-use efficiency (Chase et al., 2012; Ehleringer and Cooper, 1988; Farquhar et al., 1989; Farquhar and Richards, 1984; Pate, 2001), although a long-term enrichment in is evident across the mid- to late Holocene. This is consistent with increased water use efficiency of C_3 plants, and an increasing abundance of $\delta^{13}\text{C}$ enriched drought-resistant succulent CAM plants under drier conditions (Smith, 1972; Valsecchi et al., 2013). These data are supported by the hyraceum $\delta^{15}\text{N}$ record, which also reflects water-availability (Chase et al., 2012; Handley et al., 1999; Handley et al., 1994; Hartman, 2011; Heaton, 1987; Murphy and Bowman, 2006, 2009; Wang et al., 2010), as well as by fossil pollen data (Valsecchi et al., 2013) and derived reconstructions of relative palaeo-aridity. Each of these proxy records expresses variability similar to that observed in regional marine core records, confirming that they are reflecting variability in a tightly coupled climate system (Fig. 3).

Although De Rif lies within the core of the winter rainfall zone, our data show that changes in the duration or intensity of the summer drought season were important drivers of environmental change at this site for much of the last 19 kyr. Whereas increases in winter rainfall would result in a net increase in annual rainfall, increased precipitation in the summer drought season would have a significantly greater impact on reducing drought-stress in the region (Chase

et al., 2015). This is reflected by trends in the percentage of drought-tolerant and intolerant taxa (Valsecchi et al., 2013) and aridity index reconstructions (Fig. 2d). While a degree of variability is evident between these records as a function of their specific sensitivities, each reflects aspects of changes in drought season length and/or intensity, and corresponds well with overall changes in water availability inferred from the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ records (Fig. 2). These findings are supported by the CCSM3 TraCE-21ka general circulation model (GCM) simulations, which show qualitative agreement between austral summer precipitation in the region and the proxy records from the De Rif middens (He et al., 2013).

The De Rif data, particularly the higher resolution $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ records, highlight the impact of three major freshening events: HS1, the YD and the 8.2 ka events. We observe two primary phases in the region's deglacial climatic evolution: (1) a general increase in moisture availability and reduced seasonality from the terminal LGM to the end of HS1 at ~14.6 ka (Fig. 2b, c), and (2) a late deglacial/early Holocene period from ~14.6 to 7 ka marked by significant, clear arid episodes synchronous with the YD and 8.2 ka event (Fig. 2b, c).

Discussion

The early deglacial period is defined by the abrupt decline of the AMOC during HS1 (Fig. 3b), and the slow build-up of South Atlantic heat from ~18-14.6 ka that is registered in Antarctic ice cores (Jouzel et al., 2007; Pedro et al., 2011) (Fig. 3h), SE Atlantic sea-surface temperatures (SSTs) (Farmer et al., 2005; Kim and Schneider, 2003) and resulted in increased humidity at De Rif (Fig. 3d, e). While these changes likely affected the Subtropical Front (Barker et al., 2009) (Fig. 3h), resulting in a poleward shift of the moisture-bearing systems associated with the westerly storm track, we interpret that compensating factors such as increased flow of warm Agulhas Current waters into the SE Atlantic and reduced northward heat transport in the AMOC

110 favoured warming SSTs and increasing advection influence of tropical easterlies in the
111 southwestern Cape(Reason et al., 2006). This would have resulted in an increase in summer rain
112 and a shorter/less intense drought season (Chase et al., 2015; He et al., 2013).

113 At ~14.6 ka, the rapid increase of AMOC created an immediate cooling response in the
114 South Atlantic, as the Subtropical Front shifted equatorward at least as far as ~41°S (Barker et
115 al., 2009). In SW Africa, however, the impact was more muted (Fig. 3). In the northern (Kim et
116 al., 2002) and central (Farmer et al., 2005) Benguela system, this period marked the end of the
117 warming trend that began at ~19 ka, but the subsequent cooling was relatively slow, similar to
118 southern (Pedro et al., 2011) and northern(North Greenland Ice Core Project members, 2004)
119 polar records during the Bølling-Allerød interstadial ~14.7-12.9 ka (Lowe et al., 2008). In the
120 SW Cape, the response to these changes was a clear reduction in humidity (Fig. 3f, g). This
121 aridification, however, was short-lived, lasting only through the Bølling interstadial (14.7-14.1
122 ka (Lowe et al., 2008)), whereas the Allerød interstadial (14.1-12.9 ka (Lowe et al., 2008)) saw
123 increased water availability at De Rif. Compared to the regionally coherent HS1 signal, the
124 spatial heterogeneity of responses during this time suggests a restructuring of Earth's climate
125 system, with the increasing influence of the South Atlantic Anticyclone in the southern
126 subtropics across the last deglaciation.

127 This restructuring is underscored by the regional response to the subsequent YD. While
128 the Northern Hemisphere cooling coincides with a distinct decrease in AMOC, the SE Atlantic
129 and SW African response is in sharp contrast to the HS1 signal, with an abrupt drop in SSTs
130 (Farmer et al., 2005; Kim et al., 2002) and marked aridity at De Rif. In contrast with the slow
131 build-up of heat during HS1 the immediate response during the YD implies a dominant
132 atmospheric interhemispheric teleconnection (cf. Moreno et al., 2001), inconsistent with the

oceanic controls related to the bipolar seesaw. The De Rif records further reveals an significant (cf. Morrill and Jacobsen, 2005) and abrupt drying signal relating to the 8.2 ka event (Barber et al., 1999), indicating that even relatively small fresh water pulses in the North Atlantic (Clarke et al., 2004) produced immediate responses in SW Africa, resulting in significant aridification events in the region (Fig. 3g, h).

This dramatic contrast in response to perturbations in the North Atlantic (more humid at De Rif during the AMOC slow-down of HS1 and more arid during the slow-downs of the YD and 8.2 ka events) challenges any systematic application of the bipolar-seesaw model of north-south phasing during abrupt climate perturbations and raises questions on the relative roles of oceanic and atmospheric teleconnections in driving SE Atlantic and SW African climate. Of particular importance here is the influence of the South Atlantic Anticyclone, which dominates atmospheric circulation in the SE Atlantic basin and has a significant impact on oceanographic conditions through its regulation of upwelling intensity (Farmer et al., 2005; Kim and Schneider, 2003; Kim et al., 2003).

Either alternatively or in concert with the potential influence of a bipolar seesaw, a weakening South Atlantic Anticyclone during HS1 would have reduced upwelling intensity and raised SSTs in the Benguela system. After ~14 ka, however, the impact of the bipolar seesaw as a driver of climatic variability was apparently not expressed in the SE Atlantic. The rapidity and direction of the SW African responses to the changes in the North Atlantic that induced the YD and 8.2 ka events (Barber et al., 1999; Murton et al., 2010) strongly implicate an atmospheric rather than oceanic interhemispheric teleconnection, and the acute sensitivity of this relationship is illustrated by the influence of the relatively minor freshwater outburst that triggered the cooling of the 8.2 ka event (Barber et al., 1999). This implies that an early dominance of the

bipolar seesaw was replaced - perhaps as a result of diminishing high latitude ice sheets and a related reduction in the intensity and impact of declines in AMOC (McManus et al., 2004; Ritz et al., 2013) - in favour of more immediate atmospheric teleconnections, (Fig. 3b). This effectively displaced the boundary between positive and negative SST anomalies related to perturbations in the North Atlantic by at least 24°, shifting the ‘fulcrum’ of the bipolar seesaw poleward of the African continent, where at 41°S, and contrary to conditions in the SE Atlantic, warmer SSTs occurred during the YD (Barker et al., 2009). These findings pose an exciting challenge as they call for closer consideration of the spatiotemporal influence of the bipolar seesaw, and identify areas for refinement of Earth System Models, which may lead to a more complete understanding of global climate dynamics.

Acknowledgments: Funding was received from the European Research Council (ERC) under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Starting Grant “HYRAX”, grant agreement no. 258657. The South African National Biodiversity Institute is thanked for the use of data/information supplied by SANBI from digitized collections.

References

- Barber, D.C., Dyke, A., Hillaire-Marcel, C., Jennings, A.E., Andrews, J.T., Kerwin, M.W., Bilodeau, G., McNeely, R., Southon, J., Morehead, M.D., Gagnon, J.M., 1999. Forcing of the cold event of 8,200 years ago by catastrophic drainage of Laurentide lakes. *Nature* 400, 344-348.
- Barker, S., Diz, P., Vautravers, M.J., Pike, J., Knorr, G., Hall, I.R., Broecker, W.S., 2009. Interhemispheric Atlantic seesaw response during the last deglaciation. *Nature* 457, 1097-1102.
- Broecker, W.S., 1998. Paleocirculation during the last deglaciation: a bipolar seesaw? *Paleoceanography* 13, 119-121.
- Carr, A.S., Boom, A., Chase, B.M., 2010. The potential of plant biomarker evidence derived from rock hyrax middens as an indicator of palaeoenvironmental change. *Palaeogeography, Palaeoclimatology, Palaeoecology* 285, 321-330.
- Chase, B.M., Boom, A., Carr, A.S., Meadows, M.E., Reimer, P.J., 2013. Holocene climate change in southernmost South Africa: rock hyrax middens record shifts in the southern westerlies. *Quaternary Science Reviews* 82, 199-205.
- Chase, B.M., Lim, S., Chevalier, M., Boom, A., Carr, A.S., Meadows, M.E., Reimer, P.J., 2015. Influence of tropical easterlies in southern Africa's winter rainfall zone during the Holocene. *Quaternary Science Reviews* 107, 138-148.
- Chase, B.M., Meadows, M.E., Scott, L., Thomas, D.S.G., Marais, E., Sealy, J., Reimer, P.J., 2009. A record of rapid Holocene climate change preserved in hyrax middens from southwestern Africa. *Geology* 37, 703-706.

Chase, B.M., Quick, L.J., Meadows, M.E., Scott, L., Thomas, D.S.G., Reimer, P.J., 2011. Late glacial interhemispheric climate dynamics revealed in South African hyrax middens. *Geology* 39, 19-22.

Chase, B.M., Scott, L., Meadows, M.E., Gil-Romera, G., Boom, A., Carr, A.S., Reimer, P.J., Truc, L., Valsecchi, V., Quick, L.J., 2012. Rock hyrax middens: a palaeoenvironmental archive for southern African drylands. *Quaternary Science Reviews* 56, 107-125.

Chevalier, M., Cheddadi, R., Chase, B.M., 2014. CREST: a *pdf*-based quantitative climate reconstruction method. *Climate of the Past*.

Clarke, G.K.C., Leverington, D.W., Teller, J.T., Dyke, A.S., 2004. Paleohydraulics of the last outburst flood from glacial Lake Agassiz and the 8200 BP cold event. *Quaternary Science Reviews* 23, 389-407.

Denton, G.H., Hendy, C.H., 1994. Younger Dryas age advance of Franz Josef Glacier in the Southern Alps of New Zealand. *Science* 264, 1434-1437.

Ehleringer, J.R., Cooper, T.A., 1988. Correlations between carbon isotope ratio and microhabitat of desert plants. *Oecologia* 76, 562-566.

Farmer, E.C., deMenocal, P.B., Marchitto, T.M., 2005. Holocene and deglacial ocean temperature variability in the Benguela upwelling region: implications for low-latitude atmospheric circulation. *Paleoceanography* 20, doi:10.1029/2004PA001049.

Farquhar, G.D., Ehleringer, J.R., Hubick, K.T., 1989. Carbon isotope discrimination and photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology* 40, 503-537.

Farquhar, G.D., Richards, R.A., 1984. Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes. *Australian Journal of Plant Physiology* 11, 539-552.

Goede, A., McDermott, F., Hawkesworth, C., Webb, J., Finlayson, B., 1996. Evidence of Younger Dryas and neoglacial cooling in a late Quaternary palaeotemperature record from a speleothem in eastern Victoria, Australia. *Journal of Quaternary Science* 11, 1-7.

Handley, L.L., Austin, A.T., Stewart, G.R., Robinson, D., Scrimgeour, C.M., Raven, J.A., Heaton, T.H.E., Schmidt, S., 1999. The $\delta^{15}\text{N}$ natural abundance ($\delta^{15}\text{N}$) of ecosystem samples reflects measures of water availability. *Functional Plant Biology* 26, 185-199.

Handley, L.L., Odee, D., Scrimgeour, C.M., 1994. $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ patterns in savanna vegetation: dependence on water availability and disturbance. *Functional Ecology* 8, 306-314.

Hartman, G., 2011. Are elevated $\delta^{15}\text{N}$ values in herbivores in hot and arid environments caused by diet or animal physiology? *Functional Ecology* 25, 122-131.

He, F., Shakun, J.D., Clark, P.U., Carlson, A.E., Liu, Z., Otto-Bliesner, B.L., Kutzbach, J.E., 2013. Northern Hemisphere forcing of Southern Hemisphere climate during the last deglaciation. *Nature* 494, 81-85.

Heaton, T.H.E., 1987. The $^{15}\text{N}/^{14}\text{N}$ ratios of plants in South Africa and Namibia: relationship to climate and coastal/saline environments. *Oecologia* 74, 236-246.

Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J.M., Chappellaz, J., Fischer, H., Gallet, J.C., Johnsen, S., Leuenberger, M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J.P., Stenni, B., Stocker, T.F., Tison, J.L., Werner, M., Wolff, E.W., 2007. Orbital and millennial Antarctic climate variability over the past 800,000 years. *Science* 317, 793-797.

Kageyama, M., Paul, A., Roche, D.M., Van Meerbeeck, C.J., 2010. Modelling glacial climatic millennial-scale variability related to changes in the Atlantic meridional overturning circulation: a review. *Quaternary Science Reviews* 29, 2931-2956.

Kaplan, M.R., Schaefer, J.M., Denton, G.H., Barrell, D.J.A., Chinn, T.J.H., Putnam, A.E., Andersen, B.G., Finkel, R.C., Schwartz, R., Doughty, A.M., 2010. Glacier retreat in New Zealand during the Younger Dryas stadial. *Nature* 467, 194-197.

Kim, J.-H., Schneider, R.R., 2003. Low-latitude control of interhemispheric sea-surface temperature contrast in the tropical Atlantic over the past 21 kyrs: the possible role of SE trade winds. *Climate Dynamics* 23, 337-347.

Kim, J.-H., Schneider, R.R., Hebbeln, D., Muller, P.J., Wefer, G., 2002. Last deglacial sea-surface temperature evolution in the southeast Pacific compared to climate changes on the South American continent. *Quaternary Science Reviews* 21, 2085-2097.

Kim, J.-H., Schneider, R.R., Mulitza, S., Müller, P.J., 2003. Reconstruction of SE trade-wind intensity based on sea-surface temperature gradients in the Southeast Atlantic over the last 25 kyr. *Geophysical Research Letters* 30, 2144.

Lowe, J.J., Rasmussen, S.O., Björck, S., Hoek, W.Z., Steffensen, J.P., Walker, M.J.C., Yu, Z.C., 2008. Synchronisation of palaeoenvironmental events in the North Atlantic region during the Last Termination: a revised protocol recommended by the INTIMATE group. *Quaternary Science Reviews* 27, 6-17.

McManus, J.F., Francois, R., Gherardi, J.-M., Keigwin, L.D., Brown-Leger, S., 2004. Collapse and rapid resumption of Atlantic meridional circulation linked to deglacial climate changes. *Nature* 428, 834-837.

Meadows, M.E., Chase, B.M., Selane, M., 2010. Holocene palaeoenvironments of the Cederberg and Swartuggens mountains, Western Cape, South Africa: Pollen and stable isotope evidence from hyrax dung middens. *Journal of Arid Environments* 74, 786-793.

Moreno, P.I., Jacobson G.L., J., Lowell, T.V., Denton, G.H., 2001. Interhemispheric climate links revealed by a Lateglacial cooling episode in southern Chile. *Nature* 409, 804-808.

Morrill, C., Jacobsen, R.M., 2005. How widespread were climate anomalies 8200 years ago? *Geophys. Res. Lett.* 32.

Murphy, B.P., Bowman, D.M.J.S., 2006. Kangaroo metabolism does not cause the relationship between bone collagen $\delta^{15}\text{N}$ and water availability. *Functional Ecology* 20, 1062-1069.

Murphy, B.P., Bowman, D.M.J.S., 2009. The carbon and nitrogen isotope composition of Australian grasses in relation to climate. *Functional Ecology* 23, 1040-1049.

Murton, J.B., Bateman, M.D., Dallimore, S.R., Teller, J.T., Yang, Z., 2010. Identification of Younger Dryas outburst flood path from Lake Agassiz to the Arctic Ocean. *Nature* 464, 740-743.

North Greenland Ice Core Project members, 2004. High-resolution record of Northern Hemisphere climate extending into the last interglacial period. *Nature* 431, 147-151.

Pate, J.S., 2001. Carbon isotope discrimination and plant water-use efficiency: case scenarios for C_3 plants, in: Unkovich, M., Pate, J., McNeill, A., Gibbs, D.J. (Eds.), *Stable Isotope Techniques in the Study of Biological Processes and Functioning of Ecosystems*. Kluwer Academic Publishers, Dordrecht, pp. 19-37.

Pedro, J.B., van Ommen, T.D., Rasmussen, S.O., Morgan, V.I., Chappellaz, J., Moy, A.D., Masson-Delmotte, V., Delmotte, M., 2011. The last deglaciation: timing the bipolar seesaw. *Climates of the Past* 7, 671-683.

Putnam, A.E., Denton, G.H., Schaefer, J.M., Barrell, D.J.A., Andersen, B.G., Finkel, R.C., Schwartz, R., Doughty, A.M., Kaplan, M.R., Schluchter, C., 2010. Glacier advance in southern middle-latitudes during the Antarctic Cold Reversal. *Nature Geoscience* 3, 700-704.

Quick, L.J., Chase, B.M., Meadows, M.E., Scott, L., Reimer, P.J., 2011. A 19.5 kyr vegetation history from the central Cederberg Mountains, South Africa: Palynological evidence from rock hyrax middens. *Palaeogeography, Palaeoclimatology, Palaeoecology* 309, 253-270.

Reason, C.J.C., Landman, W., Tennant, W., 2006. Seasonal to decadal prediction of southern African climate and its links with variability of the Atlantic ocean. *Bulletin of the American Meteorological Society* 87, 941-955.

Ritz, S.P., Stocker, T.F., Grimalt, J.O., Menviel, L., Timmermann, A., 2013. Estimated strength of the Atlantic overturning circulation during the last deglaciation. *Nature Geosci* 6, 208-212.

Scott, L., Bousman, C.B., 1990. Palynological analysis of hyrax middens from Southern Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 76, 367-379.

Scott, L., Bousman, C.B., Nyakale, M., 2005. Holocene pollen from swamp, cave and hyrax dung deposits at Blydefontein (Kikvorsberge), Karoo, South Africa. *Quaternary International* 129, 49-59.

Scott, L., Vogel, J.C., 2000. Evidence for environmental conditions during the last 20,000 years in Southern Africa from ^{13}C in fossil hyrax dung. *Global and Planetary Change* 26, 207-215.

Scott, L., Woodborne, S., 2007. Vegetation history inferred from pollen in Late Quaternary faecal deposits (hyraceum) in the Cape winter-rain region and its bearing on past climates in South Africa. *Quaternary Science Reviews* 26, 941-953.

Smith, B.N., 1972. Natural abundance of the stable isotopes of carbon in biological systems. *BioScience* 22, 226-231.

Stocker, T.F., 1998. The seesaw effect. *Science* 282, 61-62.

Stocker, T.F., Johnsen, S.J., 2003. A minimum thermodynamic model for the bipolar seesaw. *Paleoceanography* 18.

Tyson, P.D., 1986. *Climatic Change and Variability in Southern Africa*. Oxford University Press, Cape Town.

Valsecchi, V., Chase, B.M., Slingsby, J.A., Carr, A.S., Quick, L.J., Meadows, M.E., Cheddadi, R., Reimer, P.J., 2013. A high resolution 15,600-year pollen and microcharcoal record from the Cederberg Mountains, South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 387, 6-16.

Wang, L., D'Odorico, P., Ries, L., Macko, S.A., 2010. Patterns and implications of plant-soil $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in African savanna ecosystems. *Quaternary Research* 73, 77-83.

Figure captions

Fig. 1. Map of study region indicating generalised atmospheric (white arrows) and oceanic (blue arrows) circulation systems and the seasonal distribution of rainfall as indicated by the percent of the total mean annual rainfall received during the austral winter months of April-September. The convergence zones of the Congo Air Boundary (CAB) and the Intertropical Convergence Zone (ITCZ) are shown in the austral summer positions. Key sites discussed in the manuscript are indicated by number: 1) GeoB1023-5 (Kim and Schneider, 2003; Kim et al., 2003); 2) ODP1084b (Farmer et al., 2005); 3) the De Rif rock hyrax midden; and 4) TNO57-21 (Barker et al., 2009).

Fig. 2. Comparison of proxy records from the De Rif rock hyrax midden and general circulation model (GCM) simulation data for the region. Radiocarbon ages shown as triangles along x-axis (DR2010 section in red, DR-2 section in blue). Heinrich stadial 1 (HS1), the Younger Dryas cold reversal (YD) and 8.2 ka event are highlighted by blue shading, and the Bølling (B) and Allerød (A) interstadials are shaded in red. Stable nitrogen (b) and carbon (c) records from the middens primarily reflect water-availability in the environment and water-use efficiency of plants respectively. These data are confirmed by pollen analysis of the De Rif midden (Valsecchi et al., 2013) and an aridity index reconstruction using a *pdf*-based modelling technique (Chevalier et al., 2014) applied to the De Rif pollen assemblage (d; shading indicates 20% and 50% errors), which indicate the importance of drought season intensity and length in determining environmental change in the region. First-order similarities between results from the CCSM3 TraCE-21ka transient GCM simulation (He et al., 2013) of austral summer (DJF) (a), and the proxy records support these conclusions, but the lack of a significant Younger Dryas signal in the simulation suggests that the model may not be capturing certain important elements of the global deglacial climate system.

Fig. 3. Comparison of proxy records from the De Rif rock hyrax midden with independent regional and extra-regional records reflecting changes in a series of related climate systems during the last 20,000 years. Radiocarbon ages shown as triangles along x-axis (DR2010 section in red, DR-2 section in blue). Heinrich stadial 1 (HS1), the Younger Dryas cold reversal (YD) and 8.2 ka event are highlighted by blue shading, and the Bølling (B) and Allerød (A) interstadials are shaded in red. Climatic perturbations in the North Atlantic basin are recorded in the NGRIP ice core record from Greenland (a) (North Greenland Ice Core Project members, 2004) and have been observed to have a significant impact on the Atlantic Meridional Overturning Circulation (AMOC) and the northward oceanic transport of heat (b) (McManus et al., 2004), resulting in an antiphase relationship between northern (a) and southern (i) hemisphere temperatures (Broecker, 1998; Stocker, 1998). This is indicated by records from the South Atlantic (g, h) (Barker et al., 2009) and the southern polar regions (i) (Jouzel et al., 2007; Pedro et al., 2011). While from ~18-14.6 ka this trend may have been expressed in SE Atlantic (c, d) (Farmer et al., 2005; Kim and Schneider, 2003; Kim et al., 2003) and SW Africa (e, f), variability in the intensity of the South Atlantic Anticyclone (c, d) (Farmer et al., 2005; Kim and Schneider, 2003; Kim et al., 2002; Kim et al., 2003) provide a coherent complimentary (Kim et al., 2002) mechanism, and highlight the increasing importance of atmospheric teleconnections with the North Atlantic in driving SW African climate change across the deglacial period.

*Figure 1

*Figure 2 REVISED

Supplementary Data

[Click here to download Supplementary Data: Chase et al_QSR_SI_REVISED_FINAL.docx](#)

Table S1
[Click here to download Supplementary Data: TableS1.xlsx](#)

Supplementary Data

[Click here to download Supplementary Data: CHASE_FIGS1.eps](#)

Supplementary Data

[Click here to download Supplementary Data: CHASE_FIG S2.eps](#)

Supplementary Data

[Click here to download Supplementary Data: SI_DeRif_d15N-d13C_Pollen_Reconstructions data.xlsx](#)

KML File (for GoogleMaps)

[Click here to download KML File \(for GoogleMaps\): De Rif hyrax midden.kml](#)