**QUEEN'S
UNIVERSITY
BELFAST**

Overview of *Taenia solium* cysticercosis in West Africa

Weka, R., Kamani, J., Cogan, T., Eisler, M., & Morgan, E. (2019). Overview of *Taenia solium* cysticercosis in West Africa. *Acta Tropica*, 190, 329-338.

Published in:
Acta Tropica

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2018 Elsevier.

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License

(<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Overview of *Taenia solium* cysticercosis in West Africa

Rebecca Paul Weka^{a,b,*}, Joshua Kamani^a, Tristan Cogan^b, Mark Eisler^b, Eric R. Morgan^{b,c}

^aParasitology Division, National Veterinary Research Institute, PMB 01 Vom, Plateau State, Nigeria

^bBristol Veterinary School, University of Bristol, Langford, Bristol BS40 5DU, United Kingdom

^cSchool of Biological Sciences, Queen's University Belfast, 97, Lisburn Road, Belfast, BT9 7BL, United Kingdom

*Corresponding author: bekkyweka@ gmail.com

Abstract

Human and porcine cysticercosis is endemic in West Africa, where epilepsy is relatively common, but rarely reported due to fear of stigmatization. Neurocysticercosis, caused by *Taenia solium*, tends to affect the poor in developing countries and control is hampered by inadequate infrastructure and financial resources coupled with lack of adequate information about its significance and distribution. The risk factors for human cysticercosis are closely associated with the characteristics of smallholder or backyard pig farming systems prevalent in this region. Poverty, ignorance and lack of political will militate against successful eradication, while tools for effective diagnosis, prevention and treatment, including vaccines for pigs, are not yet available in many countries. Cysticercosis was targeted for control by The World Health Organization global plan for 2008-2015; however, measures for control are yet to be undertaken in a coordinated manner in West Africa. Diagnostic tools, including neuroimaging facilities, should be strategically situated and made accessible to rural populations in West Africa. Community education in combination with a multipronged approach consisting of vaccination of pigs using TSOL18 vaccine and treatment with oxfendazole along with mass chemotherapy of humans with praziquantel could eliminate taeniasis and should be considered in the region. In-depth and extensive epidemiological studies are required in the region in order to fully elucidate the prevalence of *T. solium* and to support more coordinated and effective control of human and porcine cysticercosis.

Keywords: *Taenia solium*; poverty; food safety; backyard pig farming; epilepsy; West Africa.

1. Introduction

Taenia solium is a tapeworm of humans whose larval stage is normally found in pig muscle, but can also invade the human central nervous system to cause neurocysticercosis (Lightowlers, 2013; Garcia *et al.*, 2014b), and is thus a significant cause of acquired preventable epilepsy (Winkler, 2013; WHO 2016a). Ndimubanzi *et al.* (2010) reported that neurocysticercosis causes 29% of acquired epilepsy in endemic countries, and epilepsy in less developed countries is responsible for mortality rates 3–6 times higher than in developed nations (WHO, 2016a) due to inadequate infrastructure and poor access to hospitals. This zoonotic tapeworm therefore constitutes a serious, but preventable, public health problem, as well as impacting on agriculture (Ngowi *et al.*, 2013; Braae *et al.*, 2016a), and is considered a neglected parasite (WHO, 2016a; Johansen *et al.*, 2016). The disease is an emergent and persistent problem in most under-developed areas of Sub-Saharan Africa, Asia and Latin America; except for in areas mainly populated by Muslims, who do not eat pig meat (Fleury *et al.*, 2015).

Although *T. solium* has been considered an eradicable disease since 1993 by the International Task Force on Disease Eradication (ITFDE), to date none of the endemic countries has been able to eradicate this disease, despite the advances made in the development of appropriate tools for diagnosis, treatment and prevention, neurocysticercosis still persists at disturbingly high levels in endemic areas of Africa (Johansen *et al.*, 2016). Control of the disease in such regions is hampered by poverty, inadequate infrastructure, financial resource constraints, and unsanitary conditions (Lightowlers *et al.*, 2015; Johansen *et al.*, 2016). Furthermore, in Africa, ignorance surrounding the presence, magnitude and impacts of the parasite by stakeholders and political authorities have led to the disease being often unrecognized and under reported (Assana *et al.*, 2013; WHO 2016a). This is further complicated by scarcity of information on the disease due to

lack of overt disease manifestation in many cases in both humans and pigs (Fleury *et al.*, 2015; Ron-Garrido *et al.*, 2015).

In recent years, the number of studies on *T. solium* in Africa has increased (Gabriël *et al.*, 2016), helping to further elucidate the epidemiology of the parasite in sub Saharan Africa (Assana *et al.*, 2013). The purpose of this review is to summarise this new information in West Africa in the context of existing knowledge on the parasite, and to apply it to identify gaps and opportunities in research and control of this potentially devastating parasitic disease.

2. Life cycle and disease

Taenia solium has a two-stage life cycle. The adult tapeworm lives in the human small intestine after eating viable cysticerci (“pork measles”) in raw or under-cooked pork resulting in taeniasis. The life cycle is completed when the proglottids become gravid, harboring thousands of the infective oncospheres, and are shed from the tapeworm and ingested in the faeces of infected individuals. Oncospheres (=eggs) are then dispersed and contaminate the environment.

Humans can act as accidental dead-end intermediate hosts and develop cysticercosis following ingestion of tapeworm eggs. Auto-infection can also occur when proglottids reach the stomach by reverse peristalsis, resulting in massive infection. The hatched larvae penetrate the intestinal mucosa and migrate throughout the body, forming cysticerci, which commonly lodge in skeletal muscles, eyes and subcutaneous tissues, but have a particular predilection for the central nervous system (CNS), causing neurocysticercosis (NCC) ('Gal'an-Puchades', 2016).

Pigs are the main intermediate hosts and ingest *T. solium* eggs contained in infected human feces or through ingestion of contaminated water or feed). After ingestion, the oncospheres evaginate and the hatched larvae penetrate the intestinal walls into the bloodstream and develop to

metacestodes, forming cysticerci throughout the body but mostly in the striated muscles of the pigs (WHO 2016a).

2.1 Clinical signs in pigs

Infected pigs are most often asymptomatic, but can rarely show neurological signs, which can manifest as dullness, sluggishness, loss of consciousness, and in some cases hypersensitivity to sound as well as seizures as in humans. Quivering, paralysis of the ear, ataxia, dribbling saliva, circling movement, decreased production performance and social isolation have also been reported (Mkupasi *et al.*, 2014; Trevisan *et al.*, 2016, 2017).

2.2 Disease in humans

The clinical manifestations of NCC differ, varying from asymptomatic to severe (Winkler, 2013; Trevisan *et al.*, 2016). Seizures are the most common presentation of NCC in humans (Carpio and Romo, 2014) occurring in about 60-90% of symptomatic infections (Rodrigues *et al.*, 2012; Carpio *et al.*, 2013), although incidence varies in different regions according to socioeconomic and cultural practices (Del Brutto, 2013b; Winkler, 2013). Headache (Carabin *et al.*, 2011; Johansen *et al.*, 2016), focal neurological signs (Fleury *et al.*, 2011; Flisser, 2013) and generalized weakness associated with muscle pseudohypertrophy (Fleury *et al.*, 2015) are also common presentations noticed.

It has been estimated that between 2.6 and 8.3 million individuals are affected globally by symptomatic and asymptomatic NCC, and that 0.76-2.46 million of the NCC-related epileptic individuals and 0.95-3.08 million of the asymptomatic NCC individuals are in sub-Saharan Africa (Winkler, 2013), resulting in 2–5 million lost disability-adjusted life years (WHO, 2015b).

3. Diagnosis

3.1 Diagnosis of porcine cysticercosis

Diagnosis of porcine cysticercosis is essential for control, to prevent taeniasis in humans and further onward infections in pigs as well as NCC in humans (Gilman *et al.*, 2012).

3.1.1 Tongue and carcass inspection

The simplest and most common diagnostic methods for porcine cysticercosis are visual inspection of the surface of the tongue, which has a sensitivity ranging from 16-70% (Phiri *et al.*, 2006), and carcass inspection (Lightowlers *et al.*, 2015). Most epidemiological studies on

porcine cysticercosis in West Africa have used tongue or carcass inspection (Onah and Cheinjina 1995; Gweba *et al.*, 2010; Goussanou *et al.*, 2014; Edia-Asuke *et al.*, 2014; Attawalna *et al.*, 2015; Idiaka *et al.*, 2017).

Gap: Tongue and carcass inspection are commonly applied to check meat safety in West Africa, however can inadvertently lead to the entrance of a large amount of pork infected with cysticercosis into the human food chain at a reduced price, due to the low sensitivity of the technique (Goussanou *et al.*, 2014). Exemplified by studies of Onah and Cheinjina (1995) who observed 5.5% (72/1300) of pigs were infected with cysticercosis infection based on tongue inspection while carcass inspection also showed that all 72 were infected, including the 8.5% (104) that were negative by tongue inspection. Furthermore Gweba *et al.* (2010) also observed a prevalence of 5.85% (n = 205) and 14.40% (n = 118) by lingual palpation and postmortem examination. The infected meat could be a source of infection to the public if mechanisms are not in place for the proper disposal of infected meat (Gonzalez *et al.*, 1990). Hence studies by Lightowers *et al.* (2015) indicate that slicing the tongue, masseter muscle and heart is highly specific and detects natural infection in pigs with 80% sensitivity at a low cost, especially because these are convenient anatomical sites which will not affect the carcass value hence such technique should be practiced in West Africa.

3.1.2 Serology in pigs

Serological tests for the detection of specific antibodies or specific antigens use enzyme-linked immunosorbent assay (ELISA) and electro-immuno transfer blot (EITB) (Lightowers *et al.*, 2016). Most of the few serological studies of pigs in West Africa have used B158/B160 Ag-

ELISA (Secka *et al.*, 2010a; Ganaba *et al.*, 2011) and studies of Weka *et al.* (2009) used IgG antibody ELISA.

Gap; Lack of serological diagnostic kits in West Africa especially due to cost is a serious obstacle to providing detailed epidemiological data hence, availability of serological kits in the region will help in diagnosis of porcine cysticercosis. However, detection of antibodies does not necessarily imply active infection as serologically positive pigs may not have cysticerci at necropsy (Jayashi *et al.*, 2012; Devleesschauwer *et al.* 2013; Lightowlers *et al.*, 2016). Moreover, false positive or transient positive reactions might arise from exposure to *T. solium* eggs that did not develop to cysticerci to be detected at carcass inspection (Lightowlers *et al.*, 2016). Generally, antigen detection methods are by far superior to antibody detection techniques since they indicate the presence of viable cysts. Although at population level, results can nevertheless give a useful indication of infected areas where the life cycle of the parasite is ongoing (Thomas, 2015). Thus serological test interpretation in individual animals is limited by their sensitivity and specificity (Lightowlers *et al.*, 2016) and should be confirmed by necropsy where possible although purchasing pigs for necropsy is equally expensive.

3.2 Diagnosis in humans

3.2.1 Serology in humans

Serological studies for diagnosis of human cysticercosis has been carried out in West Africa, including the use of Ag-ELISA (Nitiéma *et al.*, 2012; Carabin *et al.*, 2015), Ab- ELISA (Edia-Asume *et al.* 2015; Weka *et al.* 2013) and EITB (Secka *et al.*, 2010a; 2011). The EITB assay developed by Tsang *et al.* (1989), though about 20 times more expensive than ELISA is highly specific (near-100%) and sensitive (70-90%). Antibody seropositive individuals, however, might have been exposed to the parasite or naturally cured and do not necessarily have an established

infection; hence, results should be interpreted in line with the clinical presentation (Gilman *et al.*, 2012). However, antigen-ELISA is highly sensitive, cost effective and technically simpler to implement in resource limited settings and will be an invaluable tool for hospital diagnosis, particularly in the absence of imaging facilities.

Gap; ELISA kits for antibody detection in human cysticercosis are commercially available (Rodriguez *et al.*, 2012) but they are reported to have low sensitivity and frequent cross-reactions (Garcia *et al.*, 2018) hence, the results should be interpreted with caution. Therefore, research laboratories in West Africa should give priority to the development of ELISA using local cysticercal antigens to enhance diagnosis and facilitate the much needed epidemiologic studies in most of Africa. and the financial burden of running the EITB precludes its routine use.

3.2.2 Diagnosis of NCC

Diagnosis of NCC is mainly pursued by neuroimaging techniques, including computerised tomography (CT) and magnetic resonance imaging (MRI), (Gilman *et al.*, 2012; Fleury *et al.*, 2013) which detects the presence of parasitic lesions/ calcified cysts and extraparenchymal lesions, and can give a sensitive and accurate diagnosis of NCC (Nash and Garcia, 2011; Gilman *et al.*, 2012). Only few surveys in West Africa have used imaging techniques (Secka *et al.*, 2010b; Milogo *et al.*, 2012).

Gaps; Neuroimaging facilities are expensive and not easily available for studies in West Africa and hence such diagnostic facilities should be made available in research institutes or government hospitals in the region where it can easily be assessed by researchers.

3.2.3. Diagnosis of taeniasis

193 The presence of adult *T. solium* tapeworms may be detected by stool microscopy, and
194 observation of *Taenia* sp. tapeworm eggs or sometimes adult tapeworm segments (Lightowlers
195 *et al.*, 2016). Coproscopy has been used in human surveys in West Africa (Gweba *et al.*, 2010;
196 Secka *et al.*, 2011).

197 Gap; This test has poor sensitivity (Lightowlers *et al.*, 2016), and misses 60-70% of cases (Mayta
198 *et al.*, 2000). Moreover, it is not possible to differentiate between *T. solium* and *T. saginata* eggs,
199 which are morphologically similar (Garcia *et al.*, 2003; Lightowlers *et al.*, 2016). Therefore
200 results of stool microscopy should be interpreted with caution and more sensitive diagnostic
201 techniques should be made available in the region.

202 Coproantigen detection is more sensitive than stool microcopy as reported by Allan *et al.* (1996)
203 where the detection rate was 5% and 1% for the former and the latter respectively. A species-
204 specific coproantigen ELISA developed by Guezala *et al.* (2009) detects *Taenia*-specific
205 molecules in faecal samples. Although the test has a reported sensitivity of 96.4% and specificity
206 of 100% for *T. solium* carriers, it is important not to base a diagnosis solely on this test.
207 Moreover, the test is no longer commercially available. Studies of Edia-Asuke *et al.* (2014) in
208 West Africa used coproantigen ELISA which involved the use of polyclonal antibodies to
209 identify copro antigens in faeces, hence such kits if made available to researchers in West Africa
210 will give a better picture of the prevalence of taeniais in the region.

211 Other diagnostic methods include the PCR and restriction enzyme analysis (REA) test developed
212 by (Mayta *et al.*, 2000), which has several advantages, it avoids the use of scarce, expensive and
213 radioactive reagents and specialized equipment and uses only two steps to differentiate the
214 tapeworms. A multiplex PCR for differential diagnosis of *T. solium*, *T. saginata* and *T. asiatica*
215 was also developed by Yamasaki *et al.* (2004). More recently, loop-mediated isothermal

amplification (LAMP) for *T. solium* DNA which does not need complicated equipment was developed (Nkouawa *et al.* 2010). This method could differentially diagnose *T. saginata*, *T. solium* and *T. asiatica* in 37 of 43 (86%) parasitologically diagnosed cases of taeniasis (Nkouawa *et al.*, 2010; 2012).

Gaps; These tests, however, are technically demanding and most of them have not been used in studies carried out in West Africa, however its availability and use by researchers in the region will make it possible for elaborate estimate figures of taeniasis to be generated which will invariably reduce under reporting/under estimation of results in the region .

4. Current status of *Taenia solium* infections in West Africa

4.1 Human taeniasis and NCC

Taenia solium cysticercosis is present in most West African countries since favorable conditions for parasite transmission in both humans and pigs occur widely in the region, such as defecation in the open field, illicit slaughtering of pigs and unqualified meat inspectors (Gweba *et al.*, 2010; Weka *et al.*, 2013; Carabin *et al.*, 2015). The prevalence of cysticercosis / NCC and taeniasis in humans is shown below (Tables 1 and 2). Stubbornly persistent high prevalence of neurocysticercosis has been reported in countries such as Burkina Faso and Senegal (Fleury *et al.*, 2013). In Burkina Faso, NCC is common, with 12-17 % of people suffering from epilepsy testing positive for *T. solium* (Millogo *et al.*, 2012; Nitiéma *et al.*, 2012). In Nigeria, a study by Edia-Asuke, *et al.* (2015) showed an association between epilepsy and cysticercosis, such that individuals with epilepsy were twice as likely to test seropositive compared to non-epileptics. Cases of taeniasis have also been widely reported in the region (Table 2). There are important

238 gaps in knowledge of prevalence, for example no recent published data on human cysticercosis
239 in Guinea-Bissau or Liberia.

Country	Prevalence (%)	Diagnosis	Test method	Subjects	References
Burkina Faso	0 - 11.5 (120/3609)	B158/B60 Ag-ELISA	NS	Villagers	Carabin <i>et al.</i> , 2015
	17 (10/60)	B158/B60 Ag-ELISA	NS	PWE	Milogo <i>et al.</i> , 2012
	29.4 (20/68)	CT-Scan	Specific	PWE	Milogo <i>et al.</i> , 2012
	12.8 (5/39)	B158/B60 Ag-ELISA	NS	PWE	Nitiéma <i>et al.</i> , 2012
Nigeria	9.6 (12/125)	ELISA (IgG)	NS	Villagers	Weka <i>et al.</i> , 2013
	14.3 (43/300)	ELISA (IgG)	NS	Butchers	Edia-Asuke <i>et al.</i> , 2015
Senegal	7.7 (31/403)	B158/B60 Ag-ELISA	NS	Villagers	Secka <i>et al.</i> , 2011
	7.7 (31/403)	EITB	Specific	Villagers	Secka <i>et al.</i> , 2011
	23.3 (10/43)	CT-scan	Specific	Seropositive	Secka <i>et al.</i> , 2011
Gambia	0	EITB, CT-Scan	Specific	630 PWE and controls	Secka <i>et al.</i> , 2010b

Table 1. Prevalence of cysticercosis in both people with epilepsy (PWE) and general villagers in West Africa by serology and imaging methods. *Prevalence is given in percent, followed by positive cases / sample size in brackets). Ag-ELISA = Antigen ELISA=Enzyme linked immunosorbent assay; CT = Computed Tomography. NS = method is not specific for T. solium.*

Country	Prevalence % (N)	Target group	Test	References
Ghana	0 (1080)	Schoolchildren	Stool microscopy	Nkrumah and Nguah, 2011
	1.1 (3/292)	Villagers	Stool microscopy (Kato-Katz)	Humphries <i>et al.</i> , 2011
	13 (65/494)	Villagers	Stool microscopy (Kato-Katz)	Bimi <i>et al.</i> , 2012
Nigeria	30 (3/10)	Adult butchers	Copro-Ag-ELISA (<i>T. solium</i>)	Edia-Asuke <i>et al.</i> , 2014
	8 (4/50)	Villagers	Stool microscopy (<i>Taenia</i> spp.)	Gweba <i>et al.</i> , 2010
Gambia	4.7 (2/43)	Villagers	Stool microscopy (<i>Taenia</i> spp.)	Secka <i>et al.</i> , 2011

Table 2. Prevalence of taeniasis in human faecal samples in West Africa. Prevalence is given in percent, followed by positive cases / sample size in brackets.

4.2 Porcine cysticercosis

Reports indicate that the West African region has the largest pig population on the African continent (Ngowi *et al.*, 2013), having increased by 23 % between 1985 and 2005 (FAO, 2012). Available data on the current status of porcine cysticercosis in West Africa, determined by serological or carcass inspection, is presented in Table 3.

Gaps; Although there are no recent data on porcine cysticercosis in several countries in West Africa, e.g. Togo (Dumas *et al.*, 1989, 1990), *T. solium* is highly prevalent in the pig raising parts of the wider region, including in Senegal, Gambia (Secka *et al.*, 2010a, b; Secka *et al.*, 2011) and Nigeria (Gweba *et al.*, 2010). There is a need for more epidemiological studies using suitable diagnostic technologies and neuroimaging to produce a more comprehensive picture.

Country	Prevalence % (sample size)	Diagnostic method	Test method	Reference
Burkina Faso	32.5- 48.2 (336)	B158/B60 Ag-ELISA	NS	Ganaba <i>et al.</i> , 2011
Gambia	4.8-13.2 (1705)	B158/B60 Ag-ELISA	NS	Secka <i>et al.</i> , 2010a
Ghana	2.3 (4121)	Carcass inspection	Specific	Atawalna <i>et al.</i> , 2015
	18.8 (117)	Carcass inspection	Specific	Bimi <i>et al.</i> , 2012
	11.7(60)	Carcass inspection	Specific	Permin <i>et al.</i> , 1999
Nigeria	2.4 (379)	Carcass inspection	Specific	Idiaka <i>et al.</i> , 2017
	5.9 (205)	Tongue inspection	Specific	Gweba <i>et al.</i> , 2010
	14.4 (118)	Carcass inspection	Specific	Gweba <i>et al.</i> , 2010
	20 (2358)	Carcass inspection	Specific	Onah and Cheinjina 1995
	9.3 (43)	Carcass inspection	Specific	Edia-Asuke <i>et al.</i> , 2014
	46 (115)	IgG antibodies	NS	Weka and Ikeh, 2009
Senegal	6.4-13 (1334)	B158/B60 Ag-ELISA	NS	Secka <i>et al.</i> , 2010a

Table 3. Prevalence of porcine cysticercosis in pigs in West Africa. Prevalence is given in percent, followed by sample size in brackets. AgELISA = Antigen Enzyme linked immunosorbent assay; IgG = Immunoglobulin G. NS = not specific for *T. solium*.

4.3 Economic burden of human disease

Neurocysticercosis resulting in epilepsy is a significant health problem in many developing countries (Carabin *et al.*, 2015; Johansen *et al.*, 2016), leading to disproportionate economic suffering in underprivileged populations (Ferrer and Garate 2014; Gabriël *et al.*, 2016; WHO 2016a). Reports from most parts of West Africa show that epileptics suffer discrimination, stigmatization, joblessness and disability (Nitiéma *et al.*, 2012; Komolafe *et al.*, 2012; Osakwe *et al.*, 2014). Negative attitudes and behaviours toward epileptics by the general population are due to ignorance and mistaken perceptions. In Nigeria some people believe that epilepsy is caused by witchcraft or evil spirits or that it is directly contagious, and sufferers therefore often seek treatment from herbalists, faith healers or witch doctors (Osakwe *et al.*, 2014). Hence detailed epidemiological surveys in order to generate data on communities' perception of epilepsy especially in relation to prevalence of epilepsy in NCC patients should be carried out in the region. Such a study will be used to serve as a guide in community treatment and social intervention packages (Osakwe *et al.*, 2014) to help change the negative perceptions that the populace have towards the epileptics.

4.4 Economic impacts of porcine cysticercosis

Porcine cysticercosis significantly impacts pig production in sub-Saharan Africa (Gabriël *et al.*, 2016; Trevisan *et al.*, 2017), resulting in nutritional and financial losses for smallholder farmers (Johansen *et al.*, 2016; Ngowi *et al.*, 2013). Previous studies by Zoli *et al.* (2003) indicate that annual losses resulting from porcine cysticercosis in 10 West and Central African countries is estimated at 25 million Euro, no recent region-wide economic information or reports are however, available for West Africa except for studies of Atawalna *et al.* (2015) in Ghana where

the authors gave an estimate of 104,528.11 Ghana Cedis was lost directly annually due to porcine cysticercosis

5. Risk factors for transmission in West Africa

The human populations considered to be at the highest risk of infection are those who earn their livelihood wholly or partially through livestock rearing, including pigs, and have limited access to good sanitation (Carabin *et al.*, 2015; WHO, 2016a). Here we discuss risk factors especially relevant to transmission of *T. solium* in West Africa.

5.1 Changing societal context

Increasing demand for meat and urbanization of populations in Africa are driving emerging livestock systems, including backyard pig rearing without sanitary precautions. The scenario is further complicated due to lack of potable water and poor sanitary conditions, which were reported as risk factors for taeniasis in a study conducted in Nigeria, in which *Taenia* spp. eggs were detected in soil and water (Gweba *et al.*, 2010). Unhygienic practices like dumping faecal matter in the surroundings, eating unwashed fruit and vegetables, and drinking unsafe (i.e. not boiled or otherwise treated) water contaminated with *Taenia* spp. eggs, are important risk factors for infection with *T. solium* in Burkina Faso (Carabin *et al.*, 2015). Hence provision of portable drinking water should be made to communities which will prevent not only taeniasis but includes other diseases and will also improve sanitation and hygiene in the region.

5.2 Animal husbandry systems

A strong link between various husbandry and sanitary practices and risk of infection in pigs and humans in West Africa has been established. The traditional system of pig management where most of the pigs are confined or tethered at night, and usually allowed to roam freely and scavenge during the day, is a common practice in West Africa and is likely to influence the prevalence of porcine or human cysticercosis, as observed in Burkina Faso (Ganaba *et al.*, 2011). In that study, porcine cysticercosis was associated with rearing practices, especially in the rainy season, in which pigs left to roam briefly during the rainy season were more likely to be seropositive than those kept confined in pens. Seropositivity for human cysticercosis in a village population in Senegal was also linked to free roaming pigs having access to human defecation sites (Secka *et al.*, 2011). In Nigeria, Gweba *et al.* (2010) observed that pigs had access to farmlands that were used as defecation sites by farmers. Weka *et al.* (2013), also in Nigeria, showed that extensively raised pigs had significantly higher seroprevalence of cysticercosis compared to those that were raised under more intensive management systems. The traditional system of pig management is therefore a risk factor for transmission of the disease; hence farmers should be advised to confine their pigs if possible.

5.3 Human dietary behavior

Changing human behavior is a crucial component of solving the problem of cysticercosis in West Africa. For example, there is a lack of coordinated meat inspection and illegal slaughtering is commonly practised, resulting in infected carcasses being marketed and consumed even when under cooked, especially during festivities exposing humans in the area at risk of taeniasis and cysticercosis. In some instances, there are no abattoirs or slaughter slabs in the locality for meat inspection to be carried out, and if present are often unregistered. This was a common scenario

observed in studies conducted in Nigeria, Senegal and Gambia (Gweba *et al.*, 2010; Secka *et al.*, 2010a, Edia-Asuke *et al.*, 2014).

5.4 Butcher and inspector practices in abattoirs

Meat inspection regulations vary from one country to another (Dorny *et al.*, 2005; Goussanou *et al.*, 2014). Usually, butchers carry out basic visual inspection, sometimes supplemented by incisions of parasite preference sites, as for example in Benin (Goussanou *et al.*, 2014). Butchers may allow only one incision to avoid disfigurement of meat as reported in Nigeria (Gweba *et al.*, 2010); a lack of multiple incisions renders the technique of low value, as cysts may be missed, thereby reducing sensitivity and observed prevalence (Goussanou *et al.*, 2014). Some inspectors in Benin also allow butchers to sell meat from carcasses with immature cysts, with the recommendation to cook thoroughly before consumption (Goussanou *et al.*, 2014). Confounding the issue also are the middle men who exert pressure to get the meat away from the slaughter slabs and quickly to the market place, thereby preventing thorough meat inspection as observed in Nigeria (Gweba *et al.*, 2010). Although existing legislation in many African countries requires that cysticercotic pig carcasses be condemned at meat inspection, this does not usually occur. Rather, infected pig carcasses are sold to consumers at reduced prices in Nigeria (Weka, personal observation) as commonly observed in other African countries (Ngowi *et al.*, 2013) which promotes spread of infection and further pushes the burden of disease onto the poor, who cannot pay the premium for safe meat, therefore proper meat inspection should be carried out to curb spread of the disease.

363

364 5.5 *Pork preparation and consumption*

365 There are indications that pork consumption is increasing in sub-Saharan Africa, while
366 preparation methods in the region often do not effectively kill the cysticerci (Assana *et al.*,
367 2013). Heat resulting from boiling and frying the pork passes through the meat to a greater extent
368 and is more likely to kill the cysts, compared to heat from roasting or barbecue grilling, as
369 reported in Nigeria (Edia-Asuke, *et al.*, 2015). Studies conducted in Ghana, Nigeria and Burkina
370 Faso indicated that those who ate pork with cysticerci and those who ate lightly cooked pork
371 soups in the markets had higher probabilities of taeniasis (Bimi *et al.*, 2012; Weka *et al.*, 2013;
372 Carabin *et al.*, 2015). In Benin it is frequently eaten in form of a meat product named “kpete”
373 (kind of pudding) which are not always sufficiently heated to kill all cysticerci (Goussanou
374 (2010). Consumers should be encouraged to cook meat thoroughly before consumption to halt
375 spread of infection.

376

377 5.6 *Hygiene practices*

378 Open defecation, poor personal hygiene and low environmental sanitation including improper
379 faecal disposal are common practices in West Africa and are habits that promote the spread of *T.*
380 *solium* (Gweba *et al.*, 2010; Secka *et al.*, 2011; Weka *et al.*, 2013). It has been reported that
381 persons who do not use toilet facilities or did not wash their hands after defecation were 8.3 and
382 5.5 times respectively to test seropositive compared to those who used water closet toilet
383 facilities and regularly washed their hands after defecation (Weka *et al.*, 2013). A similar report
384 had been made in Burkina Faso (Ganaba *et al.*, 2011; Carabin *et al.*, 2015).

385 5.7 *Lack of knowledge*

In endemic areas, poverty, ignorance and superstitious beliefs about *T. solium* are associated with increased risk of disease transmission and stakeholders in endemic areas may know about tapeworm infections in humans but could not relate them to porcine cysticercosis and NCC (Thys *et al.*, 2015). In a study in Nigeria, 80 % of the butchers had poor knowledge of *T. solium* cysticercosis and its public health significance (Edia-Asuke *et al.*, 2014), a likely explanation for the persistently high-risk practices as described above.

6. Treatment of *Taenia solium* infections

6.1 Treatment of neurocysticercosis/Taeniasis

There is no universally accepted gold standard for the treatment of NCC but several suggestions have been made (Winkler, 2013; Garcia *et al.*, 2014b). Pharmacological therapy succeeds in eliminating active cysts in only one third of patients (Carpio and Romo, 2014).

Albendazole is one of the cysticidal drugs of choice because of its ability to penetrate the central nervous system (Garcia *et al.*, 2014a). Another drug of choice is praziquantel, which is widely accessible in sub-Saharan Africa and is effective against cestodes and schistosomes (Evans *et al.*, 2013; Braae *et al.*, 2015c). Mass praziquantel treatment has been used widely in schistosomiasis control in Africa. There are no data available yet on widespread treatment for *T. solium* infections in humans in West Africa; however, several countries in the region have carried out MDA against schistosomiasis using praziquantel (Garba *et al.*, 2009; Evans *et al.*, 2011; Leslie *et al.*, 2012). The drug is very effective in the treatment of taeniasis at a single oral dose rate of 5–10 mg/kg and therefore can be used to treat against both parasites (Braae *et al.*, 2015a). Unfortunately, the recommended dose for the treatment of schistosomiasis (40mg//Kg); (WHO 2002) stimulates seizures in people with NCC (Johansen *et al.*, 2016). It has been observed that

integration with schistosomiasis control programmes is a good way forward for control of *T. solium*, and will also control other co-endemic infections (Braae *et al.*, 2015a).

The safety of praziquantel for mass drug administration (MDA) in West Africa is yet to be evaluated in the light of high prevalence of NCC. Niclosamide is also another drug of choice for eliminating adult tapeworm burdens (Gilman *et al.*, 2012), but must be used with caution especially in people with NCC. Adult worms eliminated by the patients constitute a serious hazard to the populace and the environment and must be safely disposed off. It may also be necessary to treat any newcomers to the region that might be carriers, and those returning from regions where transmission of *T. solium* is still ongoing (Geerts, 2016).

6.2 Treatment of cysticercosis in pigs

The benzimidazole drug, oxfendazole is very effective against muscle cysts and gastrointestinal helminths at a single dose of 30 mg/kg body weight, but has limited effect on brain cysts (Pondja *et al.*, 2012; Mkupasi *et al.*, 2013a, b). Additional benefit of the use of oxfendazole in pigs is that it increases the growth rate and general condition by eliminating other gastro intestinal nematodes.

The use of oxfendazole alone is often unable to control *T. solium*, as shown in a high endemic area in Mozambique (Pondja *et al.*, 2012). This is because some animals are likely to get reinfected following the last treatment and prior to slaughter. The role of drug treatment in the development of immunity against reinfection in endemic areas is uncertain. Further, in heavily infected pigs, inflammatory reactions arise in reaction to the anthelmintic-mediated death of cysticerci in the muscles (Assana *et al.*, 2010), and can make the meat unfit for marketing and subsequent consumption (Lightowers, 2010). The prolonged treatment time and withdrawal

period following the use of oxfendazole before the animals are sold for consumption constitutes a major constraint to the effective control of *T. solium* in most rural communities where the pigs are the main source of income for the farmers. In Morocco north Africa, oxfendazole is marketed under the trademark Paranthic[®] at a price of 0.5 US dollars per 30 kg pig, which is motivating to small-scale pig farmers (Lightowlers, 2013; Donadeu and Lightowlers, 2014; Geerts, 2016), however, data on the availability, acceptability and affordability of the drug in West Africa is lacking.

7. Control

Taenia solium cysticercosis is still endemic in West Africa despite the availability of tools to disrupt the life cycle (Braae *et al.*, 2015b). The World Health Organization (WHO) included *T. solium* cysticercosis as one of the major neglected tropical diseases (NTDs), and has adopted a resolution for intensified control efforts (WHO, 2010; Braae *et al.*, 2016b). The resolution recommended a scaled-up control and eradication strategy in selected countries to last from 2016 to 2020 with the aim to control and eradicate the disease by 2020. However; no large scale taeniasis control programme, has yet been implemented in sub-Saharan Africa (WHO, 2015c). Below, we consider the approaches available and how they have been and might be more effectively applied in West Africa.

7.1 Use of latrines to reduce open defecation

The availability and use of toilets or latrines will decrease the spread of *Taenia* spp. eggs and other soil-transmitted helminths (STH) in the environment (Bethony *et al.*, 2006; Pruss-Ustun *et*

al., 2014). Moreso, emphasis should not be on the provision and use of latrines or access to sufficient clean water only but also on the proper management of such infrastructures by the population (Gabriël *et al.*, 2016). The assessments of community-led total sanitation (CLTS) as an intervention measure for the control of porcine cysticercosis has not yet been fully evaluated as part of *T. solium* control in West Africa, hence the communities should be encouraged to stop open defecation and practice CLTS. However, studies in The Gambia and Senegal reported that 93% of the residents had toilets and only 5% used open defecation and that the 5% were likely to be responsible for considerable environmental contamination with *Taenia* spp. eggs (Secka *et al.*, 2010a). A study from Nigeria showed that although 93% of households in the study location had toilets, majority of them still defecate on farmlands thereby contaminating the environment with *Taenia* spp. eggs (Gweba *et al.*, 2010).

7.2 Personal and household hygiene

Individual and community hygiene should also be emphasized and encouraged in order to decrease the risk of NCC, including the availability of sufficient clean water which is crucial to allow improved hygiene (Gabriël *et al.*, 2016). A study in Nigeria showed higher risk (31.7 times) of exposure to taeniasis among respondents who do not wash their hands after each toilet use compared to those who wash their hand with soap and water (Weka *et al.*, 2013). Enough water for washing of fresh foods- fruits and vegetables is also necessary to reduce the spread of *Taenia* spp. eggs as reported from studies carried out in Bukina faso (Carabin *et al.*, 2015) .

7.3 Confinement of pigs

Traditional systems of pig management is commonly observed in West Africa and is likely to influence the prevalence of porcine or human cysticercosis (Ganaba *et al.*, 2011; Secka *et al.*, 2011; Weka *et al.*, 2013). The ability of farmers to provide pig housing is hindered by poverty, since the farmer is also expected to provide feed to the pigs under the confined systems of management (Assana *et al.*, 2013). This runs in opposition to the motives of the smallholder farmer, to keep pigs as a source of income without the need to invest in feed (Gilman *et al.*, 2012) which renders pig confinement an unworkable approach to *T. solium* control in the short term and changes in pig management systems should be considered a long term strategy (Thomas, 2015). However pig farmers should also be informed that apart from the advantage of a decrease in porcine cysticercosis, confined pigs also have a reduced risk of acquiring African Swine Fever (Geerts 2016), and decreased burden of other parasites (Gabriël *et al.*, 2016).
Gap; Changes in pig rearing system are likely to be driven by economic more than sanitary considerations, but research on the constraints and consequences of adopting different systems, and related trade-offs with time and funds available for other activities, resilience to internal and external disruptions, and animal welfare, remain un-studied in West Africa.

7.4 Meat and lingual inspection

Both meat and tongue inspection are carried out in West Africa in locations where slaughter slabs and abattoirs for pigs exist (Onah and Chiejina, 1995; Coulibaly and Yameogo, 2000; Gweba *et al.*, 2010; Secka *et al.*, 2010b; Edia Asuke *et al.*, 2014; Goussanou *et al.*, 2014; Idiaka *et al.*, 2017). These techniques are relatively cheap, rapid and easy to conduct (Goussanou *et al.*, 2013), their sensitivity, however, is low therefore, lightly infected pig carcasses undetected by

this method are likely to go into the food chain and serve as a source of infection in the community as reported in Benin (Goussanou *et al.*, 2014). Furthermore, farmers and traders in Benin frequently carry out tongue examinations to detect cysts prior to marketing in order to avoid condemnation, and then often use the infected carcase for their own consumption or sell it illegally (Goussanou *et al.*, 2013).

Gap; The lack of comprehensive and satisfactory meat inspection in West Africa is a significant risk factor for the transmission of cysticercosis to humans. In Nigeria, the situation is further complicated by the lackadaisical attitude of inspectors, apathy from the farmers, insincerity of policy makers, and consumers who prefer to buy infected carcasses at a cheap price without regard for the health implications (Weka, personal observation). Strict meat inspection and condemnation of infected carcasses runs the risk of exacerbating the divide between safe but expensive meat and cheaper but riskier ‘unofficial’ meat, further pushing the burden of disease onto the poor. Research is needed that encompasses the economic, social and behavioural context of meat safety as well as biological and veterinary processes.

7.5 Meat processing

Proper meat processing at household and community level is an important defence against NCC (Ertel *et al.*, 2015). Freezing for 10 days at -10°C , gamma-radiation, cooking and salt pickling of pork meat infected with *T. solium* all decrease cyst viability (OIE, 2015; Geerts 2016).

Gap; Implementation in rural areas is often difficult due to cost and unavailability of equipment (Geerts, 2016). Lack of freezers, unreliable power supply, and impatience of consumers militate against prolonged freezing of meat as a control measure in West Africa. Although meat is

normally well cooked in endemic areas (Geerts, 2016), as temperature of 80 °C will also kill *T. solium* cysticerci in infected meat (OIE, 2015), studies in including West Africa indicated that the pork meat it is frequently eaten in forms which are not always sufficiently heated to kill all cysticerci (Goussanou 2010; Edia-Asuke, *et al.*, 2015).

7.6 Anthelmintic treatment of pigs and humans

Treatment of cysticercosis in pigs and taeniasis in people can be usefully applied as MDA at population level using a single dose of 5 mg/kg praziquantel or nitazoxanide (2g) as an effective control measure. A 56% decrease of prevalence of taeniasis after 42 months was reported in Mexico while 1-3.5 % decrease in cases of taeniasis and a 7-55 % decrease in porcine cysticercosis in two villages in Guatemala 10 months after evaluation was reported (Allan *et al.*, 1997). Gap; Studies using combined treatment of pigs and humans in West Africa have not yet been reported.

7.7 Vaccination of pigs

Vaccination is a recently available intervention for the control and elimination of *T. solium* (Lightowlers, 2013; Gabriël *et al.*, 2016), and has conferred high levels of protection in both experimental and field trials (WHO, 2015a, 2016a). Three candidate vaccines currently exist, based on: a crude antigen developed by Molinari *et al.* (1993); a recombinant oncosphere antigen-based TSOL16 or TSOL18 vaccine by Lightowlers *et al.* (2010); and a peptide-based S3PVac developed by Huerta *et al.* (2001). The most effective vaccine produced to date is the recombinant protein TSOL18, which achieved 99.5–100% protection of vaccinated pigs in a field trial in Cameroon when combined with a single oxfendazole treatment (Assana *et al.*,

2010). In the study, the first vaccine dose was given at 2-3 months of age, and a second vaccine dose given with oxfendazole 4 weeks later. A third vaccine dose was given 3 months later. The combination of TSOL18 vaccination and oxfendazole in pigs has the potential to control transmission in endemic areas and indirectly decrease new cases of NCC (Assana *et al.*, 2010). The oxfendazole eliminates any *T. solium* infection that is already present in the vaccinated pigs, prior to the animals being fully vaccinated and protected (Lightowlers, 2013). The regimen allows adequate time to pass after chemotherapy for any lesions in the pork arising from killed and necrotic cysticerci to be resolved prior to the animals being slaughtered. During this period, all animals (previously infected or otherwise) are protected against new *T. solium* infection by the vaccine. A similar result was obtained in a field trial conducted in Peru (Jayashi *et al.*, 2012). The effectiveness of *T. solium* taeniasis-cysticercosis elimination by combined vaccination (TSOL18) plus treatment of pigs with oxfendazole, combined with mass or targeted treatment of human tapeworm carriers (Geerts, 2016) led to marked reduction in *T. solium* transmission in the study populations (Garcia *et al.*, 2010; Gilman *et al.*, 2012). Challenges remain, however, in achieving optimal protection of traditionally managed pigs under field conditions (Lightowlers, 2013; Pawloski, 2016). The protocol requiring three vaccinations and oxfendazole increases costs to the farmer and likely decreases feasibility (Jayashi *et al.*, 2012; Thomas, 2014). Therefore, the vaccine may be unaffordable for poor communities unless subsidized and supported by provision of infrastructure to deliver the vaccine, including an effective cold chain. Currently, the TSOL18 vaccine is produced commercially at scale by Indian Immunologicals (Thomas, 2015; Geerts, 2016). Although the vaccine has been registered for use since May 2016 and can be purchased by any country it is not registered in any West African country probably due to the cost implications associated with the registration process. Therefore, there is no data

on the vaccine trial in West Africa, even though the authors' personal interview with farmers in North central Nigeria indicates their willingness to purchase the vaccine.

7.8 Health education

Health education is a significant part of control strategies for *T. solium* (Garcia *et al.*, 2010; Fleury *et al.*, 2013) and could play a significant role by informing consumers about the risks related to the consumption of infected pork; subsequent refusal to purchase infected meat might assist in a change of management practices, as farmers will acknowledge a clearer economic cost of high-risk practices (Thomas, 2015; Gabriël *et al.*, 2016). Impact could extend throughout the meat supply chain and to health workers and the general population (Thomas, 2015), although like other interventions, education has limitations as a stand-alone approach (Gabriël *et al.*, 2016). A free computer-based health education tool called "The Vicious Worm" was developed to support efforts to control *T. solium* infections (Johansen *et al.*, 2014), targeting stakeholders across from all professions and sectors and providing information on transmission, diagnosis, risk factors, and prevention and control of the disease (Johansen *et al.*, 2016).

Gap; Recently effectiveness of community-based (EFECAB) educational programme carried out in Burkina Faso showed a reduction in the cumulative incidence of active cysticercosis (adjusted cumulative incidence ratio 0·65, 95% Bayesian credible interval [95% CrI] 0·39–1·05) and a reduction in active cysticercosis prevalence adjusted prevalence proportion ratio 0·84; 95% CrI 0·59–1·18) both from baseline to after randomisation (Carabin *et al.*, 2018). Such studies can be repeated in other countries of West Africa so that health education can be used as a control tool for *T. solium* cysticercosis in West Africa.

7.9 Integrated approach

Several predictive models have been developed to help design the most effective and feasible intervention strategies for the control of *T. solium* cysticercosis in both pigs and human populations (Braae *et al.*, 2016b; Winskill *et al.*, 2017). The recently developed CystiSim model indicated that combined intervention strategies in both pigs and humans, such as concurrent mass drug administration in humans and vaccination and treatment of pigs, have a high likelihood of success, given a 75% coverage rate sustained for more than four years (Braae *et al.*, 2016b).

Collaborative networks for *T. solium* control already exist nationally and internationally among researchers, and this effort should be sustained for interaction and transfer of knowledge and findings (Gabriël *et al.*, 2016, WHO, 2016a). A multisectorial and One Health approach across all disciplines including agricultural, food and human health sectors comprising medical, veterinary, environmental, governmental and nongovernmental and social actors is necessary for the control and eradication of cysticercosis and *T. solium* in both human and pigs (Braae *et al.*, 2016a; Gabriël *et al.*, 2016). Furthermore, it is vital to integrate control of *T. solium* cysticercosis with other NTDs or within national primary health care systems (Bockarie *et al.*, 2013; Nakagawa *et al.*, 2015) and should be encouraged in West Africa. Efforts to improve scientific understanding and design of optimal control strategies should be cognisant of the fact that the success of any intervention is largely dependent on the level of societal and political acceptance, commitment and engagement of the stakeholders (Gabriël *et al.*, 2016).

8. Conclusions

Diverse intervention methods have been recommended and tried in order to control *T. solium* cysticercosis (Fleury *et al.*, 2013). A multipronged approach consisting of vaccination of pigs

using TSOL18 vaccine and treatment with oxfendazole along with mass chemotherapy of humans is promising and with praziquantel could eliminate taeniasis. However, the fact remains that long-term and sustainable control, and potentially elimination, of *T. solium* should preferably include combinations of methods that center on both human and animal hosts. Selected combinations will depend on practicality and economic cost (Mwape *et al.*, 2015). Alas, at present there is inadequate evidence for the best combinations for cost-effective intervention packages in endemic countries (Johansen *et al.*, 2016), thus integration of economics into disease control models might help to rectify this important deficit. Identifying and addressing key obstacles to sustained control efforts, as set out in this review for West Africa, is an essential step in designing rational approaches suited to target regions, and making real progress towards eradication. Diagnostic tools, including neuroimaging facilities, should be strategically situated and made accessible to rural populations in West Africa. This will support efforts to more accurately estimate the burden of disease, highlight the problem to policymakers with competing priorities, and monitor the effectiveness of interventions.

Acknowledgement

Competing interests

The authors declare no conflict of interests of any kind.

Declaration of interest- None

Funding:

References

- Allan J. C., Velasquez-Tohom M., Torres-Alvarez R., Yurrita P., Garcia-Noval, J. (1996). Field trial of the coproantigen-based diagnosis of *Taenia solium* taeniasis by Enzyme-Linked Immunosorbent Assay. The Am. J. Trop. Med. Hyg. 54, 352–356.
- Assana, E., Craig T. K., Charles G. G., Geerts, S., Dorny, P., 'De Deken, R., Garry, A. A., Zoli P. A., 'Lightowlers M.W. (2010). "Elimination of *Taenia solium* transmission to pigs in a field trial of the TSOL18 vaccine in Cameroon. Internat. J. Parasitol. 40, 515–519.
- Assana, E., Lightowlers, M. W., Zoli, A. P., Geerts, S. (2013). *Taenia solium* taeniosis/cysticercosis in Africa: risk factors, epidemiology and prospects for control using vaccination. Vet. Parasitol. 195, 14-23.
- Atawalna, J., Ewura, S., Mensah. M. (2015). Prevalence and financial losses associated with porcine cysticercosis in the Kumasi metropolis of Ghana. Internat. J. Livestock Res. 5, 21–26.
- Bethony, J., Brooker, S., Albonico, M., Geiger, S. M., Loukas, A., Diemert, D., Hotez, P. J. (2006). Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. The Lancet 367, 1521–1532
- Bimi, L., Laar, A. K., Anto, F. (2012). Prevalence and Risk Factors of Taeniasis in the Bunkpurugu-Yunyoo District of Northern Ghana. J. Bacteriol. Parasitol. 3, 129.
- Bockarie, M. J., Kelly-Hope, L. A., Rebollo, M., Molyneux, D. H. (2013). Preventive chemotherapy as a strategy for elimination of neglected tropical parasitic diseases: endgame challenges. Philosop. Transac. Roy. Soc B: Biol. Sci. 368, 20120144.

- Braae, U. C., Magnussen, P., Lekule, F., Harrison, W., Johansen, M. V. (2014). Temporal fluctuations in the sero-prevalence of *Taenia solium* cysticercosis in pigs in Mbeya Region, Tanzania. *Parasit. Vectors.* 7, 574.
- Braae, U. C., Saarnak, C. F., Mukaratirwa, S., Devleesschauwer, B., Magnussen, P., Johansen, M. V. (2015a). *Taenia solium* taeniosis/cysticercosis and the co-distribution with schistosomiasis in Africa. *Parasit. Vectors.* 8, 323
- Braae, U. C., Harrison, W., Lekule, F., Magnussen, P., Johansen, M. V. (2015b). Feedstuff and poor latrines may put pigs at risk of cysticercosis—A case-control study. *Vet. Parasitol.* 214, 187–191.
- Braae, U. C., Magnussen, P., Ndawi, B., Harrison, W., Lekule, F., Johansen, M. V. (2015c). Effect of repeated mass drug administration with praziquantel and track and treat of taeniosis cases on the prevalence of taeniosis in *Taenia solium* endemic rural communities of Tanzania. *Acta Trop.* 165, 246–251
- Braae, U. C., Magnussen, P., Harrison, W., Ndawi, B., Lekule, F., Johansen, M. V. (2016a). Effect of national Schistosomiasis control programme on *Taenia solium* taeniosis and porcine cysticercosis in rural communities of Tanzania. *Parasit. Epid. Control.* 1, 245–251
- Braae, U.C., Devleesschauwer, B., Gabriël, S., Dorny, P., Speybroeck, N., Magnussen, P., Torgerson, P., Johansen, M.V. (2016b). CystiSim—an agent-based model for *Taenia solium* transmission and control. *PLoS NTDs.* 10, e0005184.
- Carabin, H., Ndimubanzi, P.C., Budke, C.M., Nguyen, H., Qian, Y., Cowan, L.D., Stoner, J.A., Rainwater, E., Dickey, M. (2011). Clinical manifestations associated with neurocysticercosis: a systematic review. *PLoS NTDs* 5, e1152.
- Carabin, H., Millogo, A., Cissé, A., Gabriël, S., Sahlu, I., Dorny, P., Bauer, C., Tarnagda, Z., Cowan, L.D., Ganaba, R. (2015). Prevalence of and factors associated with human cysticercosis in 60 villages in three provinces of Burkina Faso. *PLoS NTDs* 9, e0004248.
- Carabin, H., Millogo, A., Ngowi, H.A., Bauer, C., Dermauw, V., Koné, A.C., Sahlu, I., Salvator, A.L., Preux, P.M., Somé, T. and Tarnagda, Z., 2018. Effectiveness of a community-based educational programme in reducing the cumulative incidence and prevalence of human *Taenia solium* cysticercosis in Burkina Faso in 2011–14 (EFECAB): a cluster-randomised controlled trial. *The Lancet Global Health*, 6(4), pp.e411-e425.
- Carpio, A., Romo, M. L. (2014). The relationship between neurocysticercosis and epilepsy: an endless debate. *Arquivos de Neuro-Psiquiatria.* 72, 383–390.
- Carpio, A., Fleury, A., Hauser, W. A. (2013). Neurocysticercosis five new things. *Neurol. Clin. Pract.* 3, 118–125.
- Coulibaly, N.D. and K.R. Yameogo, 2000. Prevalence and control of zoonotic diseases: Collaboration between public health workers and veterinarians in Burkina Faso. *Acta Trop.*, 76: 53-57
- Del Brutto, O.H. (2013b). Human cysticercosis (*Taenia solium*). *Trop. Parasitol.* 3, 100–103.
- Devleesschauwer, B., Aryal, A., Tharmalingam, J., Joshi, D.D., Rijal, S., Speybroeck, N., Gabriël, S., Victor, B., Dorny, P. (2013). Complexities in using sentinel pigs to study

- Taenia solium* transmission dynamics under field conditions. Vet. Parasitol. 193, 172–178.
- Donadeu, M., Lightowers, M. W. (2014). RE: Commercialisation of oxfendazole and TSOL18. Type in Thomas, L.F., 2015. Landscape analysis: control of *Taenia solium*. http://apps.who.int/iris/bitstream/10665/164359/1/9789241508643_eng.pdf (accessed on 3 June 2017).
- Dorny, P., Brandt, J., Geerts, S. (2005). Detection and diagnosis. In: Murrell, K.D. (Ed.), WHO/FAO/OIE Guidelines for the Surveillance, Prevention and Control of Taeniosis/Cysticercosis. OIE, Paris, France, pp: 45–55.
- Dumas, M., Grunitzky, E., Deniau, M., Dabis, F., Bouteille, B., Belo, M., Pestre-Alexandre, M., Catanzano, G., Darde, M.L., d'Almeida, M. (1989). Epidemiological study of neurocysticercosis in northern Togo (West Africa). Acta Leidensia. 57, 191–196.
- Dumas, M., Grunitzky, K., Belo, M., Dabis, F., Deniau, M., Bouteille, B., Kassankogno, Y., Catanzano, G., Pestre-Alexandre, M. (1990). Cysticercosis et neurocysticercose: enquête épidémiologique dans le nord du Togo. Bull. Soc. Path. Exot. 83, 263/274.
- Edia-Asuke, A. U., Inabo, H. I., Umoh, V. J., Whong, C. M., Asuke, S., Edeh, R. E. (2014). Assessment of sanitary conditions of unregistered pig slaughter slabs and post mortem examination of pigs for *Taenia solium* metacestodes in Kaduna metropolis, Nigeria. Infect. Dis. Poverty. 3, 45.
- Edia-Asuke, A. U., Inabo, H. I., Mukaratirwa, S., Umoh, V. J., Whong, C. M., Asuke, S., Ella, E. (2015). Seroprevalence of human cysticercosis and its associated risk factors among humans in areas of Kaduna metropolis, Nigeria. The J. Infect. Develop. Count. 9, 799–805.
- Ertel, R.L., Braae, U.C., Ngowi, H.A., Johansen, M.V. (2015). Assessment of a computer-based *Taenia solium* health education tool ‘The vicious worm’ on knowledge uptake among professionals and their attitudes towards the program. Acta Trop. 165, 240–245.
- Evans, D., McFarland, D., Adamani, W., Eigege, A., Miri, E., Schulz, J., Pede, E., Ubugadu, C., Ogbu-Pearse, P. and Richards, F.O. (2011). Cost-effectiveness of triple drug administration (TDA) with praziquantel, ivermectin and albendazole for the prevention of neglected tropical diseases in Nigeria. Ann. Trop. Med. Parasitol. 105, 537–547.
- Evans, D.S., King, J.D., Eigege, A., Umaru, J., Adamani, W., Alphonsus, K., Sambo, Y., Miri, E.S., Goshit, D., Ogah, G., Richards, F.O. (2013). Assessing the WHO 50% prevalence threshold in school-aged children as indication for treatment of urogenital schistosomiasis in adults in central Nigeria. The Am. J. Trop. Med. Hyg. 88, 441–445.
- FAO (2012) Secteur porcin Burkina Faso. Revues Nationales del’ élevage dela division dela production et de la santé animals dela FAO. Rome Microbiological Risk Assessment Series 23, FAO/WHO. Accessed November 2015.
- Ferrer, E., Gárate, T. (2014). Taeniosis and Cysticercosis. In: Helminth Infections and their Impact on Global Public Health (201–227). Springer Vienna.
- Fleury, A., Carrillo-Mezo, R., Flisser, A., Sciutto, E., Corona, T. (2011). Subarachnoid basal neurocysticercosis: a focus on the most severe form of the disease. Expert Rev. Anti. Infect. Ther. 9, 123–133.
- Fleury, A., Sciutto, E., De Aluja, A.S., Larralde, C., Agudelo, S., Garcia, G.M., Fandiño, J., Guerra, R., Nunes, C., de Aragão, S.C., Sato, M. (2013). Control of *Taenia solium* transmission of taeniosis and cysticercosis in endemic countries: the roles of continental

- networks of specialists and of local health authorities. In: Novel Aspects on Cysticercosis and Neurocysticercosis. InTech.
- Fleury, A., Sciutto, E., de Aluja, A.S., Carpio, A. (2015). Cysticercosis: A preventable, but embarrassing neglected disease still prevalent in non-developed countries. In: Zoonoses-Infections affecting humans and animals. Springer Netherlands 335–354.
- Flisser, A. (2013). Epidemiology of Neurocysticercosis in Mexico: From a public health problem to its control. Chapter 9. http://dx.doi.org/10.5772/53839_255-274. InTech.
- Gabriël, S., Dorny, P., Mwape, K.E., Trevisan, C., Braae, U.C., Magnussen, P., Thys, S., Bulaya, C., Phiri, I.K., Sikasunge, C.S., Makungu, C. (2016). Control of *Taenia solium* taeniasis/cysticercosis: The best way forward for sub-Saharan Africa?. Acta Trop. 165, 252–260
- Galan-puchades, M. T. (2016) Taeniasis vs cysticercosis infection routes Asian pacific Journal Tropical Medicine 9(6)619-620
- Ganaba, R., Praet, N., Carabin, H., Millogo, A., Tarnagda, Z., Dorny, P., Hounton, S., Sow, A., Nitiéma, P., Cowan, L.D. (2011). Factors associated with the prevalence of circulating antigens to porcine cysticercosis in three villages of Burkina Faso. PLoS NTDs. 5, e927.
- Garba, A., Toure ' S., Dembele ' R. , Boiesier P., Tohon, Z., Bosque ' -Oliva, E., Koukounari A., Fenwick A. P. (2009), Present and future schistosomiasis control activities with support from the Schistosomiasis Control Initiative in West Africa. Parasitol.136, 1731–1737.
- García, H. H., Gonzalez, A. E., Evans, C. A., Gilman, R. H. Cysticercosis Working Group in Peru. (2003). *Taenia solium* cysticercosis. The Lancet. 362, 547–556.
- Garcia, H.H., Gonzalez, A.E., Rodriguez, S., Gonzalvez, G., Llanos-Zavalaga, F., Tsang, V.C. and Gilman, R.H. (2010). Epidemiología y control de la cisticercosis en el Perú [Epidemiology and control of cysticercosis in Peru]. Revista Peruana de Medicina Experimental y Salud Publica 27, 592–597.
- Garcia, H. H., Gonzales, I., Lescano, A. G., Bustos, J. A., Pretell, E. J., Saavedra, H., Nash, T. E. (2014a). Enhanced steroid dosing reduces seizures during antiparasitic treatment for cysticercosis and early after. Epilepsia. 55, 1452–1459.
- Garcia, H. H., Nash, T. E., Del Brutto, O. H. (2014b). Clinical symptoms, diagnosis, and treatment of neurocysticercosis. The Lancet Neurol. 13, 1202–1215.
- Garcia H H, Castillo Y, Gonzales I, Bustos JA, Saavedra H, Jacob L, Del Brutto OH, Wilkins PP, Gonzalez AE, Gilman RH (2018). Cysticercosis Working Group in Peru. Low sensitivity and frequent cross-reactions in commercially available antibody detection ELISA assays for *Taenia solium* cysticercosis. Trop Med Int Health. 23(1):101-105.
- Geerts, S. (2016) Elimination of *Taenia solium* cysticercosis through vaccination of pigs: a realistic option?. [www.kaowarsom.be/documents/PDF Bulletin/Geerts.pdf](http://www.kaowarsom.be/documents/PDF%20Bulletin/Geerts.pdf) (accessed on 2 August 2017).
- Gilman, R. H., Gonzalez, A. E., Llanos-Zavalaga, F., Tsang, C. W., Garcia, H. H. (2012). The Cysticercosis Working Group in Peru. Special issue article prevention and control of *Taenia solium* taeniasis/cysticercosis in Peru. Path. Glob. Hlth. 106, 5.
- Gonzalez, A.E., Cama, V., Gilman, R.H., Tsang, V.C., Pilcher, J.B., Chavera, A., Castro, M., Montenegro, T., Verastegui, M., Miranda, E., Bazalar, H. (1990). Prevalence and comparison of serologic assays, necropsy, and tongue examination for the diagnosis of porcine cysticercosis in Peru. The Am. J. Trop. Med.Hyg. 43, 194–199.

- Goussanou, J.S.E., 2010. Evaluation des procédés d'abattage et de la qualité microbiologique des carcasses de porcs locaux des Abattoirs de Cotonou-Porto-Novo et des tueries de Cotonou et de Godomey. Mémoire de fin de formation de master en normes, Contrôle de Qualité et Technologie Alimentaire. UAC. Bénin
- Goussanou, J.S., Kpodekon, T.M., Saegerman, C., Azagoun, E., Youssao, A.K., Farougou, S., Praet, N., Gabriël, S., Dorny, P., Korsak, N. (2013). Spatial distribution and risks factors of porcine cysticercosis in southern Benin based meat inspection records. *Internat. Res. J. Microbiol.* 4, 188–196.
- Goussanou, J.S.E., Korsak, N., Saegerman, C., Youssao, A.K.I., Azagoun, E., Farougou, S., Gabriël, S., Dorny, P., Kpodekon, M.T. (2014). Assessment of routine inspection method for diagnostic of porcine cysticercosis in South East Benin by using meat inspection records and Ag-ELISA test. *Internat. J. Anim. Vet. Adv.* 6, 80–86.
- Guezala, M.C., Rodriguez, S., Zamora, H., Garcia, H.H., Gonzalez, A.E., Tembo, A., Allan, J.C., Craig, P.S. (2009). Development of a species-specific coproantigen ELISA for human *Taenia solium* taeniasis. *The Am.J. Trop. Med. Hyg.* 81, 433–437.
- Gweba, M., Faleke, O.O., Junaidu, A.U., Fabyi, J.P., Fajinmi, A.O. (2010). Some risk factors for *Taenia solium* cysticercosis in semi intensively raised pigs in Zuru, Nigeria. *Vet. Italiana*, 46, 57–67.
- <http://www.oie.int/en/international-standard-setting/terrestrial-code/access-online/>
- Huerta, M., De Aluja, A.S., Fragoso, G., Toledo, A., Villalobos, N., Hernandez, M., Gevorkian, G., Acero, G., Diaz, A., Alvarez, I., Avila, R. (2001). Synthetic peptide vaccine against *Taenia solium* pig cysticercosis: successful vaccination in a controlled field trial in rural Mexico. *Vacc.* 20, 262–266.
- Humphries, D., Mosites, E., Otchere, J., Twum, W.A., Woo, L., Jones-Sanpei, H., Harrison, L.M., Bungiro, R.D., Benham-Pyle, B., Bimi, L., Edoh, D. (2011). Epidemiology of hookworm infection in Kintampo North Municipality, Ghana: patterns of malaria coinfection, anemia, and albendazole treatment failure. *The Am. J. Trop. Med. Hyg.* 84, 792–800.
- Idika, I. K. , Njoga, U. J., Ezech, I. O., Iheagwam, C. N., Ezenduka, E.V., Njoga, E., Onah, D. N. (2017) Re-evaluation of porcine cysticercosis in Nsukka area of Enugu State, Nigeria. *Asian Pac. J. Trop. Dis.* 7, 519–522.
- Jayashi, C. M., Arroyo, G., Lightowlers, M. W., Garcí'a, H. H., Rodri'guez, S., Gonzalez, A. E. (2012). Seroprevalence and risk factors for *Taenia solium* cysticercosis in rural pigs of northern Peru. *PLoS NTDs.* 6, e1733.
- Johansen, M. V., Trevisan, C., Braae, U. C., Magnussen, P., Ertel, R. L., Mejer, H., Saarnak, C. F. (2014). The Vicious Worm: a computer-based *Taenia solium* education tool. *Trends Parasitol.* 30, 372–374.
- Johansen, M. V., Trevisan, C., Gabriël, S., Magnussen, P., Braae, U. C. (2016). Are we ready for *Taenia solium* cysticercosis elimination in sub-Saharan Africa? *Parasitol.* 144, 59–64.
- Komolafe, M.A., Sunmonu, T.A., Afolabi, O.T., Komolafe, E.O., Fabusiwa, F.O., Groce, N., Kett, M., Disu, J.O., Ajiboye, J.K., Olaniyan, S.O. (2012). The social and economic impacts of epilepsy on women in Nigeria. *Epilep. Behav.* 24, 97–101.
- Leslie, J., Garba, A., Oliva, E.B., Barkire, A., Tinni, A.A., Djibo, A., Mounkaila, I., Fenwick, A. (2012). Correction: Schistosomiasis and soil-transmitted helminth control in Niger: Cost effectiveness of school based and community distributed mass drug administration. *PLoS NTDs.* 6, 10–1371.

856 Lightowlers, M.W. (2010). Eradication of *Taenia solium* cysticercosis: a role for vaccination of
857 pigs. *Internat. J. Parasitol.* 40, 1183–92.

858 Lightowlers, M. W. (2013). Control of *Taenia solium* taeniasis/cysticercosis: past practices and
859 new possibilities. *Parasitol.* 140, 1566–1577.

860 Lightowlers, M. W., Assana, E., Jayashi, C. M., Gauci, C. G., Donadeu, M. (2015). Sensitivity of
861 partial carcass dissection for assessment of porcine cysticercosis at necropsy. *Internat. J.*
862 *Parasitol.* 45, 815–818.

863 Lightowlers, M. W., Garcia, H. H., Gauci, C. G., Donadeu, M., Abela-Ridder, B. (2016).
864 Monitoring the outcomes of interventions against *Taenia solium*: options and
865 suggestions. *Parasite Immunol.* 38, 158–169.

866 Mayta, H., Talley, A., Gilman, R.H., Jimenez, J., Verastegui, M., Ruiz, M., Garcia, H.H.,
867 Gonzalez, A.E. (2000). Differentiating *Taenia solium* and *Taenia saginata* infections by
868 simple Hematoxylin-Eosin staining and PCR-Restriction Enzyme Analysis. *J. Clin.*
869 *Microbiol.* 38, 133–137.

870 Millogo, A., Nitiéma, P., Carabin, H., Boncoeur-Martel, M.P., Rajshekhar, V., Tarnagda, Z.,
871 Praet, N., Dorny, P., Cowan, L., Ganaba, R., Hounton, S. (2012). Prevalence of
872 neurocysticercosis among people with epilepsy in rural areas of Burkina Faso. *Epilepsia.*
873 53, 2194–2202.

874 Mkupasi, E.M., Ngowi, H.A., Sikasunge, C.S., Leifsson, P.S., Johansen, M.V. (2013a). Efficacy
875 of ivermectin and oxfendazole against *Taenia solium* cysticercosis and other parasitoses
876 in naturally infected pigs. *Acta Trop.* 128, 48–53.

877 Mkupasi, E. M., Sikasunge, C. S., Ngowi, H. A., Johansen, M. V. (2013b). Efficacy and safety of
878 anthelmintics tested against *Taenia solium* cysticercosis in pigs. *PLoS NTDs.* 7, e2200.

879 Mkupasi, E.M., Ngowi, H.A., Sikasunge, C.S., Leifsson, P.S., Johansen, M.V. (2014).
880 Distribution and histopathological changes induced by cysts of *Taenia solium* in the brain
881 of pigs from Tanzania. *J. Helminthol.* 89, 559–564.

882 Molinari, J.L., Soto, R., Tato, P., Rodriguez, D., Retana, A., Sepulveda, J., Palet, A. (1993).
883 Immunization against porcine cysticercosis in an endemic area in Mexico: a field and
884 laboratory study. *The Am. J. Trop. Med. Hyg.* 49, 502–512.

885 Mwape, K.E., Blocher, J., Wiefek, J., Schmidt, K., Dorny, P., Praet, N., Chiluba, C., Schmidt,
886 H., Phiri, I.K., Winkler, A.S., Gabriël, S.(2015). Prevalence of neurocysticercosis in
887 people with epilepsy in the Eastern province of Zambia. *PLoS NTDs* 9, e0003972.

888 Nakagawa, J., Ehrenberg, J.P., Nealon, J., Fürst, T., Aratchige, P., Gonzales, G., Chanthavisouk,
889 C., Hernandez, L.M., Fengthong, T., Utzinger, J., Steinmann, P. (2015). Towards
890 effective prevention and control of helminth neglected tropical diseases in the Western
891 Pacific Region through multi-disease and multi-sectoral interventions. *Acta Trop.* 141,
892 407–418.

893 Nash, T.E., Garcia, H.H. (2011). Diagnosis and treatment of neurocysticercosis. *Nature Rev.*
894 *Neurol.* 7, 584–594.

895 Ndimubanzi, P.C., Carabin, H., Budke, C.M., Nguyen, H., Qian, Y.J., Rainwater, E., Dickey, M.,
896 Reynolds, S., Stoner, J.A. (2010). A systematic review of the frequency of
897 neurocyticercosis with a focus on people with epilepsy. *PLoS NTDs* 4, e870.

898 Ngowi, H.A., Mukaratirwa, S., Lekule, F.P., Maingi, N., Waiswa, C., Sikasunge, C., Afonso, S.,
899 Sumbu, J., Ramiandrasoa, S., Penrith, M.L., Willingham, A.L. (2013). Agricultural
900 Impact of Porcine Cysticercosis in Africa: A Review. In *Novel Aspects on Cysticercosis*
901 *and Neurocysticercosis. InTech.*

- Nitiéma, P., Carabin, H., Hounton, S., Praet, N., Cowan, L.D., Ganaba, R., Kompaoré, C., Tarnagda, Z., Dorny, P., Millogo, A. (2012). Prevalence case control study of epilepsy in three Burkina Faso villages. *Acta Neurolog. Scandinavica*. 126, 270–278.
- Nkouawa, A., Sako, Y., Li, T., Chen, X., Wandra, T., Swastika, I.K., Nakao, M., Yanagida, T., Nakaya, K., Qiu, D., Ito, A. (2010). Evaluation of a loop-mediated isothermal amplification method using fecal specimens for differential detection of *Taenia* species from humans. *J Clin Microbiol*, 48, 3350–3352.
- Nkouawa, A., Sako, Y., Li, T., Chen, X., Nakao, M., Yanagida, T., Okamoto, M., Giraudoux, P., Raoul, F., Nakaya, K., Xiao, N. (2012). A loop-mediated isothermal amplification method for a differential identification of *Taenia* tapeworms from human: application to a field survey. *Parasitol. Int.* 61, 723–725.
- Nkrumah, B., Nguah, S.B. (2011). *Giardia lamblia*: a major parasitic cause of childhood diarrhoea in patients attending a district hospital in Ghana. *Parasit. Vectors* 4, 163.
- Onah, D.N. and S.N. Chiejina, 1995. *Taenia solium* cysticercosis and human taeniasis in the Nsukka area of Enugu State, Nigeria. *Ann. Trop. Med. Parasitol.*, 89: 399–407.
- OIE, 2015. Terrestrial Animal health Code. Chapter 15.3. Infection with *Taenia solium*.
- Osakwe, C., Otte, W. M., Alo, C. (2014). Epilepsy prevalence, potential causes and social beliefs in Ebonyi State and Benue State, Nigeria. *Epilep. Res.* 108, 316–326.
- Pawlowski, Z. (2016). Taeniosis/Neurocysticercosis control as a medical problem—A discussion paper. *World J. Neurosci.* 6, 2.
- Permin, A., L. Yelifari, P. Bloch, N.P. Steenhard and P. Nansen, 1999. Parasites in cross-bred pigs in the upper East region of Ghana. *Vet. Parasitol.*, 87: 63–71.
- Phiri, I.K., Dorny, P., Gabriël, S., Willingham, A.L., Sikasunge, C., Siziya, S., Vercruysse, J., (2006). Assessment of routine inspection methods for porcine cysticercosis in Zambian village pigs. *J. Helminthol.* 80, 69–72.
- Pondja, A., Nerves, L., Mlangwa, J., Afonso, S., Fafetine, J., Willingham III, A.L., Thamsborg, S.M., (2012). Use of oxfendazole to control porcine cysticercosis in a high endemic area of Mozambique. *PLoS NTDs*. 6, e1651.
- Prüss-Ustün, A., Bartram, J., Clasen, T., Colford Jr, J.M., Cumming, O., Curtis, V., Bonjour, S., Dangour, A.D., De France, J., Fewtrell, L., Freeman, M.C. (2014). Burden of disease from inadequate water, sanitation and hygiene in low and middle income settings: a retrospective analysis of data from 145 countries. *Trop. Med. Internat. Hlth.* 19, 894–905.
- Rodrigues, C.L., De Andrade, D.C., Livramento, J.A., Machado, L.R., Abraham, R., Massaroppe, L., Lucato, L.T., Caramelli, P. (2012). Spectrum of cognitive impairment in neurocysticercosis. Differences according to disease phase. *Neurol.* 78, 861–866.
- Ron-Garrido, L., Coral-Almeida, M., Gabriël, S., Benitez-Ortiz, W., Saegerman, C., Dorny, P., Berkvens, D., Abatih, E.N. (2015). Distribution and potential indicators of hospitalized cases of neurocysticercosis and epilepsy in Ecuador from 1996 to 2008. *PLoS NTDs* 9, e0004236.
- Secka, A., Grimm, F., Victor, B., Marcotty, T., De Deken, R., Nyan, O., Herera, O., Van Marck, E., Geerts, S. (2010a). Epilepsy is not caused by cysticercosis in The Gambia. *Trop. Med. Internat. Hlth.* 15, 476–479.

- Secka, A., Marcotty, T., De Deken, R., Van Marck, E., Geerts, S. (2010b). Porcine cysticercosis and risk factors in The Gambia and Senegal. *J. Parasitol. Res.* <http://dx.doi.org/10.1155/2010/823892>.
- Secka, A., Grimm, F., Marcotty, T., Geysen, D., Niang, A.M., Ngale, V., Boutche, L., Van Marck, E., Geerts, S. (2011). Old focus of cysticercosis in a Senegalese village revisited after half a century. *Acta Trop.* 119, 199–202.
- Thomas, L. F. (2014). Epidemiology of *Taenia solium* Cysticercosis in western Kenya. A Thesis submitted for the degree of Doctor of Philosophy University of Edinburgh.
- Thomas, L.F., 2015. Landscape analysis: control of *Taenia solium*. http://apps.who.int/iris/bitstream/10665/164359/1/9789241508643_eng.pdf (accessed on 3 June 2017).
- Thys, S., Mwape, K.E., Lefèvre, P., Dorny, P., Marcotty, T., Phiri, A.M., Phiri, I.K., Gabriël, S. (2015). Why latrines are not used: communities' perceptions and practices regarding latrines in a *Taenia solium* endemic rural area in Eastern Zambia. *PLoS NTDs* 9, e0003570.
- Trevisan, C., Mkupasi, E. M., Ngowi, H. A., Forkman, B., Johansen, M. V. (2016). Severe seizures in pigs naturally infected with *Taenia solium* in Tanzania. *Vet. Parasitol.* 220, 67–71.
- Trevisan, C., Devleesschauwer, B., Schmidt, V., Winkler, A. S., Harrison, W., Johansen, M. V. (2017). The societal cost of *Taenia solium* cysticercosis in Tanzania. *Acta Trop.* 165, 141–154.
- Tsang, V.C., Brand, J.A., Boyer, A.E. (1989). An enzyme-linked immunoelectrotransfer blot assay and glycoprotein antigens for diagnosing human cysticercosis (*Taenia solium*). *J. Infect. Dis.* 159, 50–59.
- Weka, R.P., Ikeh, E.I. (2009). Seroprevalence of cysticercosis and intestinal parasitism in pigs in Jos metropolis. *J. Anim. Vet. Adv.* 8, 883–887.
- Weka, R. P., Ikeh, E. I., Kamani, J. (2013). Seroprevalence of antibodies (IgG) to *Taenia solium* among pig rearers and associated risk factors in Jos metropolis, Nigeria. *The J. Infect. Dis. Dev. Countries.* 7, 067–072.
- WHO. Prevention and control of schistosomiasis and soil-transmitted helminthiasis: Report of a WHO Expert Committee. World Health Organization; 2002.
- WHO (2010). Working to overcome the global impact of neglected tropical diseases - First WHO report on neglected tropical diseases. Geneva: World Health Organization; 2010.
- WHO (2016a). A rationale for investment and action World Health Organization 2016 WHO/HTM/NTD/NZD/2016.
- WHO (2016b). *Taenia solium* taeniasis/cysticercosis diagnostic tools report of a stakeholder meeting Geneva, 17–18 December 2015 WHO/HTM/NTD/NZD/2016.4.
- WHO (2015a). Investing to Overcome the Global Impact of Neglected Tropical Diseases: Third WHO Report on Neglected Tropical Diseases 3.
- WHO (2015b). Assembling a framework for intensified control of taeniasis and neurocysticercosis caused by *Taenia solium*: report of an informal consultation, WHO Headquarters, Geneva, 17-18 July 2014.
- WHO(2015c). *Taenia Solium* Taeniasis/cysticercosis diagnostic tools. Report of a stakeholder meeting, Geneva, 17-18 December 2015. 1ra Ed. Génova: WHO UNICEF. 2015.

- Winkler, A.S. (2013). Epilepsy and neurocysticercosis in sub-Saharan Africa. In: Foyaca-Sibat H (ed). Novel aspects on cysticercosis and neurocysticercosis. InTech, Croatia, 2013; 307-340. <http://www.intechopen.com/books/show/title/novel-aspects-on-cysticercosis-and-neurocysticercosis> accessed June 2014.
- Winkler, A. S., Richter, H. (2015). Landscape analysis: management of neurocysticercosis with an emphasis on low- and middle-income countries. World Health Organization WHO/HTM/NTD/NZD/2015.05.
- Winskill, P., Harrison, W. E., French, M. D., Dixon, M. A., Abela-Ridder, B., Basáñez, M. G. (2017). Assessing the impact of intervention strategies against *Taenia solium* cysticercosis using the EPICYST transmission model. *Parasit.Vectors* 10, 73.
- Yamasaki, H., Allan, J.C., Sato, M.O., Nakao, M., Sako, Y., Nakaya, K., Qiu, D., Mamuti, W., Craig, P.S., Ito, A., (2004). DNA differential diagnosis of taeniasis and cysticercosis by multiplex PCR. *J. Clin.Microbiol*, 42,548-553.
- Zoli, A., Shey-Njila, O., Assana, E., Nguekam, J. P., Dorny, P., Brandt, J., Geerts, S. (2003). Regional status, epidemiology and impact of *Taenia solium* cysticercosis in Western and Central Africa. *Acta Trop.* 87, 35–42.