**QUEEN'S
UNIVERSITY
BELFAST**

Evidence for co-selection of antibiotic resistance genes and mobile genetic elements in metal polluted urban soils

Zhao, Y., Cocerva, T., Cox, S., Tardif, S., Su, J. Q., Zhu, Y. G., & Brandt, K. K. (2019). Evidence for co-selection of antibiotic resistance genes and mobile genetic elements in metal polluted urban soils. *Science of the Total Environment*, 656, 512-520. <https://doi.org/10.1016/j.scitotenv.2018.11.372>

Published in:
Science of the Total Environment

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2018 Elsevier.

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License

(<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

1		
2		
3		
4	1	Evidence for co-selection of antibiotic resistance genes and mobile genetic elements in metal polluted urban
5		
6	2	soils
7		
8		
9	3	Yi Zhao ^{a,*} , Tatiana Cocerva ^b , Siobhan Cox ^b , Stacie Tardif ^a , Jian-Qiang Su ^c , Yong-Guan Zhu ^{c,d} , Kristian Koefoed
10		
11	4	Brandt ^{a,e} *
12		
13		
14	5	^a Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Thorvaldsensvej
15		
16	6	40, 1871 Frederiksberg, Denmark
17		
18		
19	7	^b School of Natural and Built Environment, Queen’s University of Belfast, David Keir Building, Stranmillis Road
20		
21	8	Belfast, BT9 6AX, United Kingdom
22		
23		
24	9	^c Key Laboratory of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences,
25		
26	10	1799 Jimei Road, Xiamen 361021, China
27		
28		
29	11	^d State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese
30		
31	12	Academy of Sciences, Beijing 100085, China
32		
33		
34	13	^e Sino-Danish Centre for Education and Research, Huairou District, Beijing, China
35		
36		
37	14	
38		
39		
40	15	*Corresponding Authors
41		
42	16	Yi Zhao, Section for Microbial Ecology and Biotechnology, Department of Plant and Environmental Sciences,
43		
44	17	University of Copenhagen, Thorvaldsenvej 40, DK-1871 Frederiksberg C, Denmark
45		
46		
47	18	E-mail: y.zhao@plen.ku.dk, Phone: +45-35334549
48		
49		
50	19	Kristian Koefoed Brandt, Section for Microbial Ecology and Biotechnology, Department of Plant and
51		
52	20	Environmental Sciences, University of Copenhagen, Thorvaldsenvej 40, DK-1871 Frederiksberg C, Denmark
53		
54		
55	21	E-mail: kkb@plen.ku.dk, Phone: +45-35332612
56		
57		
58	22	
59		
60		
61		
62		
63		
64		
65

Highlights:

1. Fifty archived urban soils from the Belfast area Tellus Survey were analyzed.
2. Antibiotic resistance determinants were profiled by high throughput qPCR chip.
3. Evidence for metal-induced co-selection of antibiotic resistance genes (ARGs).
4. Total ARGs were positively correlated with total mobile genetic elements.
5. Metals may confer persistent selection pressures for ARGs in urban soils.

1 **Abstract**

2 Antibiotic resistance genes (ARGs) constitute emerging environmental pollutants and pose risks to public
3 health. Toxic metals are known to select for metal-resistant bacteria in metal-contaminated soils, but there
4 is growing concern that metal contaminants can also act as co-selective agents thereby causing
5 environmental proliferation of antibiotic resistance. In this study, we quantified ARGs and selected
6 mobile genetic elements (MGEs) known to constitute potential ARG hosts in 50 archived urban and
7 suburban soils from the Belfast metropolitan area using a high-throughput qPCR ARG chip. ARG
8 prevalence was linked to concentrations of individual metals and a soil metal toxicity index calculated
9 based on the relative toxicity of different metals to soil microbial processes. A total of 164 ARGs were
10 detected across the 50 soils analyzed with an average absolute abundance of 3.4×10^7 ARG gene copies
11 per gram of soil. A significant correlation between abundance of ARGs and MGEs was observed,
12 suggesting the importance of horizontal gene transfer for ARG dissemination. Network analysis revealed
13 significant co-occurrence patterns between specific metals (As, Cd, Co, Cr, Cu, Hg, Ni and Zn) and
14 associated ARGs. Path analysis further indicated that the soil metal toxicity index significantly affected
15 the number of detected ARGs ($\lambda = 0.32$, $P < 0.001$) and the abundance of metal co-occurring ARGs ($\lambda =$
16 0.612 , $P < 0.001$) via effects on MGEs. Collectively, our results indicate a role of soil metals in co-
17 selection of ARGs and MGEs in urban and semi-urban soils and suggest a risk for environmental ARG
18 dissemination via horizontal gene transfer.

19
20 **Keywords**

21 Antibiotic resistance genes, mobile genetic elements, co-selection, metal toxicity, qPCR chip, urban soils

22

23

1. Introduction

The rapid emergence of antibiotic-resistant bacteria is occurring worldwide, posing threats to global public health, food security and development (Ventola, 2015). According to the World Health Organization (WHO), a post-antibiotic era is emerging, in which antibiotic resistance threatens the effective prevention and successful treatment of an ever-increasing range of bacterial infections. The critical role of the environment for development and dissemination of antibiotic resistance has now been recognized (Ashbolt et al., 2013). Consequently, antibiotic resistance genes (ARGs) and mobile genetic elements carrying these genes can be regarded as emerging environmental pollutants (Gillings et al., 2008; Pruden et al., 2006).

The widespread use of antibiotics is generally considered to be the primary cause for elevated levels of ARGs in pathogenic bacteria (Ventola, 2015), but there is growing concern that contaminants such as metals and biocides may also co-select for antibiotic resistance (Baker-Austin et al., 2006; Berendonk et al., 2015; Hoffman et al., 2005; Pal et al., 2017; SCENIHR, 2009). Co-selection of antibiotic and metal(loid) resistance have been associated with arsenic (As), cadmium (Cd), cobalt (Co), chromium (Cr), copper (Cu), mercury (Hg), nickel (Ni), lead (Pb) and zinc (Zn) (Knapp et al., 2011; Pal et al., 2017; Pal et al., 2015; Seiler and Berendonk, 2012; Yazdankhah et al., 2014). Several co-selection mechanisms are known (Baker-Austin et al., 2006). The genes encoding resistance to antibiotics and metals may for instance be found on the same mobile genetic elements (plasmid, integron or transposon), and this physical linkage results in co-resistance. Cross-resistance is another co-selection mechanism which occurs when single genes encode resistance to both antibiotics and metals.

The relative importance of antibiotics and co-selecting agents for the selection of antibiotic resistance is likely to differ between different environments. Antibiotic residues primarily accumulate to toxic levels

able to strongly select bacterial communities in habitats where antibiotics are used by humans (e.g., human or animal gut), whereas antibiotics only rarely accumulate to toxic levels in soil or water (Brandt et al., 2015). By contrast, metals frequently accumulate to toxic levels in some environmental compartments including both agricultural and urban soils (Brandt et al., 2010; Berg et al., 2012; Imfeld et al., 2011; McLaughlin and Smolders, 2001). Indeed, toxic metals may in some cases provide stronger and more persistent selective pressures for environmental selection of antibiotic resistance as compared to antibiotic residues (Song et al., 2017). Consequently, metal-induced co-selection for ARGs in metal contaminated environments represents a risk factor for the expansion of the soil bacterial resistome and may thus represent a barrier for reversal of antibiotic resistance even if antibiotic residues are prevented from reaching the environment.

Continuous accumulation of metal(loid) contaminants widely occurs in urban soils (Luo et al., 2012). Over half of the world's population currently lives in urban areas, and the urban population continues to grow (United Nations, 2014) suggesting increasingly important linkages between urban environmental quality and human health (Li et al., 2018). Due to rapid urbanization and intensive anthropogenic activity, a massive volume of potential selective agents (such as heavy metals) and microbes carrying ARGs swarm into urban soils and successfully persist, increasing the level of ARG pollution in urban environments (Wang et al., 2014). These ARGs spread amongst humans and in the environment by horizontal gene transfer (HGT), developing into pathogenic antibiotic-resistant bacteria (pARB) and thereby raising the risk of failure of antibiotic treatments (Zhu et al., 2018). The linkage between soil metals and ARGs from urban and industrially polluted soils has been observed in previous studies (Knapp et al., 2017, Berg et al. 2010). However, only a few metals and a limited number of ARGs were targeted and comprehensive evidence for the ability of metals to co-select antibiotic-resistance in urban soils is still largely lacking (Larsson et al., 2018). Therefore, monitoring ARG distribution in urban environments and its association with metals as co-selective agents is necessary.

73

74 In this study, we profiled ARGs in metal-contaminated soils within an urban area to improve our
75 understanding of the role of metal contaminations in developing ARGs. To this end, 50 archived soils
76 across the urban area of Belfast, Northern Ireland with a gradient of various metal contaminations were
77 selected and examined using a high-throughput qPCR based ARG chip. The ARG profile was
78 subsequently linked to MGEs and metal contamination by multivariable statistical analyses and network
79 analysis. Cluster analysis, permanova test with land use and distance-decay analysis was applied to depict
80 the influence of geographic factors (spatial location) on ARG distribution. The path analysis was further
81 performed to test our hypotheses of metal-induced co-selection on ARGs.

82

83 **2. Materials and Methods**

84 **2.1. Study area**

85 Belfast is the capital and the largest city in Northern Ireland with a population of approximately 700 000
86 (NISRA, 2016). During the 18th and 19th centuries, Belfast grew to be the leading industrial city in
87 Ireland, with thriving linen and shipbuilding industries. Belfast continued to play an important role
88 throughout the industrial revolution in the 19th century and is also historically recognized for tobacco-
89 processing, rope making, glass manufacturing, tobacco production and distilleries (Royle et al., 2007).
90 Concentrations of metal(loid)s in the Belfast area have been shown to act as an ‘urbanization tracer’
91 (McIlwaine et al., 2017) and differential metal(loid) concentrations can, therefore, be largely attributed to
92 anthropogenic contamination. By contrast, anthropogenic contamination of the Belfast area soils with
93 antibiotics or ARGs can generally be considered negligible. Human fecal wastes are almost exclusively
94 released to aquatic recipients through wastewater collection and treatment processes (EMEA, 2006).
95 Leakage of untreated wastewater from the collection network into soil could potentially provide a route
96 for antibiotics and ARGs to enter the soil. However, the collection network is at a greater depth than the

soils investigated in this study (5-20 cm depth), and significant top soil contamination is therefore highly unlikely. We therefore assume that ARGs present in the sampled top soils of the Belfast area were primarily derived from the natural, ‘background’ antibiotic resistome (e.g. ARGs in soil-dwelling bacteria producing or degrading antibiotics; (D’Costa et al., 2007; Dantas et al., 2008)) with potential contributions from other diffuse sources (e.g. ARGs in fecal matter from wildlife, rain water, or particles deposited via air). Hence, the metropolitan area of Belfast provides an opportunity to evaluate the ability of soil metal(loid)s to co-select antibiotic resistance. The boundaries of the study area have been defined using the Corine land cover data (European Environment Agency, 2012) satellite images and the spatial distribution of the available urban soil samples.

2.2. Soil collection

The Tellus geochemical and geophysical survey was undertaken across the entire region of Northern Ireland by the Geological Survey of Northern Ireland (GSNI) between 2004 and 2007. Shallow soil samples (5-20 cm) from 1166 sampling locations were collected and archived at room temperature (< 25°C). Full details of the sampling strategy are described in a previous study (Smyth, 2007). In this study, a total of 50 sampling locations were selected (Figure 1a) and their archived urban soils (Knights, 2006) were retrieved from the GSNI Tellus survey archive in October 2016. Original sampling coordinates are listed in Table S1. Sample selection targeted a broad spectrum of toxic elements, with a wide range of concentrations, and included a variety of different land uses while ensuring samples were spatially distributed across the city.

2.3. Soil chemical characterization

Archived metal concentration data (As, Cd, Co, Cr, Cu, Hg, Ni, Pb and Zn) were retrieved from the Tellus database for downstream analyses. The spatial distribution of these elements across Belfast has

been reported previously (McIlwaine et al., 2017). Concentrations of As in soils were found to be controlled by anthropogenic inputs. By contrast, concentrations of Cu and Zn were influenced by both anthropogenic and geogenic inputs, while concentrations of Ni, Co and Cr were mainly attributed to geogenic sources (McIlwaine et al., 2017).

A metal toxicity index (TI_{metals}) was calculated for each soil sample to provide a normalized measure of the bacterial selection pressure posed by the toxic metals present in each sample. TI_{metals} was calculated based on previously recorded effects of individual metal(loid)s (As, Cd, Co, Cr, Cu, Hg, Ni, Pb and Zn) on soil microbial processes (Welp, 1999) following a previously established procedure (Azarbad et al., 2015; Stefanowicz et al., 2008): $TI_{\text{metals}} = \sum (C_i/EC_{50i})$, where C_i equals the total concentration of the element i in the soil and EC_{50i} equals the half-maximal effective concentration for that particular element i (Welp, 1999).

Archived soil pH and land use data were also retrieved from the Tellus database (Table S1). Full details of the analytical methods used and quality assurance/quality control procedures adopted in the Tellus geochemical survey can be found in a previous study (Smyth, 2007).

2.4. DNA extraction

The 50 selected soil samples were retrieved from the Tellus soil archive and aseptically weighed into prepared sterile plastic bags. A total of 250 - 400 mg of dry soil was transferred into PowerBead Tubes (MoBio) and incubated for rehydration at room temperature for 20-30 minutes. DNA was extracted from rehydrated soil using MoBioPowerSoil® DNA Isolation Kit according to the instruction manual. The concentration and purity of DNA were checked using ultraviolet absorbance (ND1000, Nanodrop,

Thermo Fisher Scientific). DNA was eluted in 100 µl elution buffer and stored at -20 °C for downstream analysis. Long-term storage has been proven to not cause bias in DNA results (Tzeneva et al., 2009, Knapp et al., 2010).

2.5. High-throughput qPCR

A total of 296 primer sets (Table S2) were used to examine Belfast urban soils. These 296 primer sets targeted antibiotic resistance genes (285 primer sets), mobile genetic elements (10 primer sets) and a 16S rRNA gene as a reference gene (Zhao et al., 2018; Zhu et al., 2017). Collectively, the targeted ARGs confer resistance to all major classes of antibiotics including aminoglycoside, beta-lactamase, chloramphenicol, macrolide-lincosamide-streptogramin B (MLSB), multidrug, sulfonamide, tetracycline and vancomycin. Targeted mobile genetic elements included transposase genes (8 primer sets) and Class 1 integron-integrase gene (2 primer sets). The HT-qPCR was performed with an HT-qPCR based ARG chip using the WaferGen SmartChip Real-time PCR system. Negative controls were included. The thermal cycle consisted of 10 min at 95 °C, followed by 40 cycles of denaturation at 95 °C for 30 s and annealing at 60 °C for 30 s. Melting curve analyses were automatically conducted by Wafergen SmartChip qPCR software.

All HT-qPCR were performed in technical triplicates with negative control. The data from reaction wells with r^2 smaller 0.99 were discarded. Only data for samples with three technical replicates that all generated amplification products were regarded as positive detection and used in further data analysis. Relative copy number was calculated based on previously published method (Looft et al., 2012): relative gene copy number = $10^{(31-C_T)/(10/3)}$, where C_T refers to the qPCR results and 31 refers the cut-off point. The normalized abundance of a gene (copies per 16S rRNA) was calculated by dividing relative gene copy number of the gene by relative copy number of reference gene 16S rRNA.

168

169 Absolute 16S rRNA copy numbers (copies per gram of soil) were determined using the standard curve
170 method on a Roche 480 system. Each 20 μ l qPCR reaction mixture consisted of 10 μ l 2 $\times$ LightCycle 480
171 SYBR Green I Master, one μ g μ l⁻¹ bovine serum albumin, one μ M of each primer, one ng μ l⁻¹ DNA as
172 template and six μ l nuclease-free PCR-grade water. The thermal cycle consisted of a 10 min enzyme
173 activation at 95 °C, followed by 40 cycles of denaturation at 95 °C for 30 s, annealing at 60 °C for 30 s
174 and extension at 72 °C for 15 s. A plasmid control containing a cloned and sequenced 16S rRNA gene
175 fragment (1.39×10^{10} copies per liter) was used to generate calibration curves from a tenfold dilution for
176 standard calculation. All qPCRs were performed in technical triplicates with negative controls. Absolute
177 ARG copy numbers were calculated by transforming relative copy numbers by normalization from
178 absolute 16S rRNA gene copy number.

179

180 2.6. Statistical analysis

181 All HT-qPCR data filtration and calculations were performed using Microsoft Excel 2010. Bar charts and
182 scatter diagrams were generated using Origin Pro 9.1. Correlation analyses and significance tests were
183 performed using IBM SPSS Statistics 22. Heatmaps were generated using HemI 1.0 (Deng et al., 2014).
184 Shannon H index was determined using PAST Statistics Software (Hammer et al., 2001). The distance-
185 decay analysis and permanova test were conducted using R 3.4.1 with the vegan package (Oksanen et al.,
186 2007). The co-occurrence patterns between ARGs (normalized abundance) and MGEs (normalized
187 abundance)/metals (total concentration) were explored using network analysis. To visualize the
188 correlations in the network interface, a correlation matrix was constructed using all pairwise Pearson's
189 rank correlations. Only correlations with Pearson's $r > 0.7$ (or < -0.7), a significance level of $P < 0.05$,
190 and p-values lower than the critical value calculated by the Benjamini-Hochberg procedure (false
191 discovery correction) were considered robust and used for forming the co-occurrence networks. The

network structure was confirmed by randomization procedure from randomized data. Network visualization was performed in Cytoscape 3.6.0 (Shannon et al., 2003). For spatial data representation, maps were produced using the ArcGIS software ArcMap version 10 (ESRI, 2010).

The path analysis, as a special case of structural equation model (SEM), has been proven useful as a statistical tool to explore the complex networks of causal relationships ecosystems (Eisenhauer et al., 2015). In this study, it was performed to evaluate the overall effect of soil metal contamination (represented by TI_{metals}) on ARG patterns as represented by both the number of detected ARGs and the normalized abundance of all co-occurring ARGs using SPSS AMOS. The path model was established based on the following theoretical assumptions: (i) metal contamination may directly influence ARG patterns by acting as a direct selective agent; (ii) metal contamination may indirectly affect ARG patterns through MGE associated co-selection processes, with co-resistance as a mechanism. The data were fitted to the estimated model using a maximum-likelihood estimation method. The model fit was tested and an overall goodness-of-fit of the model was indicated by satisfaction of the following criteria: (i) non-significant Chi-square value ($P > 0.05$); (ii) low root mean square error of approximation as absolute fit index (RMSEA < 0.08); (iii) high increment fit index (CFI, GFI, TLI, NFI > 0.95 ; CFI as comparative fit index, GFI as goodness of fit, TLI as Tucker-Lewis index and NFI as Normed-fit index). The disturbance terms (also called the residual error terms) were added into the model to reflect the unexplained variance and measurement error. The path coefficients (standardized regression weights) and significance were calculated in an SPSS AMOS, showing the effect of an independent variable on a dependent variable in the path model. The standardized direct, indirect and total effects were automatically calculated using SPSS AMOS following the method published previously (Finney, 1972).

3. Results

3.1. Diversity and abundance of ARGs in Belfast urban area

A total of 175 genes (164 ARGs, eight transposase genes, two Class 1 integron-integrase genes and the 16S rRNA gene) were detected by the HT-qPCR chip in the 50 studied urban soils (Figure 1). The detected ARGs represented most major resistance mechanisms including antibiotic deactivation (43%), efflux pumps (34%) and cellular protection (19%) (Figure 1b) and confer resistance to most major classes of antibiotics administered to humans and animals, including aminoglycosides, beta-lactams, chloramphenicols, MLSB, sulfonamides, tetracyclines, vancomycin and multidrug. Almost half of the detected ARGs confer resistance to beta-lactams (23%) and multidrug (23%) (Figure 1c). The number of detected ARGs in the Belfast urban area ranged from 8 to 137 with the average at 67 (Figure 1a). The Shannon diversity H index of ARGs varied from 1.2 to 3.4 in Belfast urban samples (Figure S1). To compare the spatial similarity of ARG patterns, we examined the presence and absence of ARGs across all samples. Only two shared core ARGs (*mphA-02* and *cphA-01*) conferring resistance to macrolide and beta-lactam antibiotics were found in all samples.

The absolute abundance of ARGs in Belfast urban soils varied over six orders of magnitude (6.8×10^2 to 1.7×10^8 copies per gram of soil) with an average of 3.4×10^7 (Figure 1a). The predominant ARGs encoded beta-lactam, multidrug, aminoglycoside or chloramphenicol resistance (Figure S2). The three most abundant ARGs were *mexF*, *cphA-01* and *cmx(A)*. The *mexF* gene, which encodes a multidrug resistance efflux pump for chloramphenicol and fluoroquinolone, was detected in 49 soil samples at $5 - 229 \times 10^7$ copies per gram of soil. The *cphA* gene confers resistance to different beta-lactams antibiotics (penicillin, cephalosporin and carbapenem) via hydrolysis of the beta-lactam ring and was detected in all 50 soils with an average abundance of 6.6×10^6 copies per gram of soil. The *cmx(A)* gene encodes a chloramphenicol exporter and was found in 49 samples, varying in abundance from $2 - 43 \times 10^7$ copies per gram of soil.

240

241 To better explore the prevalence of ARGs within the studied soil bacterial communities, the absolute
242 abundance of ARGs and MGEs was normalized relative to the abundance of the 16S rRNA gene.
243 Bacterial abundances in soils were in the range of 8.7×10^3 to 9.0×10^8 with an average of 2.2×10^8
244 copies per gram. The normalized ARG abundance in 50 soils varied from 0.06 to 0.77 with an average of
245 0.15 ARG copies per 16S rRNA gene.

246

247 We further investigated the spatial distribution of ARGs with distance-decay analysis and cluster analysis.
248 However, our results show that ARG distribution in the Belfast urban area was unlikely to have been
249 driven by geographic factors. The geographic distance did not show any significant correlation with the
250 similarity of ARG communities between samples (Pearson's $r = -0.026$, $P = 0.355$) (Table S3). Likewise,
251 land use at the time of soil sampling could not explain the observed ARG patterns. Combining the cluster
252 analysis with a heatmap to visualize the ARG profiles in different land uses, we thus did not find any
253 significant effect of land use (cluster) on the ARG distribution in Belfast urban soils (Figure S3&4). This
254 conclusion was further confirmed by permanova test ($R^2 < 0.02$, $P > 0.05$; Table S4). Both results
255 indicated the inconsequential role that geographic factors played in determining the fate of ARGs in
256 Belfast urban soils.

257

258 3.2. MGEs and their associations with ARGs

259 A total of two integron genes and eight transposon genes were targeted and detected in this study. The
260 total absolute abundance of these MGEs ranged from $2-38 \times 10^7$ copies per gram of soil with an average
261 of 3.4×10^6 . The normalized abundance of MGEs ranged from 0.004 to 0.069 copies per 16S rRNA gene
262 with average at 0.014. Class 1 integron, *intI-1*, was found in all 50 soils. Significant correlations were
263 found between MGEs and ARGs (Figure 2). The normalized abundance of MGEs was positively

correlated with the number of ARGs detected (Pearson's $r = 0.57$, $P < 0.001$), as well as normalized abundance of ARGs (Pearson's $r = 0.66$, $P < 0.001$) (Figure 2a). Positive correlations were also found between the absolute abundance of ARGs and the MGEs: Class 1 integrons (Pearson's $r = 0.97$, $P < 0.001$) and transposons (Pearson's $r = 0.66$, $P < 0.001$) (Figure S5).

The co-occurrence pattern between specific ARGs and MGEs were revealed by network analysis based on Pearson correlations ($r > 0.7$, $P < 0.05$) (Figure 2b). The network consists of 130 nodes corresponding to 8 MGEs and 122 ARGs. A total of 358 strong correlations between these MGEs and ARGs were found, including 334 positive correlations and 24 negative correlations. MGEs including integrons and transposons both exhibited a co-occurrence pattern with different types of ARGs. A total of 81 ARGs were positively correlated with the class 1 integron, *intI-1*, while 77 ARGs were positively correlated with the clinical class 1 integron, *cIntI-1*. For transposons, 66 ARGs were found positively correlated to *tnpA* and *IS613* genes. Among the 358 correlations between ARGs and MGEs, 31% were contributed by multidrug-resistant genes, while beta-lactam, MLSB, vancomycin, aminoglycoside, tetracycline and chloramphenicol resistant genes accounted for 20%, 12%, 12%, 7%, 6% and 2%, respectively.

3.3. Co-occurrence pattern between metals and ARGs

The soil samples profiled for ARGs and MGEs were contaminated to varying degrees by metals. The metals with the potential of co-selection for ARGs and their concentration ranges in the soil samples were shown in Table 1. The co-occurrence pattern between metals and ARGs was further explored by network analysis based on Pearson's correlation ($r > 0.7$, $P < 0.05$) (Figure 3). The network consists of 25 nodes including eight metals and 17 ARGs. A total of 24 significant positive correlations were found between metals and ARGs. No significant negative correlation was found between metals and ARGs. As, Cd, Co, Cr, Cu, Hg, Ni and Zn all exhibited co-occurrences with specific ARGs. Zinc was found to co-occur with

eight resistance genes conferring resistance to aminoglycosides (4), multidrug (3) and beta-lactam antibiotics (1). Copper co-occurred with aminoglycoside resistance genes (*aadA* and *aac*) and MLSB resistance genes (*mefA*). The number of ARGs that co-occurred with Cd, Co, Ni, Hg, Cr and As were 3, 3, 2, 2, 2 and 1, respectively. These metal(loid)s all co-occurred with 17 specific ARGs conferring resistance to aminoglycosides, beta-lactams, MLSBs, multidrug, tetracycline and vancomycin. Among the 24 connections, almost half were contributed by a combination of aminoglycoside (25%) and multidrug resistance genes (21%).

3.4. The effects of metal toxicity index and MGEs on ARGs

To further assess the effects of metals and MGEs on ARGs, structural equation model (SEM) based path analysis was performed with a multiple-pathways model based on the theoretical assumptions outlined in section 2.6 (Figure 4a). The path analysis indicated that the degree of soil metal contamination (metal toxicity index) had a significant direct positive association on normalized abundance of co-occurring ARGs (copies per 16S rRNA gene) ($\lambda = 0.187$, $P < 0.01$) and a significant indirect positive effect on number of detected ARGs ($\lambda = 0.251$, $P < 0.001$) (Figure 4b) and normalized abundance of co-occurred ARGs (copies per 16S rRNA gene) ($\lambda = 0.426$, $P < 0.001$) (Figure 4c). The total standardized effects of metal toxicity index on the number of detected ARGs and normalized abundance of co-occurring ARGs (copies per 16S rRNA gene) were 0.323 and 0.612, respectively. This suggests that increase of one unit of one metal toxicity index resulted in 0.323 more numbers of ARGs detected and 0.612 more copies of co-occurred ARGs per 16S rRNA gene. Metal toxicity changed MGE abundances in soil ($\lambda = 0.521$, $P < 0.001$) and subsequently influenced the number of detected ARGs and abundance of co-occurring ARGs ($\lambda = 0.483$, $P < 0.001$ and $\lambda = 0.817$, $P < 0.001$) (Figure 4a). Metal toxicity index and MGE abundance both had a total positive effect on the number of detected ARGs and abundance of co-occurred ARGs. The path analysis results were further tested and confirmed by correlation analysis. The correlation analysis showed significant positive correlations among metal toxicity index, the abundance of MGE,

number detected ARGs and abundance of co-occurring ARGs (Figure S6), which further confirmed the positive association between metals and ARGs via MGEs. Significant positive correlations were also found between soil pH and the number of detected ARGs (Pearson's $r = 0.329$, $P < 0.05$), and between the absolute abundance of the 16S rRNA gene and total ARGs (copies per gram) (Pearson's $r = 0.905$, $P < 0.001$).

4. Discussion

4.1. Evidence for metal-induced co-selection of ARGs in urban soils

To the best of our knowledge, this present study provides the most comprehensive investigation of the ability of metal contamination to affect the distribution of ARGs in urban soils to date. The co-occurrence between specific metal contaminants and specific ARGs together with the significant positive effect of overall soil metal loading (metal toxicity index) on soil ARGs are key findings and suggests co-selection of metals and ARGs (Figure 3 & 4). Not only was a diverse range of ARGs detected, many of which displayed significant co-occurrence patterns with both specific metals and overall metal load (metal toxicity index), but we were also able to identify a potential causal link between soil metals and ARGs. Hence, observed ARG patterns could not be explained by current land use or geographic location and with the possible exception of the two included pasture soils (Table S1), the studied soils were generally unlikely to have received significant recent point source inputs of fecal materials from humans or animals treated with antibiotics. Soil pH also affected observed ARG patterns; this could most likely be attributed to the known abilities of pH to modulate bacterial community composition (Rousk et al., 2010) and the bioavailability/toxicity of metals (Smolders et al., 2009).

4.2. Co-selection mechanisms and potential for horizontal gene transfer of ARGs

Although our study was not specifically designed to compare the relative importance of different co-selection mechanisms (Baker-Austin et al., 2006), it clearly demonstrated co-selection of ARGs and MGEs (Figure 4). The strong linkages between metal toxicity index, MGEs and ARGs indicate a significant metal impact on both the diversity and abundance of ARGs via MGEs (Figure 4). According to our results of path analysis, 78% of the total effect of metal toxicity on the number of detected ARGs and 69% of the total effect of metal toxicity on the abundance of co-occurring ARGs were observed to occur via MGEs. The majority of co-occurring ARGs with metals were found to also co-occur with MGEs, suggesting an underlying metal-driven co-selection process with co-resistance (i.e. linkage of genes conferring resistance to metals and antibiotics on the same genetic element) as the major mechanism for most studied ARGs that do not have any known roles in bacterial metal resistance. However, the resistance genes *acrF*, *adeA*, *ttgB*, *qacEAI*, *rarD*, *tetPA* and *mefA* encode efflux pumps as their resistance mechanism and cross-resistance with other classes of antimicrobial agents via efflux cannot be ruled out. For instance, the multi-drug resistance pumps encoded by the genes *acrF*, *adeA*, *qacEAIrarD* and *ttgB* can export both metals and antibiotics for detoxification purposes (Mata et al., 2000).

The ability of soil bacteria to transfer ARGs to pathogenic bacteria of clinical relevance is of considerable public health concern (Forsberg et al., 2012, Ashbolt et al., 2013, Graham et al., 2016). Importantly, our study indicates that urban soil metal pollution co-selected ARGs that may be prone to horizontal gene transfer between different species of bacteria. Results of correlation analysis, network analysis and path analysis all showed that an increase in MGE abundance was strongly correlated with an increase in ARG diversity and abundance (Figure 2, 4 & S6). The co-occurrence pattern of ARGs and MGEs revealed by network analysis showed several clusters within the network. Resistance genes such as *aac* and *aad* genes, known to be carried within integron gene cassettes (An et al., 2018; Partridge et al., 2009), were strongly correlated with the class 1 integron gene *IntI-1*, clinical class 1 integron gene *cIntI-1* and transposon

genes *tnpA* and *IS613* (Figure 2). The clinical class 1 integron-integrase gene, *cIntI-1*, exhibits rapid responses to various environmental pressures (including toxic metals) and thus has been proposed as a marker for anthropogenic pollution and as an emerging pollutant (Gillings et al., 2015; Gillings, 2018). The significant correlation and co-occurrence pattern between the clinical class 1 integron-integrase gene (*cIntI-1*) and ARGs (Table S5 & Figure 2) may therefore suggest that metal contamination increases environmental selection of bacteria containing clinical Class 1 integrons conferring resistance to both metals and antibiotics even in the absence of a selection pressure exerted by antibiotic residues.

4.3. Conclusions and perspectives for the environmental dissemination of ARGs.

Our findings provide evidence that metal contamination (As, Cd, Co, Cr, Cu, Hg, Ni and Zn) could significantly affect the diversity, abundance and mobility potential of a broad spectrum of ARGs in urban soils. Collectively, our results suggest that urban soil metal contamination increases the potential for horizontal gene transfer of ARGs via co-selection of ARGs and MGEs thereby generating a pool of high-risk mobile ARGs (Martínez et al., 2015). Indeed, it is likely that metals in many soils confer more important selective agents than antibiotic residues because metals, as opposed to antibiotics, frequently accumulate to persistent toxic levels in contaminated soils (Song et al., 2017). Based on the ongoing recruitment of resistance genes by MGEs (Gillings et al., 2015; Gillings, 2018), we further hypothesize that soil metals may become an even more important public health risk factor for the evolution (e.g. selection, mobilization, transfer and persistence) of ARGs in the future. Hence, we call for additional research into metal-induced co-selection of ARGs and MGEs in metal contaminated soils in the interest of public health (Larsson et al., 2018).

Urban soils are commonly contaminated with toxic metals (McIlwaine et al., 2017) and our study thus has implications for environmental regulation and management of urban soils worldwide. Hence, urban residents may be exposed to ARGs via ingestion of metal-contaminated urban soils (especially children)

or vegetables (urban gardening) unless adequate precautions are taken. Metals and co-selected ARGs may also be leached to groundwater or transported by surface runoff to aquatic receptors implying subsequent human ARG exposure via drinking water (Frey et al., 2015, Ma et al., 2017) or recreational activities (Leonard et al., 2018). Specific metals will probably co-select for different ARGs in different soils and it will therefore not be relevant to monitor only a few ARGs when evaluating risks posed by toxic soil metals. We therefore propose that metal susceptibility in indicator soil bacteria or absence of pollution-induced community tolerance to metals (Lekfeldt et al., 2014) might serve as additional protection goals for environmental risk assessment of toxic metals in soil. In conclusion, serious consideration is needed to set minimum standards for retarding ARGs and for mitigating the accumulation of toxic metals in urban soils.

Supporting Information

Identity of the 50 Tellus soil samples selected for this study (Table S1); primers used in this study (Table S2); relationships between geographical distance and distributions of antibiotic resistance genes (ARGs) and metal(loid) contamination as investigated by distance-decay analysis (Table S3); the ability of different land use categories to explain variation in ARG distribution by permanova test (Table S4); correlations between normalized abundance of clinical class 1 integron-integrase gene and total ARGs (Table S5) (in Supporting Information.docx).

Shannon H index of 50 soil samples showing ARG diversity (Figure S1); stacked bar chart showed the relative percentage of absolute abundance of ARGs (Figure S2); heatmap showing normalized abundance of ARGs in 50 Tellus soils and their land use (Figure S3); heatmap showing absolute abundance of ARGs in 50 Tellus soils and their land use (Figure S4); correlation between absolute abundances of class 1 integron/transposases and ARGs (Figure S5); correlations among metal toxicity index, MGEs and ARGs (Figure S6) (in Supporting Information.docx).

409

410 **Conflict of interests**

411 The authors declare no conflicts of interests.

412

413 **Acknowledgments**

414 The authors thank statistical advisory service provided by University of Copenhagen and Tianwei Xu for
415 the statistical advice on path analysis. Geochemistry data were provided by the Geological Survey of
416 Northern Ireland as part of the Tellus Project, which was funded by the Department of Enterprise, Trade
417 and Investment of Northern Ireland and by the EU ‘Building Sustainable Prosperity’ programme of the
418 Department of Agriculture and Rural Development. This work is supported financially by REMEDIATE
419 (Improved decision-making in contaminated land site investigation and risk assessment) Marie-Curie
420 Innovation Training Network from the European Union's Horizon 2020 Programme (grant number
421 643087).

422

423 **References**

424 An XL, Chen QL, Zhu D, Zhu YG, Gillings MR, Su JQ, 2018. Impact of Wastewater Treatment on the
425 Prevalence of Integrons and the Genetic Diversity of Integron Gene Cassettes. *Appl Environ*
426 *Microbiol*, 84 (9): e02766-17
427 Ashbolt NJ, Amezquita A, Backhaus T, Borriello P, Brandt KK, Collignon P, et al, 2013. Human Health
428 Risk Assessment (HHRA) for environmental development and transfer of antibiotic resistance.
429 *Environ Health Perspect*, 121: 993-1001.

430 Azarbad H, Niklinska M, Laskowski R, van Straalen NM, van Gestel CA, Zhou J, et al, 2015. Microbial
 431 community composition and functions are resilient to metal pollution along two forest soil
 432 gradients. *FEMS Microbiol Ecol*, 91: 1-11.

433 Baker-Austin C, Wright MS, Stepanauskas R, McArthur JV, 2006. Co-selection of antibiotic and metal
 434 resistance. *Trends Microbiol*, 14: 176-82.

435 Berendonk TU, Manaia CM, Merlin C, Fatta-Kassinos D, Cytryn E, Walsh F, et al, 2015. Tackling
 436 antibiotic resistance: the environmental framework. *Nat Rev Microbiol*, 13: 310-7.

437 Berg J, Brandt KK, Al-Soud WA, Holm PE, Hansen LH, Sørensen SJ, et al, 2012. Selection for Cu-
 438 tolerant bacterial communities with altered composition, but unaltered richness, via long-term Cu
 439 exposure. *Appl Environ Microbiol*, 78: 7438-7446.

440 Berg J, Thorsen MK, Holm PE, Jensen J, Nybroe O, Brandt KK, 2010. Cu exposure under field
 441 conditions coselects for antibiotic resistance as determined by a novel cultivation-independent
 442 bacterial community tolerance assay. *Environ Sci Technol*, 44(22):8724-8.

443 Brandt KK, Amézquita A, Backhaus T, Boxall A, Coors A, Heberer T, et al, 2015. Ecotoxicological
 444 assessment of antibiotics: a call for improved consideration of microorganisms. *Environ Int*,
 445 85:189-205.

446 Brandt KK, Frandsen RJ, Holm PE, Nybroe O, 2010. Development of pollution-induced community
 447 tolerance is linked to structural and functional resilience of a soil bacterial community following a
 448 five-year field exposure to copper. *Soil Biol Biochem*, 42(5):748-57.

449 Dantas G, Sommer MOA, Oluwasegun RD, Church GM, 2008. Bacteria subsisting on antibiotics. *Science*
 450 320 (5872):100-103.

451 D'Costa VM, Griffiths E, Wright GD, 2007. Expanding the soil antibiotic resistome: exploring
 452 environmental diversity. *Curr Opin Microbiol* 10 (5):481-489.

453 Deng W, Wang Y, Liu Z, Cheng H, Xue Y, 2014. HemI: a toolkit for illustrating heatmaps. *PLoS One*, 9:
 454 e111988.

455 Eisenhauer N, Bowker MA, Grace JB, Powell JR, 2015. From patterns to causal understanding: Structural
 456 equation modeling (SEM) in soil ecology. *Pedobiologia*, 58: 65-72.
 457 EMEA, 2006. European Medicines Agency, Guideline on the Environmental Risk Assessment of
 458 Medicinal Products for Human Use.
 459 ESRI, 2010. Environmental Systems Resource Institute, ArcMap 10.0, Redlands, California.
 460 European Environment Agency, 2012. Corine Land Cover 2012 seamless vector data.
 461 Finney JM, 1972. Indirect effects in path analysis. *Sociological Methods & Research*, 1: 175-186.
 462 Forsberg KJ, Reyes A, Wang B, Selleck EM, Sommer MO, Dantas G, 2012. The shared antibiotic
 463 resistome of soil bacteria and human pathogens. *Science*, 337: 1107-1111.
 464 Frey SK, Topp E, Khan IUH, Ball BR, Edwards M, Gottschall N, et al, 2015. Quantitative
 465 *Campylobacter spp.*, antibiotic resistance genes, and veterinary antibiotics in surface and ground
 466 water following manure application: Influence of tile drainage control. *Sci Total Environ*
 467 532:138-153.
 468 Gillings M, Boucher Y, Labbate M, Holmes A, Krishnan S, Holley M, et al, 2008. The evolution of class
 469 1 integrons and the rise of antibiotic resistance. *J Bacteriol*, 190: 5095-5100.
 470 Gillings MR, Gaze WH, Pruden A, Smalla K, Tiedje JM, Zhu YG, 2015. Using the class 1 integron-
 471 integrase gene as a proxy for anthropogenic pollution. *ISME J*, 9: 1269-1279.
 472 Gillings MR, 2018. DNA as a Pollutant: the Clinical Class 1 Integron,. *Current Pollution Reports*,
 473 4(1):49-55.
 474 Graham DW, Knapp CW, Christensen BT, McCluskey S, Dolfing J, 2016. Appearance of β -lactam
 475 resistance genes in agricultural soils and clinical isolates over the 20th century. *Sci Rep* 6:21550.
 476 Hammer Ø, Harper D, Ryan P, 2001. Paleontological statistics software: package for education and data
 477 analysis. *Palaeontol Electron*, 4(1):9-178
 478 Hoffman LR, D'Argenio DA, MacCoss MJ, Zhang Z, Jones RA, Miller SI, 2005. Aminoglycoside
 479 antibiotics induce bacterial biofilm formation. *Nature*, 436: 1171-5.

480 Imfeld G, Bringel F, Vuilleumier S, 2011. Bacterial tolerance in contaminated soils: potential of the PICT
 481 approach in microbial ecology. *Tolerance to Environmental Contaminants*, CRC, Boca Raton,
 482 pp341-370.

483 Knapp CW, Callan AC, Aitken B, Shearn R, Koenders A, Hinwood A, 2017. Relationship between
 484 antibiotic resistance genes and metals in residential soil samples from Western Australia. *Environ*
 485 *Sci Pollut Res Int*, 24: 2484-2494.

486 Knapp CW, McCluskey SM, Singh BK, Campbell CD, Hudson G, Graham DW, 2011. Antibiotic
 487 resistance gene abundances correlate with metal and geochemical conditions in archived Scottish
 488 soils. *PLoS One*, 6: e27300.

489 Knapp CW, Dolfing J, Ehlert PA, Graham DW, 2010. Evidence of increasing antibiotic resistance gene
 490 abundances in archived soils since 1940. *Environ Sci Technol*, 44(2):580-587.

491 Knights KV, 2006. A report on the Tellus urban field campaigns of Belfast Metropolitan areas and
 492 Londonderry. Commissioned Report CR/07/006N. British Geological Survey.

493 Larsson DGJ, Andremon A, Bengtsson-Palme J, Brandt KK, de Roda Husman AM, Fagerstedt P, et al,
 494 2018. Critical knowledge gaps and research needs related to the environmental dimensions of
 495 antibiotic resistance. *Environ Int* 117:132-138.

496 Lekfeldt JDS, Magid J, Holm PE, Nybroe O, Brandt KK, 2014. Evaluation of the leucine incorporation
 497 technique for detection of pollution-induced community tolerance to copper in a long-term
 498 agricultural field trial with urban waste fertilizers. *Environ Pollut* 194:78-85.

499 Leonard AFC, Zhang L, Balfour AJ, Garside R, Hawkey PM, Murray AK, et al, 2018. Exposure to and
 500 colonisation by antibiotic-resistant *E. coli* in UK coastal water users: Environmental surveillance,
 501 exposure assessment, and epidemiological study (Beach Bum Survey). *Environ Int* 114:326-333.

502 Li G, Sun GX, Ren Y, Luo XS, Zhu YG, 2018. Urban soil and human health: a review. *Eur J Soil Sci*, 69:
 503 196-215.

504 Looft T, Johnson TA, Allen HK, Bayles DO, Alt DP, Stedtfeld RD, et al, 2012. In-feed antibiotic effects
 505 on the swine intestinal microbiome. *Proc Natl Acad Sci USA*, 109: 1691-6.

506 Luo XS, Yu S, Zhu YG, Li XD, 2012. Trace metal contamination in urban soils of China. *Sci Total*
 507 *Environ*, 421-422: 17-30.

508 Ma L, Li B, Jiang XT, Wang YL, Xia Y, Li AD, Zhang T, 2017. Catalogue of antibiotic resistome and
 509 host-tracking in drinking water deciphered by a large scale survey. *Microbiome* 5 (1):154, 2017.

510 Martínez JL, Coque TM, Baquero F, 2015. What is a resistance gene? Ranking risk in resistomes. *Nat*
 511 *Rev Microbiol*, 13(2):116.

512 Mata M, Baquero F, Perez - Diaz J, 2000. A multidrug efflux transporter in *Listeria monocytogenes*.
 513 *FEMS Microbiol Lett*, 187: 185-188.

514 McIlwaine R, Doherty R, Cox SF, Cave M, 2017. The relationship between historical development and
 515 potentially toxic element concentrations in urban soils. *Environ Pollut*, 220: 1036-1049.

516 McLaughlin MJ, Smolders E, 2001. Background zinc concentrations in soil affect the zinc sensitivity of
 517 soil microbial processes—A rationale for a metalloregion approach to risk assessments. *Environ*
 518 *Toxicol Chem*, 20: 2639-2643.

519 NISRA, 2016. Population Projections for areas within Northern Ireland: 2016-based. Northern Ireland
 520 Statistics and Research Agency.

521 Oksanen J, Kindt R, Legendre P, O'Hara B, Stevens MHH, Oksanen MJ, et al, 2007. The vegan package.
 522 *Community ecology package*, 10: 631-637.

523 Pal C, Asiani K, Arya S, Rensing C, Stekel DJ, Larsson DGJ, et al, 2017. Metal Resistance and Its
 524 Association With Antibiotic Resistance. *Adv Microb Physiol*, 70: 261-313.

525 Pal C, Bengtsson-Palme J, Kristiansson E, Larsson DG, 2015. Co-occurrence of resistance genes to
 526 antibiotics, biocides and metals reveals novel insights into their co-selection potential. *BMC*
 527 *Genomics*, 16: 964.

528 Partridge SR, Tsafnat G, Coiera E, Iredell JR, 2009. Gene cassettes and cassette arrays in mobile
 529 resistance integrons. *FEMS Microbiol Rev*, 33: 757-84.

530 Pruden A, Pei R, Storteboom H, Carlson KH, 2006. Antibiotic resistance genes as emerging contaminants:
 531 studies in northern Colorado. *Environ Sci Technol*, 40: 7445-7450.

532 Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, Fierer N, 2010. Soil
 533 bacterial and fungal communities across a pH gradient in an arable soil. *ISME J*, 4(10):1340.

534 Royle SA, Simms A, Clarke HB, Gillespie R, 2007. Belfast, Part II, 1840-1900. Royal Irish Academy.

535 SCENIHR, 2009. SCoEaNIHR. Assessment of the Antibiotic Resistance Effects of Biocides.

536 Seiler C, Berendonk TU, 2012. Heavy metal driven co-selection of antibiotic resistance in soil and water
 537 bodies impacted by agriculture and aquaculture. *Front Microbiol*, 3: 399.

538 Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al, 2003. Cytoscape: a software
 539 environment for integrated models of biomolecular interaction networks. *Genome Res*, 13: 2498-
 540 2504.

541 Smolders E, Oorts K, Van Sprang P, Schoeters I, Janssen CR, McGrath SP, McLaughlin MJ, 2009.
 542 Toxicity of trace metals in soil as affected by soil type and aging after contamination: using
 543 calibrated bioavailability models to set ecological soil standards. *Environ Toxicol Chem*,
 544 28(8):1633-42.

545 Smyth D, 2007. Methods used in the Tellus geochemical mapping of Northern Ireland. British Geological
 546 Survey open report OR/07/022, <http://nora.nerc.ac.uk/14008/>.

547 Song J, Rensing C, Holm PE, Virta M, Brandt KK, 2017. Comparison of Metals and Tetracycline as
 548 Selective Agents for Development of Tetracycline Resistant Bacterial Communities in
 549 Agricultural Soil. *Environ Sci Technol*, 51: 3040-3047.

550 Stefanowicz AM, Niklińska M, Laskowski R, 2008. Metals affect soil bacterial and fungal functional
 551 diversity differently. *Environ Toxicol Chem*, 27(3):591-8.

552 Tzeneva VA, Salles JF, Naumova N, de Vos WM, Kuikman PJ, Dolfing J, et al, 2009. Effect of soil
 553 sample preservation, compared to the effect of other environmental variables, on bacterial and
 554 eukaryotic diversity. *Res Microbiol*, 160: 89-98.

United Nations, 2014. Department of Economic and Social Affairs Population Division. World Urbanization Prospects: the 2014 Revision. Highlights (ST/ESA/SER.A/352), United Nations.

Ventola CL, 2015. The antibiotic resistance crisis: part 1: causes and threats. *Pharm Ther*, 40: 277.

Wang FH, Qiao M, Su JQ, Chen Z, Zhou X, Zhu YG, 2014. High throughput profiling of antibiotic resistance genes in urban park soils with reclaimed water irrigation. *Environ Sci Technol*, 48: 9079-85.

Welp G, 1999. Inhibitory effects of the total and water-soluble concentrations of nine different metals on the dehydrogenase activity of a loess soil. *Biol Fertil Soils*, 30(1-2):132-9.

Yazdankhah S, Rudi K, Bernhoft A, 2014. Zinc and copper in animal feed - development of resistance and co-resistance to antimicrobial agents in bacteria of animal origin. *Microb Ecol Health Dis*, 25:25862.

Zhao Y, Su JQ, An XL, Huang FY, Rensing C, Brandt KK, et al, 2018. Feed additives shift gut microbiota and enrich antibiotic resistance in swine gut. *Sci Total Environ*, 621: 1224-1232.

Zhu YG, Gillings M, Simonet P, Stekel D, Banwart S, Penuelas J, 2018. Human dissemination of genes and microorganisms in Earth's Critical Zone. *Glob Chang Biol*, 24: 1488-1499.

Zhu YG, Zhao Y, Li B, Huang CL, Zhang SY, Yu S, et al, 2017. Continental-scale pollution of estuaries with antibiotic resistance genes. *Nat Microbiol*, 2: 16270.

Figures Legend

Figure 1 Antibiotic resistance gene (ARG) profile in Belfast urban soils. The map (a) reveals the number of different ARGs detected (dot size scale) and their absolute abundance (copies per gram of soil; color scale) in 50 urban soils from the Belfast metropolitan area. Pie charts depict (b) the percentage of detected ARGs corresponding to different resistance mechanisms and (c) their classification based on the antibiotics they confer resistance to.

Figure 2 Correlation between mobile genetic elements (MGEs) and antibiotic resistance genes (ARGs) across all 50 Belfast urban soils. (a) The normalized abundance of all targeted MGEs (2 integrons and 8 transposons) significantly correlated to the total number of detected ARGs and the normalized abundance of ARGs (copies per 16S rRNA gene) based on Pearson's correlation ($P < 0.001$). (b) Network analysis showing the co-occurrence pattern between individual MGEs and ARGs. A connection represents a strong (Pearson's $r > 0.7$) and significant ($P < 0.05$) correlation. The nodes with different colors represent MGEs and different ARG types. The edges with different colors correspond to positive (grey) and negative (red) correlations between nodes. The size of node is proportional to the number of connections between nodes. The width of edge is proportional to the degree of correlation.

Figure 3 Network analysis showing the co-occurrence pattern between metals (As, Cd, Co, Cr, Cu, Hg, Ni, Pb, and Zn) and antibiotic resistance genes (ARGs) based on Pearson correlation analysis. A connection represents a strong (Pearson's $r > 0.7$) and significant ($P < 0.05$) correlation. The color of each node represents each metal or the type of ARG. The node size is proportional to the number of the connections. The edge width represents the degree of correlation plotted with Pearson's r . All significant correlations in the data set were positive.

Figure 4 Path analysis showing the modelled effect of soil metal contamination (metal toxicity index) and normalized abundance of mobile genetic elements (MGEs) on (a) the number of detected antibiotic resistance genes (ARGs) and the normalized abundance of co-occurring ARGs (copies per 16S rRNA

598 gene). Path arrows correspond to positive (red) and negative (green) effects with significance level of $P <$
599 0.05 (solid line), and $P > 0.05$ (dotted line) with path directions. Numbers adjacent to the path arrows are
600 path coefficients (standardized regression weights), and the arrow width is proportional to the strength of
601 path coefficients. Bar charts show the standardized direct effect, indirect effect and total effect of metal
602 toxicity index and MGEs on (b) number of detected ARGs and (c) normalized abundance of co-occurring
603 ARGs (copies per 16S rRNA gene) derived from path modelling (a).

Table 1 Total concentrations of metals in Belfast urban soils.

Symbol	Name	Average (mg/kg)	Min - Max (mg/kg)	Category
Pb	Lead	354.5	22.9 - 2910	Post-transition metal
Cd	Cadmium	0.8	0.08 - 3.27	Transition metal
Co	Cobalt	22.3	6.1 - 48	Transition metal
Cr	Chromium	64.1	24 - 345	Transition metal
Cu	Copper	160.0	19 - 954	Transition metal
Hg	Mercury	0.4	0.06 - 1.86	Transition metal
Ni	Nickel	82.3	20.1 - 244	Transition metal
Zn	Zinc	458.4	32 - 5550	Transition metal
As	Arsenic	14.1	2.2 - 51.3	Metalloid

*Analyzed with *aqua regia* digestion followed by Inductively Coupled Plasma Mass Spectrometry (ICP-MS) (McIlwaine et al., 2017).

Figure

[Click here to download Figure: Figure 1.pdf](#)

a

No. detected ARGs

Abundance of total ARGs
(copies per gram of soil)

b

c

Figure
Click here to download Figure- Figure 2.pdf

Figure
[Click here to download Figure: Figure 3.pdf](#)

Figure

[Click here to download Figure 4.pdf](#)

Supplementary material for on-line publication only

[Click here to download Supplementary material for on-line publication only: Supporting Information.docx](#)