**QUEEN'S
UNIVERSITY
BELFAST**

Place versus response learning in fish: a comparison between species.

McAroe, C. L., Craig, C. M., & Holland, R. A. (2015). Place versus response learning in fish: a comparison between species. *Animal Cognition*, 1-9. <https://doi.org/10.1007/s10071-015-0922-9>

Published in:
Animal Cognition

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
<http://link.springer.com/article/10.1007%2Fs10071-015-0922-9>

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

1 **Place versus Response Learning in Fish: A Comparison between**
2 **Species**

3 Claire L. McAroe^{1,2} cmcaroe01@qub.ac.uk, +44 (0) 2892664256

4 Cathy M Craig² and Richard A Holland¹

5 ¹School of Biological Sciences, Queen's University Belfast, Medical Biology
6 Centre, 97 Lisburn Road, Belfast, BT9 7 BL, Northern Ireland

7 ²School of Psychology, Queen's University Belfast, University Road, Belfast,
8 BT7 1NN, Northern Ireland

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26 **Abstract**

27 Place learning is thought to be an adaptive and flexible facet of navigation.
28 Due to the flexibility of this learning, it is thought to be more complex than the
29 simpler strategies such as learning a particular route or navigating through the
30 use of cues. Place learning is crucial in a familiar environment as it allows an
31 individual to successfully navigate to the same endpoint, regardless of where
32 in the environment the journey begins. Much of the research to date focusing
33 on different strategies employed for navigation has used human subjects or
34 other mammals such as rodents. In this series of experiments, the spatial
35 memory of four different species of fish (goldfish, killifish, zebrafish and
36 Siamese fighting fish) was analysed using a plus maze set-up. Results
37 suggest that three of the species showed a significant preference for the
38 adoption of a place strategy during this task whereas zebrafish showed no
39 significant preference. Furthermore, zebrafish took significantly longer to learn
40 the task than the other species. Finally, results suggest that zebrafish took the
41 least amount of time (seconds) to complete trials both during training and
42 probe.

43 **Keywords:** Spatial memory, Spatial cognition, Fish, Animal model, Navigation

44 **Introduction**

45 In the animal kingdom, spatial memory is important for many reasons
46 including avoiding predation, prey detection and finding a mate (Wolbers &
47 Hegarty 2010; White & Brown 2014). There are two categories associated
48 with spatial memory and navigation (van Gerven et al. 2012). The first are
49 called “egocentric processes” and involve encoding features of the
50 environment in relation to where the individual is currently located (Shelton &

51 McNamara 2001). These processes involve learning to navigate to a location
52 using landmarks in the environment as beacons or by learning a series of
53 turning responses (O'Keefe & Nadel 1978; van Gerven et al. 2012).
54 Egocentric navigation is relatively simple and therefore allows spatial
55 memories to be formed easily (van Gerven et al. 2012). Such methods do,
56 however, lack flexibility, particularly in response to changes within the
57 environment, meaning the individual will be less likely to reach their desired
58 location from a novel start point (Rodriguez et al. 1994).

59 O'Keefe and Nadel argued that animals are also able to navigate using
60 another set of processes which, although more cognitively demanding, are
61 able to adapt more readily to changes within the environment (O'Keefe &
62 Nadel 1978; Rodriguez et al. 1994). These are called "allocentric processes".
63 Cognitive mapping, first proposed by Tolman, is the best known example of
64 an allocentric process and is described as a complex mental representation of
65 a familiar environment that can be used to influence spatial behaviour and
66 decision-making (Tolman 1948; O'Keefe & Nadel 1978). One of the key
67 features of such an allocentric process is that an individual is able to navigate
68 based on the spatial relationships between multiple landmarks, sensory
69 features and possible routes within a particular environment (Iaria et al. 2009).
70 As the individual moves through the environment, its cognitive map is
71 continuously updated, making for more robust spatial memory that is also
72 more relevant and current (Cheeseman et al. 2014). This means that changes
73 within the environment, such as removal of specific landmarks or attempting
74 to reach the same goal location from a novel start point, will not significantly
75 affect the individual's spatial memory (Wolbers & Hegarty 2010).

Many experiments have been conducted using maze tests to analyse spatial memory in a variety of species, with mammals and birds being the most popular animals to study (Tolman et al. 1946; O'Keefe & Nadel 1979; Clayton & Krebs 1995; Shettleworth & Westwood 2002; Hamilton et al. 2009). Both taxa possess a hippocampal structure in the brain which is thought to be a key area associated with complex spatial memory abilities (see: Pravosudov & Roth II 2013). Some studies have also shown that individuals who navigate through complex environments on a regular basis, or perform complex spatial memory tasks, are likely to have an enlarged hippocampus (Clayton & Krebs 1995; Maguire et al. 2000). However, the lack of a hippocampal structure in other animal classes does not necessarily mean that such species are incapable of flexible navigation.

Fish are the most numerous of all vertebrate species (Casebolt et al. 1998) and many live in complex environments, some of which are subject to instability and therefore require the animals to have powerful spatial memory capabilities (Brown 2014). Rodriguez and colleagues showed that, like mammals and birds, goldfish are capable of navigating using both egocentric and allocentric processes during maze tests (Rodriguez et al. 1994). The same authors have also been able to manipulate navigational strategy by ablating brain regions of fish, directly suggesting that there are specific areas of the fish brain responsible for different navigational processes (Salas et al. 1996; López et al. 2000; Broglio et al. 2010). Despite these suggestions, a recent search on *Web of Knowledge* (January 2015) using “spatial memory” as the search criterion reported 54,109 articles with only 216 citing fish as the study species. Furthermore, many of these articles focus on the large-scale

navigation abilities (i.e. migration) of particular species rather than on which strategies fish use to solve spatial tasks (Dittman & Quinn 1996; Saito & Watanabe 2005) . Finally, there have been limited studies comparing the similarities and differences in spatial memory across species of fish with many experiments focusing on only one study species (e.g. Braithwaite & De Perera 2006; Shapiro & Jensen 2009; Lamb et al. 2012). Such research would highlight that complex spatial memory is an attribute of fish as a taxonomic group rather than just of specific species.

 This study used four different species of fish to assess the length of time each took to learn a spatial task. It also analysed which strategy each species preferred (egocentric or allocentric) for spatial memory. The four species were goldfish, Siamese fighting fish, zebrafish and a type of killifish. Each of these species is teleost (from the class Actinopterygii) with three from the order Cypriniformes and Siamese fighting fish from the order Perciformes (www.fishbase.org). It is important that they are all from the same class, as the teleost brain is thought to be very similar across fish species, and is comparable on some levels to the mammalian brain (Tropepe & Sive 2003). As mentioned previously, the hippocampus is thought to be crucial for allocentric spatial memory performance in mammals, the area associated with similar spatial memory abilities in teleost fish has been identified as the lateral pallium area of the telencephalon, and impairment to this area results in a marked reduction in allocentric spatial learning in these animals (see: Broglio et al. 2003 for a review).

 All but one of these species (killifish) are thought of as domesticated fish (Gordon & Axelrod 1968; Andrews 2002; Spence et al. 2011) and popular

laboratory species in the studies of a variety of both physiological and cognitive experiments (Roitblat et al. 1982; Kishi 2004; Portavella & Vargas 2005). The final species was chosen as it is gaining popularity within similar fields (Herrera & Jagadeeswaran 2004). Goldfish are the most domesticated of all, and therefore there is limited available information about their natural habitat. However as descendants of carp, it is assumed they tend to occupy ponds and other such bodies of water (Andrews 2002). More is known about the natural habitats of the other three species, each of which also prefer still or slow-moving pools of water (Gordon & Axelrod 1968; Genade et al. 2005; Spence et al. 2011). This is important in a comparison, as a previous study has indicated that even within the same species, fish from different environments (fast flowing vs. still) may use different strategies for navigation (Odling-Smee & Braithwaite 2003). Finally, the species of fish range in sociality with zebrafish having an innate tendency to shoal (Spence et al. 2011), through to the killifish and Siamese fighting fish where it is recommended that the males are housed separately.

Materials and Methods

Subjects

Twenty fish from the following four species were used, giving a total of 80 individuals; *Carassius auratus auratus* (common goldfish), *Nothobranchius guentheri* (killifish), *Betta splendens* (Siamese fighting fish), and *Danio rerio* (zebrafish). The mean length of each species (with standard deviation) was 7.63±2.47cm, 3.58±1.08cm, 8.63±1.12cm, and 4.14±0.88cm, respectively. All Siamese fighting fish were male, six of the killifish were male and fourteen female, sex of goldfish and zebrafish was unknown. All animals were

commercially sourced (*goldfish, Siamese fighting fish and zebrafish from Exotic Aquatics, Belfast, N. Ireland; killifish from Maidenhead Aquatics, Newtownabbey, N. Ireland*). All animals were experimentally naïve.

Housing Conditions

All apparatus was commercially sourced (*Maidenhead Aquatics, Newtownabbey, N. Ireland and Exotic Aquatics, Belfast, N. Ireland*). Goldfish were kept in 80cm x 30cm x 40cm tanks with a stocking density of five in each. Temperatures were maintained at an average of 20° C. During experimentation, killifish, Siamese fighting fish and zebrafish were housed individually (for identification purposes) at an average temperature of 25 ° C. Individual housing jars were made of transparent glass and were kept beside each other allowing fish to have visual access to conspecifics. When not completing experiments, all fish were fed commercial flaked food. pH and waste levels in all tanks were monitored regularly using *API Freshwater Master Test Kit* and water changes were carried out on a regular basis. Waste levels were kept within safe ranges (0ppm ammonia & nitrite; <40ppm nitrate). pH range for goldfish was maintained at a range of 7.6 ± 0.2 ; 7.7 ± 0.3 for killifish and zebrafish; and 8.1 ± 0.1 for Siamese fighting fish. All fish were maintained in a 13: 11 h light: dark cycle at all times during the laboratory.

Experimental Design

Apparatus

Fig. 1 Layout of experimental T-maze formed by blocking arm directly opposite start arm with a piece of Perspex. Reverse layout would be used as the T-maze for probe trials

A square tank measuring 63cm x 63 cm x 43 cm was used for all trials. Water temperatures were similar to housing conditions for each species and waste levels were kept within the same safe limits. pH was maintained at a range of 7.8 ± 0.4 (depending on species). A plus maze (similar to that used in Odling-Smee & Braithwaite 2003) was made using pieces of white opaque Perspex, 21cm in length with each arm attached to the inner walls of the tank (see Fig. 1).

Experimental Design

Fish were placed into the arm of the maze designated as the “start arm” (see Fig 1). Both the start arm and the arm opposite were blocked with removable pieces of Perspex. Trials would begin by unblocking the start arm to allow fish access to the remaining two arms of the maze (a T would be formed by keeping the opposite arm blocked; see Fig 1). All trials (training and probe) were timed using a standard stopwatch and recorded using a *Sony HDR-CX190E handycam* video camera. The position of the camera and that of the experimenter was varied during experiments to help eliminate any unwanted cues. Similarly, the source of illumination was not in the vicinity of the experimental tank.

Training Trials

194 Fish were randomly assigned to receive a bloodworm reward in the arm either
195 to the left or to the right of the start arm ($n=10$ for each side within each
196 species). A trial was considered complete when the tail fin of the fish had
197 passed fully into either arm of the maze. If the fish swam to its assigned
198 rewarded arm, it would receive bloodworm immediately and be moved back to
199 the start arm for the next trial. If the fish swam to the unrewarded arm, exit
200 from that arm would be blocked and the fish would receive a two-minute “time
201 out” (no reward given) before being moved back to the start arm for the next
202 trial. Bloodworm was the recommended food reward for each of the four
203 species by the commercial suppliers and has been used for feeding and
204 reward in a variety of experiments involving fish (e.g. Miklósi & Andrew 1999;
205 Saito & Watanabe 2005; Thomas et al. 2008; Pike et al. 2010). Each fish
206 would receive 10 trials per day (one block). Training criterion was a minimum
207 of 8 out of 10 trials correct for three consecutive blocks. For each incorrect
208 block (7 or fewer trials correct), the training was reset to Block One. The
209 water, along with any substrate, in the tank was disturbed between each trial
210 and the tank filtered for a minimum of 20 minutes between each fish to help
211 reduce the risk of olfactory cues. Potential cues within the maze were
212 controlled where possible, e.g. by mirroring the layout so that the external
213 filter tube and the heater (both turned off during experiments) were at
214 opposing corners of the tank and a tube was placed at the opposite wall of the
215 tank from the output tube of the filter. Outside the maze, there was a wall at
216 the end of the left hand arm while there was no wall at the end of the right
217 arm. Potential global cues (i.e. those external to the maze) were not controlled
218 in order to allow fish to avail of cues external to the tank. Such cues consisted

of features on the surrounding walls (such as paper record sheets) and housing tanks.

Probe Trial

When a fish had reached training criteria, it was immediately transferred to the previously blocked arm, which would now become the start arm; the training start arm would now be blocked to form a “T”. The probe trial would begin and end in the same manner as in training trials. During the probe trial, all fish were rewarded regardless of choice of arm. If a fish swam to the same location as before, a “place-strategy” was recorded; if they chose to swim using the same turning-direction as they were trained, a “response-strategy” was recorded. Following the probe trial, each fish was returned to its housing tank and experimentation for that individual animal was complete.

Statistical Analyses

Data were analysed with the SPSS statistical package (20.0 version) and Microsoft Excel 2010. Generalized linear models were used to assess the effect of species on number of blocks to reach training criteria (Poisson loglinear Model) and to assess the effect of species on time taken to complete the training and probe trials (linear scale response). Pairwise comparisons (Fisher’s LSD method) were performed to analyse differences between species following each generalized linear model. Paired-sample t-tests were used to assess the difference in time taken to complete the first day versus last day of training within each species. Individual binomial tests and Bayesian inferences were carried out to assess whether the number of fish within each species adopting a place versus response strategy differed from chance.

Ethical note

No invasive procedures were performed and no animals were harmed. Fish numbers were the minimum required for sufficient data collection and analyses. Strict procedures were followed in accordance with the 'Guidelines for the treatment of animals in behavioural research and teaching' (2012). Complete water changes were avoided as these can be harmful and stressful to animals (see above for more information regarding water changes). Furthermore, fish were always transferred in water via containers. Laboratory conditions were inspected by the Veterinary Services Division of the DHSSPS, Northern Ireland who deemed no licence was required for this series of experiments.

Results

Acquisition Time

Acquisition time was based on the number of blocks it took each fish to reach training criteria.

Fig. 2 Boxplot showing task acquisition time for each species. Rectangular boxes display 25th & 75th quartiles and the median. Whiskers display 90th percentile of the data with outliers outside this

range marked with an X on the plot. The dotted line displays the minimum number of blocks to reach training criteria within each species (3 blocks in all species). On the graphical display, asterisks are used to show significant differences between species (a single asterisk (*) indicates $P \leq .05$ and double asterisks (**) indicate $P < .001$)

There was a significant main effect of species on task acquisition time: $Wald \chi^2(df = 3) = 25.59; P < 0.001$. Further analyses suggest that zebrafish required significantly more blocks to learn the task than the other three species (goldfish, $P = .003$; killifish, $P = .02$; Siamese fighting fish $P < .001$) and that Siamese fighting fish also took significantly fewer blocks than both goldfish ($P = .05$) and killifish ($P = .013$). There was no significant difference between goldfish and killifish ($P = .57$) (see Fig. 2).

Time Taken during Training Trials

Mean Time Taken (All Trials)

Fig. 3 Bar chart showing the mean time taken in seconds by each species to complete the training trials. Error bars indicate the standard error of the mean time taken for individual fish within each species

There was a significant main effect of species on the time taken (in seconds) to complete individual training trials (both correct and incorrect): $Wald \chi^2(df=3) = 303.52; P < .001$. Posthoc analyses showed that zebrafish took significantly less time to complete trials than the other three species ($P < .001$ in all cases). Furthermore, goldfish took significantly less time during training

trials than both killifish and Siamese fighting fish (both, $P < .001$), and killifish took less time than Siamese fighting fish ($P = .006$) (see Fig. 3).

Difference in Time Taken on First and Last Day of Training

Fig. 4 Bar chart showing mean time taken in seconds by each species to complete trials on the first day of training versus on the last day of training. Error bars indicate the standard error of the mean time taken for individual fish within each species

Goldfish ($P < .001$), Siamese fighting fish ($P = .001$) and zebrafish ($P < .001$)

each took significant more time (seconds) to complete trials on the first day of training compared to the last day of training. However, there was no significant difference in the time taken by killifish on the first and last day of training ($P = .883$). (See Fig. 4).

Navigational Strategy

Fig. 5 Bar chart showing the number of fish in each species that adopted either a place or a response strategy during probe trials. An asterisk (*) indicates that there was a significant difference in the number of fish that adopted a place strategy over a response strategy ($P < 0.5$) ($n = 20$ in all species)

Fig. 5 shows the number of individual fish in each species that adopted either a place or response strategy during the probe trial.

Goldfish, killifish and Siamese fighting fish showed a significant preference for adopting a place strategy (goldfish and Killifish, $P = 0.041$; Siamese fighting fish, $P = 0.012$) but zebrafish showed no significant preference ($P = 0.507$).

Bayesian inferences however showed no significant support for choosing the null hypothesis (performance at chance level) over alternative hypothesis (of place learning) for zebrafish as probability estimates of the null hypothesis lie within the estimates of the alternatives with probabilities of between 0.6 and 0.8 (see Table 1).

Alternative Hypothesis	Evidence Ratio for Null
0.5 (Null)	1.00
0.6	0.67
0.7	1.05
0.8	5.42

Table 1 Evidence ratio for null hypothesis against various alternative hypotheses for the proportion of fish adopting a place strategy in zebrafish species estimated using Bayesian inference. *A posteriori* probabilities suggest that there is not strong evidence for choosing the null hypothesis over some of the alternative hypotheses. Strong evidence is when the ratio of a null *a posteriori* probability to an alternative *a posteriori* probability is less than 1/3 or greater than 3. For example, if an *a priori* alternative hypothesis of 0.7 is examined, a value of 1.05 is obtained indicating no strong evidence for choosing this alternative hypothesis over the null, or vice versa.

Time Taken on Probe Trial

Fig. 6 Bar chart showing the mean time taken in seconds by each species during the probe trial. Error bars indicate the standard error of the mean time taken for individual fish within each species

There was a significant main effect of species on time taken during the probe trial: $Wald \chi^2 (df=3) = 30.92; P < .001$. Posthoc analyses showed Siamese fighting fish took significantly more time (in seconds) than the other three species to complete the probe trial ($P < .001$ in all cases). There were no other significant differences (goldfish vs. killifish; $p = .648$, goldfish vs. zebrafish; $P = .250$, killifish vs. zebrafish; $P = .108$). (See Fig. 6)

Discussion

The aims of this study were to assess the length of time it took four species of fish to learn a spatial memory task and to see what navigational strategy each species preferentially employed in doing so. Results show that zebrafish took significantly more blocks to reach training criterion than the other three species (goldfish, killifish and Siamese fighting fish). Goldfish and killifish also took significantly more blocks than Siamese fighting fish to learn the training task. Such findings correspond with previous studies (Roitblat et al. 1982; Rodriguez et al. 1994) which found that both goldfish and Siamese fighting fish are capable of learning a variety of spatial memory tasks. Although little is known about the spatial memory of *Nothobranchius guentheri*, the findings in this study do suggest that these fish are also able to learn a spatial memory task and can do so at a faster rate than a more commonly used study species (zebrafish).

Other results from this study suggest that the mean time taken (in seconds) to complete an individual trial by the Siamese fighting fish (the largest of the four species and the quickest to learn) was significantly longer in both training ($p < .01$) and probe trials ($p < .001$) than the other three species. Similarly, the mean time to complete a training trial by zebrafish (the slowest to learn) was significantly shorter ($p < .001$) than all the other fish used in these experiments. These findings suggest that taking time to explore the environment may improve the learning process, by allowing the animal to learn from both correct and incorrect trials. In other contexts it has been proposed that animals that are faster explorers and “proactive” might be less accurate whereas animals that are slower explorers and “reactive” might be more accurate in their decision making (Sih et al. 2004). Previous studies

357 have also suggested that fish which are slower to explore their environment
358 are more flexible in their behaviour and are quicker to adapt to changes (see:
359 Magnhagen 2012 for a review). However it should be noted that these studies
360 have focused on individual differences within a species and have not
361 compared differences in behavioural flexibility and plasticity across species.
362 We are cautious in our interpretation as another reason for this difference may
363 be simply that Siamese fighting fish are from a different order than the other
364 three species which may have an effect on their adaptation of spatial
365 navigation (Perciformes and Cypriniformes respectively). Further studies
366 manipulating time allowed to complete a trial and its influence on learning
367 speed would also be interesting. Likewise adopting similar analyses used in
368 this study to assess the effects of time taken to make a decision could be
369 applied to explore behavioural outcomes in a variety of species.

370 Although our findings, which show that three species (goldfish,
371 Siamese fighting fish and zebrafish) took significantly less time to complete
372 trials on the last day of training versus the first day of training, would be
373 expected during a learning task, it is unclear why there was no significant
374 difference in the killifish species. This may have been because the killifish
375 were disadvantaged by being the smallest of the four species. However
376 further research is required to fully understand these differences.

377 The findings that three of the four species showed a preference for the
378 adoption of a place strategy suggest that fish, like mammals and birds, are
379 capable of complex spatial memory and that the lack of a hippocampal
380 structure is not necessarily detrimental to the navigational abilities of fish. This
381 has also been argued by others who suggest that the telencephalon in fish, in

particular the lateral pallium, is homologous to the hippocampus of mammals and birds (Saito & Watanabe 2006; Spence et al. 2011). Zebrafish did not show a significant preference for the adoption of either a response or a place strategy despite their popularity in previous behavioural and memory studies (Williams et al. 2002; Miklosi & Andrew 2006; Sison & Gerlai 2010; Miller & Gerlai 2012b). However, results from Bayesian inference would suggest an increase in sample size may be required in order to establish whether these fish truly have a preference for either a place or a response strategy.

It is possible that the fish were using salient geometric cues external to the maze in order to navigate using what seemed to be a place strategy; this has been suggested to be the case in previous studies using rats as subjects (Ritchie 1947; Cheng 1986). The possible reason why some animals, particularly the zebrafish, did not show a preference for a place strategy may be that they were using geometrical information internal to the maze therefore causing a rotational error on the probe trial and eliciting what seemed to be a response strategy (Ritchie 1947; Cheng 1986). Further experiments could be completed in order to assess the effects of geometrical information on strategy choice.

Another possible reason for the zebrafish species performing differently could be based on zebrafish being a shoaling species which means they are more likely to navigate through their environment in groups (Wright & Krause 2006; Miller & Gerlai 2012a; Butail et al. 2013). Again, further research is needed to assess whether zebrafish show a difference in navigation or a preference for either strategy when completing the task in shoals.

Three of the four species were housed individually during experimentation (killifish, Siamese fighting fish and zebrafish). This isolation could have caused some level of stress, particularly for the shoaling zebrafish species which is highly social. Housing jars were, however, kept beside each other allowing each fish visual contact to conspecifics. Previous studies have suggested that this is satisfactory to reduce isolation stress in such animals and may also induce shoaling movements (e.g. Engeszer et al. 2004; Sison & Gerlai 2010; Karnik & Gerlai 2012). In the experimental protocol, fish would complete the trials individually which may have led to isolation stress, particularly in a novel environment. Additionally, it is possible that body size may have had an impact on the results, perhaps placing a disadvantage on the smaller species. There were, however, no observed signs of stress during experimentation with individual fish willingly eating the bloodworm reward on all correct trials, a further indicator that stress levels were minimal during experiments (Carr 2002).

As each fish would receive ten trials per block and multiple fish would be tested on each experimental day, it could be argued that olfactory cues may have been important in the learning of the task and the performance of the probe trial. These effects were controlled for where possible by disturbing the water and any substrate in the tank between each correct trial and by filtering the tank for a minimum of 20 minutes between each fish. Furthermore, a previous study using bloodworm reward and goldfish as a study species suggests that olfaction is not often used by fish during similar maze tests (Saito & Watanabe 2005). Finally, as place learning was not

employed by all fish, it is unlikely that olfactory cues played any significant role on the performance of fish during these experiments.

A previous study suggested that the number of training days has an effect on whether an animal uses a place or response strategy for navigation (Packard & McGaugh 1996). This study found that rats were more likely to choose a response over a place strategy when they had more training in the experimental set-up. This could explain why the zebrafish, who showed the greatest variation in the number of blocks of trials taken to reach training criteria, showed no significant preference for the adoption of either a place or a response strategy. However, it should be noted that the difference in the set of experiments presented here is that not all fish received the same amount of training. This differed from the previous study (Packard & McGaugh, 1996) where there was a fixed training schedule before each probe trial. The suggestion that more training could have an impact on navigational strategy also requires further investigation.

It is possible that the fish could have used the wall at the end of one of the goal arms (or the lack of wall at the end of the other) as a navigation beacon. If this was the case, then one would expect the animals to use a simpler US-CS link to learn the task rather than a more complex spatial memory strategy (Karnik & Gerlai 2012). As global cues external to the maze were not controlled for or analysed during experimentation, it is not possible to conclude what cues the fish used in order to display place learning. Indeed, it has been argued that it is difficult to exclude simpler strategies from any experiment that investigates complex spatial memory processes (Bennett 1996) However, It is surprising that the zebrafish did not show a preference

for the training location if a simple, salient feature was indicating this and could explain performance of the other species. Thus, whether it was beaconing to a cue external to the maze or a set of more complex allocentric features that explain the performance of the other three species, it is clear that the zebrafish were not able to consistently use the same strategy as the other species to learn the location of the food reward. Whilst our experimental setup cannot distinguish between beaconing to a cue and an allocentric strategy, it is worth noting that other experiments, using the same experimental setup (plus maze), have indicated that when the lateral pallium area of the telencephalon is ablated, fish are no longer able to learn place, but retain an ability to use a cue based strategy (Rodriguez et al. 1994; Salas et al. 1996; López et al. 2000; Broglio et al. 2010). Experiments using a controlled salient cue at the rewarded arm would indicate whether zebrafish can learn this task and use it from a novel start point and thus provide further insight into the strategies used in the current experiment.

The findings in this study suggest that the spatial memory of fish may be comparable to other animals as three species out of four demonstrated a preference for using a strategy that is often associated with more complex and flexible allocentric navigation when completing the plus maze task . Zebrafish did not demonstrate place learning and showed no significant preference for either a place or response strategy. Further experiments using shoals of zebrafish should be completed to assess whether navigation is facilitated by collective decision-making in this species (Couzin et al. 2005). Regardless of strategy choice, these findings suggest that all four species are able to learn a spatial memory task, reinforcing previous findings that fish are

a useful animal model for the study of cognitive behavioural tasks (Rodriguez et al. 1994; Braithwaite & De Perera 2006; White & Brown 2014).

Acknowledgements

The authors would like to acknowledge the following individuals for assistance with lab. set-up and fish husbandry: Georgina Glaser, Sinéad Smith, Chris Preshaw, Gillian Riddell and other members of the technical staff at the School of Biological Sciences, QUB. We would also like to thank the funding bodies of this research: The Department of Employment and Learning, Northern Ireland and Queen's University Belfast. Finally, we would like to thank Ken Cheng and Hannah White for advice on statistics.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Andrews, C., 2002. *An Interpet Guide to Fancy Goldfish*, Surrey: Interpet Publishing.
- Bennett, A.T., 1996. Do animals have cognitive maps? *J Exp Biol*, 199, pp.219–224.
- Braithwaite, V.A. & De Perera, T.B., 2006. Short-range orientation in fish: how fish map space. *Mar Freshw Behav Phy*, 39(1), pp.37–47.
- Broglio, C. et al., 2010. Selective involvement of the goldfish lateral pallium in spatial memory. *Behav Brain Res*, 210(2), pp.191–201.
- Broglio, C., Rodriguez, F. & Salas, C., 2003. Spatial cognition and its neural basis in teleost fishes. *Fish Fish*, 4, pp.247–255.
- Brown, C., 2014. Fish intelligence, sentience and ethics. *Anim Cogn*, 18, pp.1–17.
- Butail, S., Bartolini, T. & Porfiri, M., 2013. Collective response of zebrafish shoals to a free-swimming robotic fish. *PloS One*, 8(10), p.e76123.
- Carr, J.A., 2002. Stress, neuropeptides, and feeding behavior: a comparative perspective. *Integr Comp Bio*, 42(3), pp.582–590.

- 509 Casebolt, D., Speare, D. & Horney, B., 1998. Care and use of fish as
510 laboratory animals: Current state of knowledge. *Lab Anim Sci*, 48(2),
511 pp.124–135.
- 512 Cheeseman, J.F. et al., 2014. Way-finding in displaced clock-shifted bees
513 proves bees use a cognitive map. *P Natl A Sci USA*, 111(24), pp.8949–
514 54.
- 515 Cheng, K., 1986. A purely geometric module in the rat's spatial
516 representation. *Cognition*, 23, pp. 149–178.
- 517 Clayton, N.S. & Krebs, J.R., 1995. Memory in food-storing birds: from
518 behaviour to brain. *Curr Opin Neurobiol*, 5(2), pp.149–154.
- 519 Couzin, I.D. et al., 2005. Effective leadership and decision-making in animal
520 groups on the move. *Nature*, 433(7025), pp.513–6.
- 521 Dittman, A.H. & Quinn, T.P., 1996. Homing in pacific salmon: mechanisms
522 and ecological basis. *J Exp Biol*, 199, pp.83–91.
- 523 Engeszer, R.E., Ryan, M.J. & Parichy, D.M., 2004. Learned social preference
524 in zebrafish. *Curr Biol*, 14(10), pp.881–4.
- 525 Genade, T. et al., 2005. Annual fishes of the genus *Nothobranchius* as a
526 model system for aging research. *Aging Cell*, 4(5), pp.223–233.
- 527 Van Gerven, D.J.H. et al., 2012. Direct measurement of spontaneous strategy
528 selection in a virtual Morris water maze shows females choose an
529 allocentric strategy at least as often as males do. *Behav Neurosci*,
530 126(3), pp.465–78.
- 531 Gordon, M. & Axelrod, H., 1968. *Siamese Fighting Fish*, New Jersey: T.F.H
532 Publications, Inc.
- 533 Guidelines for the Use of Animals (2012) Guidelines for the treatment of
534 animals in behavioural research and teaching. *Anim Behav*, 83, 301-309
- 535 Hamilton, D.A. et al., 2009. Control of rodent and human spatial navigation by
536 room and apparatus cues. *Behav Process*, 81(2), pp.154–69.
- 537 Herrera, M. & Jagadeeswaran, P., 2004. Annual fish as a genetic model for
538 aging. *J Gerontol A Biol Sci Med Sci*, 59(2), pp.101–7.
- 539 Iaria, G. et al., 2009. Age differences in the formation and use of cognitive
540 maps. *Behav Brain Res*, 196(2), pp.187–91.
- 541 Karnik, I. & Gerlai, R., 2012. Can zebrafish learn spatial tasks? An empirical
542 analysis of place and single CS-US associative learning. *Behav Brain*
543 *Res*, 233(2), pp.415–421.

- 544 Kishi, S., 2004. Functional Aging and Gradual Senescence in Zebrafish. Ann
545 NY Acad Sci, 1019, pp.521–526.
- 546 Lamb, E.A., Echevarria, D.J. & Jouandot, D.J., 2012. The utility of the T-maze
547 in assessing learning, memory, and models of neurological disorders in
548 the zebrafish. Behaviour, 149(10-12), pp.1081–1097.
- 549 López, J.C. et al., 2000. Dissociation of place and cue learning by
550 telencephalic ablation in goldfish. Behav Neurosci, 114(4), pp.687–699.
- 551 Magnhagen, C., 2012. Personalities in a crowd: What shapes the behaviour of
552 Eurasian perch and other shoaling fishes? Curr Zoo, 58(1), pp.35–44.
- 553 Maguire, E. et al., 2000. Navigation-related structural change in the
554 hippocampi of taxi drivers. PNAS, 97(8), pp.4398–4403.
- 555 Miklósi, a & Andrew, R.J., 1999. Right eye use associated with decision to
556 bite in zebrafish. Behav Brain Res, 105(2), pp.199–205.
- 557 Miklosi, A. & Andrew, R., 2006. The zebrafish as a model for behavioral
558 studies. Zebrafish, 3(2), pp.227–234.
- 559 Miller, N. & Gerlai, R., 2012a. From schooling to shoaling: patterns of
560 collective motion in zebrafish (*Danio rerio*). PloS one, 7(11), p.e48865.
- 561 Miller, N. & Gerlai, R., 2012b. Automated tracking of zebrafish shoals and the
562 analysis of shoaling behaviour. In: Kalueff, A.V. & Steward, A.M.
563 Zebrafish Protocols for Neurobehavioral Research, pp. 217–230. New
564 Jersey: Humana Press.
- 565 O'Keefe, J. & Nadel, L., 1978. The Hippocampus as a Cognitive Map, Oxford:
566 Clarendon Press.
- 567 Odling-Smee, L. & Braithwaite, V.A., 2003. The influence of habitat stability on
568 landmark use during spatial learning in the three-spined stickleback.
569 Anim Behav, 65(4), pp.701–707.
- 570 Packard, M.G. & McGaugh, J.L., 1996. Inactivation of hippocampus or
571 caudate nucleus with lidocaine differentially affects expression of place
572 and response learning. Neurobiol Learn Mem, 65(1), pp.65–72.
- 573 Pike, T.W. et al., 2010. Learning by proportional observation in a species of
574 fish. Behav Ecol, 21(3), pp.570–575.
- 575 Portavella, M. & Vargas, J., 2005. Emotional and spatial learning in goldfish is
576 dependent on different telencephalic pallial systems. Eur J Neurosci, 21,
577 pp.2800–2806.

- 578 Pravosudov, V. V. & Roth II, T.C., 2013. Cognitive ecology of food hoarding:
579 The evolution of spatial memory and the Hippocampus. *Annu Rev Ecol*
580 *Evol S*, 44(1), pp.173–193.
- 581 Ritchie, B.F., 1947. Studies in spatial learning III: Two paths to the same
582 location and two paths to two different locations. *J Expl Psychol*, 37(1),
583 pp. 25–38.
- 584 Rodriguez, F. et al., 1994. Performance of goldfish trained in allocentric and
585 egocentric maze procedures suggests the presence of a cognitive
586 mapping system in fishes. *Anim Learn Behav*, 22(4), pp.409–420.
- 587 Roitblat, H., Tham, W. & Golub, L., 1982. Performance of *Betta splendens* in
588 a radial arm maze. *Anim Learn Behav*, 10(1), pp.108–114.
- 589 Saito, K. & Watanabe, S., 2006. Deficits in acquisition of spatial learning after
590 dorsomedial telencephalon lesions in goldfish. *Behav*
591 *Brain Res*, 172(2), pp.187–94.
- 592 Saito, K. & Watanabe, S., 2005. Experimental analysis of spatial learning in
593 goldfish. *Psychol Rec*, 55, pp.647–662.
- 594 Salas, C. et al., 1996. Spatial Learning and Memory Deficits After
595 Telencephalic Ablation in Goldfish Trained in Place and Turn Maze
596 Procedures. *Behav Neurosci*, 110(5), pp.965–980.
- 597 Shapiro, M.S. & Jensen, A.L., 2009. Parameters of rewards on choice
598 behavior in Siamese fighting fish (*Betta splendens*). *Behav Process*,
599 82(1), pp.30–8.
- 600 Shelton, A.L. & McNamara, T.P., 2001. Systems of spatial reference in human
601 memory. *Cognitive Psychol*, 43(4), pp.274–310.
- 602 Shettleworth, S.J. & Westwood, R.P., 2002. Divided attention, memory, and
603 spatial discrimination in food-storing and nonstoring birds, black-capped
604 chickadees (*Poecile atricapilla*) and dark-eyed Juncos (*Junco hyemalis*).
605 *J Exp Psychol Anim B*, 28(3), pp.227–241.
- 606 Sih, A., Bell, A. & Johnson, J.C., 2004. Behavioral syndromes: An ecological
607 and evolutionary overview. *Trends Ecol Evol*, 19(7), pp.372–378.
- 608 Sison, M. & Gerlai, R., 2010. Associative learning in zebrafish (*Danio rerio*) in
609 the plus maze. *Behav Brain Res*, 207(1), pp.99–104.
- 610 Spence, R., Magurran, A.E. & Smith, C., 2011. Spatial cognition in zebrafish:
611 the role of strain and rearing environment. *Anim Cogn*, 14(4), pp.607–12.
- 612 Thomas, P.O.R. et al., 2008. Does defection during predator inspection affect
613 social structure in wild shoals of guppies? *Anim Behav*, 75(1), pp.43–53.

- 614 Tolman, E., Ritchie, B. & Kalish, D., 1946. Studies in spatial learning. II. Place
615 learning versus response learning. J Exp Psychol, 36(3), pp.221–229.
- 616 Tolman, E.C., 1948. Cognitive maps in rats and men (1). Psychol Rev, 55(4),
617 pp.189–208.
- 618 Tropepe, V. & HL Sive, 2003. Can zebrafish be used as a model to study the
619 neurodevelopmental causes of autism? Genes Brain Behav, 2, pp.268–
620 281.
- 621 White, G.E. & Brown, C., 2014. A comparison of spatial learning and memory
622 capabilities in intertidal gobies. Behav Ecol Sociobiol, 68(9), pp.1393–
623 1401.
- 624 Williams, F.E., White, D. & Messer, W.S., 2002. A simple spatial alternation
625 task for assessing memory function in zebrafish. Behav Process, 58(3),
626 pp.125–132.
- 627 Wolbers, T. & Hegarty, M., 2010. What determines our navigational abilities?
628 Trends Cogn Sci, 14(3), pp.138–46.
- 629 Wright, D. & Krause, J., 2006. Repeated measures of shoaling tendency in
630 zebrafish (*Danio rerio*) and other small teleost fishes. Nat Protoc, 1(4),
631 pp.1828–31.

632 **Figure Captions**

- 633 **Fig. 1** Layout of experimental T-maze formed by blocking arm directly opposite start arm with a
634 piece of Perspex. Reverse layout would be used as the T-maze for probe trials
- 635 **Fig. 2** Boxplot showing task acquisition time for each species. Rectangular boxes display 25th &
636 75th quartiles and the median. Whiskers display 90th percentile of the data with outliers outside
637 this range marked with an X on the plot. The dotted line displays the minimum number of blocks
638 to reach training criteria within each species (3 blocks in all species). On the graphical display,
639 asterisks are used to show significant differences between species (a single asterisk (*)
640 indicates $P \leq .05$ and double asterisks (**) indicate $P < .001$)
- 641 **Fig. 3** Bar chart showing the mean time taken in seconds by each species to complete the
642 training trials. Error bars indicate the standard error of the mean time taken for individual fish
643 within each species
- 644 **Fig. 4** Bar chart showing mean time taken in seconds by each species to complete trials on the
645 first day of training versus on the last day of training. Error bars indicate the standard error of
646 the mean time taken for individual fish within each species
- 647 **Fig. 5** Bar chart showing the number of fish in each species that adopted either a place or a
648 response strategy during probe trials. An asterisk (*) indicates that there was a significant
649 difference in the number of fish that adopted a place strategy over a response strategy ($P < 0.05$)
650 ($n = 20$ in all species)
- 651 **Fig. 6** Bar chart showing the mean time taken in seconds by each species during the probe trial.
652 Error bars indicate the standard error of the mean time taken for individual fish within each
653 species
- 654 **Table 2** Evidence ratio for null hypothesis against various alternative hypotheses for the
655 proportion of fish adopting a place strategy in zebrafish species estimated using Bayesian
656 inference. A *posteriori* probabilities suggest that there is not strong evidence for choosing the
657 null hypothesis over some of the alternative hypotheses. Strong evidence is when the ratio of a

658 null *a posteriori* probability to an alternative *a posteriori* probability is less than 1/3 or greater
659 than 3. For example, if an *a priori* alternative hypothesis of 0.7 is examined, a value of 1.05 is
660 obtained indicating no strong evidence for choosing this alternative hypothesis over the null, or
661 vice versa.

662