**QUEEN'S
UNIVERSITY
BELFAST**

Airborne microbial transport limitation to isolated Antarctic soil habitats

Archer, S. D.J., Lee, K. C., Caruso, T., Maki, T., Lee, C. K., Cary, S. C., Cowan, D. A., Maestre, F. T., & Pointing, S. B. (2019). Airborne microbial transport limitation to isolated Antarctic soil habitats. *Nature Microbiology*.

Published in:
Nature Microbiology

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2019 The Authors. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Airborne microbial transport limitation to isolated Antarctic soil habitats

Stephen D.J. Archer^{1,2}, Kevin C. Lee², Tancredi Caruso³, Teruya Maki⁴, Charles K. Lee⁵, S. Craig Cary⁵, Don A. Cowan⁶, Fernando T. Maestre⁷, Stephen B. Pointing^{1,8,9}

¹ *Yale-NUS College, National University of Singapore, Singapore 138527*

² *Institute for Applied Ecology New Zealand, Auckland University of Technology, Auckland 1142, New Zealand*

³ *School of Biological Sciences and Institute for Global Food Security, Queen's University Belfast, Belfast BT9 7BL, Northern Ireland, UK*

⁴ *Department of Chemical Engineering, Kanazawa University, Kanazawa 920-1192, Japan*

⁵ *International Centre for Terrestrial Antarctic Research, University of Waikato, Hamilton 3240, New Zealand*

⁶ *Centre for Microbial Ecology and Genomics, University of Pretoria, Pretoria 0002, South Africa*

⁷ *Departamento de Biología y Geología, Física y Química Inorgánica, Escuela Superior de Ciencias Experimentales y Tecnología, Universidad Rey Juan Carlos, C/ Tulipán s/n, 28933 Móstoles, Spain*

⁸ *Department of Biological Sciences, National University of Singapore, Singapore 117558*

⁹ *Institute of Nature and Environmental Technology, Kanazawa University, Kanazawa 920-1192, Japan*

Dispersal is a critical yet poorly understood factor underlying macroecological patterns in microbial communities ¹. Airborne microbial transport is assumed to occupy a central role in determining dispersal outcomes ^{2,3} and extra-range dispersal has important implications for predicting ecosystem resilience and response to environmental change ⁴. One of the most pertinent biomes in this regard is Antarctica given its geographic isolation and vulnerability to climate change and human disturbance ⁵. Here we report microbial diversity in near-ground and high-altitude air above the largest ice-free Antarctic habitat as well as that of underlying soil microbial communities. We found that persistent airborne inputs were unable to fully explain local soil community assembly. Comparison with airborne microbial diversity from non-polar sources suggests that strong selection occurs during atmospheric transport resulting in regionally restricted airborne inputs. Highly specialized soil communities occurred where fungi displayed greater isolation than bacteria from non-polar sources. Overall, microbial communities from this isolated Antarctic ecosystem displayed limited connectivity to the non-polar microbial pool and alternative sources of recruitment are necessary to fully explain extant soil diversity. Our findings provide critical insights to forecast the potential outcomes for microbial communities of climate change-mediated shifts in air circulation to Antarctica.

Airborne microbial transport on fine particulate matter has typically been regarded as neutral and ubiquitous due to the small size and survivability of cells ⁶⁻⁸. Diverse microbial signatures have been recovered from airborne particulates after transport across inter-continental distances ^{3,8,9}. Large scale patterns in microbial diversity are therefore often viewed as developing largely due to deterministic niche-driven processes ^{10,11}. The assumption that airborne transport is a neutral and

ubiquitous process has, however, been challenged ¹² and we postulate that deterministic dispersal drivers may help explain extant diversity patterns in isolated microbial communities. The McMurdo Dry Valleys comprise the largest ice-free habitat in Antarctica and one of the most isolated soil habitats on Earth. They are devoid of vascular plants and dominated by highly specialised soil microbial communities ¹³ that display adaptations to the extreme environmental conditions ¹⁴. Some taxa, notably cyanobacteria, have been shown to display phylogenetic endemism in Antarctica at the level of rRNA gene-defined diversity ¹⁵⁻¹⁷. Also, lichenised fungi displayed patterns in diversity that suggest that they radiated from local refugia rather than from exogenous sources outside Antarctica ¹⁸. A global theoretical model for atmospheric aerosols estimated that the rate of airborne microbial exchange to Antarctica may be extremely low, with 90% of aerosols expected to be of local origin ¹⁹. In contrast, empirical studies have claimed circum-polar distribution for some soil Cyanobacteria, chlorophyte algae, and Fungi ^{20,21}. Antarctica therefore presents a paradox in microbial biogeography with regard to microbial dispersal. Evidence for airborne microorganisms in Antarctica is scarce and limited to a few preliminary biodiversity estimates indicating cosmopolitan and local taxa are recoverable from the near-surface atmosphere ²²⁻²⁴.

Here we use state-of-the art sampling, sequencing and statistical approaches to study the diversity of airborne and soil microbial communities in a typical Antarctic Dry Valley. We targeted Bacteria and Fungi because these domains are the most abundant microorganisms in the McMurdo Dry Valleys ¹³. We tested the null hypotheses that air and soil microbial communities are a random sample of the regional and global microbial pools. For doing so, we acquired massive bulk-air samples and estimated microbial diversity in near-ground air and underlying soil for low and high elevation

1 sites, as well as polar air above the boundary layer for surface interactions and non-
2 polar sources.

3 The incoming air mass to our study site in the McMurdo Dry Valleys largely
4 transited above the Antarctic Plateau during the maximum predicted residence time for
5 bacteria and fungi in air (15 days) ¹⁹, whilst the most distant air mass had a non-polar
6 origin above the coastal shelf of New Zealand (Fig. 1a). Transport was exclusively from
7 the Polar Plateau and across the Trans-Antarctic Mountains during the average
8 residence time for microorganisms in air (Fig. 1a). We thus envisage that severe
9 selection pressure should occur during airborne transit in an air mass with freezing
10 temperatures and high UV exposure at mean altitudes of 2769m (3 day transit) and
11 3034m (15 day transit).

12 In general terms, we found that alpha diversity metrics for air and elevated high-
13 altitude soils were distinct from those observed in valley soils, with taxa richness being
14 more variable in valley soils than in air and elevated soils (Fig. 1b,c). These results
15 highlight the more heterogenous nature of valley soils as a habitat compared with bulk
16 phase air or elevated altitude soils, where conditions are generally unfavourable to
17 colonisation. The elevated altitude mineral soil sites may therefore be representative of
18 near-term airborne deposition to this system. Bacterial taxa richness was similar among
19 all samples although slightly higher in air and elevated soil (Fig. 1b). Conversely, the
20 richness of fungal taxa was highest in valley soils, and lower in air and elevated soils
21 (Fig. 1c). Soils also displayed greater evenness and lower richness values than air
22 samples and this tentatively indicated that airborne fungi are under strong selective
23 pressure. Ordination analyses of weighted UniFrac distances for bacteria and fungi
24 supported these trends in alpha diversity (Fig. 1b,c). Valley soil bacterial communities

1 separated clearly from air and elevated soil communities (Fig. 1b). A similar although
2 less pronounced pattern was observed for fungi (Fig. 1c).

3 Taxonomic assignment of bacteria and fungi revealed further complexity.
4 Airborne phylum-level bacterial diversity was dominated by Proteobacteria,
5 Bacteroidetes and Firmicutes (Fig. 1d, Online Supplementary Information Fig. S2). Phyla
6 with high relative abundance comprised spore-formers and taxa with known UV and/or
7 desiccation tolerant traits viewed as advantageous during atmospheric transport and
8 survival in Antarctic soil. The airborne bacterial samples supported relatively high
9 levels of taxa associated with marine influence ⁸ suggesting recruitment during transit
10 over the Southern Ocean (Online Supplementary Information, Fig. S2). Terrestrial
11 bacteria were also transported and may have benefited from islands acting as stepping
12 stones for dispersal ⁸. Soil communities supported greater abundance of Actinobacteria
13 and other taxa typical from arid soils ¹⁰. Near-ground air supported 3-5 fold more
14 habitat-specific taxa than high-altitude air; samples from the former habitat were most
15 similar to their underlying soil communities. Valley soils supported 56.4% soil-specific
16 taxa compared with only 15.8% in elevated high-altitude soils. Valley soils shared very
17 few taxa with the total air sample pool (4.5%) whilst different air habitats (valley,
18 elevated and high-altitude) shared approximately half the taxa encountered in each
19 habitat.

20 The most abundant fungal taxa in air were basidiomycetous yeasts (Fig. 1e,
21 Online Supplementary Information Fig. S2) whereas soils were dominated by
22 unclassified fungi also including yeasts. The yeasts are thought to be well-adapted to
23 growth in Antarctic soil habitats ²⁵. Ascomycetes were also commonly encountered but
24 Chytrids occurred only in valley soil and air. Valley soils supported 48.3% habitat-
25 specific taxa while elevated soils and high-altitude air showed the lowest number of

1 habitat-specific taxa (4.9-8.7%). The different air habitats shared approximately half the
2 taxa encountered in each habitat. Fungi are well-adapted to conditions anticipated
3 during atmospheric transport due to the production of resistant spores and UV-
4 protective compounds. Local recruitment may, however, be limited to asexual states
5 since teleomorph fruiting structures are not known from Antarctic fungi.

6 We achieved near-asymptote in diversity estimation for all samples (Online
7 Supplementary Information, Supplementary Methods). Thus we further interrogated
8 the phylogenetic diversity of air and soil by generating distribution heatmaps by habitat
9 for the 1,000 most abundant taxa. This analysis captured 91% of total bacterial and
10 96% of total fungal diversity in the libraries (Fig. 2a,b). We also incorporated diversity
11 data from air originating at the nearest non-polar land mass into this analysis. A striking
12 pattern emerged where soil bacterial and fungal assemblages in the McMurdo Dry
13 Valleys were only partially recruited from local air taxa. This pattern cannot be further
14 explained by airborne recruitment from exogenously sourced aerosols. These findings
15 support the notion of a system operating in stark contrast with the long-held
16 assumption that microbial dispersal is ubiquitous and deterministic niche processes are
17 the primary driver of community assembly in terrestrial surfaces ^{1,12}.

18 We therefore further interrogated this association using Ecological Network
19 Analysis (Fig. 2c). Overall non-polar air displayed least connectivity to all other
20 Antarctic habitats as previously observed by the weak associations and greater Bray-
21 Curtis distances between them (Fig. 2c). Bacterial communities clustered by habitat
22 type (Fig. 2c). This pattern is likely to be indicative of selection pressures due to local
23 environmental filtering, which could combine a mixture of biotic and abiotic factors.
24 Conversely, fungal communities associated by geographic distance and were thus more
25 likely to be influenced by dispersal limitation. No significant distance-decay

relationships were observed for airborne or soil communities between valley and elevated sites (air bacteria $R^2 = 0.009$, soil bacteria $R^2 = 0.006$, air fungi $R^2 = 0.016$, soil fungi $R^2 = 0.024$). These results indicate that dispersal within the Dry Valleys may be limited and likely reflects the associated steep environmental gradients present in this region. These findings provide empirical support to prevailing theoretical models of emission and transport for biological particles in the atmosphere that predicts relatively low exchange between Antarctic and non-polar air as well as reduced residence time in air for fungi compared to bacteria due to allometric considerations ¹⁹.

We conducted additional analyses (Nestedness Analysis and Net Relatedness Index analysis) to reveal the extent to which taxonomic and phylogenetic structuring reflected the likelihood of exogenous recruitment (Fig. 3). Null models applied to nestedness metrics showed that bacterial and fungal communities overall were significantly anti-nested (NODF <30) (Fig. 3a). This general result implies that passive sampling from the regional pool alone was not sufficient to explain the structure of local Antarctic communities. Both Bacteria and Fungi, however, were significantly nested for taxonomic composition (NODFc) under the hypothesis that nestedness can be maximised when ordering sites from the most connected, to the least connected, to a global species pool. This suggests that species poor assemblages of the least connected sites are a proper subset of richer, more connected sites. Fungi were markedly more nested (NODFc = 62) than Bacteria (NODFc = 18) (Fig. 3a), suggesting a potential major role of dispersal limitation for this group.

The Net Relatedness Index (NRI) added phylogenetic support to the findings of our Network Analysis and Nestedness Analysis, by further demonstrating that local Antarctic communities were not a random sample of the overall species pool. Antarctic bacteria displayed greater and highly significant phylogenetic clustering compared to

1 non-polar samples, which were almost randomly structured and in some case over-
2 dispersed (Fig. 3b). Although the pattern itself does not prove any specific process, the
3 results clearly indicate that Antarctic bacterial communities both in soil and air must
4 have been selected non-randomly, which is consistent with both the taxonomic and
5 phylogenetic observations for our air and soil communities. This result is congruent
6 with observations of biodiversity for other soils in the Dry Valleys region ²⁶. The Fungi
7 were always significantly clustered but bacterial communities were always much more
8 clustered than fungi at any given airborne or soil location. The fungal data should,
9 however, be interpreted with care given current drawbacks with phylogenetic
10 reconstruction based on ITS and despite our efforts to correct for them. Nonetheless
11 even a cautious interpretation of the data suggests a limited extent of input from fungal
12 taxa not present in local reservoirs. This interpretation concurs with our other lines of
13 evidence presented here and with studies on fungal dispersal from other biomes ²⁷.

14 Contrary to the view that “everything is everywhere” in terms of airborne
15 microbial transport, our data indicates that the aerosphere is a strongly selective
16 habitat that limits dispersal, although the extent may vary between taxonomic groups
17 and spatial scales. We conclude that inter-continental microbial connectivity to the
18 McMurdo Dry Valleys of East Antarctica is limited, and this supports the hypothesis that
19 the Hadley Cell circulation acts as a dispersal barrier to the poles even during the
20 austral summer when the Polar Vortex is annually at its weakest. The Antarctic
21 continent supports other smaller ice-free soil regions and whilst we are unable to
22 directly extrapolate our data to these, it is reasonable to expect similar patterns given
23 what is known of air circulation to the continent; that is, that other Antarctic ice-free
24 areas may also be somewhat decoupled from global microbial reservoirs. An exception
25 may be the peninsula in West Antarctica due to its proximity to the South American

continent. Comparison of our soil biodiversity estimates with those for other Dry Valleys locations suggest there is a common core diversity throughout the Antarctic Dry Valleys²⁶. Hence, we expect our data to be broadly applicable to this region. The low level of airborne immigration from exogenous sources may represent an inherently low flux for Antarctica and this may help to explain the unique microbial composition of Antarctic soils compared to others globally²⁸.

We have presented multiple lines of evidence to refute the null hypothesis that local air and soil microorganisms are a random sample of phylogenetic diversity in the regional/global pools. Sources of recruitment other than persistent airborne transport are therefore necessary to fully explain the extant Antarctic soil microbial diversity patterns observed. One potential explanation is stochastic storm events where particulate matter supporting biological propagules is thought to be transported on local scales within the Dry Valleys²⁹, although we did not encounter any such events during our sampling expedition. A further source may be local dispersal from geothermal refugia as they are important reservoirs for radiative dispersal of animal, plant and lichen taxa⁵. The periodicity from which dispersal from such reservoirs occurs is, however, unknown. An additional reservoir may be the moisture-sufficient soil around lakes where microbial mats are known to persist over inter-annual periods²⁶. Local refugia may be important in facilitating resilience at the landscape scale where severe local extinction pressure occurs due to stochasticity and steep environmental gradients for abiotic variables.

Challenges remain in deciphering the relationship of diversity patterns to biomass and ecosystem function³⁰, but the revelation that airborne connectivity is largely localised rather than being an inter-continental scale process emphasises the conservation value of the McMurdo Dry Valleys as a unique ecosystem. This is

1 particularly pertinent in light of a predicted increase in stochasticity for atmospheric air
2 circulation as a result of climate change, which may led to an increased flux of foreign
3 and invasive taxa into Antarctic ecosystems. Such an increased flux acting in concert
4 with warmer temperatures could profoundly alter the unique biota inhabiting the
5 Antarctic Dry Valleys, one of the of the last pristine ecosystems on Earth.

6 7 **Methods**

8 **Sample Recovery**

9 We employed a high-volume liquid impinger apparatus (Coriolis μ , Bertin
10 Technologies, France) and developed a novel collection protocol optimised for low-
11 temperature environments. This involved collection of samples directly into *RNAlater*
12 nucleic acid preservative solution (Invitrogen, Carlsbad, CA). Evaporation was
13 compensated for using a peristaltic pump set to between 0.5 and 1.3mL per minute with
14 a mixture of phosphate-buffered saline (PBS) and 20% v/v *RNAlater*. For each sampling
15 interval a random collection cone was also assembled into the machine but not
16 activated, and these were used as the negative controls. All collection cones were
17 soaked in 1.5% sodium hypochlorite (NaClO) then washed with 70% ethanol and three
18 washes of Milli-Q H₂O before being filled with filtered *RNAlater*. All sampling equipment
19 was disassembled between locations and cleaned with NaClO, ethanol and Milli-Q H₂O.
20 Samples in *RNAlater* were stored at 4°C during transit from Antarctica and until
21 processed.

22 Air mass at near-surface (1.5m above ground) and corresponding soil sampling
23 (top 10mm surface soil after removing pebbles and rocks) was conducted from 11th –
24 23rd January 2017 from eight locations throughout the Wright Valley floor (77.518633
25 S, 161.768783 E) and high elevation locations at the valley ridge and a high elevation

inter-valley feature known as Bull Pass (7.47085 S, 161.77345 E). Air mass above the boundary layer for surface influence was also recovered by mounting the apparatus in a helicopter with an external sampling port (flightpath: 2,000m A.M.S.L., 77.440836 S, 162.657553 E to 77.524583 S, 161.690917 E). Overall, we retrieved 33 massive bulk air samples plus underlying soil samples from 18 locations (high altitude Antarctic air samples and non-polar air did not have accompanying soil samples). Massive bulk-phase air volumes (72,000L per sample) were collected for each discreet sampling location. Biotic data were retrieved for 30 air samples with a total sampled volume of 2,160,000L, and from all soil samples.

At each station and time interval air flow rates of 300L/min were employed for 4hrs into 15ml RNA*later*. This approach overcomes limitations from earlier studies where low volume pumps have necessitated long sampling durations with uncertain microbial survival and recovery and impaction techniques that are known to bias against certain phyla ^{31,32}. Non-polar air samples were collected from New Zealand's North Island (36.916153 S, 174.645760 E) during the same austral summer season and using the same method. We selected this place because our HYSPLIT back trajectory analysis (see below) indicated that this was the nearest non-polar land mass from which air mass arriving at the Dry Valleys location was derived. These samples were used to make broad diversity comparisons with possible exogenous sources although we acknowledge that additional variability is likely among non-polar aerosols given inherent uncertainties over their trajectory to the Antarctic. Each location was sampled at three discreet time intervals during the austral summer field season.

Sampling was conducted within local weather parameters as follows: Relative humidity 20-58%, Temperature -3.3-6.9°C, wind speed 0.6-9.5m/S, wind direction E-ESE (Kestrel 3500 Weather Meter, Nielson-Kellerman Co, Minnesota, USA); Total near-

ground air particulate matter 2939-6558 $\mu\text{g}/\text{m}^3$ PM_{2.5-10} (Aerotrak, TSI Incorporated, Minnesota, USA). Back-trajectories of air mass arriving at each sampling interval were generated using the National Oceanic and Atmospheric Administration (NOAA) HYSPLIT-WEB model (<https://ready.arl.noaa.gov/HYSPLIT.php>). Separate calculations were made at 3d and 15d because they represent the average minimum and maximum residence times for microorganisms in air transported to the McMurdo Dry Valleys ¹⁹. HYSPLIT back trajectories were calculated using the GDAS database and the model vertical velocity option. Three day back trajectories travelled between 598 and 2581 km with an average of 1350 km at an average altitude of 2769 m and a maximum altitude of 5174 m above mean sea level (AMSL) Fifteen day back trajectories travelled between 4673 and 11216 km with an average of 6886 km at an average altitude of 3034 m and a maximum altitude of 8211 m AMSL.

Environmental DNA sequencing and bioinformatics

Airborne microbial samples from the RNA*later* preservation solution were filtered onto a 25mm 0.2 μm polycarbonate filter and stored frozen until processed. Total DNA was directly extracted using a CTAB protocol ³³. DNA was extracted from three 0.75g $\pm$ 0.025 of soil using the same CTAB protocol. DNA yield for these ultra-low biomass samples was quantified using the Qubit 2.0 Fluorometer (Invitrogen) in the range 1.06-8.44ng. Samples were then stored at -20°C until processed. We used DNA yield as an indirect estimate for biomass. We were unable to successfully apply direct cell/fluorescent particle counting due to the extremely low cell numbers in Antarctic air, although we acknowledge the validity of this approach in studies of higher biomass aerial habitats ⁸. We did not apply real-time quantitative PCR as a further indirect estimate of biomass since the approach has significant limitations that preclude

1 meaningful estimates of biomass from environmental samples with domain-specific
2 PCR primers ^{34,35}.

3 Illumina MiSeq libraries were prepared as per manufacturer's protocol
4 (Metagenomic Sequencing Library Preparation Part # 15044223 Rev. B; Illumina, San
5 Diego, CA, USA) and as previously described with PhiX positive controls ³⁶. We targeted
6 Bacteria and Fungi since these domains are the most abundant microorganisms in the
7 McMurdo Dry Valleys ²⁶. PCR was conducted with primer sets targeting the V3-V4
8 regions of bacterial and archaeal 16S rRNA gene: PCR1 forward (5' TCGTCGGCAG
9 CGTCAGATGT GTATAAGAGA CAGCCTACGG GNGGCWGCAG 3') and PCR1 reverse (5'
10 GTCTCGTGGG CTCGGAGATG TGTATAAGAG ACAGGACTAC HVGGGTATCT AATCC 3') and
11 the internal transcribed spacer region of fungal 18S and 5.8S rRNA genes: ITS1 forward
12 (5' CTTGGTCATTTAGAGGAAGTAA 3') and ITS2 reverse (5' GCTGCGTTCTTCATCGATGC
13 3'). These primers for Bacteria and Fungi are widely accepted to capture the broadest
14 estimates of diversity ^{10,37,38} and were used according to recommended workflows for
15 the Earth Microbiome Project (<http://www.earthmicrobiome.org>). Total sequence
16 library sizes were 3,994,561 for bacteria and 2,437,256 for fungi before filtering and
17 total counts in the processed dataset were 1,333,553 bacterial and 2,220,883 fungal
18 sequences. A total of 3,636 bacterial and 5,525 fungal taxa were identified from these.
19 All sequence data generated by this study has been submitted to the NCBI Sequence
20 Read Archive under BioProject PRJEB27416 with accession numbers ERS3573837 to
21 ERS3573946.

22 Sequencing data for 16S rRNA gene amplicons was processed based on the
23 DADA2 v1.8 ³⁹ pipeline. Primers sequences were removed using cutadapt ⁴⁰ to remove
24 forward (CCTACGGGNGGCWGCAG) and reverse (GACTACHVGGGTATCTAATCC). The
25 reads were uniformly trimmed to 280 bp (forward) and 250 bp (reverse) and then

1 filtered by removing reads exceeding maximum expected error of 2 for forward reads
2 and 5 for reverse reads or reads containing ambiguity N symbol. The reads were used to
3 train the error model and then dereplicated to acquire unique sequences, which were
4 used to infer sequence variants with the trained error model. The forward and reverse
5 reads were merged and chimeric sequences were removed. For bacteria we used
6 amplicon sequence variants (ASV) to assign operational taxonomic units (OTUs) since
7 this has been shown as the most robust method currently available for bacterial 16S
8 rRNA gene-defined taxa identification ⁴¹. OTUs were given taxonomic assignment using
9 DADA2 with SILVA nr v132 database ⁴² to provide species level assignment based on
10 exact match between ASVs and known reference sequences. For fungal ITS1 amplicon
11 data, the sequences were processed using USEARCH v9.0.2132 ⁴³. The forward and
12 reverse paired-end sequences were merged and filtered to remove reads >1 maximum
13 expected error per sequence. Additionally, anomalous sequences (<200 or >500 bp in
14 length or exceeding 20 homopolymers) were also removed. After dereplication and
15 removal of singletons, the reads were clustered at 97% identity threshold to obtain
16 representative sequences as OTUs ³⁷. Unfiltered reads were mapped onto these OTUs to
17 produce an abundance table of the occurrence of these OTUs within the communities.
18 The representative sequences were given taxonomic assignment using USEARCH
19 SINTAX classifier and RDP Warcup training set v2 (rdp_its_v2) ⁴⁴.

20 The resulting OTUs were then processed as previously described ⁴⁵. The R
21 packages phyloseq ⁴⁶, DESeq2 ⁴⁷ and ggplot2 ⁴⁸ were used for downstream analysis and
22 visualisation including ordination and alpha/beta diversity calculations. Despite
23 inherent bias due to underlying differences in substrate biomass influencing species
24 richness estimates with any cross-habitat biogeographic analysis ⁴⁹, we are confident
25 that comparable yet inherently low biomass in all our air and ultra-oligotrophic mineral

soil samples minimised such influence and this was reflected in our diversity estimates. An exception was that fungi in non-polar air were markedly more taxon-rich than in Antarctic samples. Therefore we used guild analysis (FUNGuild⁵⁰) to establish the predominantly phyllosphere origin of fungi in non-polar samples which are absent in Antarctica as well as the latitudinal gradient in fungal diversity⁵¹ (lack of database depth for FUNGuild limited its value to identifying ecological guilds for non-polar fungi only). For heatmap visualisations the 1,000 most abundant OTUs in each data set were selected, and these captured 91% bacterial and 96% fungal sequences. All other analysis used the entire sequence library data. Our heatmap analysis therefore had high confidence since the unsampled 'tail' comprised only extremely rare sequence variants at very low/singleton abundance.

Statistical Treatments

We used multiple statistical approaches to test the null hypothesis that local soil and air sample communities were a random sample of the of the regional pool. Rejection of the hypothesis (i.e., non-random patterns) yields observational evidence for the alternative hypothesis that local communities are a non-random selection from the regional pool. Specifically, we expected local soil communities to be selected against the extreme conditions found at soils in the McMurdo Dry Valleys soil and thus to display clustering. At the same time, we expected air samples to be less structured or even random and thus better reflect the regional pool, although some structuring due to local influences from atmospheric stressors such as low temperatures and UV exposure were also be expected.

We employed approaches that utilised both taxonomic identity and phylogenetic structure of the communities. Ecological Network Analysis is a commonly employed tool to infer biotic interactions within and between communities by visualising links

1 between species nodes. Potential relationships pertinent to our system include
2 connectivity, clustering and nestedness which are informative to interpreting the
3 biogeographic patterns of species occurrence ⁵². We performed Ecological Network
4 Analysis on our samples using the R package phyloseq ⁴⁶ with maximum Bray-Curtis
5 distance of 0.2 for Bacteria and 0.4 for Fungi to establish connection between nodes
6 (representing communities). The nodes were positioned using the Fruchterman-
7 Reingold method ⁵³.

8 Nestedness is a widespread biogeographical pattern that emerges when species
9 composition of small assemblages is a nested subset of larger regional assemblages. We
10 quantified this pattern using the metrics NODF ⁵⁴ and its compositional (NODFc) and
11 incidence (NODFr) version. This metric is currently considered one of the most effective
12 and statistically robust, especially in relation to the null models that are used to test
13 whether observed metrics are smaller or larger than expected by chance ^{52,55}. The
14 metric ranges from 0, that is no nestedness or perfect antinestedness, to 100 (perfect
15 nestedness). In a perfectly nested assemblage, species poor communities are just a
16 subset of species richer communities, or less frequent species always occur in subsets of
17 sites where most widespread species also occur, or a combination of both. We applied
18 the NODF metrics to the matrix of the investigated sites both for Bacteria and Fungi. We
19 simulated null models using a random swap algorithm (R project, vegan package,
20 function “oecosimu” ⁵⁶) with fixed row and column sums. This combination to create a
21 random matrix is the most conservative in terms of Type I and II errors and is
22 particularly recommended when species co-occurrence is critical to the tested
23 hypothesis ⁵². We calculated Standardised Effect Sizes and tested for significance of
24 effects with 999 permutations and at $P < 0.05$. We calculated NODF and null models by a
25 specific order of sites in the matrix that reflected our main null hypothesis. This was

1 that local communities are just a passive sampling from regional species pools.
2 Specifically, we used two complementary orderings of the sites to test our hypothesis.
3 First, we created a connectivity gradient, which assumed New Zealand is the site more
4 connected to the global species pool while soils in the Antarctic Dry Valleys are the least
5 connected (the exact order was: non-polar>high-altitude air>elevated air>valley
6 air>elevated soil>valley soil). In the second test of the hypothesis local, more isolated
7 soils were assumed to select more than connected soil and air habitats. In this case, sites
8 were ordered as follows: Valley Soil>Elevated Soil>Valley Air>Elevated Air>high-
9 altitude air>non-polar air.

10 Community phylogenetic metrics were calculated using the R ⁵⁷ packages picante
11 ⁵⁸, ape ⁵⁹, phylobase ⁶⁰, adephylo ⁶¹, and phytools ⁶². Phylogenetic trees for community
12 phylogenetic structure analysis were constructed for all OTUs with FastTree v2.1.9 on
13 multiple alignment of sequences produced by MUSCLE v3.8.31. For Bacteria an
14 approximately Maximum-Likelihood approach was used whilst for Fungi an alignment-
15 free distance approach with Neighbour-Joining method was employed in order generate
16 a ITS-based phylogenetic tree for community metrics ⁶³. The distance approach was
17 used for Fungi as the hypervariability of ITS1 loci hinders multiple sequence alignment
18 required for most phylogenetic analyses. To validate this approach, we compared our
19 tree topology with the most recent whole genome phylogenies for the Fungi ^{64,65}. This
20 approach was robust at higher taxonomic levels (there is no consensus for fungal
21 phylogenies using multiple loci or whole genomes below Order rank) and has been used
22 successfully with other eukaryotic taxa as a workflow for Net Relatedness analysis ⁶⁶.
23 Although we acknowledge limitations to this approach, the advantages of using ITS loci
24 for taxonomic identification vastly outweighed its shortcomings, and we also
25 triangulated data from this test with additional analytical approaches, so overall our

inclusion of this test is justified and interpreted conservatively. Mean phylogenetic distance (MPD) ⁶⁷ was calculated to measure phylogenetic distance between ASVs and OTUs in each sample. A null model algorithm based on independent swap (999 randomisation) was used to test the extent of phylogenetically clustering (positive values) or overdispersion (negative values) ⁶⁸. Results for NRI were expressed as effects size, $(MPD - MPD_{null}) / SD(MPD_{null})$. Distance Decay of phylogenetic versus geographic distance for bacteria and fungi was estimated using the R package Vegan ⁵⁶.

Acknowledgements

Field and logistical support was provided by Antarctica New Zealand and the United States Antarctic Program. The authors thank Craig Cary (University of Waikato) for facilitating access to the McMurdo Dry Valleys. The research was funded by a grant from the New Zealand Ministry of Business, Innovation & Employment (UOWX1401) and the Yale-NUS College Start-Up Fund. F.T.M. is supported by the European Research Council (BIODESERT project, ERC Grant agreement n° 647038).

Author contributions

S.D.J.A. and S.B.P. conceived the study; S.D.J.A and C.K.L. conducted fieldwork; T.M. developed and validated the helicopter sampling method; S.D.J.A. performed laboratory experiments; S.D.J.A., K.C.L., T.C., and S.B.P. performed data analysis and interpretation; D.A.C., F.T.M. and S.B.P. critically assessed and interpreted the findings; S.B.P. wrote the manuscript.

Materials & Correspondence

Correspondence and requests for materials should be addressed to S.B.P. Email:

stephen.pointing@yale-nus.edu.sg.

Data Availability

All sequence data generated by this study has been submitted to the EMBL European Nucleotide Archive (ENL) under BioProject PRJEB27416 with accession numbers ERS3573837 to ERS3573946. PLEASE CHECK PUBLIC AVAILABILITY

References

1. Hanson, C. A., Fuhrman, J. A., Horner-Devine, M. C. & Martiny, J. B. H. Beyond biogeographic patterns: processes shaping the microbial landscape. *Nat. Rev. Microbiol.* **10**, 497–506 (2012).
2. Burrows, S. M., Elbert, W., Lawrence, M. G. & Pöschl, U. Bacteria in the global atmosphere – Part 1: Review and synthesis of literature data for different ecosystems. *Atmos. Chem. Phys.* **9**, 9263–9280 (2009).
3. Kellogg, C. A. & Griffin, D. W. Aerobiology and the global transport of desert dust. *Trends Ecol. Evol.* **21**, 638–644 (2006).
4. Wilson, J. R. U., Dormontt, E. E., Prentis, P. J., Lowe, A. J. & Richardson, D. M. Something in the way you move: dispersal pathways affect invasion success. *Trends Ecol. Evol.* **24**, 136–44 (2009).
5. Chown, S. L. *et al.* The changing form of Antarctic biodiversity. *Nature* **522**, 431 (2015).
6. de Wit, R. & Bouvier, T. ‘Everything is everywhere, but, the environment selects’;

what did Baas Becking and Beijerinck really say? *Environ. Microbiol.* **8**, 755–758 (2006).

7. Finlay, B. J. & Clarke, K. J. Ubiquitous dispersal of microbial species. *Nature* **400**, 828–828 (1999).

8. Mayol, E. *et al.* Long-range transport of airborne microbes over the global tropical and subtropical ocean. *Nat. Commun.* **8**, 201 (2017).

9. Favet, J. *et al.* Microbial hitchhikers on intercontinental dust: catching a lift in Chad. *ISME J.* **7**, 850–67 (2013).

10. Delgado-Baquerizo, M. *et al.* A global atlas of the dominant bacteria found in soil. *Science* (80-.). **359**, 320–325 (2018).

11. Wang, J. *et al.* Phylogenetic beta diversity in bacterial assemblages across ecosystems: deterministic versus stochastic processes. *ISME J.* **7**, 1310–1321 (2013).

12. Lowe, W. H. & McPeck, M. A. Is dispersal neutral? *Trends Ecol. Evol.* **29**, 444–50 (2014).

13. Pointing, S. B. *et al.* Highly specialized microbial diversity in hyper-arid polar desert. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 19964–19969 (2009).

14. Chan, Y., Van Nostrand, J. D., Zhou, J., Pointing, S. B. & Farrell, R. L. Functional ecology of an Antarctic Dry Valley. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 8990–5 (2013).

15. Bahl, J. *et al.* Ancient origins determine global biogeography of hot and cold desert cyanobacteria. *Nat. Commun.* **2**, 163 (2011).

16. Jungblut, A., Lovejoy, C. & Vincent, W. Global distribution of cyanobacterial ecotypes in the cold biosphere. *ISME J.* **4**, 191–202 (2010).

17. Vyverman, W. *et al.* Evidence for widespread endemism among Antarctic micro-

- organisms. *Polar Sci.* **4**, 103–113 (2010).
18. Fraser, C. I., Terauds, A., Smellie, J., Convey, P. & Chown, S. L. Geothermal activity helps life survive glacial cycles. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 5634–9 (2014).
19. Burrows, S. M. *et al.* Bacteria in the global atmosphere – Part 2: Modeling of emissions and transport between different ecosystems. *Atmos. Chem. Phys.* **9**, 9281–9297 (2009).
20. Kleinteich, J. *et al.* Pole-to-Pole Connections: Similarities between Arctic and Antarctic Microbiomes and Their Vulnerability to Environmental Change. *Front. Ecol. Evol.* **5**, 137 (2017).
21. Cox, F., Newsham, K. K., Bol, R., Dungait, J. A. J. & Robinson, C. H. Not poles apart: Antarctic soil fungal communities show similarities to those of the distant Arctic. *Ecol. Lett.* **19**, 528–536 (2016).
22. Kobayashi, F. *et al.* Atmospheric bioaerosols originating from Adélie penguins (*Pygoscelis adeliae*): Ecological observations of airborne bacteria at Hukuro Cove, Langhovde, Antarctica. *Polar Sci.* **10**, 71–78 (2016).
23. Pearce, D. A., Hughes, K. A., Lachlan-Cope, T., Harangozo, S. A. & Jones, A. E. Biodiversity of air-borne microorganisms at Halley station, Antarctica. *Extremophiles* **14**, 145–159 (2010).
24. Bottos, E. M. E. M., Woo, A. C. A. C., Zawar-Reza, P., Pointing, S. B. S. B. & Cary, S. C. Airborne Bacterial Populations Above Desert Soils of the McMurdo Dry Valleys, Antarctica. *Microb. Ecol.* **67**, 120–128 (2013).
25. Rao, S. *et al.* Low-diversity fungal assemblage in an Antarctic Dry Valleys soil. *Polar Biol.* **35**, 567–574 (2011).
26. Cowan, D. A., Makhalanyane, T. P., Dennis, P. G. & Hopkins, D. W. Microbial ecology and biogeochemistry of continental Antarctic soils. *Front. Microbiol.* **5**, 154

(2014).

27. Brown, S. P. & Jumpponen, A. Phylogenetic diversity analyses reveal disparity between fungal and bacterial communities during microbial primary succession. *Soil Biol. Biochem.* **89**, 52–60 (2015).
28. Fierer, N. *et al.* Cross-biome metagenomic analyses of soil microbial communities and their functional attributes. *Proc. Natl. Acad. Sci. USA* **109**, 21390–21395 (2012).
29. Atkins, C. B. & Dunbar, G. B. Aeolian sediment flux from sea ice into Southern McMurdo Sound, Antarctica. *Glob. Planet. Change* **69**, 133–141 (2009).
30. Pointing, S. B., Fierer, N., Smith, G. J. D., Steinberg, P. D. & Wiedmann, M. Quantifying human impact on Earth’s microbiome. *Nat. Microbiol.* **1**, 16145 (2016).
31. Dybwad, M., Skogan, G. & Blatny, J. M. Comparative testing and evaluation of nine different air samplers: End-to-end sampling efficiencies as specific performance measurements for bioaerosol applications. *Aerosol Sci. Technol.* **48**, 282–295 (2014).
32. Šantl-Temkiv, T. *et al.* High-Flow-Rate Impinger for the Study of Concentration, Viability, Metabolic Activity, and Ice-Nucleation Activity of Airborne Bacteria. *Environ. Sci. Technol.* **51**, 11224–11234 (2017).
33. Archer, S. D. J., McDonald, I. R., Herbold, C. W., Lee, C. K. & Cary, C. S. Benthic microbial communities of coastal terrestrial and ice shelf Antarctic meltwater ponds. *Frontiers in Microbiology* **6**, 485 (2015).
34. Fierer, N., Jackson, J. A., Vilgalys, R. & Jackson, R. B. Assessment of Soil Microbial Community Structure by Use of Taxon-Specific Quantitative PCR Assays. *Appl. Environ. Microbiol.* **71**, 4117–4120 (2005).

35. Props, R. *et al.* Absolute quantification of microbial taxon abundances. *ISME J.* **11**, 584–587 (2017).
36. Warren-Rhodes, K. *et al.* Subsurface microbial habitats in an extreme desert Mars-analogue environment. *bioRxiv* 269605 (2018). doi:10.1101/269605
37. Schoch, C. L. *et al.* Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 6241–6 (2012).
38. Thompson, L. R. *et al.* A communal catalogue reveals Earth’s multiscale microbial diversity. *Nature* **551**, 457–463 (2017).
39. Callahan, B. J., Mcmurdie, P. J., Rosen, M. J., Han, A. W. & A, A. J. DADA2: High resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583 (2016).
40. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10 (2011).
41. Callahan, B. J., McMurdie, P. J. & Holmes, S. P. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J.* **11**, 2639–2643 (2017).
42. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res.* **41**, 590–596 (2013).
43. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461 (2010).
44. Deshpande, V. *et al.* Fungal identification using a Bayesian classifier and the Warcup training set of internal transcribed spacer sequences. *Mycologia* **108**, 1–5 (2016).
45. Maki, T. *et al.* Variations in the structure of airborne bacterial communities in

- 1 Tsogt-Ovoo of Gobi desert area during dust events. *Air Qual. Atmos. Heal.* **10**, 249–
2 260 (2017).
- 3 46. McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive
4 Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217 (2013).
- 5 47. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
6 dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
- 7 48. Wickham, H. *ggplot2*. (Springer New York, 2009). doi:10.1007/978-0-387-98141-
8 3
- 9 49. Pointing, S. B. *et al.* Biogeography of photoautotrophs in the high polar biome.
10 *Front. Plant Sci. Funct. Plant Ecol.* **6**, 692 (2015).
- 11 50. Nguyen, N. H. *et al.* FUNGuild: An open annotation tool for parsing fungal
12 community datasets by ecological guild. *Fungal Ecol.* **20**, 241–248 (2016).
- 13 51. Tedersoo, L. *et al.* Global diversity and geography of soil fungi. *Science (80-.)*. **346**,
14 (2014).
- 15 52. Ulrich, W., Almeida-Neto, M. & Gotelli, N. J. A consumer's guide to nestedness
16 analysis. *Oikos* **118**, 3–17 (2009).
- 17 53. Fruchterman, T. M. J. & Reingold, E. M. *Graph Drawing by Force-directed*
18 *Placement*. **21**, (1991).
- 19 54. Almeida-Neto, M., Guimarães, P., Guimarães, P. R., Loyola, R. D. & Ulrich, W. A
20 consistent metric for nestedness analysis in ecological systems: reconciling
21 concept and measurement. *Oikos* **117**, 1227–1239 (2008).
- 22 55. Ulrich, W. *et al.* A comprehensive framework for the study of species co-
23 occurrences, nestedness and turnover. *Oikos* **126**, 1607–1616 (2017).
- 24 56. Oksanen, J. *et al.* R package: Vegan. (2017).
- 25 57. Swenson, N. G. *Functional and Phylogenetic Ecology in R*. (Springer New York,

2014). doi:10.1007/978-1-4614-9542-0

58. Kembel, S. W. *et al.* Picante: R tools for integrating phylogenies and ecology.

Bioinformatics **26**, 1463–1464 (2010).

59. Paradis, E. Ape package. (2018). Available at:

<https://www.rdocumentation.org/packages/ape/versions/5.2>. (Accessed: 24th

October 2018)

60. Michonneau, F. Phylobase package. Available at:

<https://www.rdocumentation.org/packages/phylobase/versions/0.8.4>.

(Accessed: 24th October 2018)

61. Dray, S. Adephylo package. Available at:

<https://www.rdocumentation.org/packages/adephylo/versions/1.1-11>.

(Accessed: 24th October 2018)

62. Zhang, J. Phylotools package. Available at:

<https://www.rdocumentation.org/packages/phylotools/versions/0.2.2>.

(Accessed: 24th October 2018)

63. Höhl, M., Rigoutsos, I. & Ragan, M. A. Pattern-based phylogenetic distance

estimation and tree reconstruction. *Evol. Bioinform. Online* **2**, 359–75 (2007).

64. Choi, J. & Kim, S.-H. A genome Tree of Life for the Fungi kingdom. *Proc. Natl. Acad.*

Sci. U. S. A. **114**, 9391–9396 (2017).

65. Ebersberger, I. *et al.* A Consistent Phylogenetic Backbone for the Fungi. *Mol. Biol.*

Evol. **29**, 1319–1334 (2012).

66. Boyle, E. E. & Adamowicz, S. J. Community phylogenetics: Assessing tree

reconstruction methods and the utility of DNA barcodes. *PLoS One* **10**, (2015).

67. Kembel, S. W. & Hubbell, S. P. The phylogenetic structure of a neotropical forest

tree community. *Ecology* **87**, 86–99 (2006).

- 1 68. Webb, C. O., Ackerly, D. D. & Kembel, S. W. Phylocom: software for the analysis of
2 phylogenetic community structure and trait evolution. *Bioinformatics* **24**, 2098–
3 2100 (2008).
- 4 69. Horner-Devine, M. C. & Bohannan, B. J. M. Phlogenetic lcustering and
5 overdispersion in bacterial communities. *Ecology* **87**, S100–S108
6
7

Display Items (2 Figures, 1 table)

Fig. 1. REVISE FORMAT OF LEGEND AND IMAGE Antarctic air and soil habitats

support distinct bacterial and fungal communities. a) Route predictions for average minimum and maximum modelled residence time for microorganisms in air based on HYSPLIT back trajectory analyses. Back trajectories indicate distance travelled for sampled air mass at 3 d (598 - 2581 km distance, average altitude of 2769 m, maximum altitude of 5174 m A.M.S.L.) and 15 d (4673 - 11216 km distance average altitude 3034m, maximum altitude 6886 km). **b) and c)** Alpha diversity estimates (Chao1 richness and Pielou's relative evenness) and visualisation of community dissimilarity using Principal Co-ordinate Analysis of weighted UniFrac distance by habitat for **b)** Bacteria and **c)** Fungi. Boxplot whiskers represent 1.5 times the interquartile range from the first to the third quartiles or the maximum/minimum data point within the range. **d) and e)** Distribution and relative abundance of **d)** Bacteria and **e)** Fungi in Antarctic air and soil. Each stack bar represents data from three pooled replicates for each substrate location. Diversity is shown at phylum level as this is the highest taxonomic rank at which between-substrate differences are noticeable. Venn diagrams show amplicon sequence variants (ASVs) and operational taxonomic units (OTUs) count and percentage occurrence within and between each habitat. The high altitude samples, i.e., those without underlying soil, do not have corresponding soil samples. Sampling locations: valley (soil and 1.5m above ground), elevated (soil and 1.5m above ground at higher altitude locations at Bull Pass and valley ridges), 2000m (helicopter samples). Interactive graphics identifying taxonomic composition to lower taxonomic ranks within each sample are presented in the Online Supplementary Information (Fig. S2). Comparison with non-polar samples is given in the Online Supplementary Information Fig. S2.

Fig. 2. REVISE FORMAT OF LEGEND AND IMAGE Comparison of bacterial and

fungal diversity from Antarctic and non-polar sources. a) Distribution and relative

abundance for the 1,000 most abundant bacterial amplicon sequence variants (ASVs)

and fungal operational taxonomic units (OTUs). **b)** Rarefaction curves are shown for

Bacteria and Fungi for each Antarctic habitat to illustrate sampling depth to near-

asymptote. **c)** Co-occurrence associations derived from Ecological Network Analysis.

We enforced maximum Bray-Curtis distance of 0.2 for Bacteria and 0.4 for Fungi to

establish connection between nodes (representing communities). The nodes were

positioned using the Fruchterman-Reingold method. Sampling locations: valley (soil and

1.5m above ground), elevated (soil and 1.5m above ground at higher altitude locations

at Bull Pass and valley ridges), 2000m (helicopter samples) and New Zealand (non-

polar). Full taxonomic comparison for all polar and non-polar samples is given in the

Online Supplementary Information Fig. S2.

Fig. 3. CAN WE CONVERT THE NRI TO A TABLE TOO? PLEASE CAN YOU PROVIDE

THE DAT IN TALE FORM AND CHECK LEGENDS BELOW

Phylogenetic structuring of local and global pools for bacterial and fungal

diversity. a) Nestedness estimates made using the NODF model (where 0 = no

nestedness, 100 = perfect nestedness). Fungi were more nested (NODFc = 62) than

Bacteria (NODFc = 18). Bacteria and Fungi were significantly nested for taxa

composition under the hypothesis that nestedness can be maximised by ordering sites

from the most connected to the least connected. Least connected sites are demonstrated

as a proper subset of richer, more connected sites. **b)** Net Relatedness Index analysis of

phylogenetic structure within each sample type for Bacteria (b) and Fungi (f). Error

1 bars show the standard error of the mean for all samples in a given substrate type.
2 Values for highly dispersed non-polar bacterial communities associated with forest soil
3 are given for comparison (bn) and indicated by an asterisk ⁶⁹. Sampling locations: valley
4 (soil and 1.5m above ground), elevated (soil and 1.5m above ground at higher altitude
5 locations at Bull Pass and valley ridges), 2000m (helicopter samples) and New Zealand
6 (non-polar).

7