**QUEEN'S
UNIVERSITY
BELFAST**

The distribution of novel bacterial laccases in alpine paleosols is directly related to soil stratigraphy

Dandare, S., Young, J. M., Kelleher, B. P., & Allen, C. (2019). The distribution of novel bacterial laccases in alpine paleosols is directly related to soil stratigraphy. *Science of the Total Environment*, 671, 19-27.

Published in:
Science of the Total Environment

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2019 Elsevier B. V.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>, which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

The distribution of novel bacterial laccases in alpine paleosols is directly related to soil stratigraphy.

S.U. Dandare¹, J.M. Young¹, B.P. Kelleher², C.C.R. Allen^{*1,3}

¹School of Biological Sciences, Queen's University Belfast, UK

²School of Chemical Sciences, Dublin City University, Ireland

³Institute for Global Food Security, Queen's University Belfast, UK

**Corresponding author*

ABSTRACT

Bacterial laccases are now known to be abundant in soil and to function outside of the cell facilitating the bacterial degradation of lignin. In this study we wanted to test the hypotheses that: i) Such enzymes can be identified readily in stratified paleosols using metagenomics approaches, ii) The distribution of these genes as potential 'public good' proteins in soil is a function of the soil environment, iii) Such laccase genes can be readily retrieved and expressed in *E.coli* cloning systems to demonstrate that *de novo* assembly processes can be used to obtain similar metagenome-derived enzyme activities. To test these hypotheses, *in silico* gene-targeted assembly was employed to identify genes encoding novel typeB two-domain bacterial laccases from alpine soil metagenomes sequenced on an Illumina MiSeq sequencer. The genes obtained from different strata were heterologously cloned, expressed and the gene products were shown to be active against two classical laccase substrates. The use of a metagenome-driven pipeline to obtain such active biocatalysts has demonstrated the potential for gene mining to be applied systematically for the discovery of such enzymes. These data ultimately further demonstrate the application of soil pedology methods to

environmental enzyme discovery. As an interdisciplinary effort, we can now establish that paleosols can serve as a useful source of novel biocatalytic enzymes for various applications. We also, for the first time, link soil stratigraphy to enzyme profiling for widespread functional gene activity in paleosols.

Keywords paleosols, stratigraphy, laccase, metagenome

1. INTRODUCTION

Laccases are oxidoreductase enzymes which can act upon a broad range of phenolic and non-phenolic compounds *via* the reduction of molecular oxygen to H₂O. Consequently, they have many applications as 'green' biocatalysts in varied industries - including textile and dye decolourisation, bioremediation, pulp bio-bleaching, microbial fuel cells and biosensors (Couto et al., 2006). The advent of high throughput sequencing and associated metagenomics has revealed that bacteria possess a huge diversity of laccase-like multicopper-oxidases (LMCOs) genes. Many of these appear to possess N terminal signal peptide sequences (Ausec et al., 2011), suggesting they are secreted by the cell, potentially enabling oxidation of lignin-derived compounds outside the cell. Only a few bacterial LMCOs have been purified and characterised to date; however, data from these studies suggests unique and biotechnologically advantageous properties, including chloride tolerance (Guan et al., 2014), increased stability at extremes of pH and temperature (Chauhan and Jha, 2018, Fang et al., 2011) and ionic liquid tolerance (Chauhan et al., 2018). Soils are the primary site of terrestrial organic carbon degradation and are some of the most genetically diverse environments on earth (Torsvik 1990). Despite this, very little is known about both the abundance and distribution of Bacterial LMCO's in the soil.

Bacterial laccases can be divided into several categories based upon sequence similarity, the number of cupredoxin domains they possess and their tertiary structure (Ausec et al., 2011). Type B and C two domain laccases are homotrimeric proteins with each monomer consisting of two cupredoxin domains (Martins et al., 2015). By contrast, three-domain bacterial laccases are monomeric and can be

subdivided into three categories based upon sequence similarity, ‘small’ three domain laccases, ‘big’ three domain laccases and CotA type laccase (similar to cotA from *B. subtilis*). Most Bacterial laccases which have been heterologously expressed and characterised to date are three domain type laccases (Martins et al., 2015), and recombinant three-domain Laccases have found widespread industrial usage.

There is presently considerable interest in the degradation of lignin in diverse environments, where different microorganisms compete for lignin and lignin breakdown products (Cragg et al., 2015). However, fundamental questions about the distribution of lignin-degrading bacteria are evident: Do different types of lignin degrading bacteria exist in competing or cooperative bacterial communities? How do these bacteria exist alongside the lignin-degrading fungi? Do various types of extracellular lignin-degrading ‘common good’ proteins (such as peroxidases and laccases) exhibit functional variation that is related to the environmental niche where they are found as opposed to their phylogenetic origin? Analysis of soil metagenomes may help us to answer some of these questions (Jacquiod et al., 2014). Further gene-mining approaches can be used to retrieve functional laccases from soil metagenomes also (Yang et al., 2018).

In this study, we used a metagenome analysis of genes for bacterial laccase enzymes in an alpine paleosol. In bacteria, laccases have differing functionalities: whereas some enzymes are clearly associated with non-catabolic roles, such as endospore germination in Firmicutes and are not known to be secreted from the cell (Francisco et al., 2004); others (e.g. the two-domain group of bacterial laccases) appear to play an extracellular function.

Here we took samples from Alpine paleosols (Mahaney *et al.*, 2016) that we know have well-defined stratigraphy, with the inceptisols harbouring distinct Ah, Bw and Cox soil horizons. We then tested the hypothesis that there is a differential distribution of the major well-defined groups of functional bacterial laccases across these horizons, which are therefore affected by changing physical and chemical environmental conditions. By investigating the demarcation of different classes of laccase

gene across paleosol soil horizon boundaries, we sought to show for the first time that the different environments present in these soil environments could affect the molecular ecology of such genes and therefore that their functionality could be linked to their ecological niche.

In validating these core questions, we also wanted to show that the two-domain group of bacterial laccases can be cloned and expressed for biochemical analysis when directly derived from such soil metagenome sources.

2. MATERIALS AND METHODS

Materials

The substrates 2, 2' – Azino-bis (3-ethylbenzothiazoline-6-sulfonate) (ABTS) and syringaldazine (SGZ) were products obtained from Sigma-Aldrich (St. Louis, MO, USA). Mo Bio Powersoil DNA extraction kit was obtained from Qiagen. Gel Extraction Kit, Plasmid Mini-Prep Kit, Dream Taq DNA polymerase, Phusion green DNA polymerase, ClonJET cloning kit, aLicator LIC cloning and expression system, SYBR Green qPCR Master Mix, prestained protein ladder, and the TOP10 chemically competent *E. coli* cells were all purchased from Thermo Fisher Scientific (UK). The BL21 (DE3) chemically competent cells were purchased from New England Biolabs, while the restriction enzymes were obtained from Fermentas. All other chemicals used in this study were of analytical reagent grade.

Sampling sites

Soils samples used in the study were as previously described: vertical profiles of soil from a series of glacial moraines in the Guil river valley in the French Alps, (G1, G2, G3a, G9, and G11) and the Po river valley in the Italian Alps (V10) were sampled by their soil horizon classification (Mahaney et al., 2016).

Soil sample collection

Samples were collected as previously described in Mahaney et al. (2016). A map of the sampling site is reproduced in the supplementary section (S6) from Young et al. (2019). Briefly, samples were collected from the mid-point depth of soil horizon profiles for specific horizons (e.g. Ah, Bw, Cox). So for a soil horizon of 5cm depth, samples were collected 2.5 cm into the horizon. The aim was to obtain samples that were representative of each horizon. Samples were collected using sterile implements and storage vials. They were immediately stored on ice, in the dark, throughout transport to the laboratory. They were then frozen at -20°C before further analysis.

Metagenomic library construction

DNA was extracted from alpine paleosol samples using a PowerSoil DNA extraction kit (Mo Bio). For each extraction 0.25g of soil was used, extractions were performed in triplicate for each sample and pooled. Extractions were performed according to the manufacturer's protocol with the following modifications: samples were homogenised using a FastPrep 120 cell disrupter system (Thermo-Fisher) at 5.5 m.s⁻¹ for 2 minutes, rather than a standard benchtop vortex. The eluted DNA was further purified *via* two rounds of ethanol precipitation; DNA solution was suspended in 3 volumes of ice-cold 100% ethanol, 0.1 volumes 5 M sodium acetate solution (pH 5.2) and 2 µL of linear polyacrylamide (LPA), the solution was then incubated overnight at -20°C and centrifuged at 18000 x g at 4°C for 30 minutes. The supernatant was discarded, and the pellet washed in 70% ice-cold ethanol and again centrifuged at 18000 x g at 4°C for 5 minutes. Finally, the supernatant was discarded and the pellet allowed to air dry for 15 minutes before resuspension in 50 µL molecular grade H₂O. Final DNA concentrations were measured using a Quantus Fluorometer (Promega) in conjunction with the Quantiflour dsDNA dye system (Promega).

DNA sequencing libraries for Illumina sequencing prepared using a Nextera NX library preparation kit (Illumina), DNA libraries were sequenced on an Illumina MiSeq DNA sequencer in the paired-end mode for 300 cycles. Library preparation and sequencing were performed at the University of Cambridge

DNA sequencing facility. Raw Sequence data files are available in the NCBI sequence read archive under Bioproject number PRJNA39461.

Metagenomic library screening and sequence analysis

Raw reads were quality trimmed to a minimum mean quality of 20 over a sliding window of 10 bases using bbdut from the bbmap package (Bushnell, 2015). Reads from all 18 samples were co-assembled using Megahit (Li et al., 2016), contigs < 1kbp were discarded, and open reading frames were identified and extracted using Prodigal (Hyatt et al., 2010) with the `–meta` flag. Resulting in a library of 1154087 ORFs. Putative bacterial laccase sequences were classified from this library of ORFs using previously published profile hidden Markov models for bacterial laccases (Ausec et al., 2011). Only hits with an e-value < 1e-10 were retained, these were manually checked against the NCBI conserved domain database to ensure they possessed the cupredoxin domain characteristic of laccases. The final library of laccases was taxonomically classified using Kaiju (Menzel et al., 2016), against a database of all proteins from all bacteria, archaea, single-celled eukaryotes and viruses in the NCBI non-redundant protein database (NCBI Resource Coordinators, 2017). Raw reads were mapped onto the library of putative laccase genes using bbmap with the flags `k=13 vslow=t`. Normalised relative abundances were calculated as TPMS (transcripts per million) as this is the most robust and appropriate method for comparisons of gene sets between samples described by Wagner et al. (2012) and is applicable to metagenomic as well as RNA-Seq data.

In order to isolate novel type B two-domain laccase for downstream expression and characterisation, primers were designed for three type-B two-domain laccases for which an entire ORF including upstream ribosomal binding site was captured in the assembly (**supplementary data S1**). However, only one of these primer sets yielded PCR products (mg_lacc_fw: 5'-ATGAGCAATAGACGTGGCTT-3', mg_lac_rev: 5'-ATTCATGTTTGTGCTCCATCTC-3').

PCR was performed using these primers with total extracted DNA from the three Viso 10 site samples, with the following conditions: initial denaturation 95°C for 3 minutes; denaturation 95°C for 30

seconds; annealing 53°C for 90 seconds; extension 72°C for 1 minute; final extension 72°C for 7 min; 30 cycles. DNA gel electrophoresis confirmed fragments of the expected size (1500bp), these were excised from the gel and purified using a GeneJET gel extraction kit (Thermo-Fisher). Purified PCR products were ligated into PJET 1.2/blunt vector using a CloneJET PCR cloning kit (Thermo-Fisher), and transformed into *E.Coli* BL21 (DE3) competent cells by heat shock transformation. Transformed clones were grown overnight on 100 µg mL⁻¹ ampicillin LB agar plates at 30°C. Individual colonies were selected and grown overnight in 100 µg mL⁻¹ ampicillin LB broth, Plasmids were extracted from each broth culture using a GeneJET plasmid extraction kit (Thermo-scientific), extracted plasmids were sequenced using an Applied Biosystems 3730 DNA analyser at the University of Dundee DNA sequencing facility, yielding a final library of 8 sequenced PCR products of the expected insert size.

Statistical analysis of data

Metagenome sequence data were analysed using the same methods to those described in our earlier report – the identical metagenome dataset was used in the present study (Young et al., 2019). Statistical analysis were carried out in R version 3.4.1 (R CoreTeam, 2011), running on an Ubuntu 16.04 LTS PC with 16 GB RAM and an Intel core i7

6700K processor. The reader is referred to our earlier study for details of methodology used.

Laccase sequence dissimilarity was compared using Krustal-Wallace ANOVA analysis performed using the BASE R: kruskal test package (R Core Team, 2011, Kruskaland Wallis, 1952).

The analysis of 26 abiotic variables – previously described by Mahaney et al. (2016) –a Euclidean distance dissimilarity matrix of soil geochemical variables was produced using the package Vegan (Dixon, 2003) in R. v3.4.1 [R Core Team, 2011] using permutational analysis of variance. Principle component analysis was performed on this matrix using the vegan function cap scale, two-dimensional ordination plots were produced using the first two principle components as axes. For statistical comparison of laccase, the laccase genes were identified as being of 5 distinct types (see

Figure 3 and references in the legend). The laccase genes identified in each sample were annotated using the package Kaiju (Menzel et al., 2016). Statistical analysis of each enzyme subgroup across the metagenome datasets was performed using Kruskal-Wallis one way ANOVA (Kruskal & Wallis, 1952). Maximum likelihood phylogenetic trees were drawn using Mega 7.0 (Kumar et al., 2016). The reference sequences shown were downloaded from the NCBI database.

Cloning and heterologous expression of novel laccase genes

The laccase genes were then sub-cloned into pLATE11 (untagged) ligation independent cloning (LIC) and expression vector without their signal peptides. Genes without their signal peptides were obtained by designing new forward primers that excluded the 26 amino acids N-terminal signal peptide during amplification. The pJET 1.2/blunt plasmids containing laccase genes were used as the templates for amplification of laccase genes without their signal peptides. Each Lac-pLATE11 recombinant was transformed into *E. coli* BL21 (DE3), and transformants were selected on LB agar plates supplemented with 100 $\mu\text{g mL}^{-1}$ ampicillin.

For the expression of recombinant laccases, LB (5 g L⁻¹ yeast extract, 10 g L⁻¹ tryptone, 5 g L⁻¹ NaCl) medium was used (Bertani, 1951). Expression studies were performed in shake flasks; each culture was inoculated with individual overnight pre-cultures (grown at 30 °C, 140 rpm) and incubated at 30 °C and 160 rpm until the cells grew to a mid-exponential phase ($\text{OD}_{600} \approx 0.6$). At mid-growth phase, the temperature and rotation speed were downshifted to 25 °C and 100 rpm respectively; and the addition of 0.1 mM IPTG induced the expression of recombinant laccases. The culture was also supplemented with copper ions in the form of CuSO_4 to a final concentration of 0.25 mM. Oxygen-limiting condition (microaeration) was introduced by switching off rotation (0 rpm) 4 h after induction; static incubation has been reported to increase the amount of copper-loaded bacterial laccases in *E. coli* (Durão et al., 2008). The static incubation was maintained for 20 h, and then cells were harvested by centrifugation (6000 rpm, 15 min). The recovered cell pellets were frozen at -80 °C.

In order to recover the proteins, cells were thawed and lysed by resuspending in CelLytic B solution (Sigma-Aldrich) containing 0.5 U mL⁻¹ Benzonase Nuclease (Novagen) and 0.2 mg mL⁻¹ lysozyme. The mixture was incubated at room temperature with shaking (200 rpm) for 20 min according to manufacturer's instructions. After the soluble fraction was separated from the cell debris by centrifugation (18000 x g, 15 min), it was heated at 70 °C for 15 min to precipitate most of the *E. coli* proteins as a form of partial purification. The thermal activation of laccases has been reported in several studies (Mollania et al., 2018; Papinutti et al., 2008; Koroleva et al., 2001). Samples collected throughout the expression studies including the cell-free extract (CFE) were mixed 1:1 with SDS-PAGE loading buffer, denatured at 95 °C for 5 min, and electrophoresed on a 10% SDS-PAGE resolving gel. The concentration of protein was estimated using the Bradford assay with bovine serum albumin (BSA) as the standard.

Enzyme assay

The activity of laccases was measured at 25 °C using ABTS ($\epsilon_{420\text{nm}} = 36 \text{ mM}^{-1} \text{ cm}^{-1}$) and syringaldazine ($\epsilon_{530\text{nm}} = 65 \text{ mM}^{-1} \text{ cm}^{-1}$) as substrates with a 6705 UV/Vis spectrophotometer (Jenway). Each assay mixture was made up to 1 mL and contained 2 mM ABTS (dissolved in phosphate buffer) or 0.05 mM SGZ (dissolved in absolute methanol), 100 μL of cell-free extract (CFE) and phosphate buffer at pH 4.0 or pH 6.5 for ABTS and SGZ assays respectively. The fungal laccase from *Trametes versicolor* (Sigma Aldrich) was used as a positive control while cell-free extract prepared from an expression experiment with an empty vector was used as a negative control. The oxidation of both ABTS and SGZ was followed by a change in absorbance at 420 nm and 530 nm respectively for 10 min. One unit (U) of laccase activity was defined as the amount of laccase required to oxidise 1 μmol of substrate per minute (1 $\mu\text{mol min}^{-1}$). All assays were performed in triplicate.

3. RESULTS

3.1 THE STRATIGRAPHY OF SAMPLING SITES IS CONSISTENT WITH SOIL HORIZONS REPRESENTING DISTINCT SETS OF ENVIRONMENTAL PARAMETERS THAT ARE INDEPENDENT OF GEOGRAPHIC SEPARATION.

Previous studies at this site in the Guil valley and Mount Viso region have classified the sampled paleosols as inceptisols with specific soil pedons as defined in the geological literature (Mahaney et al., 2016; Young et al., 2019). While there is a long established convention of describing glacial sediment deposits in terms of Ah, Bw, Cox, Cu, and D horizons - based upon variation in physical and chemical distinctions down the soil profile - there is no detailed attempt to date, to link differences in extra-cellular microbial enzyme distribution to this classification.

We initially used data gathered in an earlier paper (Mahaney et al., 2016) to show that the 27 chemical and physical parameters previously measured in each of the 18 samples – 6 each being designated as Ah, Bw and Cox – did indeed indicate that soil Horizons represent distinct physical/chemical environments. Soil horizons were chemically and physically distinct as indicated by Principle Coordinates Analysis, wherein the first two principle co-ordinate axes explained over 70% of the total variation between samples (Figure 1).

3.2 THE DISTRIBUTION OF BACTERIAL LACCASE GENES IS RELATED TO SOIL STRATIGRAPHY

We searched a library of 1154088 ORFs from all the paleosol samples sequenced from a metagenomic co-assembly of 135365644 raw reads for putative bacterial laccases using previously published profile hidden Markov models for bacterial laccases (Ausec et al., 2011), resulting in 323 new putative laccase-like ORFs being identified. Of these, there were many that include both start and stop codons. The amino acid translation of these sequences were additionally aligned against the NCBI nr protein database revealing that the majority of these sequences had low sequence identity but high sequence

similarity to known LMCO genes [supplementary data table S2]. This analysis highlighted the abundance of novel LMCO sequences in these soils.

We hypothesised that the abundances of different classes of bacterial laccases would vary significantly between stratified soil horizons, as their functional role as extracellular biocatalysts would inevitably be linked to localised environmental parameters. In order to test this hypothesis, we applied Kruskal Wallace one way ANOVA (Kruskal & Wallis, 1952) to the relative abundance of putative laccases grouped by soil horizon (**Figure 2**), revealing that all but two classes of bacterial laccases vary significantly in abundance between soil horizon. The type B two domain laccases appear to increase in abundance down a soil profile and show the strongest variation with soil horizon, while the big three domain, small three-domain and CotA –type laccases all decrease in abundance with soil depth. It should be noted that, in terms of relative abundance, the two most abundant classes of enzyme in these samples are the small three-domain and type B two domain laccases (**Figure 3**). These two groups comprise >70% of all laccase-like genes identified.

The predicted taxonomic affiliations for putative laccase ORFs shows strikingly different community structures between laccase types (**Supplementary Figure S4**). Additionally, principal coordinate analysis of the predicted taxonomic affiliations shows clear differences between the Ah Horizon and Bw/Cox horizons for the type B two domain, small three domain and CotA type laccases (**Figure 3**). Indicating that the distribution of laccase genes in the soil horizons is affected by the soil horizon location as well as by the phylogeny of any expressing bacterial cell. Critically, the two most abundant groups of identified genes show the greatest demarcation between soil horizons in these PCA plots. Where relatively few genes were identified in the horizons (e.g. the ‘Big 3D’ group) there is no clear demarcation. It is important to note here that given the fragmented nature of the assembly from which these ORFs were derived, that we cannot deduce which organisms may be harbouring multiple laccase gene types, although such organisms should certainly exist.

3.3. Further analysis of typeB two-domain laccases, including expression and activity.

Three sets of primers were designed to amplify novel type-B two-domain laccases for which a complete ORF including start and stop codons, as well as a signal peptide (indicating potential extracellular secretion), could be identified. These primers were all tested against multiple samples from the paleosols, to see if laccases could be readily amplified. We found only one set of primers gave reliable amplification and only when using DNA extracted from the V10 Bw and Cox samples. These primers were designated [V10_lac]. Cloning and sequencing of these PCR products yielded eight high-quality sequences of the expected insert size; these showed high homology with the assembled parent sequence; these sequences are compared in a phylogenetic tree (**Figure 5**). Interestingly, none of the PCR amplified sequences were 100% identical to the parent sequence, this may be attributed to the fact that a contiguous sequence derived from a typical metagenomic assembly is often a consensus sequence of closely related strains and variants present in the dataset, other factors may be involved including PCR error, variants introduced during exponential growth of the cloning vectors and sequencing error (both Illumina and Sanger sequencing). Despite this, in this case, it appears that the assembled gene sequence was sufficiently accurate to allow the design of PCR primers which could be used to isolate functionally active proteins from a portion of the soil samples.

3.4 Characterisation of cloned laccases

There is a limited detailed biochemical analysis of two domain laccase enzymes from Acidobacteria in the literature, despite the importance of these bacteria in soil (Jones et al., 2009). Further, there is just one account of a metagenome laccase analysed to date (Ausec et al., 2017). To validate the metagenomic enzyme discovery pipeline used in this study, we selected four Acidobacteria two-domain laccase enzymes from our earlier gene mining activities (**Supplementary S3**) that were cloned and expressed for biochemical characterisation. The aim here was to demonstrate bona fide laccase like activity for the identified gene products.

287 In all the laccases tested, the enzymes showed higher activity against SGZ than ABTS (Figure 6). This
 288 observation has been made with other laccases (Guan et al., 2014; Lu et al., 2013; Ye et al., 2010).
 289 Also, enzymes mined from the Bw horizon (LacB3 and LacB6) showed more activity against both
 290 substrates tested than enzymes from the C horizon (LacC12 and LacC15).

291 This experiment confirmed that complete functional genes for the typeB 2D group could be isolated
 292 from the paleosols. It effectively validated both the use of our metagenomics methods for novel
 293 laccase gene discovery and complete gene isolation, from these paleosols. It also validated the
 294 designation of our metagenome analysis methods for the identification of putative laccase genes.

295 Further experimentation confirmed that these enzymes were in fact functionally distinct laccases. For
 296 example, copper dependence is a key characteristic of laccases, and we showed that one of the
 297 enzymes (LacB6, that had the highest activity with both substrate of the four clones) required CuSO₄
 298 addition for full activity (**Figure 7**). We also showed that the activity of LacB6 against syringaldazine
 299 increased with increasing concentration of copper. The addition of 10-, 50- and 100 µM CuSO₄ in the
 300 assay mix increased the enzyme activity 2.5-, 5, and 8-fold respectively. These observations confirm
 301 that LacB6 is a copper-dependent enzyme and that copper was only partially incorporated into the
 302 enzyme's active site during expression. Cloned and expressed laccases are usually partially copper-
 303 depleted as a result of the difficulty of the passage of copper ions from the medium into the cell during
 304 expression (Harris, 2000; Vulpe and Packman, 1995).

305 Molecular modelling was performed on the Laccase_B_6 amino acid sequence using three publically
 306 available protein modelling servers, ITASSER (<https://zhanglab.ccmb.med.umich.edu/I-TASSER/>),
 307 Phyre2 (<http://www.sbg.bio.ic.ac.uk/~phyre2/html/page.cgi?id=index>) and Swiss-model
 308 (<https://swissmodel.expasy.org/>). All three tools found the closest available crystal structures for
 309 homology modelling had between just 14-32% amino acid identity to the query sequence (data not
 310 shown). This suggests that homology models produced for this protein sequence are unlikely to be
 311 representative of the true structure.

4.DISCUSSION

The relationship between soil stratigraphy nomenclature and abiotic chemical/physical variables has been long established in the geological literature – and has also previously been considered in the context of Alpine glacial deposition (Birkland et al., 1979). In this study, we were able to confirm a clear differentiation between the designated soil horizon samples previously described by Mahaney et al. (2016) and the physical/chemical composition of the samples using PCA plots. The data form a framework for later comparison with the distribution of laccases in the same samples.

Our main objective in this study was to see if there was a connection between soil pedon classification and the distribution of bacterial laccases in a model soil ecosystem. We chose to look at bacterial laccases for a number of reasons: i) they are enzymes with diverse function. Some are required for intracellular endospore germination/formation processes, such as the Cot A proteins (Francisco et al., 2004), while there is clear evidence that the small three domain and typeB two-domain enzymes are extracellular enzymes are used for lignin degradation (Bugg et al., 2004); ii) The extracellular bacterial laccases are secreted by one taxon into the environment, but then can also benefit other components in a microbial ecosystem such as soil. As such, we might expect the distribution of these genes to be related to cooperative behaviours between different bacterial taxa in localised communities (West et al. 2006); iii) As potential industrial biocatalysts they are particularly useful, while also being structurally simple as single component enzymes – making them good candidates to demonstrate the potential for paleosol metagenome-mining as a source of useful functional genes (Call & Mucke, 1997).

Looking at the differences in the distribution of the genes encoding five different ‘classes’ of enzymes, that are already established in the literature, two key observations become evident: Firstly, there does appear to be present distinct sub-populations of laccase enzymes in soil pedons from our data. Not only do the relative abundance of laccase genes in the stratified soils clearly demarcate between Ah, Bw and Cox soil horizons, but we also observed that the beta diversity

337 between populations of laccase genes in soil horizons is suggestive of greater dissimilarity of gene
338 populations between soil horizons in any individual paleosol than between geographical locations –
339 even when these locations are kilometres apart (such as between the site V10C and any of the G
340 sites studied here). This is especially evident for the typeB two-domain and small three-domain
341 enzyme groups that were generally abundant in all the soil pedons (**Figure 3**). Clearly, from the
342 perspective of bacterial laccase gene function and phylogenetic diversity, we can say that Ah and Bw
343 soil horizon - though just separated physically by centimetres - are fundamentally different
344 environments . One explanation we can propose for this marked difference in laccase gene diversity
345 across soil horizons down profile is their potential importance as ‘public good’ proteins in complex
346 bacterial communities of heterotrophic bacteria that rely upon the degradation of plant-derived
347 lignin as a carbon and energy source. The principal component analysis of the 27 physical
348 parameters measured in the soil horizons (**Figure 1**) showed clear demarcation of the three types of
349 soil horizon across all 18 soil pedons. However, differences between the populations of the most
350 abundant typeB two-domain laccases and the small three-domain laccases across the boundary of
351 the Ah and Bw/Cox pedons follow a different distribution: clear dissimilarity is evident between Ah
352 and Bw gene populations for PCoA1, whereas no dissimilarity is seen between the populations of
353 these genes in the Bw and Cox horizons. This would support a hypothesis that biological and perhaps
354 even societal factors consistently lead to polarisation of these groups of enzymes between Ah and
355 Bw horizons. This conclusion is certainly speculative at this stage, but it does highlight the possibility
356 of using paleosol horizon metagenome analysis of such ‘public goods protein’ functional genes for
357 the study of developing field of bacterial sociobiology (Griffin et al., 2004). Critically, the different
358 classes of laccase studied do not partition in an identical fashion between the pedons. For example,
359 the CotA type three domain laccase gene populations differentiate as well between Bw and Cox soil
360 horizons as they do between Ah and Bw horizons from the PCA analysis (**Figure 3**). Furthermore, the
361 fact that CotA enzymes are not likely to be public goods proteins could be important. Enzymes may

have the same functionality but different roles in a bacterial community, and laccases in particular may serve a community function in soil microbial ecosystems (Freedman and Zak, 2015).

As initially noted, the use of metagenome gene-mining as a process to source new industrial enzymes is an important possibility for the generation of new biocatalysts. We wanted to demonstrate here that with our paleosol data set we could successfully obtain, clone and validate expression of novel typeB two-domain laccase enzymes. We chose to isolate genes for the most abundant 2D two component enzymes – that have not been studied extensively as biocatalytic enzymes – due to their relative novelty. The methodology developed here shows clearly that such enzymes can be readily obtained and assayed for enzyme activity. The observed activity against model substrates is consistent with other extracellular bacterial laccases of biotechnological importance (Couto et al., 2006).

5. Acknowledgements

We thank the REMEDIATE ITN project (BPK, CCRA), Invest Northern Ireland (JMY) and the Commonwealth Scholarships Commission (SUD) for financial support. We also acknowledge a reviewer of the manuscript for useful suggestions.

379 6. SUPPLEMENTARY DATA

380 **S1:** Complete laccase ORF for which PCR primers were designed. This was assembled using the
381 Megahit assembly approach described in the methods.

382 ATGAGCAATAGACGTGGCTTTTTGCGAAATATTCTCGCCGGAGCTGGCGCTATGGTTTCCGCGAGAGTTCTTT
383 CAGCCCAAGAGACGGACACGTCTCATGGGATGCACGGAATGAAAATGAAAGGTAGGCAGGAGTCGGAGCGC
384 GGCCACGGTTCTCCCGTGCTTATGGAAACGCCGGACGTTCCGCAACTCCGTGGCGCATGGATGGGGGCGTA
385 AAGGAATTCATCTCATTGCCGAGCCGGTGAAACAGGAGATTTTTCTGGCCGGATCGTTGACCTCTGGGGTT
386 ACAACGCCAGCGCTCCGGGGCCGACGATTCAAGTCAACCAGGGCGATCGCGTTCGGATCATCGTGGACAACC
387 ACCTTCCCGAAGCGACGTGCATGCACTGGCACGGCTTCGAAATCCCGAGCGAGATGGACGGAGCGCCCGGTT
388 CGAGTCAAGACCCGATCCCTCCCGGGGGCCGCTTTGTCTATGAATTACGCTCCACCAGGAAGGCACTTATTTT
389 TACCACTCGCACATGGCGATGCAGGAAATGATGGGCATGATCGGCGCGTTCATCATGCATCCCAAGGAGCCA
390 TACAAGCCGCGCGCCGATAAAGACTTCGCCATCATTATGCAGGAGTACGCCATCTTGCCGAACATCAAGGTCC
391 CGAACTCCATGAACATGGAGTTCAACTGGCTTACCTTCAACGGAAAAATCCGGTCCAGCCAACACTCCGCTCATT
392 GTGCGGCACGGAGAGCGCGTCCGCATCCGCCTTATCAATCTGGGGATGGACCACCATCCGATCCACCTTCATG
393 GGCACCAGTTCGTAGTTACAGGTACTGAAGGAGGCCGACAGCCCGAAACGACTTGGGGGGCCGGGCAATACT
394 GTTCTCGTTGGAGTCGCACAGTCCCGCGACGTGGAGTTCGTTGCCAACAACCCCGGCGACTGGATGCTGCACT
395 GCCATCTTCCGCACCACATGATGAACCAGATGTCATCGATGGTCGGCCCGATGTCACGCCGAAATGGAATGCC
396 TGCGGGTCTCGACATGGAGCGGGGCATGGGAATGCTGCGACAAGGAAGCGCGACGTCTGAGGAGAACGGT
397 CCGAGCCTGGGTGCGGGAATGGGCGTCGGTTCGACAGCGGAGCAAACCATGTGCAATTCTCCGCTCAAAGCC
398 GGGAGTCCGATGCAGCACCAAGACATGCCCCGACATGCAGCAGCAAGGGATGCAGGGGAGGCAGCCCGACAA
399 GCGCGATGACACGAAGGACGCGCATTGCTTGCCGGAGGATGCCTTGGTGAAGGTATCATGATGGCGATGG
400 ATCAAATGGTCGACAAACCTGAAACTTCGGCCTGCGTCCCGGCTGGAGCGGCTTCATGGCAGGAATGATGA
401 CGTTCGTACGCGTGCTGCCTCAAGACAAGTACGACCACAGCATGGAGCTTCGAAAAAGCAGGAAGGCAATA
402 AGCCAATGAAAATGGACATGCCGGAGATGGAGCACAAACATGAATAA

403

404 **S2:** Sequence statistics from total co-assembly and isolated laccase ORFs. Statistics are shown for both
405 the complete co-assembly and the putative Laccase ORFs. Laccase hits were identified using the HMM
406 search pipeline as described in the text.

	All_contigs	Laccase_hits
# contigs (>= 0 bp)	7171198	323
# contigs (>= 1000 bp)	540891	151
# contigs (>= 5000 bp)	2869	0
# contigs (>= 10000 bp)	165	0
Total length (>= 0 bp)	4127121895	316566
Total length (>= 1000 bp)	804814220	199008
Total length (>= 5000 bp)	19157720	0
Total length (>= 10000 bp)	2202670	0
# contigs	3220893	283
Largest contig	62040	2877
GC (%)	62.22	61.55

N50	775	1143
N75	614	912
L50	1102891	106
L75	2045807	180

407

408

409 **S3:** Complete amino acid sequences of gene-mined laccases that were shown to have activity. These
410 were obtained using PCR amplification using the V10_las primers as described in the text.

411 **>LacB3**

412 MSNRRGFLRNILAGAGAMASAKVLSAQEMHMSHGTQGMKGMKETERGHVSPTLVETPDVS
413 QLSWRMGGSVKEFHIAEPVKQEIFPGRIVDLWGYNGSAPGPTIQVNEGDRVRIIVDNHL
414 PEATSMHWHGFEIPSAMDGAPGSSQDPIPPGGRFVYEFTLHQEGTYFYHSHMAMQEMMGM
415 IGAFIMHPKEPHKPRADKDFAIIMQEYAILPNIKVPNSMNMEFNWLTfNGKAGPATTPLI
416 VRHGDRVRVRLINLGMDHHPHILHGHQFVVTGTEGGRQPQSTWSPGNTVLVGVAQSRDVE
417 FVATNPGDWMLHCHLPHHMMNQMSMVGPMSSRRNGTPAGLDMERGMGMLRQESATSEENG
418 PSLGRGMGMGSTAEQTVSN SPLKAVNPMQHKDMPNMQQPAMQMKGPDVSKDAISVPGFPQ
419 DAFMEGPMAMMDQMVDKPNFADPPGGSGPTAGTMTFVRVPPPDNYDQIMEHRKKQEGNN
420 PLQMDMPQMEHNQA

421

422 **>LacB6**

423 MSNRRGFLRNILAGAGAMVSARVLSAQETDTSHGMMHGMKMKGTRESESGHGSPMLVEMPD
424 VAQLPWRLDGDVKEFHIAEPVKQEIFPGRIVDLWGYNGSVPGPTIQVNQGDRVRIIVDN
425 HLPEATSMHWHGFEIPNEMDGAPGSSQDPIPPGGRFVYEFTLHQEGTYFYHSHMAMQEMM
426 GMIGAFIMHPKEPYKPRADKDFAIIMQEYAILPNIKVPNPMNMEFNWLTfNGKAGPATT
427 LIVRHGERVRIRLINLGMDHHPHILHGHQFVVTGTEGGRQPQSTWGPNTVLVGVAQSRD
428 VEFVASNPGDWMLHCHLPHHMMNQMSMVGPMSSRRNEMPAGLDMERGMGMLRQGSATSEE
429 NGPSLGRGMGVGSTAEQTSNSPLKAENPMQHQDMPDMQPQGMQMGNPDVSKDANSVPGF
430 PQDAFMEGPMAMMDQMVDKPNFGLRPGWSGFMAGMMTFVRVLPQDKYNQIMELRKKQEG
431 NKPMKMDMPMEHKHE

432

433 **>LacC12**

434 MSNRRGFLRNILAGAGAMASAKVLSAQEMDMSHGMQGMKMKGVKKSEHAHASRMLVETPD
435 VAQLPWRMEGNVKEFHIAEPVKQEIFPGRVVDLWGYNGSAPGPTIQVNQGDRVRIIVEN
436 RLPEATSMHWHGFEVPNEMDGAPGSSQDPIPPGGRFVYEFTLHQEGTYFYHSHMAMQEMM
437 GMIGAFVMRPKQAYRPRVDHDFAILQEYAILPNISVPNSMNMEFNWLTfNGKAGPATT
438 LIVRQGERVRIRLINLGMDHHPHILHGHQFVVTGTEGGRQPESTWGPNTVfVGVAQSRD
439 IEFVASNPGDWMLHCHLPHHMMNQMSNVGPMTRRNGIPAGLDMERGMGMLRQGSATSEE
440 HGPSLGRGMGVGSTFEQRMSNSPLKAGSPMQHQGMQMKGPDVSKDANSVPGFPQDAFMEG
441 PMMAMDQVRVDKPNFGLRPGWSGFMAGMMTFVRVLPQDKYDHIMELRKKQEGNKPMKMDM
442 PEMEHKHE

443

444 **>LacC15**

445 MSNRRGFLRNILAGAGAMASAKVLSAQEMEMSHGMQGMKMKGVKKSERDHASRMLVETPD
446 VAQLPWRMDGSVKEFHIAEPVKQEIFPGRVVDLWGYNGSVPGPTIQVNQGDRVRIIVDN
447 HLPEATSMHWHGFEIPNEMDGAPGSSQDPIPPGGRFVYEFTLHQEGTYFYHSHMAMQEMM
448 GMIGAFIMHPKEPYKPRSDKDFAIIMQEYAILPNIKVPNSMNMEFNWLTfNGKAGPATT
449 LIVRHGDRVRVRLINLGMDHHPHILHGHQFVVTGTEGGRQPQSTWGPNTVLVGVAQSRD
450 VEFVASNPGDWMLHCHLPHHMMNQMSMVGPMSSRRNGMPAGLDMERGMGMLRQGSAMSEE
451 NGPSLGRGMGVGSTAEQTSNSPLKAGNPMQHQAMPNMQQQGMKMGKPDVSKDANSVPGF

PQDAFMEGPMAMQMDKPFNGLRRGWSGFMAGMMTFVRVLPQDKYDHIMEPRKKQEG
NKPMKMDMPMEHKHEY

S4: Laccase taxonomic distribution by type Phylum level

Taxonomic profile for each class of laccase genes Isolated. Laccase gene sequences were taxonomically annotated using Kaiju (Menzel et al., 2016), as described in the main methods section of the paper. See figure 4 for origin of the five laccase type designations.

S.5. Map of sampling sites used in the study (from Young et al., 2019).

Overview of Sampling sites and soil stratigraphy. Panel A: Satellite map of sampling area. Panel B: Stratigraphy of soil profiles for each sample (from Mahaney et al., 2016). Sampling sites are shown at

the top of each profiles, depth in centimetres is displayed on the left and soil horizon classification is displayed on the right of each profile.

7. REFERENCES

- Ausec, L., Zakrzewski, M., Goesmann, A., Schlüter, A., Mandic-Mulec, I., 2011. Bioinformatic analysis reveals high diversity of bacterial genes for laccase-like enzymes. *PLoS ONE*. 6(10), e25724
- Ausec, L., Berini, F., Casciello, C., Cretoiu, M.S., Van Elsas, J.D., Marinelli, F. and Mandic-Mulec, I., 2017. The first acidobacterial laccase-like multicopper oxidase revealed by metagenomics shows high salt and thermo-tolerance. *Applied Microbiology and Biotechnology*, 101(15), pp. 6261-6276.
- Bertani, G. 1951. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. *J. Bacteriol.* 62, 293-300
- Birkland, P.W., Colman, S.M., Burke, R.M., Shroba, R.R., Meierding, T.C., 1979. Nomenclature of alpine glacial deposits, or, What's in a name? *Geology*, 7, pp. 532-536.
- Bugg, T.D.H., Ahmad, M., Hardiman, E.M., Singh, R., 2011. The emerging role for bacteria in lignin degradation and bio-product formation. *Current Opinion in Biotechnology*, 22(3), pp. 394-400.
- Bushnell B. (2015). BBMap short-read aligner, and other bioinformatics tools. Available from sourceforge net/projects/bbmap.
- Call, H.P., Mucke, I., 1997. History, overview and applications of mediated lignolytic systems, especially laccase-mediator-systems (Lignozym®-process). *Journal of biotechnology*. 53 (2–3), pp. 163-202
- Chauhan, P.S., Goradia, B. And Jha, B., 2018. Optimization and up scaling of ionic liquid tolerant and thermo-alkali stable laccase from a marine *Staphylococcus arlettae* S1-20 using tea waste. *Journal of the Taiwan Institute of Chemical Engineers*, 86, pp. 1-8.
- Chauhan, P.S. And Jha, B., 2018. Pilot scale production of extracellular thermo-alkali stable laccase from *Pseudomonas* sp. S2 using agro waste and its application in organophosphorous pesticides degradation. *Journal of Chemical Technology and Biotechnology*, 93(4), pp. 1022-1030.
- Couto, S. R., Herrera, J.L.T., 2006. Industrial and biotechnological applications of laccases: a review. *Biotechnology advances* 24(5), 500-513.
- Cragg, S.M., Beckham, G.T., Bruce, N.C., Bugg, T.D., Distel, D.L., Dupree, P., Etxabe, A.G., Goodell, B.S., Jellison, J., McGeehan, J.E., McQueen-Mason, S.J., 2015. Lignocellulose degradation mechanisms across the Tree of Life. *Current Opinion in Chemical Biology*. 1(29), pp.108-119.
- Dixon, P., 2003. VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science*, 14(6), pp. 927-930.
- Durão, P., Chen, Z., Fernandes, A.T., Hildebrandt, P., Murgida, D.H., Todorovic, S., Pereira, M.M., Melo, E.P. and Martins, L.O., 2008. Copper incorporation into recombinant CotA laccase from *Bacillus*

subtilis: Characterization of fully copper loaded enzymes. *Journal of Biological Inorganic Chemistry*, 13(2), pp. 183-193.

Fang, Z., Li, T., Wang, Q., Zhang, X., Peng, H., Fang, W., Hong, Y., Ge, H., Xiao, Y., 2011, A bacterial laccase from marine microbial metagenome exhibiting chloride tolerance and dye decolorization ability. *Applied microbiology and Biotechnology*. 89(4), pp1103-1110.

Francisco, J.E., Marçal, D., Martins, L.O., Grenha, R., Henriques, A.O., Lindley, P.F., Carrondo, M.A., 2004. Substrate and Dioxygen Binding to the Endospore Coat Laccase from *Bacillus subtilis*. *Journal of Biological Chemistry*, 279 (22), pp. 23472–23476.

Freedman, Z.B. and Zak, B.R., 2015. Atmospheric N deposition alters connectance, but not functional potential among saprotrophic bacterial communities. *Molecular Ecology*. 24(12), pp3170

Godornes, C., Leader, B.T., Molini, B.J., Centurion-Lara, A., Lukehart, S.A. 2007, Quantitation of rabbit cytokine mRNA by real-time PT-PCR. *Cytokine* 38(1), 1-7

Griffin, A.S., West, S.A., Buckling, A. 2004, Cooperation and competition in pathogenic bacteria. *Nature* 430, 1024-1027

Guan, Z., Song, C., Zhang, N., Zhou, W., Xu, C., Zhou, L., Zhao, H., Cai, Y. and Liao, X., 2014. Overexpression, characterisation, and dye-decolorizing ability of a thermostable, pH-stable, and organic solvent-tolerant laccase from *Bacillus pumilus* W3. *Journal of Molecular Catalysis B: Enzymatic*, 101, pp. 1-6

Harris, E.D., 2000. Cellular copper transport and metabolism. *Annual Review of Nutrition*, 20, pp. 291-310

Hyatt, D., Chen, G-L., Locascio, P.F., Land, M.L., Larimer, F.W., Hauser, L.J., 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11, pp119.

Ihssen, J., Reiss, R., Luchsinger, R., Thöny-Meyer, L. and Richter, M., 2015. Biochemical properties and yields of diverse bacterial laccase-like multicopper oxidases expressed in *Escherichia coli*. *Scientific Reports*, 5.

Jacquiod, S., Demaneche, S., Franqueville, L., Ausec, L., Xu, Z., Delmont, T.O. et al., 2014. Characterization of new bacterial catabolic genes and mobile genetic elements by high throughput genetic screening of a soil metagenomic library. *Journal of Biotechnology*, 190(20), pp.18-29.

Jones, R.T., Robeson, M.S., Lauber, C.L., Hamady, M., Knight, R., Fierer, N., 2009. A comprehensive survey of soil acidobacterial diversity using pyrosequencing and clone library analyses. *The ISME Journal*. 3, pp. 442–453

Koroleva, O.V., Stepanova, E.V., Binukov, V.I., Timofeev, V.P. and Pfeil, W., 2001. Temperature-induced changes in copper centers and protein conformation of two fungal laccases from *Coriolus hirsutus* and *Coriolus zonatus*. *Biochimica et Biophysica Acta - Protein Structure and Molecular Enzymology*, 1547(2), pp. 397-407.

Kruskal, W.H., Wallis, W.A., 1952. Use of ranks in one-criterion variance analysis. *Journal of the American Statistical Association*. 47: 583–621

541 Kumar,S., Stecher, G., Tamura, K., 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version
542 7.0 for Bigger Datasets. *Molecular Biology and Evolution*. 33 (7), pp 1870–1874

543 Li, D., Luo, R., Liu, C.M., Leung, C.M., Ting, H.F., Sadakane, K., Yamashita, H., and Lam, T.W., 2016.
544 MEGAHIT v1. 0: A fast and scalable metagenome assembler driven by advanced methodologies and
545 community practices. *Methods*, 102, pp3-11

546 Lu, L., Wang, T.-., Xu, T.-., Wang, J.-., Wang, C.-. and Zhao, M., 2013. Cloning and expression of thermo-
547 alkali-stable laccase of *Bacillus licheniformis* in *Pichia pastoris* and its characterization. *Bioresource*
548 *technology*, 134, pp. 81-86.

549 Machczynski , M.C., Vijgenboom, E., Samyn, B., Canters, G.W., 2009. Characterization of SLAC: A small
550 laccase from *Streptomyces coelicolor* with unprecedented activity. *Protein Science*. 13(9), pp 2388-
551 2397

552 Mahaney WC, Somelar P, Dirszowsky RW, Kelleher B, Pentlavalli P, McLaughlin S, et al. (2016). A
553 Microbial Link to Weathering of Postglacial Rocks and Sediments, Mount Viso Area, Western Alps,
554 Demonstrated through Analysis of a Soil/Paleosol Bio/Chronosequence. *Journal of Geology*, 124, 149–
555 169.¶

556 Martins, L.O., Durão, P., Brissos, V., Lindley , P.F., 2015. Laccases of prokaryotic origin: enzymes at the
557 interface of protein science and protein technology. *Cellular and molecular life sciences*. 72(5) pp.911-
558 22.

559 McKnight, C.J., Briggs, M.S. and Gierasch, L.M., 1989. Functional and nonfunctional LamB signal
560 sequences can be distinguished by their biophysical properties. *Journal of Biological Chemistry*,
561 264(29), pp. 17293-17297.

562 Menzel, P., Ng, K.L., Krogh, A., Marth, G., Lipman, D., 2016. Fast and sensitive taxonomic classification
563 for metagenomics with Kaiju. *Nature Communications*. 7: pp11257.

564 Mohammadian, M., Fathi-Roudsari, M., Mollania, N., Badoei-Dalfard, A. and Khajeh, K., 2010.
565 Enhanced expression of a recombinant bacterial laccase at low temperature and microaerobic
566 conditions: Purification and biochemical characterization. *Journal of Industrial Microbiology and*
567 *Biotechnology*, 37(8), pp. 863-869.

568 Mollania, N., Heidari, M. and Khajeh, K., 2018. Catalytic activation of *Bacillus* laccase after temperature
569 treatment: Structural & biochemical characterization. *International journal of biological*
570 *macromolecules*, 109, pp. 49-56.

571 Muyzer, G., De Waal, E.C. & Uitterlinden, A.G. 1993. Profiling of complex microbial populations by
572 denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding
573 for 16S rRNA. *Applied and Environmental Microbiology*, 59(3), pp. 695-700.

574 Nakamura, K., Kawabata, T., Nobuhiro Go, K.Y., 2003. Novel types of two-domain multi-copper
575 oxidases: possible missing links in the evolution. *FEBS Letters*, 553, pp1873-3468

576 NCBI Resource Coordinators, 2017. Database Resources of the National Center for Biotechnology
577 Information. *Nucleic Acids Research*. 45: D12–D17

578 Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara RB, Simpson GL, Solymos P, Stevens
579 MH, Wagner H (2011). *vegan: Community ecology package*. R package version. 2011, pp.117-118.

580 Papinutti, L., Dimitriu, P. and Forchiassin, F., 2008. Stabilization studies of *Fomes sclerodermeus*
581 laccases. *Bioresource technology*, 99(2), pp. 419-424.

582 R Development CoreTeam, 2011. A language and environment for statistical computing. R Foundation
583 for Statistical Computing, Vienna. <http://www.R-project.org>

584 Suzuki, M.T., Taylor, L.T. & DeLong, E.F. 2000, "Quantitative analysis of small-subunit rRNA genes in
585 mixed microbial populations via 5'-nuclease assays", *Applied and Environmental Microbiology*, 66(11),
586 pp. 4605-4614.

587 Torsvik, V., Jostein, G., Daae, F.L., 1990. "High diversity in DNA of soil bacteria." *Applied and*
588 *environmental microbiology* 56(3), pp. 782-787.

589 Vulpe, C.D. and Packman, S., 1995. Cellular copper transport. *Annual Review of Nutrition*, 15, pp. 293-
590 322

591 Wagner, G.P., Kin, K., Lynch, V.J., 2012. Measurement of mRNA abundance using RNA-seq data: RPKM
592 measure is inconsistent among samples. *Theory Biosciences*. 131, pp.281–285.

593 West, S.A., Griffin, A.S., Gardner, A., Diggle, S.P., 2006. Social evolution theory for microorganisms.
594 *Nature Reviews Microbiology*, 4, pp. 597-607.

595 Yang, Q., Zhang, M., Zhang, M., Wang, C., Liu, Y., Fan, X., Li, H., 2018. Characterization of a Novel, Cold-
596 Adapted, and Thermostable Laccase-Like Enzyme With High Tolerance for Organic Solvents and Salt
597 and Potent Dye Decolorization Ability, Derived From a Marine Metagenomic Library. *Frontiers in*
598 *Microbiology*. 9, pp.2998

599

600 Ye, M., Li, G., Liang, W.Q. and Liu, Y.H., 2010. Molecular cloning and characterization of a novel
601 metagenome-derived multicopper oxidase with alkaline laccase activity and highly soluble expression.
602 *Applied Microbiology and Biotechnology*, 87(3), pp. 1023-1031.

603 Young, J.M., Skvortsov, T., Kelleher, B.P., Mahaney, W.C., Somelar, P., Allen, C.C.R, 2019. Effect of soil
604 horizon stratigraphy on the microbial ecology of alpine paleosols. *Science of The Total Environment*.
605 657, pp. 1183-1193

606

607

Figure 1. Principal coordinates analysis, of Alpine soil physical-chemical characteristics. A total of 26 Soil Abiotic variables were compared, (supplementary table 1.). Variables were log transformed and normalised such that the sum of squares for each sample = 1. Sample dissimilarities were calculated by their Euclidean distances. PcoA was performed using the capscale function in the package vegan (Oksanen et al., 2016) in R version 3.4.1 (R Core Team, 2011), The first two principle co-ordinate axes were used to ordinate the samples in 2d space. Samples are coloured by their soil horizon classification and convex hulls are displayed for each horizon.

608

609

Figure 2. Laccase genes by Type Comparison of relative abundances of laccases between stratified soil horizons. Relative abundances of metagenome derived type B 2 domain, type C 2 domain, Big 3 domain, small 3 domain, cot A 3 domain respectively. Metagenomic proteins were classified using previously published Hidden Markov Models (HMMs) (Ausec et al., 2011). Abundances are expressed as TPMs. The statistical significance of differences in mean abundances between soil horizon was assessed by Kruskal-Wallis one way ANOVA (Kruskal & Wallis, 1952), the resulting p values are displayed on each plot.

Figure 3. Total relative abundance for each of the 323 putative identified laccase genes was performed across all sample microbiomes. Error bars represent standard error of the mean abundance across all samples. The laccase genes identified in each sample were annotated using the package Kaiju (Menzel et al., 2016). Statistical analysis of each enzyme subgroup across the metagenome datasets was performed using Kruskal-Wallis one way ANOVA (Kruskal & Wallis, 1952). The distinction between type B and C 2-domain laccase is from Nakuamura et al. (2003). The designation of small and large 3-domain laccases comes from Machczynski et al. (2009). The CotA 3-Domain laccases were described by Martin et al. (2015).

632

633

634

635

636

637

638

639

640

641

Figure 4. PCA plots for each laccase class abundance plus summed abundances (Phylum level): Laccase genes were taxonomically classified at the Phylum level using the lowest common ancestor algorithm. For each sub-class, Percentage abundances were normalized by Hellinger transformation, sample dissimilarities were calculated by their Euclidean distances. PcoA was performed using the capscale function in the package vegan (Oksanen et al., 2016) in R version 3.4.1 (R Core Team, 2011), The first two principle co-ordinate axes were used to ordinate the samples in 2d space. Samples are coloured by their soil horizon classification and convex hulls are displayed for each horizon.

642

643

644 **Figure 5.** Maximum likelihood phylogenetic tree of alpine metagenome-derived LMCOs with reference
645 sequences of bacterial two domain laccases that demonstrated activity. The sequences are type B 2
646 domain laccases amplified using the V10_lac PCR primers as described in the text, against DNA
647 extracted from separate V10 Bw and Cox samples (hence designation B or C). Sample LacM (Ausec et
648 al., 2017) was also included as the first acidobacterial metagenome-derived LMCO, though it is a three
649 domain laccase. The tree was drawn using Mega 7.0 (Kumar et al., 2016). The reference sequence
650 shown were downloaded from the NCBI database. The GSLac sequences are representatives of the
651 metagenome-assembled putative laccase genes referred to in 3.2, while the LacB and C samples are
652 sequences obtained after PCR amplification from Bw and Cox horizons I section 3.3.

653

655

656 **Figure 6:** Specific activities of heat treated partially purified metagenome-derived laccase-like multi
 657 copper oxidases (LMCOs) from *E.coli* clone cell free extracts. The enzyme activity against these
 658 substrates was also compared with a vector control for the *E.coli* expression system (data not shown)
 659 where little activity was detectable (<0.01 U/mg for both substrates). Activity measurements are
 660 measurements of triplicate analysis (error bars show standard deviation). See methods section for
 661 assay details.

662

663

664

665 **Figure 7:** Effect of increasing concentrations of CuSO₄ on the cloned laccase LacB6. Values are fold
666 increase at different concentrations relative to the specific activity of 2.5 U/mg recorded in the
667 absence of copper, where relative activity is set as 1. Activity measurements are measurements of
668 triplicate analysis (error bars show standard deviation). See methods section for assay details.

669

670