



**QUEEN'S
UNIVERSITY
BELFAST**

Survival for oesophageal, stomach and small intestine cancers in Europe 1999-2007: Results from EUROCARE-5

Anderson, L. A., Tavilla, A., Brenner, H., Luttmann, S., Navarro, C., Gavin, A. T., Holleczech, B., Johnston, B. T., Cook, M. B., Bannon, F., & Sant, M. (2015). Survival for oesophageal, stomach and small intestine cancers in Europe 1999-2007: Results from EUROCARE-5. *European Journal of Cancer*, 51(15), 2144-2157.
<https://doi.org/10.1016/j.ejca.2015.07.026>

Published in:
European Journal of Cancer

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2015 Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>, which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

Survival for oesophageal, stomach and small intestine cancers: results from EUROCARE-5.

Key words: oesophageal, stomach, small intestine, survival, Europe

L. A. Anderson^a, A. Tavilla^b, H. Brenner^{c,d}, S. Luttmann^e, C. Navarro^{f,g,h}, A.T. Gavin^{a,i},
B. Hollecze^j, B.T. Johnston^k, M.B. Cook^l, F. Bannon^a, M. Sant^m and the EUROCARE-5
Working Group¹⁴.

^a Centre for Public Health, School of Medicine, Dentistry and Biomedical Science,
Queen's University Belfast, Northern Ireland.

^b National Center of Epidemiology, Italian National Institute of Health, Rome, Italy.

^c Division of Clinical Epidemiology and Aging Research and Division of Preventive
Oncology, German Cancer Research Center (DKFZ), Heidelberg, Germany.

^d German Cancer Consortium (DKTK), German Cancer Research Center (DKFZ),
Heidelberg, Germany.

^e Bremen Cancer Registry, Leibniz-Institute for Prevention Research and Epidemiology –
BIPS, Bremen, Germany.

^f Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca, Murcia,
Spain.

^g CIBER Epidemiología y Salud Pública (CIBERESP), Spain.

^h Department of Health and Social Sciences, Universidad de Murcia, Murcia, Spain.

ⁱ Northern Ireland Cancer Registry, Queen's University Belfast, Northern Ireland.

^j Saarland Cancer Registry, Präsident Baltz Straße 5, 66119 Saarbrücken, Germany.

^k Belfast Health and Social Care Trust, Belfast, Northern Ireland.

^l Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Maryland, USA.

^m Analytical Epidemiology and Health Impact Unit. Department of Preventive and Predictive Medicine, Fondazione IRCCS Istituto Nazionale dei Tumori Via Venezian 1, 20133, Milan, Italy.

¹⁴**EUROCARE-5 Working Group:** **Austria:** M. Hackl, N. Zielonke (*Austrian CR*); W. Oberaigner (*Tyrol CR*); **Belgium:** E. Van Eycken, K. Henau (*Belgian CR*); **Bulgaria:** Z. Valerianova, N. Dimitrova (*Bulgarian CR*); **Croatia:** M. Sekerija (*Croatian CR*); **Czech Republic:** M. Zvolský, L. Dušek (*Czech National CR*); **Denmark:** H. Storm, G. Engholm (*Danish Cancer Society*); **Estonia:** M. Mägi (*Estonian CR*); T. Aareleid (*National Institute for Health Development, Tallinn*); **Finland:** N. Malila, K. Seppä (*Finnish CR*); **France:** M. Velten (*Bas Rhin CR*); X. Troussard (*Basse Normandie Haematological malignancies CR*); V. Bouvier, G. Launoy (*Calvados Digestive Tract Registry*); A. V. Guizard (*Calvados, General CR*); J. Faivre*, A. M. Bouvier (*Côte d'Or Digestive Tract Registry, Burgundy*); P. Arveux (*Côte d'Or Gynaecologic CR*); M. Maynadié (*Côte d'Or Haematological Malignancies CR*); A. S. Woronoff (*Doubs CR*); M. Robaszkiewicz (*Finistère Digestive Tract Registry*); I. Baldi (*Gironde CNS CR*); A. Monnereau (*Gironde Haematological Malignancies CR*); B. Tretarre (*Hérault CR*); N. Bossard* (*Hospices Civils de Lyon*); A. Belot (*Hospices Civils de Lyon, FRANCIM*); M. Colonna* (*Isère CR*); F. Molinié (*Loire-Atlantique CR*); S. Bara (*Manche CR*); C. Schwartz (*Marne & Ardennes, Thyroid CR*); B. Lapôtre-Ledoux (*Somme CR*); P. Grosclaude (*Tarn CR*); **Germany:** M. Meyer (*Bavaria CR*); R. Stabenow (*Berlin CR; Brandenburg CR; Mecklenburg-West Pomerania CR; Saxony CR; Saxony-Anhalt CR; Thüringen CR*); S. Luttmann, A. Eberle (*Bremen CR, Leibniz Institute for Prevention Research and Epidemiology*); H. Brenner (*German Cancer Research Center*); A. Nennecke (*Hamburg CR*); J. Engel, G. Schubert-Fritschle (*Munich CR*); J. Kieschke (*Niedersachsen CR*); J. Heidrich (*North Rhine-Westphalia CR*); B. Holleczeck (*Saarland CR*); A. Katalinic* (*Schleswig-Holstein CR*); **Iceland:** J. G. Jónasson, L. Tryggvadóttir (*Icelandic CR*); **Ireland:** H. Comber (*National Cancer Registry Ireland*); **Italy:** G. Mazzoleni, A. Bulatko (*Alto Adige CR*); C. Buzzoni (*Associazione Italiana Registri Tumori*); A. Giacomini (*Biella CR*); A. Suter Sardo, P. Mancuso (*Catanzaro CR*); S. Ferretti (*Ferrara CR*); E. Crocetti, A. Caldarella (*Firenze-Prato CR*); G. Gatta*, M. Sant*, H. Amash, C. Amati, P. Baili*, F. Berrino*, S. Bonfarnuzzo, L. Botta, F. Di Salvo, R. Foschi, C. Margutti, E. Meneghini, P. Minicozzi,

A. Trama (*Fondazione IRCCS Istituto Nazionale dei Tumori, Milan*); D. Serraino, L. Dal Maso (*Friuli Venezia Giulia CR, CRO Aviano National Cancer Institute*); R. De Angelis*, M. Caldora, R. Capocaccia*, E. Carrani, S. Francisci*, S. Mallone, D. Pierannunzio, P. Roazzi, S. Rossi, M. Santaquilani, A. Tavilla (*Istituto Superiore di Sanità, Rome*); F. Pannozzo, S. Busco (*Latina CR*); L. Bonelli, M. Vercelli (*Liguria CR, IRCCS AOU SM-IST*); V. Gennaro (*Liguria Mesothelioma CR*); P. Ricci (*Mantova CR*); M. Autelitano, G. Randi (*Milano CR*); M. Ponz De Leon (*Modena colorettali CR*); C. Marchesi, C. Cirilli (*Modena CR*); M. Fusco, M. F. Vitale (*Napoli 3 South CR*); M. Usala (*Nuoro CR*); A. Traina, R. Staiti (*Palermo Breast CR*); F. Vitale, B. Ravazzolo (*Palermo CR*); M. Michiara (*Parma CR*); R. Tumino (*Ragusa CR*); P. Giorgi Rossi, E. Di Felice (*Reggio Emilia CR*); F. Falcini (*Romagna CR*); A. Iannelli (*Salerno CR*); O. Sechi, R. Cesaraccio (*Sassari CR*); S. Piffer (*Servizio Epidemiologia Clinica e Valutativa, Trento*); A. Madeddu, F. Tisano (*Siracusa CR*); S. Maspero, A. C. Fanetti (*Sondrio CR*); R. Zanetti, S. Rosso (*Torino CR*); P. Candela, T. Scuderi (*Trapani CR*); F. Stracci, F. Bianconi (*Umbria CR*); G. Tagliabue, P. Contiero (*Varese Province CR, Fondazione IRCCS Istituto Nazionale dei Tumori*); A. P. Dei Tos, S. Guzzinati (*Veneto CR*); **Latvia**: S. Pildava (*Latvian CR*); **Lithuania**: G. Smailyte (*Lithuanian CR*); **Malta**: N. Calleja, D. Agius (*Malta National Cancer Registry, Health Information and Research*); **Norway**: T. B. Johannesen (*Norwegian CR*); **Poland**: J. Rachtan (*Cracow CR*); S. Gózdź, R. Mężyk (*Kielce CR*); J. Błaszczyk, M. Bębenek (*Lower Silesia CR, Wroclaw*); M. Bielska-Lasota* (*National Institute of Public Health- NIH, Warszawa*); **Portugal**: G. Forjaz de Lacerda (*Açores CR*); M. J. Bento, C. Castro (*Northern Portugal CR*); A. Miranda, A. Mayer-da-Silva (*Southern Portugal CR*); **Romania**: F. Nicula, D. Coza (*The Oncology Institute "I. Chiricuta", Cluj-Napoca*); **Slovakia**: C. Safaei Diba (*Slovakian National CR*); **Slovenia**: M. Primic-Zakelj* (*Cancer Registry of Republic of Slovenia*); **Spain**: E. Almar, C. Ramírez (*Albacete. Castilla-La Mancha CR*); M. Errezola, J. Bidaurrezaga (*Basque Country CR*); A. Torrella-Ramos (*Castellón-Valencia CR, breast*); J. M. Díaz García, R. Jimenez-Chillaron (*Cuenca CR*); R. Marcos-Gragera*, A. Izquierdo Font (*Girona CR*); M. J. Sanchez, D. Y. L. Chang (*Granada CR, CIBERESP, ibs.Granada*); C. Navarro, M. D. Chirlaque (*Murcia CR, CIBERESP, IMIB-Arrixaca*); C. Moreno-Iribas, E. Ardanaz (*Navarra CR, CIBERESP*); J. Galceran, M. Carulla (*Tarragona CR*); **Sweden**: M. Lambe* (*Regionalt cancercentrum and Karolinska Institutet*); S. Khan (*Swedish CR*); **Switzerland**: M. Mousavi (*Basel CR*); C. Bouchardy, M. Usel (*Geneva CR*); S. M. Ess, H. Frick (*Grisons-Glarus CR*); M. Lorez (*NICER*); S. M. Ess, C. Herrmann (*St. Gallen CR*); A. Bordoni, A. Spitale (*Ticino CR*); I. Konzelmann (*Valais CR*); **The Netherlands**: O. Visser, R. Verhoeven (*The Netherlands CR*); **UK-England**: M. Coleman*, C. Allemani, B. Rachet (*London School of Hygiene and Tropical Medicine*); J. Verne*, N. Easey, G. Lawrence, T. Moran, J. Rashbass, M. Roche, J. Wilkinson (*Public Health England*); **UK-Northern Ireland**: A. Gavin, C. Donnelly (*Northern Ireland CR*); **UK-Scotland**: D. H. Brewster (*Scottish CR*); **UK-Wales**: D. W. Huws, C. White (*Welsh Cancer Intelligence and Surveillance Unit*); R. Otter*.

*EUROCARE Steering Committee

Corresponding author details:

Dr Lesley Anderson

Centre for Public Health

Queen's University Belfast

Institute for Clinical Sciences Block B

Grosvenor Road

Belfast

BT12 6BJ

Tel: ++44 (0)28 9063 2315

Fax: ++44 (0)28 9023 5900

E-mail: l.anderson@qub.ac.uk

Acknowledgements: We thank Chiara Margutti, Simone Bonfarnuzzo and Camilla Amati for secretarial assistance.

Role of funding source

The study was funded by the Compagnia di San Paolo, the Fondazione Cariplo Italy, the Italian Ministry of Health (Ricerca Finalizzata 2009, RF-2009-1529710) and the European Commission (European Action Against Cancer, EPAAC, Joint Action No20102202). The Northern Ireland Cancer Registry is supported by the Public Health Agency for N. Ireland. Dr Michael Cook is funded by US Federal Funds. The Compagnia di San Paolo, the Fondazione Cariplo Italy, the Italian Ministry of Health (Ricerca Finalizzata 2009, RF-2009-1529710) and the European Commission (European Action Against Cancer, EPAAC, Joint Action No20102202). The funding sources had no role in study design, the collection, analysis or interpretation of data, the writing of the report, or the decision to submit the article for publication.

Abstract

Background

European regional variation in cancer survival was reported in the EUROCARE-4 study for patients diagnosed in 1995-1999. Relative survival (RS) estimates are here updated for patients diagnosed with cancer of the oesophagus, stomach, and small intestine from 2000 to 2007. Trends in RS from 1999-2001 to 2005-2007 are presented to monitor and discuss improvements in patient survival in Europe.

Materials and Methods

EUROCARE-5 data from 29 countries (87 cancer registries) were used to investigate 1- and 5-year RS. Using registry-specific life-tables stratified by age, gender, and calendar year, age-standardised 'complete analysis' RS estimates by country and region were calculated for Northern, Southern, Eastern and Central Europe, and for Ireland and United Kingdom (UK). Survival trends of patients in periods 1999-2001, 2002-2004, and 2005-2007 were investigated using the 'period' RS approach. We computed the 5-year RS conditional on surviving the first year (5-year conditional survival), as the ratio of age-standardised 5-year RS to 1-year RS.

Results

Oesophageal cancer 1- and 5-year RS (40% and 12%, respectively) remained poor in Europe. Patient survival was worst in Eastern (8%), Northern (11%), and Southern Europe (10%). Europe-wide, there was a 3% improvement in oesophageal cancer 5-year survival by 2005-2007, with Ireland and the UK (3%), and Central Europe (4%) showing large improvements.

Europe-wide, stomach cancer 5-year RS was 25%. Ireland and UK (17%) and Eastern Europe (19%) had the poorest 5-year patient survival. Southern Europe had the best 5-year survival (30%), though only showing an improvement of 2% by 2005-2007.

Small intestine cancer 5-year RS for Europe was 48%, with Central Europe having the best (54%), and Ireland and UK the poorest (37%). Five-year patient survival improvement for Europe was 8% by 2005-2007, with Central, Southern, and Eastern Europe showing the greatest increases ($\geq 9\%$).

Conclusions

Survival for these cancer sites, particularly oesophageal cancer, remains poor in Europe with wide variation. Further investigation into the wide variation, including analysis by histology and anatomical sub-site, will yield insight to better monitor and explain the improvements in survival observed over time.

Introduction

This article focuses on European relative survival (RS) estimates and trends for oesophageal, stomach and small intestine cancer patients, diagnosed up to 2007, with follow-up to December 31st 2008, as part of EURO CARE-5. Regional variation in RS estimates throughout Europe has been consistently reported for cancer patients, including upper gastrointestinal tract cancers, diagnosed in 1990-1994 [1], 1995-1999 [2] and 1999-2007 [3].

Oesophageal cancer ranks as the eighth most common cancer worldwide with approximately 5 cases per 100,000 diagnosed in Europe annually [4]. Two main histological subtypes, adenocarcinoma (OAC) and squamous cell carcinoma (OSCC), display regional variation in incidence across Europe [5]. Stomach cancer is the third most common cause of cancer death globally [6]. Wide variation in stomach cancer incidence across Europe has been reported with recent declines in most European countries as a result of lifestyle changes, *Helicobacter pylori* detection and cancer treatment. Incidence of non-cardia tumors is high in Southern Europe [7] which, correspondingly, has the best 5-year patient survival [3]. While the small intestine comprises 90% of the length of the bowel, small intestine cancers are rare with an age-standardised incidence rate of 2 per 100,000 person-years in the USA [8] with lower incidence rates reported within Europe [9]. Small intestine cancers exhibit a diverse histology with adenocarcinomas, carcinoid (now classified as neuroendocrine), lymphomas and sarcomas most common [10]. Incidence of small intestine cancers, particularly neuroendocrine malignancies, have increased in the USA [11,12] and Sweden [13], likely as a result of improved detection and classification. Neuroendocrine small intestine cancers are the most common histological subtype and confer superior prognosis

compared to other small intestine entities [12]. Incidence of epithelial small intestine cancers is reportedly highest in Northern and lowest in Eastern Europe [14]; possibly due to geographic differences in diagnostic testing and variable capture by cancer registries.

Methods

Methods used for the analysis of EUROCare-5 data are described in a dedicated paper in this EJC issue [15]. Briefly, survival data were obtained from 29 countries, 21 with 100% national coverage, from 87 cancer registries. Countries were grouped into Northern, Central, Southern and Eastern Europe and Ireland and UK.

All patients diagnosed with a primary and malignant oesophageal, stomach or small intestine cancer, as identified by topography codes C15, C16 (cardia C16.0 and non-cardia C16.1-C16.6) and C17, respectively, of the International Classification of Diseases for Oncology, 3rd edition (ICD-O-3), diagnosed from 2000-2007 were included. Patients with morphology codes 9590-9989 (ICD-O-3), or who were diagnosed by death certificate only (DCO), autopsy only, or censored with null survival time, were excluded. Patients were not excluded if they had a previous primary tumour. All the registries with less than 13% of DCO (for all cancers combined) were included in the analysis.

One-year RS, 5-year RS and 5-year RS conditional on surviving the first year after diagnosis (5-year conditional) were estimated using the ‘complete’ cohort approach for patients diagnosed 2000-2007 (with follow-up to 2008) stratified by gender and age-group (i.e. 15-44, 45-54, 55-64, 65-74, 75 years or older) as previously described [15]. Age standardised survival [16] and European average estimates [15] are also provided. Survival trends were estimated for countries with cases diagnosed between 1999 and

2007 (n=24 countries) with follow-up to 2008, using the 'period' approach [17] to reliably predict 5-year survival in the years, 1999-2001, 2002-2004, and 2005-2007.

Results

Oesophageal, stomach and small intestine cancers were more common in men than women, Table 1. Some countries in Eastern Europe had a high percentage of DCO cases. Elsewhere in Europe the highest DCO rates were reported in Germany. Mean age at diagnosis for oesophageal, stomach and small intestine cancers ranged from 60.7-71.6, 66.8-73.1 and 60.5-68.9 years, respectively, Table 1.

Oesophageal cancer

European average 1-year age-standardised RS was 39.9%, with 12.4% of patients surviving 5-years, Figure 1. Patients in the Central Europe region, particularly Belgium, had the best survival in Europe while survival was poorest in Eastern Europe. Lithuania and Bulgaria had the lowest 5-year RS estimates. Conditional 5-year survival displayed less heterogeneity across Europe, Figure 1.

Survival, at all follow-up time points investigated, decreased with increasing age, Figure 1. One-, 3- and 5-year age-standardised RS was higher in women than men across all follow-up time points, Figure 1.

Overall oesophageal cancer 5-year age-standardised patient survival improved from 9.9% to 12.6% between 1999-2001 and 2005-2007. Graphs of 5-year RS by region and Europe

overall are presented in Supplement 1. The largest regional improvements in 5-year RS were observed in Ireland and UK and Central Europe with limited improvements observed in Eastern or Southern Europe, (Table 2 and Supplement 1). Similar improvements in patient survival were noted between 1999-2001 and 2002-2004, and between 2002-2004 and 2005-2007 for most regions.

Stomach Cancer

One-year age-standardised RS for stomach cancer patients reached almost 50% with substantial regional variation, see Figure 2. While the Eastern Europe region had the poorest 1-year RS (38.4%), the 5-year RS was lowest in Ireland and UK (17.2%) region, with similar survival across all UK countries. Southern Europe had the best 5-year patient survival (29.6%) in Europe. While Eastern Europe had low 1- and 5-year RS, 5-year conditional survival was better than in Northern Europe, and Ireland and UK. Wide variation among countries was identified in 5-year RS estimates from 11.9% in Bulgaria to 34.5% in Iceland. Survival, at all follow-up time points investigated, decreased with increasing age, and women appeared to fare better than men.

Overall 5-year patient survival increased absolutely by less than 2% points across Europe between 1999-2001 and 2005-2007 (Table 3 and Supplement 2). The most marked improvement in patient survival was in Slovenia from 1999-2001 (RS 20.8%) to 2002-2004 (RS 27.1%), Table 3. Although no change was observed in 5-year RS in Northern Europe, improved patient survival was evident in Denmark and Sweden with a decrease

in 5-year RS observed in Finland. The Netherlands had low RS compared to the rest of Central Europe across all periods.

Southern and Central Europe had better patient survival for cardia and non-cardia cancers than other regions, Table 4. Survival for non-cardia cancer patients was significantly higher than for cardia cancer patients, Table 4. In Eastern Europe, as in Southern and Central Europe, patients with non-cardia cancer predominated, Table 4.

Small Intestine Cancer

Small intestine cancer 1- and 5-year RS was 67.9% and 47.9%, respectively, see Figure 3. Ireland and UK was the region with the worst 1-year patient survival at 58.8%. Croatia was the country with the poorest 1-year RS (53.3%). The Central Europe region had the best 5-year RS for small intestine cancer (53.9%) with the poorest in the Ireland and UK region (36.9%). Wide country variation was identified in 5-year RS from 23.5% in Malta to 58.6% in Switzerland. Five-year conditional survival in patients in Ireland and UK remained significantly below the European average, Figure 3.

European patient survival declined with increasing age. Overall 1-, 3- and 5-year age-standardised RS were slightly higher in women compared to men; particularly evident in younger patients, Figure 3.

Overall 5-year RS increased from 40.5% to 48.7% from 1999-2001 until 2005-2007 (Table 5 and supplement 3). The largest improvements (>10% points) in patient survival

were observed in Italy, Austria, Czech Republic and Finland. All regions, except Ireland and UK, showed a significant increase in survival from 1999-2001 to 2005-2007.

Discussion

European wide variation in patient survival was observed for all three cancer sites investigated between regions. Country-specific patient survival also displayed wide variation with several countries showing inconsistent estimates to their region, including Denmark, the Netherlands, Bulgaria and Croatia. Survival of patients improved modestly from 1999-2001 until 2005-2007 for all cancer sites. Oesophageal and stomach cancer 5-year RS for Europe remained very poor. Small intestine cancer had the best overall 5-year RS in Europe and displayed the largest improvement in patient survival.

Oesophageal cancer

European 1- and 5-year RS for oesophageal cancer patients remained poor (35.8% and 10.6%, respectively). With the exception of Central Europe, which maintained the highest patient survival compared with other European regions as reported in EURO CARE-4 [18], RS in other European regions remained below that reported in the USA [19]. Eastern Europe, where OSCC predominates, continued to have the worst RS. Geographical differences in the proportion of oesophageal cancer patients with histology 'not otherwise specified' between regions may account for some of these disparities (data not shown). Additionally, differences in diagnostic accuracy may also account for regional variation with potential misclassification of gastro-oesophageal tumours [20,21]. Cancer stage is a major predictor of cancer patient survival and differences in stage distribution between countries and regions, as a result of early detection and/or diagnostic

practices, could also account for some of the observed disparity seen in Eastern Europe [22,23].

Five-year RS for oesophageal cancer patients, for Europe as a whole, increased marginally from 9.8% in 1999-2001 to 12.6% in 2005-2007. Central Europe and Ireland and UK demonstrated the most marked improvement. This may be explained by improvements in surgical techniques, adjuvant therapy, earlier diagnosis and/or centralisation of treatment. The trends in Europe in mortality [24] and incidence [25] in oesophageal cancer vary markedly across the countries in the study, but generally there is tight correlation between them, suggesting that improvements in survival are not due to over-diagnosis arising from increased surveillance. Variation in incidence trends may be caused by regional changes in the risk-factor prevalence [26]. Obesity may be increasing the incidence of OAC particularly in northern and western Europe, while reduction in tobacco and alcohol consumption is reducing the incidence of OSCC [26]. The generally better prognosis of patients diagnosed with EAC is not consistent across Europe [18].

Centralisation of treatment has produced a marked improvement in oesophageal cancer patient survival with many European countries introducing such strategies in recent years. Ireland and UK demonstrated comparatively better patient survival improvements for oesophageal cancer than most Northern European countries in both time frames investigated and in line with the centralisation of cancer services for oesophagogastric cancer surgery implemented in the UK in 2001. While hospitals performing more than 40 oesophagectomies annually had lower 30-day postoperative mortality, this may not fully explain regional differences in oesophageal or gastric cancer patient survival [27]. Other factors, as highlighted by the International Benchmarking Partnership, may be important such as late diagnosis, differences in public awareness of cancer symptoms, cancer stage,

morphology and topography, presence of co-morbidities, lifestyle factors such as cigarette smoking, and access to optimum care [28]. Body mass index has also been shown to be a prognostic marker for OSCC [29]. The fact that 5-year conditional patient survival is rather similar across Europe indicates relevant differences in short term mortality and points towards early diagnosis and access to care as important areas to consider with regards to improvement of oesophageal cancer patient treatment and standardisation of care.

Stomach cancer

One- and 5-year RS for stomach cancer patients remained low particularly in comparison to 5-year survival of around 69% achieved in Asia [30]. Compared to Europe, stomach cancer incidence in Asia is high, with a predominance of non-cardia tumours which have better patient survival [31]. Screening programs and more aggressive treatment undoubtedly contribute to the superior survival of patients seen in Asia but similar strategies are unlikely to be cost-effective in comparatively low incidence countries within Europe. Histological and staging variability across Europe may account for some of the differences in stomach cancer patient survival observed between countries. Patient survival improved overall in Europe from 1999-2001 to 2005-2007 particularly in Denmark and the Czech Republic. Both mortality [20] and incidence [32] rates for stomach cancer continue to fall for most countries during the period of this study, suggesting no appreciable surveillance-driven over-diagnosis that could compromise estimated survival improvement. A recent report using data from the World Health Organisation reported lower stomach cancer mortality from 2000 onwards in the UK, the USA, Japan and several European countries [33]. Centralisation of treatment for gastric cancer was implemented in several European countries, including the UK, Denmark and

the Netherlands, in recent years despite reports of no survival benefit [27,34] for patients. While 5-year RS was worst in Ireland and UK, improvements in the most recent time period were observed particularly in Wales and England. While delayed diagnosis, first line treatment, or post-operative mortality could explain the patient survival disadvantage in Ireland and UK, other factors appear to be important given the poor 5-year conditional patient survival. Lifestyle differences such as smoking behaviour, co-morbidities, cancer stage and/or subtype could explain the variability observed across countries.

The decreasing 5-year RS in Finland and Norway could be related to the marked decrease in incidence, mainly affecting distal stomach cancer [35], in these countries. Patients with distal stomach cancer have better prognosis, as presented in this report, and this cancer is more responsive to preventative measures than cancers arising in the cardia or proximal stomach. As an effect of this selective incidence decrease, patient with proximal cancers, who carry a worse prognosis, may have become relatively more frequent over time.

Small intestine cancer

European 1- and 5-year RS for patients with epithelial small bowel carcinomas diagnosed from 1978-2002 were comparatively lower than those reported here for all small intestine cancers, excluding lymphomas [14]. Incidence of epithelial small intestine cancers are similar in Ireland and UK and Northern and Southern Europe [14] despite variation in RS. Differences in cancer stage at diagnosis and subtype throughout Europe could explain the reported variations in patient survival. The EURO CARE-5 data encompasses all small intestine cancer histologies with the exception of lymphomas. Small intestine sarcomas reportedly have worse prognosis than neuroendocrine cancers which have a more favourable outcome [8,36]. Small intestine cancers are notoriously difficult to diagnose

due to their vague symptoms. Delays in diagnosis and treatment of small intestine cancer patients are associated with poorer prognosis [37]. One-year RS was lower in Ireland and UK as previously reported [14], and also in Denmark and several Eastern European countries, suggesting that delayed diagnosis, at patient, primary care or referral stages, might be an important factor. This would not however explain the poorer 5-year conditional survival estimates in Ireland and UK, Denmark and Malta for those patients who survived the first year post diagnosis.

Improved survival is reported across all European regions particularly in Northern, Central and Eastern Europe for small intestine cancer patients. Increasing trends in small intestine cancer incidence has been reported [11,12,13,38,39] but mortality rates have remained stable or slightly increasing [38,39]. Given the low incidence and mortality rates, and the heterogeneity of tumour types, it is difficult to say whether effective therapy has increased patient survival [40]. Recent improvements in treatment of small intestine sarcomas, with the use of tyrosine kinase inhibitors since 2001 [41] may have influenced patient survival. Due to the low incidence of gastrointestinal stromal tumours [42], a rare sarcoma sub-type, the effect on patient survival in large datasets like EUROCORE is difficult to measure without ad hoc analyses.

Detailed discussion of the strengths and limitations of the EUROCORE-5 data are available in the article by Rossi et al. in this issue [15]. Increasing survival trends after 5 years of follow-up were found in patients with poor prognosis cancer and aged 75 year and older for Austria, Croatia, Germany, Poland and Slovakia, and may be related to difficulties in the ascertainment of life status [43] or to DCO proportions [15]. Survival estimates from these countries should be interpreted with caution. However, comparing

individual countries may provide more meaningful assessment of reasons for disparities in patient survival; this is limited, however, for cancers with low incidence estimates such as small intestine and oesophageal cancer as the standard errors become large. In addition, the % DCO statistic for each country and cancer are available in Table 1, and should inform comparisons being made between individual countries' patient survival estimates [44].

Conclusions

This article presents overall patient survival for three anatomical sub-sites: oesophagus, stomach and small intestine. They provide some indication of areas that need further investigation to determine the drivers of the variation in survival of cancer patients across Europe. More in-depth investigation by anatomic sub-site and histology could explain the variability observed and are planned using additional data from EURO CARE-5. The historic nature of these large collaborative studies means that recent developments in early detection, routes to treatment, changes to service provision and new treatment modalities for patients will have had insufficient time to have a visible effect. Continued monitoring of cancer survival across Europe will allow further evaluation of survival differences to further promote the widespread application of effective diagnosis and treatment modalities [45]. In summary, although improvements in survival have been reported for cancers of the oesophagus, stomach and small intestine, survival remains poor with wide variation across Europe.

References

- [1] Sant M, Aareleid T, Berrino F, Bielska Lasota M, Carli PM, Faivre J, et al. EUROCORE-3: survival of cancer patients diagnosed 1990-94--results and commentary. *Ann Oncol* 2003;14 Suppl 5:v61–118.
- [2] Berrino F, De Angelis R, Sant M, Rosso S, Bielska-Lasota M, Lasota MB, et al. Survival for eight major cancers and all cancers combined for European adults diagnosed in 1995-99: results of the EUROCORE-4 study. *Lancet Oncol* 2007;8:773–83. doi:10.1016/S1470-2045(07)70245-0.
- [3] De Angelis R, Sant M, Coleman MP, Francisci S, Baili P, Pierannunzio D, et al. Cancer survival in Europe 1999-2007 by country and age: results of EUROCORE--5-a population-based study. *Lancet Oncol* 2014;15:23–34. doi:10.1016/S1470-2045(13)70546-1.
- [4] Bosetti C, Levi F, Ferlay J, Garavello W, Lucchini F, Bertuccio P, et al. Trends in oesophageal cancer incidence and mortality in Europe. *Int J Cancer* 2008;122:1118–29. doi:10.1002/ijc.23232.
- [5] Arnold M, Soerjomataram I, Ferlay J, Forman D. Global incidence of oesophageal cancer by histological subtype in 2012. *Gut* 2015;64:381–7. doi:10.1136/gutjnl-2014-308124.
- [6] Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2014;136:E359–86. doi:10.1002/ijc.29210.

- [7] Verdecchia A, Corazziari I, Gatta G, Lisi D, Faivre J, Forman D. Explaining gastric cancer survival differences among European countries. *Int J Cancer* 2004;109:737–41. doi:10.1002/ijc.20047.
- [8] Qubaiah O, Devesa SS, Platz CE, Huycke MM, Dores GM. Small intestinal cancer: a population-based study of incidence and survival patterns in the United States, 1992 to 2006. *Cancer Epidemiol Biomarkers Prev* 2010;19:1908–18. doi:10.1158/1055-9965.EPI-10-0328.
- [9] Haselkorn T, Whittemore AS, Lilienfeld DE. Incidence of small bowel cancer in the United States and worldwide: geographic, temporal, and racial differences. *Cancer Causes Control* 2005;16:781–7. doi:10.1007/s10552-005-3635-6.
- [10] Schottenfeld D, Beebe-Dimmer JL, Vigneau FD. The epidemiology and pathogenesis of neoplasia in the small intestine. *Ann Epidemiol* 2009;19:58–69. doi:10.1016/j.annepidem.2008.10.004.
- [11] Tsikitis VL, Wertheim BC, Guerrero MA. Trends of incidence and survival of gastrointestinal neuroendocrine tumors in the United States: a seer analysis. *J Cancer* 2012;3:292–302. doi:10.7150/jca.4502.
- [12] Bilimoria KY, Bentrem DJ, Wayne JD, Ko CY, Bennett CL, Talamonti MS. Small bowel cancer in the United States: changes in epidemiology, treatment, and survival over the last 20 years. *Ann Surg* 2009;249:63–71. doi:10.1097/SLA.0b013e31818e4641.

- [13] Lu Y, Fröbom R, Lagergren J. Incidence patterns of small bowel cancer in a population-based study in Sweden: increase in duodenal adenocarcinoma. *Cancer Epidemiol* 2012;36:e158–63. doi:10.1016/j.canep.2012.01.008.
- [14] Faivre J, Trama A, De Angelis R, Elferink M, Siesling S, Audisio R, et al. Incidence, prevalence and survival of patients with rare epithelial digestive cancers diagnosed in Europe in 1995-2002. *Eur J Cancer* 2012;48:1417–24. doi:10.1016/j.ejca.2011.10.038.
- [15] Rossi S, Baili P, Caldora M, Carrani E, Minicozzi P, Pierannunzio D et al. The EURO CARE-5 database, quality checks and methods of statistical analysis. *Eur J Cancer* 2015.
- [16] Corazziari I, Quinn M, Capocaccia R. Standard cancer patient population for age standardising survival ratios. *Eur J Cancer* 2004;40:2307–16. doi:10.1016/j.ejca.2004.07.002.
- [17] Brenner H, Gefeller O. An alternative approach to monitoring cancer patient survival. *Cancer* 1996;78:2004–10.
- [18] Gavin AT, Francisci S, Foschi R, Donnelly DW, Lemmens V, Brenner H, et al. Oesophageal cancer survival in Europe: a EURO CARE-4 study. *Cancer Epidemiol* 2012;36:505–12. doi:10.1016/j.canep.2012.07.009.
- [19] Ries LAG, Young JL, Keel GE, Eisner MP, Lin YD HM-J, editor. SEER Survival Monograph: Cancer Survival Among Adults: U.S. SEER Program, 1988-2001, Patient and Tumor Characteristics. NIH Pub. N. National Cancer Institute, SEER Program; 2007.

- [20] Buas MF, Vaughan TL. Epidemiology and risk factors for gastroesophageal junction tumors: understanding the rising incidence of this disease. *Semin Radiat Oncol* 2013;23:3–9. doi:10.1016/j.semradonc.2012.09.008.
- [21] Marsman WA, Tytgat GNJ, ten Kate FJW, van Lanschot JJB. Differences and similarities of adenocarcinomas of the esophagus and esophagogastric junction. *J Surg Oncol* 2005;92:160–8. doi:10.1002/jso.20358.
- [22] Walters S, Maringe C, Butler J, Brierley JD, Rachet B, Coleman MP. Comparability of stage data in cancer registries in six countries: lessons from the International Cancer Benchmarking Partnership. *Int J Cancer* 2013;132:676–85. doi:10.1002/ijc.27651.
- [23] Maringe C, Walters S, Butler J, Coleman MP, Hacker N, Hanna L, et al. Stage at diagnosis and ovarian cancer survival: evidence from the International Cancer Benchmarking Partnership. *Gynecol Oncol* 2012;127:75–82. doi:10.1016/j.ygyno.2012.06.033.
- [24] Bosetti C, Bertuccio P, Malvezzi M, Levi F, Chatenoud L, Negri E, et al. Cancer mortality in Europe, 2005-2009, and an overview of trends since 1980. *Ann Oncol*. 2013 Oct;24(10):2657-71.
- [25] Lortet-Tieulent J, Renteria E, Sharp L, Weiderpass E, Comber H, Baas P, et al. Convergence of decreasing male and increasing female incidence rates in major tobacco-related cancers in Europe in 1988-2010. *Eur J Cancer*. 2015 Jun;51(9):1144-63.

- [26] Castro C, Bosetti C, Malvezzi M, Bertuccio P, Levi F, Negri E, et al. Patterns and trends in esophageal cancer mortality and incidence in Europe (1980-2011) and predictions to 2015. *Ann Oncol*. 2014 Jan;25(1):283-90.
- [27] Dikken JL, van Sandick JW, Allum WH, Johansson J, Jensen LS, Putter H, et al. Differences in outcomes of oesophageal and gastric cancer surgery across Europe. *Br J Surg* 2013;100:83–94. doi:10.1002/bjs.8966.
- [28] Forbes LJL, Simon AE, Warburton F, Boniface D, Brain KE, Dessaix A, et al. Differences in cancer awareness and beliefs between Australia, Canada, Denmark, Norway, Sweden and the UK (the International Cancer Benchmarking Partnership): do they contribute to differences in cancer survival? *Br J Cancer* 2013;108:292–300. doi:10.1038/bjc.2012.542.
- [29] Watanabe M, Ishimoto T, Baba Y, Nagai Y, Yoshida N, Yamanaka T, et al. Prognostic impact of body mass index in patients with squamous cell carcinoma of the esophagus. *Ann Surg Oncol* 2013;20:3984–91. doi:10.1245/s10434-013-3073-8.
- [30] Nashimoto A, Akazawa K, Isobe Y, Miyashiro I, Katai H, Kodera Y, et al. Gastric cancer treated in 2002 in Japan: 2009 annual report of the JGCA nationwide registry. *Gastric Cancer* 2013;16:1–27. doi:10.1007/s10120-012-0163-4.
- [31] Yako-Suketomo H, Katanoda K. Comparison of time trends in stomach cancer mortality (1990-2006) in the world, from the WHO mortality database. *Jpn J Clin Oncol* 2009;39:622–3. doi:10.1093/jjco/hyp107.

- [32] Arnold M, Karim-Kos HE, Coebergh JW, Byrnes G, Antilla A, Ferlay J, et al. Recent trends in incidence of five common cancers in 26 european countries since 1988: Analysis of the european cancer observatory. *Eur J Cancer*. 2015 Jun;51(9):1164-87.
- [33] Matsuda A, Matsuda T. Time trends in stomach cancer mortality (1950-2008) in Japan, the USA and Europe based on the WHO mortality database. *Jpn J Clin Oncol* 2011;41:932–3. doi:10.1093/jjco/hyr093.
- [34] Van de Poll-Franse L V, Lemmens VEPP, Roukema JA, Coebergh JWW, Nieuwenhuijzen GAP. Impact of concentration of oesophageal and gastric cardia cancer surgery on long-term population-based survival. *Br J Surg* 2011;98:956–63. doi:10.1002/bjs.7493.
- [35] Schmassmann A, Oldendorf M-G, Gebbers J-O. Changing incidence of gastric and oesophageal cancer subtypes in central Switzerland between 1982 and 2007. *Eur J Epidemiol* 2009;24:603–9. doi:10.1007/s10654-009-9379-y.
- [36] Zeeneldin AA, Saber MM, Seif El-Din IA, Frag SA. Small intestinal cancers among adults in an Egyptian district: a clinicopathological study using a population-based cancer registry. *J Egypt Natl Canc Inst* 2013;25:107–14. doi:10.1016/j.jnci.2013.01.004.
- [37] Bauer RL, Palmer ML, Bauer AM, Nava HR, Douglass HO. Adenocarcinoma of the small intestine: 21-year review of diagnosis, treatment, and prognosis. *Ann Surg Oncol* 1994;1:183–8. 1.

- [38]. Shack LG, Wood HE, Kang JY, Brewster DH, Quinn MJ, Maxwell JD, et al. Small intestinal cancer in England & Wales and Scotland: Time trends in incidence, mortality and survival. *Aliment Pharmacol Ther.* 2006 May 1;23(9):1297-306.
- [39]. Klint A, Engholm G, Storm HH, Tryggvadottir L, Gislum M, Hakulinen T, et al. Trends in survival of patients diagnosed with cancer of the digestive organs in the nordic countries 1964-2003 followed up to the end of 2006. *Acta Oncol.* 2010 Jun;49(5):578-607.
- [40]. Karim-Kos HE, Kiemeny LA, Louwman MW, Coebergh JW, de Vries E. Progress against cancer in the netherlands since the late 1980s: An epidemiological evaluation. *Int J Cancer.* 2012 Jun 15;130(12):2981-9.
- [41]. Demetri GD. Identification and treatment of chemoresistant inoperable or metastatic GIST: experience with the selective tyrosine kinase inhibitor imatinib mesylate (STI571). *Eur J Cancer* 2002;38 Suppl 5:S52-9.
- [42]. Goodman MT, Matsuno RK, Shvetsov YB. Racial and ethnic variation in the incidence of small-bowel cancer subtypes in the United States, 1995-2008. *Dis Colon Rectum* 2013;56:441-8. doi:10.1097/DCR.0b013e31826b9d0a.
- [43]. Andersen MR, Storm HH, Eurocourse Work Package 2 Group. Cancer registration, public health and the reform of the European data protection framework: Abandoning or improving european public health research? *Eur J Cancer.* 2015 Jun;51(9):1028-38.
- [44]. Robinson D, Sankila R, Hakulinen T, Moller H. Interpreting international comparisons of cancer survival: The effects of incomplete registration and the presence of death certificate only cases on survival estimates. *Eur J Cancer.* 2007 Mar;43(5):909-13.

- [45] Baili P, Di Salvo F, Marcos-Gragera R, Siesling S, Mallone S et al. Survival for all cancer patients diagnosed between 1999 and 2007 in Europe: results of EURO CARE-5, a population-based study. *Eur J Cancer* 2015.

Table 1: Number of cases, percentage Death Certificate Only (DCO) cases and mean age at diagnosis (years) for oesophageal, stomach and small intestine cancers by country/region before exclusion of autopsy and DCO cases.

	Oesophagus					Stomach					Small Intestine				
	All				Mean	All				Mean	All				Mean
	cases	Men	Women	DCO ^a	age	cases	Men	Women	DCO ^a	age	cases	Men	Women	DCO ^a	age
Northern EU															
Denmark	3,177	2,242	935	0	68.2	4,200	2,700	1,500	0	68.7	602	303	299	0	66.7
Finland	1,859	1,218	641	1.1	69.2	5,812	3,213	2,599	0.7	70.1	691	367	324	1.4	65.3
Iceland	121	87	34	0	71.6	281	171	110	0	72.5	47	27	20	0	66.2
Norway	1,466	1,041	425	0.8	70.1	4,521	2,717	1,804	0.6	72.5	836	441	395	0.2	66.9
Sweden	3,203	2,300	903	0	70.1	7,863	4,740	3,123	0	72.1	1,819	1,012	807	0	68.9
Ireland and UK															
Ireland	2,706	1,707	999	0.8	69.7	3,701	2,297	1,404	1.6	69.8	392	228	164	1.5	65.3
UK-England	50,610	32,299	18,311	2.6	71.5	55,973	36,023	19,950	3.5	72.9	5,501	2,985	2,516	2.5	68.1
UK-Northern															
Ireland	1,294	817	477	0.7	69.9	1,882	1,129	753	1.3	71.5	189	109	80	1.1	65.7

[illegible]

Bulgaria	1,478	1,152	326	22.5	64.6	14,616	9,005	5,611	20.5	68.2	345	197	148	24.1	63.5
Czech Republic	3,680	3,090	590	5.1	63.6	13,760	7,996	5,764	4.6	69.3	1,019	559	460	3.9	65.4
Estonia	434	355	79	0	64.9	3,277	1,776	1,501	0.2	66.8	90	36	54	0	64.2
Latvia	881	739	142	6.5	63.8	5,324	2,948	2,376	6.6	67.0	114	52	62	11.4	65.8
Lithuania	1,180	1,022	158	4.8	63.1	7,047	4,095	2,952	4.2	67.2	176	83	93	8.5	65.8
Poland ^b	1,353	1,070	283	1.7	63.5	6,253	3,938	2,315	1.6	67.0	230	122	108	0	64.3
Slovakia	1,937	1,732	205	12.5	60.7	6,826	4,111	2,715	12.3	68.1	397	212	185	10.8	64.3

^a Also includes ‘autopsy-only’ basis of diagnosis. ^b Pooled rates as these countries did not have national coverage.

Table 2: Five-year relative survival (RS) and corresponding 95% confidence interval (CI) of oesophageal cancer in three periods (1999-2001, 2002-2004, 2005-2007) by country, European region and European average, with p-values of differences^a between periods.

	Number of cases analysed across all time periods	1999-2001		2002-2004		2005-2007		2005-2007 vs 1999-2001	
		% RS	95% CI	% RS	95% CI	% RS	95% CI	Abs diff	<i>p-value</i>
Europe	111006	9.9	(9.3-10.5)	11.7	(11.0-12.3)	12.6	(12.0-13.2)	2.7	<0.001
Northern EU	10471	9.1	(8.0-10.4)	11.3	(10.1-12.8)	10.8	(9.7-12.1)	1.7	0.023
Denmark	3401	4.6	(3.3-6.4)	9.0	(7.0-11.5)	9.7	(7.9-11.8)	5.1	<0.001
Finland	1959	9.6	(7.2-12.7)	12.9	(9.9-16.7)	12.1	(9.5-15.2)	2.5	0.108
Iceland ^b	129	-		-	-	-	-	-	-
Norway	1572	8.4	(5.7-12.4)	12.5	(9.5-16.5)	10.9	(8.0-14.8)	2.5	0.150
Sweden	3411	13.3	(11.0-16.0)	12.3	(10.0-15.2)	10.6	(8.8-12.8)	2.7	0.052
Ireland and UK	67862	10.3	(9.8-10.8)	11.9	(11.4-12.4)	13.5	(13.0-14.1)	3.2	<0.001
Ireland	2816	11.9	(9.6-14.7)	15.3	(12.7-18.3)	16.7	(14.2-19.6)	4.8	0.005
England	52786	9.9	(9.4-10.6)	11.5	(10.9-12.1)	13.7	(13.1-14.3)	3.7	<0.001

Northern Ireland ^b	1389	9.6	(7.1-13.0)	14.6	(11.3-18.7)	-	-	-	-
Scotland	7142	10.0	(8.6-11.6)	11.7	(10.1-13.5)	11.1	(9.7-12.7)	1.1	0.157
Wales	3727	14.1	(11.7-17.1)	13.8	(11.6-16.4)	12.6	(10.5-15.0)	1.6	0.188
Central EU	18139	10.8	(9.9-11.8)	13.9	(12.8-15.0)	15.2	(14.2-16.2)	4.3	<0.001
Austria	2711	11.7	(9.4-14.6)	17.9	(15.1-21.2)	17.1	(14.6-20.0)	5.4	0.002
France	3365	13.0	(11.2-15.0)	11.2	(9.6-13.0)	-	-	-	-
Germany ^b	1804	16.1	(12.8-20.4)	13.7	(10.6-17.7)	-	-	-	-
Switzerland	1075	15.3	(11.4-20.5)	18.3	(14.3-23.4)	18.9	(14.6-24.4)	3.6	0.145
The Netherlands	11744	9.6	(8.6-10.8)	13.1	(11.9-14.5)	14.4	(13.3-15.7)	4.8	<0.001
Southern EU	4474	9.7	(8.2-11.6)	10.6	(9.0-12.5)	10.9	(9.2-12.7)	1.1	0.186
Italy	3278	10.7	(8.8-13.2)	12.4	(10.3-14.8)	11.0	(9.1-13.3)	0.3	0.432
Malta ^b	67	-	-	-	-	-	-	-	-
Slovenia ^b	805	7.1	(4.2-12.1)	-	-	8.6	(5.5-13.6)	1.5	0.289
Spain	1792	7.9	(6.1-10.2)	7.9	(6.1-10.2)	-	-	-	-
Eastern EU	10063	7.3	(6.1-8.6)	7.4	(6.3-8.6)	8.1	(7.0-9.3)	0.8	0.175
Bulgaria ^b	1172	-	-	-	-	6.7	(4.2-10.7)	-	-

Czech Republic	3496	7.3	(5.4-9.8)	9.2	(7.0-12.2)	11.4	(9.5-13.7)	4.2	0.003
Estonia ^b	485	-	-	-	-	-	-	-	-
Lithuania ^b	1348	8.4	(5.6-12.5)	4.7	(2.9-7.5)	-	-	-	-
Poland ^b	1474	8.1	(5.3-12.4)	7.7	(5.3-11.3)	6.2	(4.0-9.7)	1.9	0.205
Slovakia ^b	111006	6.4	(3.9-10.3)	10.1	(6.4-15.6)	-	-	-	-

Abs = absolute, Diff = Difference.

^a Survival differences between periods have been assessed by the Z-test.

^b Standardized Survival rates could not be calculated where one or more age specific rates are absent due to small number of cases.

Note: % difference is the relative difference.

Note: Empty fields of RS in France and Spain in 2007 are due to a limitation of analysis to periods 1999-2001 and 2002-2004 only.

Table 3: Five-year relative survival (RS) and corresponding 95% confidence interval (CI) of stomach cancer in three periods (1999-2001, 2002-2004, 2005-2007) by country, European region and European average, with p-values of differences^a between periods.

	Number of cases analysed across all time periods	1999-2001		2002-2004		2005-2007		2005-2007 vs 1999-2001	
		% RS	95% CI	% RS	95% CI	% RS	95% CI	Abs diff	p-value
Europe	232452	23.3	(22.9-23.8)	23.8	(23.4-24.3)	25.1	(24.6-25.6)	1.8	<0.001
Northern EU	26201	22.4	(21.4-23.5)	21.7	(20.7-22.7)	22.7	(21.6-23.8)	0.3	0.360
Denmark	4691	14.0	(12.2-16.2)	14.7	(12.7-16.9)	18.3	(16.3-20.6)	4.3	0.002
Finland	6667	28.5	(26.5-30.8)	25.0	(23.0-27.1)	25.2	(23.1-27.5)	-3.3	0.016
Iceland ^b	341	-	-	-	-	-	-	-	-
Norway	5341	23.5	(21.2-26.1)	21.8	(19.6-24.3)	23.5	(21.1-26.2)	0.0	0.499
Sweden	9152	21.4	(19.8-23.2)	22.6	(20.8-24.5)	22.5	(20.7-24.4)	1.0	0.214
Ireland and UK	83908	16.1	(15.6-16.7)	16.5	(16.0-17.1)	18.2	(17.6-18.8)	2.0	<0.001
Ireland	4056	19.4	(17.1-22.0)	19.4	(17.0-22.0)	21.9	(19.4-24.6)	2.5	0.086
England	64533	16.1	(15.5-16.7)	16.3	(15.7-16.9)	18.0	(17.3-18.7)	1.9	<0.001

Northern Ireland	2189	17.7	(14.7-21.4)	18.8	(15.8-22.4)	18.4	(15.0-22.5)	0.6	0.400
Scotland	7992	14.7	(13.1-16.5)	16.8	(15.1-18.7)	16.3	(14.5-18.3)	1.6	0.104
Wales	5144	16.1	(13.9-18.6)	16.5	(14.3-18.9)	20.0	(17.5-22.9)	4.0	0.015
Central EU	39365	24.0	(23.2-24.9)	24.7	(23.9-25.6)	26.2	(25.3-27.1)	2.1	<0.001
Austria	12740	30.7	(29.1-32.3)	29.8	(28.2-31.5)	33.6	(31.8-35.4)	2.9	0.009
France ^b	4997	25.4	(23.4-27.5)	28.1	(26.1-30.3)	-	-	-	-
Germany	4486	27.2	(24.7-30.0)	27.0	(24.5-29.7)	27.5	(24.9-30.3)	0.3	0.439
Switzerland	2019	25.0	(21.6-29.0)	29.3	(25.5-33.7)	31.4	(27.4-36.0)	6.4	0.014
The Netherlands	18808	18.9	(17.8-20.1)	20.6	(19.4-21.8)	21.1	(19.9-22.3)	2.2	0.005
Southern EU	29234	30.5	(29.4-31.6)	30.4	(29.4-31.5)	32.1	(31.0-33.3)	1.6	0.021
Italy pool	23784	32.7	(31.5-34.0)	31.6	(30.4-32.9)	33.8	(32.5-35.1)	1.1	0.126
Malta ^b	398	-	-	-	-	-	-	-	-
Slovenia	4116	20.8	(18.5-23.5)	27.1	(24.4-30.1)	27.9	(25.3-30.7)	7.0	<0.001
Spain	6569	25.1	(23.5-26.8)	25.9	(24.2-27.7)	-	-	-	-
Eastern EU	53747	17.6	(16.9-18.2)	19.0	(18.3-19.6)	18.8	(18.2-19.5)	1.3	0.004
Bulgaria	12555	10.9	(9.8-12.1)	12.5	(11.3-13.8)	12.8	(11.7-14.0)	2.0	0.010

Czech Republic	14449	18.1	(16.9-19.4)	21.3	(19.9-22.7)	22.6	(21.2-24.0)	4.4	<0.001
Estonia	3852	21.8	(19.3-24.7)	24.8	(22.0-27.8)	22.2	(19.6-25.1)	0.3	0.432
Lithuania	8614	22.0	(20.4-23.8)	23.4	(21.7-25.2)	23.7	(21.7-25.8)	1.7	0.112
Poland	7164	15.2	(13.5-17.0)	16.7	(15.0-28.5)	15.6	(14.0-17.4)	0.4	0.367
Slovakia	7186	21.2	(19.3-23.2)	20.3	(18.6-22.3)	21.1	(19.2-23.1)	-0.1	0.471

Abs = absolute, Diff = Difference.

^a Survival differences between periods have been assessed by the Z-test.

^b Standardized Survival rates could not be calculated where one or more age specific rates are absent due to small number of cases.

Note: % difference is the relative difference.

Note: Empty fields of RS in Spain in 2007 are due to a limitation of analysis to periods 1999-2001 and 2002-2004 only.

Table 4: Age-standardised 1-year, 5-year relative survival, and 5-year relative survival conditional on surviving 1 year, with 95% confidence intervals, for cardia and non-cardia stomach cancers.

	Cardia							Non-cardia						
	No. of	1-year		5-year		Conditional		No. of	1-year		5-year		Conditional	
	cases	% RS	95% CI	% RS	95% CI	% RS	95% CI	cases	% RS	95% CI	% RS	95% CI	% RS	95% CI
Europe	48611	46.0	45.5-46.4	16.0	15.5-16.4	34.0	33.2-34.9	96020	54.6	54.3-54.9	30.5	30.1-30.9	66.3	65.9-66.8
Northern EU	5299	43.3	41.9-44.7	14.1	12.9-15.3	32.5	30.0-35.1	6027	55.2	53.8-56.5	28.6	27.1-30.2	51.9	49.4-54.5
Denmark	1687	41.0	38.6-43.4	12.8	10.8-15.0	31.2	26.8-36.2	1149	48.0	44.9-51.0	25.9	22.8-29.2	53.9	48.4-60.1
Finland	936	45.9	42.5-49.2	16.0	13.2-19.1	34.9	29.5-41.4	272	57.8	51.3-63.7	-	-	-	-
Iceland	41	31.0	18.0-44.8	10.3	3.4-19.1	33.2	15.2-72.5	70	74.7	62.7-83.0	-	-	-	-
Norway	967	43.9	40.6-47.1	15.4	12.7-18.5	35.2	29.6-42.0	1831	58.0	55.4-60.5	30.0	27.2-32.9	51.8	47.7-56.2
Sweden	1668	44.2	41.7-46.6	13.6	11.6-15.7	30.7	26.6-35.5	2705	55.9	53.7-58.0	28.0	25.6-30.4	50.1	46.3-54.1
Ireland and UK	19244	46.6	45.8-47.3	14.4	13.8-15.1	31.0	29.7-32.4	17457	48.4	47.5-49.3	23.2	22.3-24.0	47.9	46.3-49.5
Ireland	986	42.2	39.0-45.3	17.0	14.1-20.0	40.2	34.3-47.1	1705	46.2	43.8-48.7	24.4	21.9-27.0	52.8	48.3-57.7
England	14510	47.4	46.5-48.2	14.8	14.1-15.5	31.2	29.8-32.70	11932	49.2	48.1-50.3	23.1	22.0-24.2	46.9	45.1-48.9
Northern Ireland	462	46.4	41.6-51.1	16.2	12.3-20.7	35.0	27.5-44.5	534	42.2	37.4-47.0	21.2	16.9-25.8	50.1	41.9-59.9
Scotland	1983	42.9	40.5-45.3	11.3	9.6-13.2	26.4	22.7-30.7	1787	47.6	44.9-50.2	23.4	20.8-26.1	49.2	44.6-54.4
Wales	1303	46.7	43.8-49.6	13.2	10.9-15.8	28.4	23.8-33.8	1499	47.3	44.2-50.4	22.3	19.4-25.4	47.2	41.9-53.1
Central EU	13230	49.2	48.4-50.1	18.2	17.4-19.0	37.0	35.4-38.5	26709	60.9	60.3-61.5	36.0	35.2-36.7	59.1	58.0-60.1
Austria	1297	50.5	47.7-53.3	22.6	19.8-25.5	44.7	39.9-50.1	1895	65.9	63.5-68.1	40.2	37.3-43.1	61.0	57.3-65.0
Belgium	1264	55.9	53.1-58.7	20.5	17.7-23.5	36.7	32.1-41.9	1504	63.0	60.3-65.7	35.6	32.3-38.8	56.4	52.0-61.2

France	1384	50.6	47.9-53.3	14.7	12.6-17.0	29.0	25.3-33.4	2949	58.6	56.6-60.6	32.8	30.6-35.0	56.0	52.9-59.2
Germany	4506	52.7	51.2-54.2	22.3	20.7-23.9	42.3	39.7-45.1	11906	64.4	63.5-65.3	40.1	39.0-41.3	62.3	60.8-63.9
Switzerland	494	52.0	47.4-56.5	-	-	-	-	1145	64.1	61.1-67.0	40.8	37.2-44.3	63.6	59.1-68.4
The Netherlands	4285	42.4	40.9-43.9	13.1	11.9-14.5	31.0	28.3-34.0	7310	53.7	52.5-54.9	28.9	27.5-30.2	53.8	51.6-56.0
Croatia	476	42.43	37.7-47.1	27.0	21.9-32.4	63.7	54.2-74.9	446	56.6	51.6-61.4	37.4	31.4-43.5	66.1	57.6-75.8
Southern EU	5793	48.6	47.2-49.9	20.2	18.9-21.5	41.5	39.2-44.0	24960	60.1	59.4-60.7	36.2	35.5-36.9	60.2	59.2-61.3
Italy	3193	52.2	50.3-54.0	20.9	19.1-22.7	40.0	37.0-43.3	15728	61.3	60.5-62.1	37.1	36.2-38.1	60.6	59.2-61.9
Malta	65	35.9	25.5-46.4	-	-	-	-	107	49.2	38.8-58.7	25.4	16.9-34.8	51.7	38.6-69.3
Portugal	890	45.5	42.0-48.9	20.6	17.6-23.9	45.4	39.7-51.9	4188	59.4	57.8-60.9	36.1	34.4-37.9	60.8	58.4-63.4
Slovenia	473	46.2	41.4-50.8	18.3	14.1-22.9	39.6	31.8-49.3	1424	61.2	58.6-63.8	41.0	37.8-44.2	67.0	62.8-71.5
Spain	696	44.7	40.8-48.5	16.9	13.9-20.2	37.9	32.0-44.8	3067	55.4	53.5-57.2	30.8	28.9-32.7	55.6	52.7-58.5
Eastern EU	5045	36.4	35.1-37.8	13.1	12.0-14.4	36.1	33.2-39.2	20867	45.6	44.9-46.3	23.7	23.0-24.4	52.0	50.6-53.4
Bulgaria	1273	25.3	22.9-27.8	7.6	5.8-9.9	30.2	23.5-38.7	6460	33.8	32.6-35.0	14.2	13.1-15.4	42.0	39.1-45.2
Czech Republic	1596	41.4	38.9-43.9	15.6	13.4-18.0	37.7	32.9-43.2	5927	49.5	48.2-50.9	27.5	26.0-29.0	55.5	53.0-58.1
Estonia	262	41.2	35.0-47.3	18.4	12.9-24.7	44.7	33.6-59.3	1760	50.9	48.4-53.2	29.1	26.3-31.8	57.2	52.7-62.0
Latvia	266	32.8	27.0-38.7	15.9	10.8-21.9	48.4	35.8-65.5	1012	44.1	40.9-47.3	22.8	19.4-26.3	51.7	45.2-59.0
Lithuania	312	42.2	36.3-47.9	15.4	11.2-20.3	36.6	28.0-47.6	2218	54.6	52.4-56.7	31.3	28.8-33.7	57.3	53.5-61.3
Poland	627	41.6	37.7-45.5	12.8	9.6-16.5	30.8	23.9-39.6	398	47.9	42.8-52.9	17.4	12.6-22.9	36.4	27.5-48.1
Slovakia	709	38.9	35.1-42.7	13.6	10.5-17.0	34.8	28.0-43.3	3092	53.9	52.0-55.7	28.7	26.7-30.7	53.2	50.1-56.5

Table 5: Five-year relative survival (RS) and corresponding 95% confidence interval (CI) of small intestine cancer in three periods (1999-2001, 2002-2004, 2005-2007) by country, European region and European average, with p-values of differences* between periods.

	Number of cases analysed across all time periods	1999-2001		2002-2004		2005-2007		2005-2007 vs 1999-2001	
		% RS	95% CI	% RS	95% CI	% RS	95% CI	Abs diff	p-value
Europe	18116	40.5	(38.5-42.7)	45.8	(43.9-47.9)	48.7	(46.9-50.5)	8.1	<0.001
Northern EU	4021	49.9	(46.7-53.3)	50.7	(47.7-53.9)	55.8	(53.0-58.8)	6.0	0.004
Denmark	626	37.4	(30.3-46.1)	37.7	(30.4-46.6)	39.6	(33.2-47.1)	2.2	0.341
Finland	678	51.7	(43.7-61.2)	55.4	(48.8-62.9)	62.1	(54.9-70.3)	10.4	0.040
Iceland ^b	49	-	-	-	-	-	-	-	-
Norway	834	52.6	(45.7-60.6)	51.6	(45.4-58.6)	56.6	(50.5-63.3)	3.9	0.216
Sweden	1835	52.0	(47.5-57.0)	53.3	(48.9-58.1)	59.5	(55.3-64.0)	7.5	0.011
Ireland and UK	7178	35.3	(33.1-37.6)	36.1	(34.1-38.3)	37.7	(35.7-39.8)	2.4	0.058
Ireland	376	35.5	(26.2-48.1)	44.7	(36.3-55.2)	42.8	(34.7-52.9)	7.3	0.154
England	5539	34.4	(31.9-37.0)	35.1	(32.8-37.5)	37.7	(35.5-40.2)	3.4	0.027

Northern Ireland	230	37.4 (27.9-50.2)	33.9 (23.0-50.0)	43.5 (30.6-61.9)	6.1	0.263
Scotland	639	39.6 (32.7-48.0)	37.5 (30.1-46.7)	38.4 (31.9-46.1)	-1.3	0.405
Wales	401	38.9 (28.6-52.7)	38.2 (29.6-49.3)	33.3 (25.9-42.8)	-5.6	0.227
Central EU	3399	44.1 (40.7-47.8)	47.8 (44.5-51.2)	53.0 (50.0-56.3)	9.0	<0.001
Austria	899	43.6 (37.3-50.9)	52.3 (46.0-59.4)	55.7 (50.1-61.9)	12.1	0.004
France	572	45.3 (39.3-52.3)	48.3 (42.2-56.2)	- -	-	-
Germany	323	42.0 (31.6-56.0)	45.6 (34.9-59.6)	50.1 (41.2-61.0)	8.1	0.154
Switzerland	294	54.7 (45.0-66.5)	59.9 (50.4-71.0)	55.4 (45.4-67.5)	0.7	0.467
The Netherlands	1737	44.5 (39.6-50.0)	43.3 (39.0-48.2)	51.5 (47.2-56.3)	7.1	0.022
Southern EU	1570	39.5 (34.7-44.9)	49.0 (44.3-54.2)	49.7 (45.5-54.3)	10.2	0.001
Italy	1338	38.7 (33.5-44.6)	48.7 (43.7-54.4)	51.1 (46.4-56.3)	12.5	<0.001
Malta ^b	34	- -	- -	- -	-	-
Slovenia ^b	153	- -	48.5 (33.8-69.6)	- -	-	-
Spain	347	46.3 (38.4-56.0)	42.4 (35.9-50.1)	- -	-	-
Eastern EU	1951	34.5 (29.4-40.4)	43.4 (38.9-48.4)	43.5 (39.6-47.8)	9.1	0.005
Bulgaria	248	41.9 (26.0-67.6)	35.8 (25.1-51.0)	- -	-	-

Czech Republic	849	35.8 (28.3-45.4)	46.1 (39.4-54.0)	46.9 (41.0-53.6)	11.0	0.020
Estonia ^b	95	- -	- -	- -	-	-
Lithuania ^b	186	- -	- -	32.9 (22.0-49.4)	-	-
Poland ^b	225	- -	- -	44.7 (33.7-59.1)	-	-
Slovakia	368	51.4 (39.4-67.1)	42.0 (32.2-54.7)	46.2 (37.6-56.8)	-5.2	0.269

Abs = absolute, Diff = Difference.

^a Survival differences between periods have been assessed by the Z-test.


^b Standardized Survival rates could not be calculated where one or more age specific rates are absent due to small number of cases.

Note: % difference is the relative difference.

Note: Empty fields of RS in France and Spain in 2007 are due to a limitation of analysis to periods 1999-2001 and 2002-2004 only.


Supplement 1: Age-standardised 5-year RS for oesophageal cancer in a) Northern Europe, b) UK and Ireland, c) Central Europe, d) Southern Europe, e) Eastern Europe and f) European average.

a)




Note: Iceland removed as no points are available due to small number of cases


b)



c)




d)




Note: Malta removed as no points are available due to small number of cases

e)




f)




Supplement 2: Age-standardised 5-year RS for stomach cancer in a) Northern Europe, b) Ireland and UK, c) Central Europe, d) Southern Europe, e) Eastern Europe and f) European average.

a)




Note: Iceland removed as no points are available due to small number of cases


b)



c)




d)




Note: Malta removed as no points are available due to small number of cases.

e)



Note: Estonia removed as no points are available due to small number of cases.


f)



Ref supplement 2f previously presented de Angelis *et al.*, 2014[3]


Supplement 3: Age-standardised 5-year RS for small intestine cancer in a)Northern Europe,
b) Ireland and UK, c) Central Europe, d) Southern Europe, e) Eastern Europe and f)
European average.

a)




Note: Iceland removed as no points are available due to small number of cases


b)



c)




d)




Note: Malta removed as no points are available due to small number of cases.

e)



Note: Estonia removed as no points are available due to small number of cases

f)




Oesophagus

Age-standardised 1-year, 5-year relative survival, and 5-year relative survival conditional to surviving 1 year, with 95% confidence intervals in parentheses

	Number of cases	1-year			5-year			Conditional		
Northern Europe	9,654	35.8(34.8-	36.9)	10.6(9.8-	11.5)	29.6(27.5-	31.9)
Denmark	3,165	30.0(28.4-	31.8)	8.9(7.7-	10.3)	29.6(25.8-	33.8)
Finland	1,798	38.1(35.6-	40.6)	12.0(10.1-	14.2)	31.4(26.8-	36.8)
Iceland	120	43.1(34.2-	54.2)	-(-)	-(-)
Norway	1,442	38.4(35.7-	41.2)	10.0(7.9-	12.6)	25.9(20.7-	32.5)
Sweden	3,129	39.2(37.3-	41.2)	11.7(10.2-	13.3)	29.7(26.3-	33.5)
Ireland and UK	63,179	39.8(39.4-	40.2)	12.5(12.1-	12.8)	31.4(30.5-	32.2)
Ireland	2,673	41.0(39.1-	43.0)	15.6(13.9-	17.4)	37.9(34.2-	42.0)
UK, England	49,287	39.7(39.2-	40.2)	12.4(12.0-	12.8)	31.2(30.3-	32.2)
UK, Northern Ireland	1,283	39.0(36.3-	42.0)	16.1(13.7-	18.9)	41.3(35.8-	47.6)
UK, Scotland	6,492	40.0(38.6-	41.3)	11.0(10.0-	12.1)	27.5(25.1-	30.2)
UK, Wales	3,444	40.4(38.6-	42.2)	13.1(11.7-	14.8)	32.5(29.1-	36.3)
Central Europe	32,860	45.5(44.9-	46.1)	15.3(14.8-	15.9)	33.7(32.7-	34.7)
Austria	2,567	42.8(40.8-	44.9)	16.3(14.5-	18.3)	38.0(34.3-	42.2)
Belgium	3,936	51.8(50.2-	53.5)	21.8(20.2-	23.5)	42.0(39.2-	45.0)
France	4,499	46.7(45.2-	48.3)	13.9(12.7-	15.2)	29.7(27.3-	32.2)
Germany	9,038	46.1(45.0-	47.2)	16.2(15.1-	17.3)	35.1(33.0-	37.3)
Switzerland	1,190	49.6(46.6-	52.7)	18.4(15.7-	21.7)	37.2(32.1-	43.2)
The Netherlands	11,630	42.7(41.8-	43.7)	13.0(12.2-	13.8)	30.4(28.7-	32.3)
Southern Europe	12,222	35.5(34.6-	36.4)	10.1(9.5-	10.8)	28.5(27.0-	30.2)
Croatia	1,676	26.4(24.2-	28.8)	10.2(8.3-	12.6)	38.7(32.0-	46.9)
Italy	5,488	40.3(39.0-	41.7)	11.7(10.7-	12.8)	29.0(26.7-	31.5)
Malta	86	32.3(24.2-	43.2)	-(-)	-(-)
Portugal	2,530	33.5(31.6-	35.5)	10.0(8.7-	11.6)	29.9(26.1-	34.1)
Slovenia	712	29.6(26.3-	33.4)	8.3(6.2-	11.2)	28.1(21.4-	36.8)
Spain	1,730	38.2(35.9-	40.7)	9.3(7.8-	11.1)	24.5(20.8-	28.8)
Eastern Europe	9,770	27.0(26.0-	28.0)	7.7(7.0-	8.6)	28.7(26.1-	31.5)
Bulgaria	1,145	17.8(15.4-	20.4)	6.1(4.3-	8.6)	34.4(25.0-	47.3)
Czech Republic	3,283	31.2(29.5-	33.0)	10.6(9.2-	12.2)	34.1(30.0-	38.7)
Estonia	426	28.1(23.7-	33.3)	7.0(4.4-	11.2)	24.9(16.0-	38.7)
Latvia	809	28.4(24.9-	32.4)	-(-)	-(-)
Lithuania	1,121	24.0(21.3-	27.0)	5.7(3.9-	8.3)	23.7(16.6-	33.9)
Poland	1,328	27.2(24.7-	30.1)	6.5(4.8-	8.7)	23.8(18.0-	31.5)
Slovakia	1,658	25.7(23.3-	28.4)	6.9(5.2-	9.2)	26.8(20.4-	35.2)
Europe	127,685	39.9(39.4-	40.4)	12.4(12.0-	12.8)	31.0(30.1-	31.9)

Age-standardised 5-year relative survival (%)




European age-specific and age-standardised observed (obs, %) and relative (rel, %) survival

Age group	Number of cases		1-year	3-year	5-year
15-44	2,995	obs	51.1	26.8	20.9
		rel	51.2	27.0	21.1
45-54	14,298	obs	46.6	19.4	14.4
		rel	46.9	19.7	14.8
55-64	30,054	obs	45.2	18.6	13.5
		rel	45.6	19.2	14.3
65-74	36,782	obs	39.3	16.1	11.2
		rel	40.3	17.5	12.9
75+	43,556	obs	27.0	8.3	4.7
		rel	29.2	10.5	7.2
All cases	127,685	obs	38.8	15.6	10.9
		rel	39.9	16.8	12.4
Men (71%)					
15-44	2,401	obs	49.9	25.7	19.7
		rel	50.0	25.9	19.9
45-54	11,784	obs	46.0	18.6	13.6
		rel	46.2	18.9	14.0
55-64	24,050	obs	44.3	17.9	12.7
		rel	44.8	18.6	13.6
65-74	27,393	obs	39.1	15.5	10.5
		rel	40.2	16.9	12.3
75+	24,766	obs	27.4	8.1	4.6
		rel	29.9	10.5	7.2
All cases	90,394	obs	38.5	15.0	10.3
		rel	39.7	16.3	11.9
Women (29%)					
15-44	594	obs	56.6	32.2	27.6
		rel	56.7	32.3	27.8
45-54	2,514	obs	50.6	24.6	20.8
		rel	50.7	24.8	21.1
55-64	6,004	obs	50.6	23.1	18.8
		rel	50.9	23.5	19.4
65-74	9,389	obs	40.0	19.0	14.1
		rel	40.6	19.8	15.4
75+	18,789	obs	26.4	8.6	5.0
		rel	28.4	10.6	7.1
All cases	37,290	obs	40.9	18.5	14.3
		rel	41.8	19.5	15.5

Age-standardised 1-year, 5-year relative survival, and 5-year relative survival conditional to surviving 1 year, with 95% confidence intervals in parentheses

	Number of cases	1-year			5-year			Conditional		
Northern Europe	22,186	46.9 (46.2-	47.6)	21.9 (21.2-	22.6)	46.6 (45.3-	47.9)
Denmark	4,188	39.4 (37.9-	41.0)	16.0 (14.7-	17.5)	40.6 (37.6-	43.9)
Finland	5,616	49.1 (47.8-	50.5)	25.3 (23.9-	26.7)	51.4 (49.0-	54.0)
Iceland	275	54.5 (48.5-	61.3)	34.5 (28.3-	42.1)	63.3 (53.9-	74.2)
Norway	4,452	49.0 (47.4-	50.7)	22.7 (21.1-	24.4)	46.3 (43.4-	49.3)
Sweden	7,655	48.3 (47.1-	49.6)	21.7 (20.5-	22.9)	44.9 (42.7-	47.2)
Ireland and UK	70,341	42.1 (41.7-	42.5)	17.2 (16.8-	17.5)	40.8 (40.0-	41.6)
Ireland	3,616	42.3 (40.6-	44.0)	20.3 (18.7-	22.1)	48.1 (44.7-	51.6)
UK, England	53,998	42.3 (41.8-	42.8)	17.0 (16.6-	17.4)	40.2 (39.3-	41.2)
UK, Northern Ireland	1,854	41.5 (39.0-	44.0)	18.1 (15.9-	20.5)	43.6 (38.9-	48.8)
UK, Scotland	6,713	40.4 (39.1-	41.7)	16.1 (15.0-	17.3)	39.9 (37.3-	42.6)
UK, Wales	4,160	42.3 (40.6-	44.1)	17.8 (16.2-	19.5)	42.0 (38.7-	45.7)
Central Europe	68,234	53.3 (52.9-	53.7)	28.1 (27.6-	28.5)	52.7 (52.0-	53.4)
Austria	10,569	54.8 (53.8-	55.8)	31.0 (29.9-	32.2)	56.6 (54.9-	58.5)
Belgium	6,650	56.2 (54.9-	57.5)	30.5 (29.1-	32.0)	54.4 (52.2-	56.7)
France	6,084	54.8 (53.5-	56.2)	26.3 (25.0-	27.7)	47.9 (45.8-	50.1)
Germany	26,662	56.3 (55.7-	56.9)	31.3 (30.6-	32.0)	55.6 (54.5-	56.7)
Switzerland	2,146	57.7 (55.5-	59.9)	31.6 (29.3-	34.2)	54.9 (51.3-	58.7)
The Netherlands	16,123	45.2 (44.4-	46.0)	20.4 (19.7-	21.3)	45.3 (43.7-	46.8)
Southern Europe	68,213	52.2 (51.9-	52.6)	29.6 (29.2-	30.0)	56.7 (56.0-	57.4)
Croatia	8,397	38.4 (37.3-	39.5)	21.3 (20.2-	22.5)	55.6 (53.2-	58.1)
Italy	35,428	56.6 (56.0-	57.2)	32.4 (31.7-	33.0)	57.2 (56.2-	58.1)
Malta	340	41.0 (35.7-	47.1)	18.7 (14.5-	24.0)	45.5 (36.9-	56.2)
Portugal	14,127	53.9 (53.0-	54.7)	31.8 (30.9-	32.7)	59.0 (57.6-	60.5)
Slovenia	3,617	46.0 (44.3-	47.7)	26.6 (24.9-	28.5)	57.9 (54.7-	61.2)
Spain	6,304	49.4 (48.1-	50.7)	25.6 (24.3-	26.9)	51.8 (49.7-	54.0)
Eastern Europe	50,775	38.4 (38.0-	38.9)	18.8 (18.4-	19.2)	48.9 (48.0-	49.9)
Bulgaria	11,614	28.8 (27.9-	29.6)	11.9 (11.2-	12.7)	41.4 (39.1-	43.9)
Czech Republic	12,354	41.8 (40.9-	42.8)	22.0 (21.1-	23.0)	52.7 (50.8-	54.6)
Estonia	3,242	44.4 (42.7-	46.2)	22.8 (21.0-	24.8)	51.4 (47.9-	55.2)
Latvia	4,842	38.3 (36.9-	39.7)	20.2 (18.8-	21.7)	52.7 (49.5-	56.1)
Lithuania	6,741	42.5 (41.3-	43.7)	23.1 (21.9-	24.4)	54.3 (51.9-	56.9)
Poland	6,112	38.1 (36.8-	39.4)	15.6 (14.4-	16.8)	40.8 (38.2-	43.7)
Slovakia	5,870	43.5 (42.2-	44.8)	20.9 (19.6-	22.2)	48.0 (45.5-	50.6)
Europe	279,749	49.7 (49.4-	50.0)	25.1 (24.8-	25.4)	50.6 (50.1-	51.1)

Age-standardised 5-year relative survival (%)



European age-specific and age-standardised observed (obs, %) and relative (rel, %) survival


Age group	Number of cases		1-year	3-year	5-year
15-44	10,313	obs	60.3	37.4	32.0
		rel	60.4	37.5	32.3
45-54	22,471	obs	59.0	35.6	29.8
		rel	59.3	36.1	30.5
55-64	47,246	obs	55.8	33.6	27.5
		rel	56.4	34.7	29.1
65-74	84,324	obs	48.9	27.6	21.9
		rel	50.1	29.8	25.1
75+	115,395	obs	34.2	16.6	11.7
		rel	37.3	21.3	18.1
All cases	279,749	obs	48.3	27.4	21.9
		rel	49.7	29.8	25.1
Men (61%)					
15-44	5,712	obs	59.1	35.9	30.4
		rel	59.2	36.1	30.8
45-54	14,758	obs	58.0	34.4	28.3
		rel	58.3	35.0	29.2
55-64	32,479	obs	54.6	31.8	25.8
		rel	55.3	33.0	27.6
65-74	55,667	obs	47.9	25.7	19.9
		rel	49.4	28.2	23.4
75+	61,560	obs	33.7	15.4	10.5
		rel	36.9	20.4	17.0
All cases	170,176	obs	47.3	25.9	20.3
		rel	48.9	28.4	23.7
Women (39%)					
15-44	4,601	obs	61.9	39.3	34.1
		rel	62.0	39.4	34.2
45-54	7,713	obs	60.9	37.9	32.6
		rel	61.1	38.2	33.1
55-64	14,767	obs	58.6	37.8	31.5
		rel	58.9	38.5	32.5
65-74	28,657	obs	50.8	31.5	26.2
		rel	51.5	33.0	28.5
75+	53,835	obs	34.9	18.0	13.1
		rel	37.7	22.4	19.2
All cases	109,573	obs	50.0	30.3	24.9
		rel	51.1	32.3	27.7

Small intestine

Age-standardised 1-year, 5-year relative survival, and 5-year relative survival conditional to surviving 1 year, with 95% confidence intervals in parentheses

	Number of cases	1-year			5-year			Conditional		
Northern Europe	3,828	73.5	(72.1- 75.0)		52.5	(50.5- 54.6)		71.5	(69.1- 73.9)	
Denmark	600	59.8	(56.0- 63.7)		36.8	(32.4- 41.9)		61.6	(55.1- 69.0)	
Finland	644	76.0	(72.7- 79.5)		56.0	(51.2- 61.2)		73.7	(68.2- 79.6)	
Iceland	44	77.5	(64.7- 92.9)		45.3	(31.2- 65.7)		58.4	(42.2- 80.8)	
Norway	821	75.1	(72.1- 78.1)		55.0	(50.8- 59.4)		73.2	(68.4- 78.3)	
Sweden	1,719	76.2	(74.2- 78.3)		55.7	(52.7- 58.9)		73.0	(69.5- 76.7)	
Ireland and UK	6,948	58.8	(57.6- 59.9)		36.9	(35.5- 38.3)		62.8	(60.7- 64.9)	
Ireland	383	66.1	(61.5- 71.0)		43.0	(37.2- 49.8)		65.1	(57.3- 73.9)	
UK, England	5,361	57.5	(56.2- 58.9)		36.4	(34.9- 38.1)		63.3	(61.0- 65.7)	
UK, Northern Ireland	187	57.1	(50.3- 64.7)		34.6	(26.9- 44.5)		60.6	(48.8- 75.3)	
UK, Scotland	612	63.8	(60.1- 67.8)		38.4	(33.6- 43.8)		60.1	(53.4- 67.6)	
UK, Wales	405	61.7	(57.1- 66.6)		37.3	(31.7- 43.8)		60.4	(52.4- 69.6)	
Central Europe	6,754	74.1	(73.1- 75.2)		53.9	(52.4- 55.4)		72.7	(70.9- 74.5)	
Austria	889	74.7	(71.8- 77.7)		54.6	(50.6- 59.0)		73.1	(68.4- 78.1)	
Belgium	841	75.1	(72.2- 78.1)		55.8	(51.7- 60.3)		74.3	(69.6- 79.4)	
France	802	74.4	(71.3- 77.5)		50.8	(47.0- 55.0)		68.4	(63.9- 73.1)	
Germany	2,151	77.2	(75.4- 79.1)		57.7	(54.9- 60.7)		74.7	(71.5- 78.1)	
Switzerland	348	78.9	(74.7- 83.4)		58.6	(52.4- 65.4)		74.2	(67.4- 81.6)	
The Netherlands	1,723	68.5	(66.3- 70.8)		48.9	(46.0- 52.1)		71.4	(67.7- 75.4)	
Southern Europe	3,677	66.2	(64.7- 67.7)		46.8	(44.9- 48.7)		70.6	(68.3- 73.1)	
Croatia	286	53.3	(47.9- 59.4)		42.3	(35.4- 50.6)		79.4	(68.8- 91.5)	
Italy	2,206	69.7	(67.8- 71.6)		49.0	(46.7- 51.6)		70.4	(67.5- 73.4)	
Malta	30	68.7	(51.2- 92.2)		23.5	(13.4- 40.9)		34.1	(21.3- 54.7)	
Portugal	626	62.3	(58.6- 66.2)		42.4	(38.0- 47.3)		68.1	(62.1- 74.6)	
Slovenia	158	64.5	(57.6- 72.2)		44.4	(36.1- 54.8)		68.9	(57.8- 82.2)	
Spain	371	61.5	(56.8- 66.6)		45.4	(40.1- 51.3)		73.7	(67.2- 81.0)	
Eastern Europe	2,071	61.4	(59.2- 63.6)		43.0	(40.1- 46.0)		70.0	(66.1- 74.2)	
Bulgaria	262	55.5	(49.1- 62.8)		36.6	(28.5- 46.8)		65.9	(53.1- 81.7)	
Czech Republic	890	64.6	(61.5- 67.9)		47.4	(43.1- 52.2)		73.4	(67.6- 79.7)	
Estonia	87	64.4	(54.6- 76.0)		42.2	(29.7- 60.1)		65.5	(48.0- 89.6)	
Latvia	101	53.7	(44.3- 65.1)		39.6	(28.7- 54.6)		73.7	(56.9- 95.3)	
Lithuania	161	56.8	(49.4- 65.2)		32.5	(24.8- 42.7)		57.3	(45.3- 72.3)	
Poland	229	58.3	(52.0- 65.3)		40.0	(32.7- 49.0)		68.7	(58.0- 81.3)	
Slovakia	341	63.2	(58.1- 68.9)		46.3	(39.8- 53.9)		73.2	(64.5- 83.0)	
Europe	23,278	67.9	(67.0- 68.9)		47.9	(46.7- 49.1)		70.5	(69.0- 72.0)	

Age-standardised 5-year relative survival (%)



European age-specific and age-standardised observed (obs, %) and relative (rel, %) survival

Age group	Number of cases		1-year	3-year	5-year
15-44	1,490	obs	86.2	72.5	67.3
		rel	86.3	72.8	67.9
45-54	2,756	obs	82.5	67.9	59.7
		rel	82.9	68.9	61.2
55-64	5,141	obs	74.6	58.8	51.0
		rel	75.3	60.6	53.8
65-74	6,577	obs	66.5	49.2	41.0
		rel	68.0	52.8	46.6
75+	7,314	obs	47.7	28.9	23.0
		rel	51.4	36.2	34.2
All cases	23,278	obs	66.2	49.4	42.2
		rel	67.9	53.1	47.9
Men (54%)					
15-44	816	obs	84.4	68.1	63.9
		rel	84.5	68.5	64.6
45-54	1,663	obs	81.3	65.8	58.3
		rel	81.7	66.9	60.1
55-64	2,977	obs	72.5	55.9	46.8
		rel	73.3	58.1	50.0
65-74	3,729	obs	65.4	47.3	39.1
		rel	67.3	51.7	46.0
75+	3,416	obs	47.7	27.3	22.6
		rel	52.0	35.5	35.9
All cases	12,601	obs	65.1	47.1	40.1
		rel	67.2	51.5	47.0
Women (46%)					
15-44	674	obs	89.1	78.4	71.7
		rel	89.2	78.6	72.1
45-54	1,093	obs	84.5	71.3	62.4
		rel	84.7	71.9	63.3
55-64	2,164	obs	77.7	63.6	57.4
		rel	78.1	64.7	59.3
65-74	2,848	obs	67.8	51.7	43.7
		rel	68.8	54.1	47.6
75+	3,897	obs	47.8	30.1	24.3
		rel	51.0	36.5	33.9
All cases	10,676	obs	67.8	52.4	45.4
		rel	69.1	55.3	49.9