**QUEEN'S
UNIVERSITY
BELFAST**

Poster abstract: Adapted Home-Based Cardiac Rehabilitation Following a TIA or Minor Stroke? A Pilot Randomised Trial of “The Healthy Brain Rehabilitation Manual.

Heron, N., Cupples, M., Donnelly, M., Mant, J., & Kee, F. (2019). Poster abstract: Adapted Home-Based Cardiac Rehabilitation Following a TIA or Minor Stroke? A Pilot Randomised Trial of “The Healthy Brain Rehabilitation Manual. *Clinical Journal of Sport Medicine*, 29(3), [e49–e69]. <https://doi.org/10.1097/JSM.0000000000000748>

Published in:
Clinical Journal of Sport Medicine

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2019 Lippincott, Williams & Wilkins. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Adapted home-based cardiac rehabilitation following a TIA or
minor stroke? A pilot randomised trial of 'The Healthy Brain
Rehabilitation Manual'

Authors: Neil Heron PhD,^{1,2,3} Frank Kee MD,^{2,3} Jonathan Mant MD,⁴
Margaret E Cupples MD,^{1,2,3} Michael Donnelly PhD^{2,3}

Affiliations:

¹Dept of General Practice and Primary Care, Queen's University, Belfast

²Centre for Public Health Research, Queen's University, Belfast

³UKCRC Centre of Excellence for Public Health Research (NI)

⁴Primary Care Unit, Department of Public Health and Primary Care,
University of Cambridge, Strangeways Research Laboratory, Cambridge,
UK

Corresponding author: Dr Neil Heron

Address: Dept of General Practice, Queen's University, 1 Dunluce
Avenue, Belfast, BT9 7HR.

Telephone: 0044(0)7808774292

Email: nheron02@qub.ac.uk

Twitter: @neilSportDoc

ClinicalTrials.gov Identifier: NCT02712385; registered, 18/03/2016

Abstract

Objective

Secondary prevention, including positive lifestyle change and exercise/physical activity, after transient ischaemic attack (TIA) or minor stroke is important but the best way to effectively implement this is uncertain. Cardiac rehabilitation (CR) delivers effective secondary prevention for coronary heart disease but its value after a TIA or minor stroke is untested despite these conditions sharing similar pathology. We aimed to pilot a trial of a novel home-based prevention programme (*'The Healthy Brain Rehabilitation Manual'*), adapted from home-based CR, for patients with a TIA or 'minor' stroke.

Design

Pilot randomised controlled trial.

Subjects

Patients within 4 weeks of their first TIA or 'minor' stroke.

Intervention

Patients received study information from clinicians in 4 hospitals and were randomly allocated to: (1) standard care (control group) (n=12); (2) standard care, manual and GP follow-up (n=14); (3) standard care, manual and stroke nurse follow-up (n=14). All participants received telephone follow-up at 1, 4 and 9 weeks. The manual included information about TIA/stroke, discussed a different lifestyle risk factor each week for 6 successive weeks and promoted physical activity, using pedometer step-count targets.

Outcome measures

We assessed eligibility, recruitment and retention rates; measured stroke/cardiovascular risk factors, including body mass index (BMI), blood pressure, physical activity (questionnaire; accelerometer), diet (validated questionnaire) and functional measures (2-minute walk test; Timed-Up-and-Go test) at baseline and 12 weeks post-enrollment; and elicited participants' views about the study via focus groups.

Results

Over a 32-week period, 28% of clinic attendees (125/443) were eligible; 35% (44/125) consented to research contact; 91% (40/44) participated; 98% (39/40) completed the study. At baseline most participants were overweight, physically inactive and had unhealthy diets. After 12 weeks, intervention groups' BMI, diet, physical activity (questionnaire and objective measures) and functional performance improved. The research methods and programme were acceptable to patients and health professionals; the programme was perceived to 'fill gaps' in current post-TIA management.

Conclusions

Our findings indicate that implementation of this novel home-based CR programme, and of a trial to evaluate its effectiveness, is feasible, with potential for clinically important benefits, increasing physical activity and improving secondary prevention following a TIA or 'minor' stroke.

Acknowledgments

We thank Dr Christopher Cardwell for statistical support and Prof Philip Reilly, our patient and public group, participants and physicians and stroke nurses in participating hospitals for their support.