**QUEEN'S
UNIVERSITY
BELFAST**

A Human Relevant Defined Mixture of Persistent Organic Pollutants (POPs) Affects In Vitro Secretion of Glucagon-Like Peptide 1 (GLP-1), but Does not Affect Translocation of its Receptor

Shannon, M., Xie, Y., Verhaegen, S., Wilson, J., Berntsen, H. F., Zimmer, K. E., Ropstad, E., Green, B., & Connolly, L. (2019). A Human Relevant Defined Mixture of Persistent Organic Pollutants (POPs) Affects In Vitro Secretion of Glucagon-Like Peptide 1 (GLP-1), but Does not Affect Translocation of its Receptor. *Toxicological Sciences*, 172(2), 359-367. <https://doi.org/10.1093/toxsci/kfz192>

Published in:
Toxicological Sciences

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© The Author(s) 2019. Published by Oxford University Press on behalf of the Society of Toxicology. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

A Human Relevant Defined Mixture of Persistent Organic Pollutants (POPs) Affects *In Vitro* Secretion of Glucagon-Like Peptide 1 (GLP-1), but Does not Affect Translocation of its Receptor

Maeve Shannon^{a*}, Yuling Xie^{a*}, Steven Verhaegen^b, Jodie Wilson^a, Hanne Friis Berntsen^{b,c}, Karin Elisabeth Zimmer^d, Erik Ropstad^b, Brian Green^a, Lisa Connolly^{a**}

^a Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, Northern Ireland, United Kingdom

^b Department of Production Animal Clinical Sciences, Faculty of Veterinary Medicine, Norwegian University of Life Sciences, Postboks 369 sentrum 0102, Oslo, Norway.

^c Department of Administration, Lab Animal Unit, National Institute of Occupational Health, P.O. Box 5330, Oslo, Norway.

^d Department of Basic Sciences and Aquatic Medicine, Faculty of Veterinary Medicine, Norwegian University of Life Sciences, Postboks 369 sentrum 0102, Oslo, Norway.

****Joint primary co-authorship***

*****Corresponding author and person to whom reprint requests should be addressed:***

Dr. Lisa Connolly

Institute for Global Food Security

Queen's University Belfast

18-30 Malone Road Belfast, BT9 5BN

Phone: +44 (0)28 90976668 Email: l.connolly@qub.ac.uk

Keywords: Metabolic disruption, Gut hormones, Mixture, Enteroendocrine cell, High Content Analysis, ELISA.

Abstract

Environmental exposure to persistent organic pollutants (POPs) has been suggested as a contributing factor for the increased rate of T2D and obesity. A complex mixture of 29 POPs (Total mixture), based on human blood concentrations was used to expose a glucagon-like peptide 1 (GLP-1) secreting enteroendocrine cell line (pGIP/neo: STC-1) in vitro, for 3 and 24 h. Significant increases of GLP-1 occurred when cells were exposed to Total Mixture at x 500 blood levels. Six sub-mixtures representing chlorinated (Cl), brominated (Br), and perfluorinated chemicals (PFAA), and their combinations (Cl + Br, Cl + PFAA, Br + PFAA) were also tested at x 500. Secretion levels seen for these, remained lower than for the Total mixture, and the Br mixture had no effect. After 24 h, increased secretion was seen with all mixtures at x 1 blood levels. Cytotoxicity was present for x 100 and x 500 blood levels. When tested in a GLP-1 receptor translocation assay (U2OS-GLP1R-EGFP), neither agonistic, nor antagonist effects on receptor internalisation were seen for any of the mixtures. We conclude individual classes of POP, alone or in combination, can affect GLP-1 secretion and might contribute as a molecular mechanism, linking environmental toxicants and diabetes.

1. Introduction

The incidence of metabolic diseases such as metabolic syndrome, obesity, and type 2 diabetes (T2D) is increasing globally. Environmental chemicals have been proposed as additional risk factors, in addition to main contributors like genetic predisposition, excess calorie intake, or lack of exercise (Jeon *et al.*, 2015). Experimental and epidemiological evidence suggests a role for early and chronic exposure to low doses of chemical pollutants with endocrine and metabolic disrupting activity, including persistent organic pollutants (POPs) (Chevalier and Fénichel, 2015). In humans, the main exposure to POPs in non-occupational settings occurs through ingestion (Darnerud *et al.*, 2006; Vestergren *et al.*, 2012). This makes the gastrointestinal tract the first organ of exposure, and thus it is conceivable that enteroendocrine (EE) cells are a potential target for endocrine disruption.

POPs constitute a diverse group of chemicals that are resistant to environmental degradation, bioaccumulate within living organisms, and biomagnify through the food web. Due to their mixing in the environment, food web, and the long-term accumulation in fatty tissues, exposure to POPs always occurs as a mixture (WHO/UNEP, 2013).

Human and experimental evidence linking chlorinated POPs with obesity and/or T2D has been extensively reviewed (Lee *et al.*, 2014). Associations for brominated POPs have been reported by Lim *et al.*, (2008). Background exposure of particular brominated POPs (PBB-153 and PBDE-153) may be closely linked to disturbance of glucose and lipid metabolism in the general population (Lim *et al.*, 2008). Perfluorinated alkylated acids (PFAAs) have been associated with disturbed lipid metabolism and shown to bind to peroxisome proliferator-activated receptors (PPARs), which are of major importance for lipid metabolism and fat storage (Zhang *et al.*, 2014, Fang *et al.*, 2015)). A study by Lind *et al.*, 2014 assessed the relationship between circulating levels of PFAAs and markers of insulin secretion and

resistance. The results revealed that perfluorononanoic acid (PFNA) is positively correlated to prevalent diabetes.

In addition to the mechanisms outlined above, it is possible that POP exposure leads to an endocrine disrupting action on gut hormones involved in controlling satiety responses and insulin release. GLP-1 (glucagon-like peptide 1) is one such hormonal system involved and vital in maintaining glucose and weight homeostasis.

GLP-1 is a peripheral humoral factor secreted by entero-endocrine (EE) L-type cells (Lim and Brubaker, 2006). After food ingestion, early-phase secretion starts within 10-15 mins as triggered by the mechano- and chemo-sensitive afferent neurons. Direct nutrient contact with L-cells starts GLP-1 secretion within 30 mins to three hours of meal intake. Exocytosis of GLP-1 secretory granules requires activation of intracellular signalling pathways such as PKA, PKC, and ion channels-dependent membrane depolarisation (Nadkarni, Chepurny, and Holz, 2014). Secreted GLP-1 either acts locally in the gut with EE cells, vagal afferents, or is drained into the circulation via the hepatic portal vein or lymph to bind to specific receptors (Steinert and Begliger, 2011).

The GLP-1 receptor (GLP-1 R), a class B G protein-coupled receptor (GPCR), is distributed in multiple tissues and has been known to mediate GLP-1 actions (Graaf et al., 2016). The pancreas is one of the major sites for GLP-1R expression. GLP-1 binding promotes the exocytosis of insulin vesicles in beta cells and suppress the secretion of hyperglycaemic hormone glucagon in alpha cells. In addition, GLP-1 increases cell proliferation, and inhibits beta cell apoptosis.

In this study, we investigated the potential for POP mixtures to modify GLP-1 secretion, and to interact with the GLP-1R, providing insight into alternative mechanisms by which POPs may be involved in the pathogenesis of diabetes and obesity.

In addition, to simulate real-life exposure scenarios relevant to humans, we utilised a synthetic mixture of 29 different POPs (Table 1) designed as relevant to the Scandinavian population, with the individual concentration of the different POPs based on mean human blood levels (Supplementary table 3). In addition to the Total mixture, 6 sub-mixtures containing the same concentration of chlorinated + brominated (Cl + Br), chlorinated + perfluorinated (Cl + PFAA), brominated + perfluorinated (Br + PFAA), or chlorinated (Cl), brominated (Br) or perfluorinated compounds (PFAA) only were constructed. Using these sub-mixtures enables investigating the effect of adding or removing one or more chemical groups (Berntsen et al., 2017). The concentration of the 29 POPs within the mixtures were based on the blood levels found in the Scandinavian population. However, a wider range of concentrations (from below and up to 500x blood levels) were tested to reflect human body burden levels in different exposure scenarios, including populations exposed to much higher POP concentrations. For example, Chinese occupational exposure to PFAAs is 100 to 500 times higher than that reported in the Scandinavian population (Fu et al., 2016). In addition, the US exposure levels to some POPs are ten to hundreds of times higher than the European population (Schechter et al., 2005). The body burden of POPs is higher also in Inuit populations (Krüger et al., 2007; Laird, Goncharov and Chan, 2013).

In summary, this study investigates seven complex POP mixtures relevant to real life human exposure scenarios (Berntsen *et al.*, 2017) for effects on:

1. Cytotoxicity on GLP-1 secreting EE gut cells (pGIP/neo: a sub-clone of the STC-1 cell line).
2. GLP-1 hormone secretion by EE gut cells.
3. Endocrine disrupting potential via (ant)agonism of the GLP-1 receptor internalization in the U2OS-GLP1R-EGFP cell line.

2. Material and Methods

2.1 Chemicals

All brominated and chlorinated chemicals were originally purchased from Chiron As (Trondheim, Norway). All PFAAs were obtained from Sigma-Aldrich (St. Louis, MO, USA) except perfluorohexanesulfonic acid (PFHxS) which was from Santa Cruz (Dallas, US). Hexabromocyclododecane (HBCD) was obtained from Sigma-Aldrich (Dorset, UK). Mitochondrial membrane potential dye (MMPD) and Hoechst nuclear stain 33342 were purchased from Perbio (Northumberland, England)). Cell culture reagents were supplied by Life Technologies (Paisley, UK). All other reagents were standard laboratory grade.

2.2 Mixtures of Persistent Organic Pollutants (POPs)

The POP mixtures used in the study were based on concentrations of relevant POPs measured in human blood of Scandinavian population. The compounds were mixed in concentration ratios relevant to average human blood levels. An extensive description and information of chemical composition and concentrations can be found in Berntsen et al., 2017. Seven mixtures were used in the study: (1) Total mixture, containing all the test compounds, (2) chlorinated mixture (Cl), (3) brominated mixture (Br), (4) perfluorinated mixture (PFAA), (5) chlorinated and brominated mixture (Cl + Br), (6) chlorinated and perfluorinated mixture (Cl + PFAA) and (7) brominated and perfluorinated mixture (Br + PFAA).

2.3 Cell culture

pGIP/neo: STC-1 cells are an EE cell model and its GLP-1 secretory ability has been extensively investigated (Gillespie *et al.*, 2015; Hand *et al.*, 2012; Jafri *et al.*, 2016). These were a gift from Dr. B. Wice (Washington University of St. Louis) with permission from Dr. D. Hanahan (University of California, San Francisco, CA). Cells were cultured in DMEM with Glutamax, 10% FBS, 1% penicillin and streptomycin, and 50 mg/ml G418. Cells were used between passages 15–50.

The U2OS-GLP1R-EGFP cell line is obtained from Thermo Scientific (UK), which stably express the human GLP-1R with a C-terminal enhanced green fluorescent protein tag. The cells were cultured routinely in Dulbecco's Modified Eagle Medium (DMEM) with Glutamax, 10% foetal bovine serum (FBS), 1% penicillin and streptomycin, and 50 mg/ml G418 at 37°C with 5% CO₂. The assay has been validated with cells up to a passage of 30 (Thermo Scientific, UK).

2.4 Analysis of cytotoxicity parameters by High Content Analysis (HCA)

Briefly, pGIP/neo: STC-1 cells were seeded 6×10^4 in 96-well plates 24 h prior to the assay. MMPD was prepared by adding 117 μ l of anhydrous DMSO to make a 1 mM stock. Following incubation (3 h and 24 h), 50 μ l of live cell stain was added to each well for 30 min at 37 °C and protected from light. The live stain was removed and cells were then fixed with a 10% formalin solution for 20 min at RT, and washed with PBS. Hoechst 33342 dye at a final concentration of 1.6 μ M was added to each well and incubated for 10 min at RT; after which cells were washed with PBS four times and evaluated on a CellInsight™ NXT High Content Screening (HCS) Platform (Thermo Fisher Scientific, UK). Data were captured at X20 objective magnification with selected filters: Hoechst dye (Ex/Em 350/461 nm) and MMPD (Ex/Em 554/576 nm).

2.5 GLP-1 secretion studies

The pGIP/neo: STC-1 cells were seeded into 24-well plates (1×10^6 per well) and cultured overnight at 37 °C in a humidified atmosphere of 5% CO₂. Medium was removed and cells were washed once with HBSS buffer and equilibrated for 1 h in 500 µl HBSS buffer. Forskolin and IBMX were used as a combined positive control at a concentration of 10 µM each. POP mixtures, positive control, and solvent control were made at a final dimethyl sulfoxide (DMSO) concentration of 0.2% in HBSS buffer. One ml of each working concentration of POP mixtures containing HBSS buffer, or 500µl of DMSO solvent control, was added to the relevant wells of the plate. For 24 h study, POP mixtures and controls were added in serum containing DMEM. Supernatant was removed after 3 and 24 h and stored at -20°C prior to GLP-1 ELISA analysis.

2.6 Hormone analysis

GLP-1 was measured using a Glucagon Like Peptide-1 (GLP-1) (Active) ELISA kit (EGLP-35K; in accordance with the manufacturer's instructions (Millipore, Watford, UK)). All experiments were performed in triplicate for each experimental point and repeated in three independent experiments. For 3 h exposure, the supernatants were diluted by HBSS buffer 1 in 4 prior to the measurement of GLP-1 hormone levels by ELISA. For 24 h exposure, the supernatants were diluted by 1 in 40 in media. Absorbance was measured using a Sunrise spectrophotometer (TECAN, Switzerland).

2.7 GLP-1 redistribution assay

U2OS-GLP1R-EGFP cells were seeded (100 µl of 6×10^4 cells/well) in 96-well black plates with clear, flat bottoms in seeding media (DMEM with Glutamax, 1% FBS, 1% penicillin and streptomycin (50 U/ml)). After 24 h, 100 µl of the POP mixtures and the GLP-1 (7-37) standards at a final DMSO concentration of 0.2% were added into plates. Test mixtures and standards were diluted in serum free media. The ability of 150 nM GLP-1 to induce

internalisation of the receptor was assessed in the presence and absence of the mixtures. Cells were incubated for 1 h then fixed with formalin (approximately 4% formaldehyde) for 20 min. Cells were washed, then followed by addition of 100 μ l of 1 μ M Hoechst staining solution. After 30 min, the plate was imaged on a HCS Platform with the filters: Hoechst dye (350/461 nm) and GFP/FITC (488/509 nm) (Thermo Fisher Scientific, UK). The assay was performed in triplicate for each experimental point and repeated in three independent exposures.

2.7 Statistical analysis

All values shown are expressed as mean $\pm$ standard error of the mean (SEM) of the three independent exposures for the POP mixtures. Data from the HCA cytotoxicity, GLP-1 secretion and GLP-1R redistribution assays were analysed using Microsoft Excel and GraphPad PRISM 5 software (GraphPad Software Inc, San Diego, CA). A one-way analysis of variance (ANOVA) followed by Dunnett's multiple comparison test was used to determine significant differences between treatments and the corresponding control. A P -value of ≤ 0.05 was considered as significant ($P \leq 0.05$ *, $P \leq 0.01$ ** and $P \leq 0.001$ ***).

3. Results

3.1. Effects of POP mixture exposure on Cellular Health and GLP-1 secretion in pGIP/neo:STC-1 cells

3.1.1 Cellular Health evaluated by HCA analysis

The general health of the exposed cell populations was evaluated by a multi-parameter approach on an HCA platform. Hoechst-staining of nuclei provided information on cell number (object count) and possible morphological changes were monitored using nuclear area (NA)

and nuclear intensity (NI). A mitochondrial specific fluorescent label was used to quantitate mitochondrial mass (MM) and the mitochondrial membrane potential (MMP). Mean values, standard error of mean, and p-values for total mixture and the 6 sub-mixtures are listed in Supplementary Table 1 (Shannon *et al.*, 2019).

In summary, exposure to the Total mixture (1:10 to x 500 blood level) for 3 h did not significantly alter the cell number. For 24 h exposure, a reduction in cell number compared to solvent control (0.2% DMSO) was seen for the two highest concentrations tested (x 100 and x 500) (Fig. 1A). No significant changes were seen for the Cl mixture after 3 h exposure (Fig. 1A), however after 24 h, all concentrations tested led to a significant reduction (Fig. 1B). For the Br mixture at 3 h exposure only x 500 led to a significant decrease (Fig. 1A), whilst for 24 h significant decreases were seen for x 50 and higher (Fig. 1B). PFAA exposure had no effect at 3 h, but reduced cell number from x 1 blood level onwards after 24 h (Fig. 1B). Cl + Br mixture did not induce any significant decrease after 3 h exposure, while 24 h treatment resulted in the significant decrease for x 50, x 100, and x 500 blood level (Fig. 1B). For the PFAA + Cl mixture, a significant decrease in cell number was seen for x 100 and x 500 after 3 h exposure (Fig. 1A). For 24h, a decrease was already seen from x 50 onwards (Fig. 1B). Br + PFAAs mixture exposure at 3 h only caused significant decrease at x 500 blood level (Fig. 1A), however, the 24 h exposure induced significant decrease at all tested concentrations (Fig. 1B).

Apart from cell number, cellular health was further evaluated using the area and fluorescent intensity of selected nuclei as pre-lethal markers. Only the Br, PFAA, and Br + PFAA mixtures induced significant nuclear area decrease at x 500 after 24 h exposure (Fig. 2). Other mixtures did not change nuclear area significantly at all tested levels for 3 or 24 h (Supp. Table 1A and B). Exposure to any of the seven mixtures tested did not induce any significant change of nuclear intensity (Supp. Table 1A and B).

Finally, markers for mitochondrial health, mitochondrial mass (MM) and mitochondrial membrane potential (MMP) were evaluated. For MM no significant changes were observed for any of the seven mixtures at the five concentrations tested. A summary of the results is shown in Supplementary Table 1A (3 h exposure) and 1B (24 h exposure) (Shannon *et al.*, 2019). Cytotoxicity results in fewer cells and a reduction in MMP. The uptake of MMPD is dependent on MMP levels. Therefore, lower MMP levels is reflected in a decrease of MMPD intensity as quantified by HCA measurement. Such changes are illustrated in Fig.3 whereby the PFAAs + Br mix at 500 times blood levels induced nuclear and mitochondrial changes as shown by a decrease in cell number and decrease in fluorescence intensity when compared to the DMSO solvent control. Visual changes in decreased MMPD dye intensity corresponding to MMP levels after 24 h exposure to the POP mixtures is illustrated in Fig 4.

3.1.2 GLP-1 secretion by pGIP/neo: STC-1 cells by ELISA analysis of Culture Medium

Exposure of the pGIP/neo: STC-1 cells were limited to non-cytotoxic ranges. Thus, for 3 h exposure, concentrations of the POP mixtures tested ranged from 1:10 to x 500 blood level. For 24 h exposure, the concentration range was limited from 1:10 to x 50. Basal levels of GLP-1 were measured in cells exposed to 0.2% DMSO only (Solvent Control - SC). Forskolin combined with 3-isobutyl-1-methylxanthine (IBMX) was used as positive control (PC) that trigger GLP-1 secretion via cAMP-elevating and intracellular calcium pathways (Simpson *et al.*, 2007). The collected media were analysed by ELISA to quantify active GLP-1. Mean values, standard error of mean, and p-values for all mixtures tested, at all concentrations are summarized in Suppl. Table 2 for 3 hr exposure (2A) and 24 hr (2B) (Shannon *et al.*, 2019).

Total mixture increased the GLP-1 secretion significantly at x 500 blood levels after 3 h exposure (Fig 5A); while the secretion enhanced significantly at all tested levels (1:10, x 1, and x 50 blood level) after 24 h exposure (Fig. 5B). For Cl mixture, 3 h exposure significant

increased GLP-1 secretion at all concentrations tested (Fig 5B); however, 24 h exposure only promoted secretion at x 1 and x 50 blood level (Fig 6). The Br mixture failed to induce any significant change in GLP-1 secretion compared to the solvent control at 3 h (Fig 5B), while the significant increase GLP-1 levels detected after 24 h exposure at x 1 and x 50 blood level (Fig. 6). The PFAAs mixture led to significant GLP-1 level increase at the highest concentration after 3 h (Fig 5B), and the pro-longed exposure resulted in the significant increase at x 1 and x 50 blood level (Fig. 6). The Cl + Br and Cl + PFAAs mixtures induced significant GLP-1 increase at x 500 blood level after 3 h exposure (Fig 5B), and at all tested concentrations after 24 h exposure (Fig 6). For the Br + PFAAs mixture, both 3 and 24 h exposure resulted in significant increase at two highest tested concentrations (Fig 5B and Fig. 6).

3.2 Effect of POP mixture exposure on GLP-1 receptor redistribution in U2OS cells using HCA.

A standard curve was generated using a concentration range of GLP-1 (1-300 nM) to drive receptor internalisation. The HCA images showed the location of fluorescent GLP-1 receptor after one hour treatment of solvent control and GLP-1 (at 300 nM) (Supplementary Fig 1). The EC₅₀ value was calculated as 37 nM GLP-1. The seven POP mixtures at all concentrations tested did not change internalisation of the GLP-1R in U2OS-GLP1R-EGFP cells with or without presence of GLP-1 (Supplementary Fig 2).

4. Discussion

Associations between POP exposure and obesity/diabetes risk have recently been highlighted without the underlying mechanisms being identified (Evangelou *et al.*, 2016;

Dirinck *et al.*, 2011; Lee *et al.*, 2014;. Lee *et al.*, 2017a: Lee *et al.*, 2017b). The present study investigated the ability of human relevant POP mixtures (Berntsen *et al.*, 2017), to potentially exert their obesogenic/diabetogenic effects via disruption to gut EE cells survival, GLP-1 secretion, and GLP-1 receptor signalling which are involved in weight and glucose homeostasis.

GLP-1 secretion from pGIP/neo STC-1 cells was determined by ELISA Five out of seven POP mixtures induced an increased GLP-1 secretion from the pGIP/neo STC-1 cells following 3 h exposure at higher concentrations (x 100 to x 500 blood levels) while all mixtures promoted GLP-1 secretion in the 24 h study at x 1:10 to x 50. At this time point, higher concentrations (x 100 and x 500 blood levels) induced significant cytotoxicity thus were not tested here.

Interestingly, at 3 h the Total mixture induced a two times higher level of measured GLP-1, than any of its sub-mixtures tested individually. The secretion level is also higher than the summed levels seen for the Cl, Br, and PFAA mixture suggestive of additive or synergistic interactions. But as summarized in Table 2A, interactions between two chemical groups are complex, with binary mixtures stimulating GLP-1 extend to a lesser degree than would be expected by summing the two constituent mixtures.

Thus, the magnitude of GLP-1 induction by the different mixture can be ranked as follows:

Total mixture > *Cl* + *PFAA* > *Br* + *PFAA* > *PFAA* > *Cl* > *Cl* + *Br* > *Br*.

The 24 h exposure induced considerable GLP-1 secretion even at concentration as low as x1 1:10 blood levels, at x 50 blood levels the Total mixture still did not plateau out GLP-1 secretion as exposure of forskolin (10 µM), combined with IBMX (10 µM) still induced greater GLP-1 secretion. Secretion of GLP-1 peptide from pGIP/neo STC-1 is triggered by several cellular pathways including cAMP/ PKA, PKC, and intracellular calcium ion levels (Hand,

Giblin, and Green, 2012). γ -HCH, PFOA, PFOS, and selective PCB congeners increased the levels of intracellular calcium in previous studies (Harada et al., 2005; Liu et al., 2011; Pessah et al., 2006; Silvestroni et al., 1997; Westerink, 2013). Only PBDE-47 increased the intracellular calcium levels at both 2 and 20 μ M, but other more highly brominated congeners did not induce significant calcium fluctuation (Dingemans et al., 2009; Dingemans et al., 2016). The different effects on GLP-1 secretion between the POP mixtures may be due to their various potential effects on intracellular Ca ion levels.

Taylor *et al.*, 2013 reviewed epidemiological studies of POPs exposure and diabetes and pointed out organochloride chemicals exposure were positively related to T2D, with nonorganochlorine POPs such as perfluoroalkyl acids and brominated compounds showed less association. Our observations indicate however that perfluorinated, as well as brominated POPs should be evaluated in mechanistic assays related to T2D, as it is clear they might have significant contribution to the effect, or affect outcome through combinatorial effects.

HCA provides the means to investigate the cellular health of pGIP/neo STC-1 cells at the level of pre-lethal cytotoxicity, whereby cells may still be alive but becoming unhealthy and jeopardising the ability to function optimally (Clarke *et al.*, 2015; Wilson *et al.*, 2016). It has been proposed that both cell number and nuclear area are among the most sensitive indicators of changes in cellular health (O'Brien *et al.*, 2006). The Total mixture only induced a decrease of cell number at the higher concentrations after 24 h exposure. Individual mixtures and sub-mixtures induced more remarkable changes (Fig 4). Cl mixtures decreased cell number after 24 h exposure at lower concentrations. Br induced MMP decrease and nuclear area decrease for both 3 and 24 h exposure. The 3 h exposure to the PFAAs mixture resulted in a significant reduction of cell number and effects on MMP at the higher concentrations. More profound toxic effects at lower concentrations were observed among these four mixtures following 24 h exposure. However, the Cl + Br mixture only led to a reduction of cell number

for 24 h at high concentrations, which suggests that the Cl compounds might compensate for the Br compounds induced cellular damage. Cl + PFAA mixture led to cell number reduction after 3 h, but did not show in Cl or PFAA mixtures, which indicates the additive effect when Cl and PFAA chemicals mixed together. However, the additive effect was not observed for 24 h exposure. Br + PFAA mixture led to MMP decrease at lower concentrations than individual Br or PFAA mixtures. Thus, the Br and PFAA compounds might lead to the additive effect on the cellular damage via mitochondrial impairment. In summary, reduction of the cell number can be ranked as: *Br* > *Br* + *PFAA* > *Cl* + *PFAA* > *Cl* + *Br* > *Cl* > *PFAA* > *Total mixture*.

The decrease in MMP along with the decrease in cell number is indicative of mitochondrial dysfunction, and in some case this is considered as an early marker for cell death pathways (Bernardi *et al.*, 1992; Gottlieb *et al.*, 2003). In some cases, a decrease in cell number was observed without changes in nuclear and mitochondrial parameters. This reduction is often the first consequence of toxic impact, even before cytotoxic parameters appear within the cell, therefore making it one of the most sensitive markers of cell health (Lucke and Mumtsidu, 2010). Abnormal mitochondrial function has been associated with diseases such as diabetes and metabolic syndrome in epidemiological studies, highlighting the significance of the results found in the present study in relation to mitochondrial effects (Park *et al.*, 2013). The Br mixture presented higher cytotoxicity than the other POP mixtures and induced a decrease in secretion of GLP-1 by the gip/neo STC-1 gut cells. In contrast, the Total mixture induced the greatest increase by the mixtures on GLP-1 secretion in the gip/neo STC-1 gut cells. Thus, gut cells cytotoxicity may reduce function and secretion on GLP-1. On the other hand, the toxic impact may result in an altered gut microbiome which has been linked to increased obesity risk (Riva *et al.*, 2017).

No receptor translocation was observed in U2OS-GLP1R-EGFP cells after exposure to any mixtures at tested concentrations. The receptor redistribution in vitro assay was validated

and applied in previous study (Shannon et al., 2017). This is the first study to investigate the interaction of POPs with GLP-1 receptor signalling. Thus, the potential activity of POPs on GLP-1 receptor signalling is difficult to predict.

5. Conclusions

To our knowledge, this is the first study to show that different classes of persistent organic pollutants can affect GLP-1 secretion. Interestingly, both ‘legacy’ POPs such as chlorinated and brominated, chemicals, as well as the emerging perfluorinated chemicals, were found to be active inducers of GLP-1 secretion. Moreover, complex mixtures combining these three groups showed combinatorial effects. Strikingly, this effect was seen with concentrations of the chemicals at levels to those observed in human blood. No effect at the level of GLP-1 receptor signalling were observed. These findings suggested POP exposure roles may contribute to the increasing rates of the pathogenesis of T2D and obesity. This study provides useful insight into the observations that POPs alter glucose homeostasis and should help guide future studies in this area.

Acknowledgments

We gratefully acknowledge PhD Studentship funding provided by the Department of Education and Learning (DEL) Northern Ireland. The POP mixtures used in this study were constructed under projects funded by the Norwegian Research Council (NFR), project 213076/H10 and project 204361/H10.

1. References

Alberts B., Johnson A., Lewis J., Raff M., Roberts K., and Watter P. (2002). *Molecular Biology of the Cell*. 4th edition. New York: Garland Science; The Mitochondrion. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK26894/>

Baggio L. and Drucker D. (2007). Biology of Incretins: GLP-1 and GIP. *Gastroenterology*, 132, 2131–2157.

Bernardi P., Vassanelli S., Veronese P., Colonna R., and Szabo I. (1992). Modulation of the Mitochondrial Permeability Transition Pore. *The Journal of Biological Chemistry*, 267(5), 2934–2939.

Berntsen H.F., Berg V., Thomsen C., Lyche J.L., Ropstad E., Zimmer K.E. (2017). The design of an environmentally relevant mixture of persistent organic pollutants for use in vivo and in vitro studies. *Journal of Toxicology and Environmental Health* 80, 16-18.

Berntsen H., Bjørklund C., Audinot J., Hofer T., Verhaegen S., Lentzen E., Gutleb A. and Ropstad, E. (2017). Time-dependent effects of perfluorinated compounds on viability in cerebellar granule neurons: Dependence on carbon chain length and functional group attached. *Neurotoxicology* 63, 70-83

Chevalier N., and Fénichel P. (2015). Endocrine disruptors: New players in the pathophysiology of type 2 diabetes? *Diabetes and Metabolism*, 41(2), 107–115.

Clarke R., Connolly L., Frizzell C., and Elliott C. T. (2015). High content analysis: A sensitive tool to detect and quantify the cytotoxic, synergistic and antagonistic effects of chemical contaminants in foods. *Toxicology Letters*, 233(3), 278–286.

Darnerud P.O., Atuma S., Aune M., Bjerselius R., Glynn A., Grawé K. P., Becker W. (2006). Dietary intake estimations of organohalogen contaminants (dioxins, PCB, PBDE

and chlorinated pesticides, e.g. DDT) based on Swedish market basket data. *Food Chem Toxicol*, 44, 1597–1606.

Dirinck E., Jorens P. G., Covaci A., Geens T., Roosens L., Neels H., and Van Gaal L. (2011). Obesity and Persistent Organic Pollutants: Possible Obesogenic Effect of Organochlorine Pesticides and Polychlorinated Biphenyls. *Obesity*, 19(4), 709–714.

Dingemans, M., Heusinkveld, H., Bergman, Å., van den Berg, M. and Westerink, R. (2009). Bromination Pattern of Hydroxylated Metabolites of BDE-47 Affects Their Potency to Release Calcium from Intracellular Stores in PC12 Cells. *Environmental Health Perspectives*, 118(4), pp.519-525.

Dingemans, M., Kock, M. and van den Berg, M. (2016). Mechanisms of Action Point Towards Combined PBDE/NDL-PCB Risk Assessment. *Toxicological Sciences*, 153(2), pp.215-224.

Fu, J., Gao, Y., Cui, L., Wang, T., Liang, Y., Qu, G., Yuan, B., Wang, Y., Zhang, A. and Jiang, G. (2016). Occurrence, temporal trends, and half-lives of perfluoroalkyl acids (PFAAs) in occupational workers in China. *Scientific Reports*, 6(1).

Evangelou E., Ntritsos G., Chondrogiorgi M., Kavvoura F. K., Hernández A. F., Ntzani E. E., and Tzoulaki I. (2016). Exposure to pesticides and diabetes: A systematic review and meta-analysis. *Environment International* 91, 60–68.

Fang X., Gao G., Zhang X., and Wang H. (2015). Perfluorononanoic acid disturbed the metabolism of lipid in the liver of streptozotocin-induced diabetic rats. *Toxicology Mechanisms and Methods* 25(8), 622–627.

Gillespie A. L., Calderwood D., Hobson L., and Green B. D. (2015). Whey proteins have beneficial effects on intestinal enteroendocrine cells stimulating cell growth and increasing the production and secretion of incretin hormones. *Food Chemistry*, 189, 120–128.

Graaf C. d., Donnelly D., Wootten D., Lau J., Sexton P. M., Miller L. J., Ahn J. M., Liao J., Fletcher M. M., Yang D., Brown A. J., Zhou C., Deng J. and Wang, M. W. (2016). Glucagon-Like Peptide-1 and Its Class B G Protein-Coupled Receptors: A Long March to Therapeutic Successes. *Pharmacological reviews*, 68(4), 954-1013.

Gottlieb E., Armour S. M., Harris M. H., and Thompson C. B. (2003). Mitochondrial membrane potential regulates matrix configuration and cytochrome-c release during apoptosis. *Cell Death and Differentiation*, 10, 709–717.

Hand K. V., Giblin L., and Green B. D. (2012). Hormone profiling in a novel enteroendocrine cell line pGIP/neo: STC-1. *Metabolism: Clinical and Experimental*, 61(12), 1683–6.

Harada, K., Xu, F., Ono, K., Iijima, T. and Koizumi, A. (2005). Effects of PFOS and PFOA on L-type Ca²⁺ currents in guinea-pig ventricular myocytes. *Biochemical and Biophysical Research Communications*, 329(2), pp.487-494.

Hölscher C., (2010). The Role of GLP-1 in Neuronal Activity and Neurodegeneration. *Vitamins and Hormones* 84, 331-354.

Jafri L., Saleem S., Calderwood D., Gillespie A., Mirza B., and Green B. D. (2016). Naturally-occurring TGR5 agonists modulating glucagon-like peptide-1 biosynthesis and secretion. *Peptides* 78, 51–58.

Jeon J., Ha K., and Kim D. (2015). New risk factors for obesity and diabetes: Environmental chemicals. *Journal of Diabetes Investigation* 6(2), 109–111.

Krüger, T., Hjelmberg, P., Jönsson, B., Hagmar, L., Giwercman, A., Manicardi, G., Bizzaro, D., Spanò, M., Rignell-Hydbom, A., Pedersen, H., Toft, G., Bonde, J. and Bonefeld-Jørgensen, E. (2007). Xenoandrogenic Activity in Serum Differs across European and Inuit Populations. *Environmental Health Perspectives*, 115(Suppl 1), pp.21-27.

Laird, B., Goncharov, A. and Chan, H. (2013). Body burden of metals and persistent organic pollutants among Inuit in the Canadian Arctic. *Environment International*, 59, pp.33-40.

Lee D. H., Lee I. K., Jin S. H., Steffes M., and Jacobs D. R. (2007). Association Between Serum Concentrations of Persistent Organic Pollutants and Insulin Resistance Among Nondiabetic Adults: Results from the National Health and Nutrition Examination Survey 1999-2002. *Diabetes Care* 30(3), 622–628.

Lee D., Porta M., and Vandenberg L. N. (2014). Chlorinated Persistent Organic Pollutants, Obesity, and Type 2 Diabetes. *Endocrine Review* 35(4), 557–601.

Lee Y.-M., Kim K.-S., Jacobs D. R., and Lee D.-H. (2017a). Persistent organic pollutants in adipose tissue should be considered in obesity research. *Obesity Reviews* 18, 129–139.

Lee Y., Ha C., Kim S., and Thoudam T. (2017b). Persistent organic pollutants impair insulin secretory function of pancreatic beta-cells: Human and in vitro evidence. *Diabetes* 66(9), 1–58.

Legler, J., Fletcher, T., Govarts, E., Porta, M., Blumberg, B., Heindel, J. J., and Trasande, L. (2015). Obesity, diabetes, and associated costs of exposure to endocrine-disrupting

chemicals in the European Union. *Journal of Clinical Endocrinology and Metabolism*, 100(4), 1278–1288.

Lim G. E. and Brubaker P. L. (2006). Glucagon-Like Peptide 1 Secretion by the L-Cell, *Diabetes* 55, S70-S77.

Lim J., Lee D., and Jacobs D. (2008). Association of Brominated Flame Retardants With Diabetes and Metabolic Syndrome in the U.S. Population, 2003–2004. *Epidemiology/Health Services Research* 31(9), 1802–1807.

Lind L., Zethelius B., Salihovic S., van Bavel B., and Lind P. M. (2014). Circulating levels of perfluoroalkyl substances and prevalent diabetes in the elderly. *Diabetologia*, 57, 473–479.

Liu, X., Jin, Y., Liu, W., Wang, F. and Hao, S. (2011). Possible mechanism of perfluorooctane sulfonate and perfluorooctanoate on the release of calcium ion from calcium stores in primary cultures of rat hippocampal neurons. *Toxicology in Vitro*, 25(7), pp.1294-1301.

Lucke J., and Mumtsidu E. (2010) Evaluation of Compound Cytotoxicity using the Opera. Mitochondrial Mass, Cell Count, Membrane Permeability and Nuclear Size. Available from: <http://www.perkinelmer.co.uk/PDFs/downloads/APP-Evaluation-Compound-Cytotoxicity.pdf>.

Murphy K.G, and Bloom S.R. (2006). Gut hormones and the regulation of energy homeostasis, *Nature* 444(7121), 854–859.

Nadkarni P., Chepurny O. G., and Holz G. G. (2014). Regulation of glucose homeostasis by GLP-1. *Progress in molecular biology and translational science*, 121, 23-65.

O'Brien P. J., Irwin W., Diaz D., Howard-Cofield E., Krejsa C. M., Slaughter M. R., and Hougham C. (2006). High concordance of drug-induced human hepatotoxicity with in vitro cytotoxicity measured in a novel cell-based model using high content screening. *Archives of Toxicology* 80(9), 580–604.

Park W., and Jeong J. H. (2013). Novel cell-based assay reveals associations of circulating serum AhR-Ligands with metabolic syndrome and mitochondrial dysfunction. *BioFactors* 39(4), 494–504.

Pessah, I., Hansen, L., Albertson, T., Garner, C., Ta, T., Do, Z., Kim, K. and Wong, P. (2006). Structure–Activity Relationship for Noncoplanar Polychlorinated Biphenyl Congeners toward the Ryanodine Receptor-Ca²⁺Channel Complex Type 1 (RyR1). *Chemical Research in Toxicology*, 19(1), pp.92-101.

Riva A., Borgo F., Lassandro C., Verduci E., Morace G., Borghi E., and Berry D. (2017). Pediatric obesity is associated with an altered gut microbiota and discordant shifts in Firmicutes populations. *Environmental Microbiology* 19(1), 95–105.

Schechter, A., Päpke, O., Tung, K., Joseph, J., Harris, T. and Dahlgren, J. (2005). Polybrominated Diphenyl Ether Flame Retardants in the U.S. Population: Current Levels, Temporal Trends, and Comparison With Dioxins, Dibenzofurans, and Polychlorinated Biphenyls. *Journal of Occupational and Environmental Medicine*, 47(3), pp.199-211.

Shannon M., Xie Y., Verhaegen S., Wilson J., Berntsen H. F., Zimmer K. E., Ropstad E., Green B., and Connolly L. (2019). Data from: A Human Relevant Defined Mixture of Persistent Organic Pollutants (POPs) Affects In Vitro Secretion of Glucagon-Like Peptide 1 (GLP 1), but Does not Affect Translocation of its Receptor. *Dryad Digital Repository*. doi:10.5061/dryad.45sk03p

Silvestroni, L., Fiorini, R. and Palleschi, S. (1997). Partition of the organochlorine insecticide lindane into the human sperm surface induces membrane depolarization and Ca^{2+} influx. *Biochemical Journal*, 321(3), pp.691-698.

Simpson A.K., Ward P.S., Wong K.Y., Collord G.J., Habib A.M., Reimann F., Gribble F.M. (2007). Cyclic AMP triggers glucagon-like peptide-1 secretion from the GLUTag enteroendocrine cell line. *Diabetologia* 50: 2181–2189.

Steinert R. E. and Begliger C. (2011). Nutrient sensing in the gut: interactions between chemosensory cells, visceral afferents and the secretion of satiation peptides, *Physiology and Behavior*. Elsevier Inc., 105(1), 62–70.

Taylor K. W., Novak R. F., Anderson H. A., Birnbaum L. S., Blystone C., and Lind L. (2013) Evaluation of the Association between Persistent Organic Pollutants (POPs) and Diabetes in Epidemiological Studies: A National Toxicology Program Workshop Review, *Environ Health Perspect.* 774(7), pp. 774–783.

Vestergren R., Berger U., Glynn A., Cousins I.T. (2012). Dietary exposure to per-fluoroalkyl acids for the Swedish population in 1999, 2005 and 2010. *Environ. Int.* 49, 120-127.

Westerink, R. (2013). Modulation of cell viability, oxidative stress, calcium homeostasis, and voltage- and ligand-gated ion channels as common mechanisms of action of (mixtures of) non-dioxin-like polychlorinated biphenyls and polybrominated diphenyl ethers. *Environmental Science and Pollution Research*, 21(10), pp.6373-6383.

Wilson J., Friis H., Elisabeth K., Frizzell C., Verhaegen S., Ropstad E., and Connolly L. (2016). Effects of defined mixtures of persistent organic pollutants (POPs) on multiple

cellular responses in the human hepatocarcinoma cell line, HepG2, using high content analysis screening. *Toxicology and Applied Pharmacology* 294, 21–31.

Zhang L., Ren X.M., Wan B., and Guo L.H. (2014). Structure-dependent binding and activation of perfluorinated compounds on human peroxisome proliferator-activated receptor γ . *Toxicology and Applied Pharmacology* 279(3), 275–283.

TABLE 1. Overview of the 29 different POPs constituting the Total mixture for *in vitro* evaluation. For a complete description of selection of compounds and their individual concentration, see Berntsen *et al.*, 2017.

Chlorinated		Brominated	Perfluorinated
PCB	OCP	BFR	PFAAs
PCB 28	p,p'-DDE	PBDE 47	PFHxS
PCB 52	HCB	PBDE 99	PFOS
PCB 101	α -Chlordane	PBDE 100	PFOA
PCB 118	Oxychlordane	PBDE 153	PFNA
PCB 138	trans-Nonachlor	PBDE 154	PFDA
PCB 153	α -HCH	PBDE 209	PFUnDA
PCB 180	β -HCH	HBCD	
	γ -HCH		
	(Lindane)		
	Dieldrin		

TABLE 2. The comparison of measured GLP-1 peptide levels induced by co-mixtures to theoretical sum ($\Sigma = \text{Sum}$) of corresponding sub-mixtures.

A. 3 h and x 500 blood levels				B. 24 h and x 50 blood levels			
Mixture				Mixture			
Mixture	Measured		$\Sigma = \text{Sum}$	Mixture	Measured		$\Sigma = \text{Sum}$
		Cl	161.1			Cl	2357
		Br	55.1			Br	1774
		PFAA	168.3			PFAA	1977
Total	572.6	>>	$\Sigma = 384.5$	Total	3154	<<	$\Sigma = 6108$
		Cl	161.1			Cl	2357
		Br	55.1			Br	1774
Cl + Br	105.3	<<	$\Sigma = 216.2$	Cl + Br	2577	<<	$\Sigma = 4131$
		Br	55.1			Br	1774
		PFAA	168.3			PFAA	1977
Br + PFAA	201.4	<<	$\Sigma = 223.4$	Br + PFAA	1741	<<	$\Sigma = 3751$
		Cl	161.1			Cl	2357
		PFAA	168.3			PFAA	1977
Cl + PFAA	250.8	<<	$\Sigma = 329.4$	Cl + PFAA	2587	<<	$\Sigma = 4334$

Fig. 1: Cell number changes in pGIP/neo STC-1 cells after a) 3h and b) 24h exposure to POP mixtures. Cells were exposed to five concentrations of POPs (x 1:10, x 1, x 50, x 100 and x 500 blood levels) and cytotoxicity was measured by cell number parameter. Data is expressed as a percentage of untreated control for each parameter; mean $\pm$ SEM, n = 3. $P \leq 0.05$ (*), $P \leq 0.01$ (**) and $P \leq 0.001$ (***) represent significant cytotoxic effects.

Fig. 2. Nuclear area changes in pGIP/neo STC-1 cells after exposure to POP mixtures. Cells were exposed to concentrations of POPs (1:10, X 1, X 50, X 100 and X 500 blood levels) for 24 h and cytotoxicity measured by nuclear area parameter. Data is expressed as a percentage of untreated control for each parameter; mean $\pm$ SEM, $n = 3$. $P \leq 0.05$ (*), $P \leq 0.01$ (**) and $P \leq 0.001$ (***) represent significant cytotoxic effects.

Fig. 3. Mitochondrial changes after exposure to three POP mixtures, Br, PFAA, and Br + PFAA, in pGIP/neo STC-1 cells. Cells were exposed to concentrations of POPs (1:10, X 1, X 50, X 100 and X 500 blood levels) for a) 3 h and b) 24 h, cytotoxicity measured by the HCA endpoint mitochondrial membrane potential. Data is expressed as a percentage of untreated control for each parameter; mean $\pm$ SEM, $n = 3$. $P \leq 0.05$ (*), $P \leq 0.01$ (**) and $P \leq 0.001$ (***) represent significant cytotoxic effects.

Fig. 4. HCA images for (A) negative control (DMSO) and B-H) POP mixtures after 24 h exposure. Each image was acquired at 20× objective magnification using Hoechst dye (blue; nuclear staining) and MMPD dye (pink; mitochondrial staining). The arrow in figure A shows a higher cell number and more intense stained mitochondria, arrows in figure C and D show decreased cell number and decreased MMPD dye intensity.

Fig 5. GLP-1 secretory responses of pGIP/neo: STC-1 cells during 3 h exposure to A) Total mixture; B) Cl, Br, PFAA, Cl + Br, Cl + PFAA, and Br + PFAA mixtures. Graph shows GLP-1 secretion from pGIP/neo: STC-1 cells (mean $\pm$ SEM, $n = 3$) following 3 h incubation with solvent control, positive control, and each mixture at 1:10, X 1, X 50, X 100 and X 500 blood levels. $P \leq 0.05$ (*), $P \leq 0.01$ (**), $P \leq 0.001$ (***) represent significance.

Fig. 6. GLP-1 secretory responses of pGIP/neo: STC-1 cells during 24 h exposure to Total mixture, Cl, Br, PFAA, Cl + Br, Cl + PFAA, and Br + PFAA mixtures. Graph shows GLP-1 secretion from pGIP/neo: STC-1 cells (mean $\pm$ SEM, $n = 3$) following 3 h incubation with solvent control (SC), positive control (PC), and ($\times 1:10$, $\times 1$, and $\times 50$ blood levels). $P \leq 0.05$ (*), $P \leq 0.01$ (**), $P \leq 0.001$ (***) represent significance.