

Draft genome sequence of halomonas sp. CSM-2, a moderately halophilic bacterium isolated from a triassic salt mine

Megaw, J., & Gilmore, B. F. (2018). Draft genome sequence of halomonas sp. CSM-2, a moderately halophilic bacterium isolated from a triassic salt mine. Microbiology resource announcements, 7(3), Article e00836-18. https://doi.org/10.1128/MRA.00836-18

Published in:

Microbiology resource announcements

Document Version:

Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:

Link to publication record in Queen's University Belfast Research Portal

Publisher rights
Copyright 2018 the authors.

This is an open access article published under a Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. - Share your feedback with us: http://go.qub.ac.uk/oa-feedback







Draft Genome Sequence of *Halomonas* sp. CSM-2, a Moderately Halophilic Bacterium Isolated from a Triassic Salt Mine

Julianne Megaw, a Brendan F. Gilmorea

^aSchool of Pharmacy, Queen's University Belfast, Belfast, United Kingdom

ABSTRACT Here, we report the draft genome sequence of Halomonas sp. CSM-2. This moderately halophilic bacterium was isolated from a brine sample obtained from a Triassic salt mine.

he genus Halomonas contains Gram-negative rod-shaped bacteria, first described in 1980 after the isolation of the new genus and species Halomonas elongata from a solar salt facility in the Netherlands (1). Representatives of this genus have since been isolated from diverse saline environments, such as solar salt facilities, intertidal estuaries, the open ocean, and hypersaline lakes. Halomonas species have been described as halotolerant, or slight to moderate halophiles, capable of growth in NaCl concentrations of 0.1 to 32.5% (wt/vol) (2).

Strain CSM-2 was isolated from a sample of brine collected from the Kilroot salt mine, a Triassic halite deposit located in Carrickfergus, Northern Ireland, onto LB agar containing 10% NaCl at 37°C. The strain showed no growth in the absence of salt but grew in the salinity range of 5 to 25% NaCl, with an optimum of 10%, indicating that it is a moderate halophile. Based on 16S sequence similarity, the organism was found to be a member of the genus Halomonas, with the closest neighbor deemed to be the type strain Halomonas janggokensis M24 (99% similarity) (3).

Genomic DNA was extracted from a culture freshly grown in LB broth with 10% NaCl using a GenElute bacterial genomic DNA kit (Sigma-Aldrich, UK), following the protocol for Gram-negative bacteria, yielding 37.2 µg DNA. Whole-genome sequencing was performed by MR DNA (Shallowater, TX), using the Illumina MiSeq platform, on a prepared library with an average size of 739 bp. The resulting 150-bp paired-end reads were assembled de novo using the NGen DNA assembly software by DNAStar, Inc. The assembled genome had a total size of 4,131,419 bp in 55 contigs (the largest was 325,242 bp), with an N_{50} value of 190,775 bp, an L_{50} value of 9, and a GC content of 56.9%. Annotation in Rapid Annotations using Subsystems Technology (RAST) (4) revealed 485 subsystems, 3,877 coding sequences, and 73 RNAs. Several genes were identified that were associated with resistance to heavy metals and toxic compounds (copper, cobalt, zinc, cadmium, mercury, arsenic, and chromium) and antibiotic resistance (fluoroquinolones, β -lactams, and multidrug resistance genes), as well as bacteriocin production and an omega-transaminase, which has since been cloned and expressed in Escherichia coli and shown to be active, with no decrease in activity up to 1.5 M NaCl (5).

Data availability. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number NBYR00000000. The version described in this paper is the first version, NBYR01000000.

ACKNOWLEDGMENT

This work was funded by the Irish Marine Institute under the Beaufort Marine Biodiscovery Research award, funded by the Irish Government under the National Development Plan (J.M. and B.F.G.).

Received 7 June 2018 Accepted 26 June 2018 **Published** 26 July 2018

Citation Megaw J, Gilmore BF. 2018. Draft genome sequence of Halomonas sp. CSM-2, a moderately halophilic bacterium isolated from a Triassic salt mine. Microbiol Resour Announc 7:e00836-18. https://doi.org/10.1128/MRA

Editor Jason Stajich, University of California,

Copyright © 2018 Megaw and Gilmore. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Julianne Megaw. jmegaw01@gub.ac.uk.

REFERENCES

- Vreeland RH, Litchfield CD, Martin EL, Elliot E. 1980. Halomonas elongata, a new genus and species of extremely salt-tolerant bacteria. Int J Syst Bacteriol 30:485–495. https://doi.org/10.1099/00207713-30-2-485.
- 2. Vreeland RH. 2015. *Halomonas*, p 1–19. *In* Bergey's manual of systematics of archaea and bacteria. John Wiley & Sons, Ltd, Chichester, United Kingdom.
- 3. Kim KK, Jin L, Yang HC, Lee S-T. 2007. *Halomonas gomseomensis* sp. nov., *Halomonas janggokensis* sp. nov., *Halomonas salaria* sp. nov. and *Halomonas denitrificans* sp. nov., moderately halophilic bacteria isolated from saline water. Int J Syst Evol Microbiol 57:675–681. https://doi.org/10.1099/ijs.0.64767-0.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. BMC Genomics 9:75. https://doi.org/10.1186/1471-2164-9-75.
- Kelly SA, Megaw J, Caswell J, Scott CJ, Allen CCR, Moody TS, Gilmore BF. 2017. Isolation and characterisation of a halotolerant ω-transaminase from a Triassic period salt mine and its application to biocatalysis. ChemistrySelect 2:9783–9791. https://doi.org/10.1002/slct.201701642.