**QUEEN'S
UNIVERSITY
BELFAST**

Composition and consequences of the IntCal20 radiocarbon calibration curve

Reimer, P. J. (Accepted/In press). Composition and consequences of the IntCal20 radiocarbon calibration curve. *Quaternary Research*.

Published in:
Quaternary Research

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Composition and consequences of the IntCal20 radiocarbon calibration curve: ~~composition and consequences~~

Paula J. Reimer

¹⁴CHRONO Centre for Climate, the Environment and Chronology, Queen's University
Belfast, Belfast BT7 1NN, U.K.

Abstract

Radiocarbon calibration is necessary to correct for variations in atmospheric radiocarbon over time. The IntCal working group has developed an updated and extended radiocarbon calibration curve, IntCal20, for Northern Hemisphere terrestrial samples from 0 to 55,000 cal yr BP. This paper summarizes the new datasets, changes to existing datasets and the statistical method used for constructing the new curve. Examples of the effect of the new calibration curve compared to IntCal13 for hypothetical radiocarbon ages are given. For the recent Holocene the effect is minimal but for older radiocarbon ages the shift in calibrated ages can be up to several hundred years with the potential for multiple calibrated age ranges in periods with higher resolution data. In addition, the IntCal20 curve is used to recalibrate the radiocarbon ages for the glaciation of the Puget Lowland and to recalculate the advance rate. The ice may have reached its maximum position a few hundred years earlier using the new calibration curve; the calculated advance rate is virtually unchanged from the prior estimate.

Keywords

Radiocarbon, calibration, IntCal20

Introduction

Radiocarbon ages require a correction to account for changes in ~~because the assumption that~~ atmospheric concentration of ¹⁴C ~~has been constant over time, which is essential in the age calculation, is not valid.~~ Without this correction, or calibration, radiocarbon ages cannot be directly compared to historical dates or ages measured by other methods nor can rates of change be calculated. Radiocarbon calibrations have been done through the use of a curve based on compilations of ¹⁴C measurements of known age material, such as dendrochronologically dated ~~tree-ring~~ tree rings, since the 1960s (e.g. Stuiver and Suess 1966; Clark 1975). To prevent confusion from the use of the various calibration curves available, an international working group was established to provide a consensus curve (Klein et al.

1982). Since that time updated calibration curves have been ratified by the radiocarbon community. Separate curves for the Northern and Southern Hemisphere terrestrial samples and for marine samples are now available. Numerous researchers have contributed to the calibration effort by providing high precision ^{14}C measurements from ~~tree-ring~~ tree ring ~~tree rings~~ and other archives.

From 2004, the IntCal Working Group has updated, extended and refined radiocarbon calibrations semi-regularly. As more independently dated archives have been radiocarbon dated and our understanding of the Earth systems have increased, calibration curves have been extended and refined. The most recent calibration curves for the Northern and Southern Hemisphere and the ocean surface have recently been published (Reimer et al. 2020; Hogg et al. 2020; Heaton et al. 2020a) ~~extending for 0~~ to 55,000 cal yr BP. The methods and datasets used for the IntCal20 curve are summarized in this paper. Calibrations of some hypothetical radiocarbon ages with both IntCal13 (Reimer et al. 2013a) and IntCal20 are used to highlight some of the similarities and differences between the two curves. An example of the effect of the new calibration curve on previously published radiocarbon ages for the timing and rate of advance of the Cordilleran ice sheet in the Pacific Northwest, USA (Porter and Swanson 1998) is given.

New statistical tools

The construction of the IntCal20 calibration curve was underpinned by a new statistical model. Heaton et al. (2020b) adapted Bayesian splines to provide a flexible method for curve construction suited to the complexities of the different types of data and error structures. The Bayesian method also allowed for prior knowledge to inform the construction. For instance, reported laboratory errors don't usually include all sources of uncertainty as evidenced by the Sixth International Radiocarbon Intercomparison exercise (SIRI, Scott et al. 2017). For ~~tree-ring~~ tree ring ~~tree rings~~, growing season and potentially species differences may contribute additional uncertainty (e.g. Kromer et al. 2001; Dee et al. 2010). The Bayesian model used prior information for additional radiocarbon uncertainty based on ~~tree-ring~~ tree ring data from the SIRI exercise although the resulting curve uncertainty was dominated by the high quality IntCal data. In addition, wiggle-matching of floating ~~tree-ring~~ tree ring series (i.e. not dendrochronologically linked to an absolutely dated chronology) was done internal to the model within estimated uncertainty. The Bayesian spline also allows for rapid changes in ^{14}C

to be captured in the curve. Complete details of the Bayesian spline implementation are given in Heaton et al. (2020b).

Calibration datasets

Dendrochronologically-dated ~~tree-ring~~tree ring archives are still preferred for radiocarbon calibration for terrestrial samples because they are direct recorders of atmospheric ^{14}C and there is little or no uncertainty in the calendar age. The number of calibration quality radiocarbon measurements on known age ~~tree-ring~~tree rings, many of them single year, has proliferated especially since the discovery of rapid increases in atmospheric ^{14}C at AD 774-5 and AD 993 (Miyake et al. 2012; 2013). Other time periods have been targeted for single ring measurements to search for additional ^{14}C events (e.g. Miyake et al., 2017a,b; Jull et al., 2018) and to improve calibration around a radiocarbon plateau ca. ~~400-800~~2700-2400 cal yr ~~BC-BP~~ (Park et al., 2017; Fahrni et al., 2020) as well as attempting to pinpoint the timing of the Minoan eruption of Santorini (Thera) (Friedrich et al., 2020; Kuitens et al., 2020; Pearson et al., 2018; 2020). The oldest dendrochronologically dated ~~tree-ring~~tree ring chronology in the Northern Hemisphere is the central European Preboreal Pine Chronology (PPC, Friedrich et al. 2004) which has been extended to 12,049-235 cal yr BP (Sookdeo Reinig et al. 2020a). Older ~~tree-ring~~tree rings used in calibration remain floating. Multi-laboratory ^{14}C measurements of the late glacial New Zealand kauri floating chronology (Hogg et al., 2016) pointed to an error in the link between the absolutely dated Central European PPC and the floating Swiss Late Glacial Master Chronology (Kaiser et al. 2012) used in IntCal13. Investigating the previous ~~tree-ring~~tree ring links resulted in an improved match with the Swiss floating chronology shifting it 35 ± 8 years older. This was supported by the overlap with new ^{14}C measurements from a floating chronology from the southern French Alps (Capano et al. 2018; 2020). A major discovery at a construction site in Zurich of hundreds of pine trees buried *in situ* has provided ample material for chronological replication and increased resolution ca. 13,160 – 11,950 cal yr BP (Reinig et al. 2020; Sookdeo et al. 2020b).

Three floating ~~tree-ring~~tree ring series from northern Italy, that had previously been fitted to Greenland ice core ^{10}Be to ca. 14,700 to 14,000 cal yr BP (Adolphi et al. 2017), were incorporated by ^{14}C matching to the other IntCal data. These measurements add structure to the otherwise rather smooth calibration curve in this time period. A 2000 year long floating ~~tree-ring~~tree ring series from New Zealand (Turney et al. 2016) was also incorporated using

an interhemispheric offset of 43 ± 23 ^{14}C yrs (Hogg et al. 2013). This series adds structure to the curve during Heinrich Stadial 3. Another glacial New Zealand series with 1300 rings (Turney et al. 2010) was also ^{14}C matched into the curve.

The most influential dataset in the IntCal20 curve older than ca. 14,000 cal yr BP is undoubtedly the U-Th dated Hulu cave ^{14}C record from China (Cheng et al. 2018) which extends to 53,900 cal ka-BP (~~0 BP = AD 1950~~). The correction for old carbon (dead carbon fraction) in the ^{14}C ages, estimated from the overlap with ~~tree-ring~~ tree ring data, can be assumed to be relatively constant within uncertainty because the speleothem formed in a portion of the cave where the limestone had largely been replaced by iron oxides. In addition, a short residence time for the soil carbon above the cave is presumed due to the observation of seasonal $\delta^{18}\text{O}$ values and lack of high ^{14}C levels from nuclear weapons testing observed in the dripwaters (Cheng et al. 2018). The $\delta^{18}\text{O}$ fluctuations recorded in the speleothem (Wang et al. 2001; Cheng et al. 2016) also serve as tie-points for marine foraminifera records from the Iberian margin, Pakistan margin and Cariaco basin (Bard et al. 2013; Heaton et al., 2013; Hughen and Heaton 2020). In addition, the Lake Suigetsu varved sediment record, which contains terrestrial macrofossils, has been revised and extended (Schlölaut et al., 2018) and the calendar age modeled with the Hulu cave timescale (Bronk Ramsey et al. 2020). Also included in IntCal20 were two speleothem records from an underwater cave on Grand Bahamas (Beck et al. 2001; Hoffmann et al. 2010). The Bahamas speleothem records, despite having large uncertainty on the dead carbon fractions, serve as a check on the Hulu cave data. By incorporating a range of datasets we can assess which features are likely to represent atmospheric signals and which are local features or noise.

Marine ^{14}C measurements of coral and foraminifera from marine sediments have been included with a correction for the marine reservoir age (MRA) of the ocean region where they grew. In the past the MRA was assumed to be constant in time but there is an abundance of evidence to suggest this is not a valid assumption especially for the last glacial period. For IntCal20 we used MRAs calculated with the Hamburg Large Scale Geostrophic Ocean General Circulation Model (LSG OGCM) with atmospheric input provided by a curve constructed with a Bayesian spline of the Hulu data (Butzin et al. 2020). The calculated MRAs ~~were not applied directly to the marine ^{14}C ages but~~ were only used as prior information to correct the marine ^{14}C ages in the Bayesian spline which then adjusted to best fit the data within uncertainty. The LSG OGCM modelled MRA for the Cariaco basin did not agree with the data, possibly due to the coarse resolution of the model compared to the size of

the basin, so a slowly varying spline was used instead (Heaton et al. 2020b; Hughen and Heaton 2020).

Data from aragonitic coral that grew close to the surface of the ocean were included in previous IntCal curves if the corals met previously established criteria (Reimer et al. 2013b). However, some of the coral ^{14}C data older than 25,000 cal ka-BP is highly variable regardless of meeting the criteria. It's likely there has been some undetected diagenesis due to exposure to freshwater during the Last Glacial Maximum lowstand (21,000 $\pm$ 2000 cal kBP). Therefore, no corals older than 25,000 cal ka-BP were used in IntCal20.

Results and discussion

The effect of the IntCal20 calibration curve compared to IntCal13 (Reimer et al. 2013a) is shown for hypothetical radiocarbon ages of 5000 $\pm$ 20 ^{14}C yr BP, 15,000 $\pm$ 30 ^{14}C yr BP, 30,000 $\pm$ ^{14}C yr 50 BP, and 40,000 $\pm$ 200 ^{14}C yr BP (Figure 1). For 5000 ^{14}C yr BP there is hardly any noticeable difference between the probability distributions calculated using IntCal13 and IntCal20. However, at 15,000 ^{14}C yr BP the distribution for IntCal20 has an additional younger peak compared to that for IntCal13. For 30,000 ^{14}C yr BP the distribution calculated with IntCal20 is about 400 years younger than with IntCal13 whereas for 40,000 ^{14}C yr BP the IntCal20 distribution is bimodal but the main peak is about 500 years older than with IntCal20IntCal13.

A ~~goods-an~~ example of the consistency of the overall shape of radiocarbon calibration curves over the past ~~20-yearsthree decades~~, ~~I-consideris~~ the case of the advance rate of the Puget Lobe of the Cordilleran ice sheet in Washington State during the last glaciation and the timing of the arrival of the ice in the Issaquah delta calibrated with IntCal93 (Stuiver and Reimer 1993) and with the new IntCal20 curve. Porter and Swanson (1998) presented 7 radiocarbon dates on outer wood and branches of pine taken from the top of a pro-glacial delta near Issaquah, Washington (Figure 2). The weighted mean of these radiocarbon dates was 14,546 $\pm$ 55 ^{14}C yr BP and the mean intercept with the IntCal93 calibration curve was given as 17,420 $\pm$ 90 cal yr BP. Although it is no longer recommended to use the mean intercept (Telford et al. 2004), the calibrated age range from IntCal20 is only slightly older than this at 17,455 – 18,005 cal yr BP (at 2 σ , rounding out to 5 years). The glacier advance rate was calculated from two radiocarbon dates on spruce wood from Allison Pool, southern British Columbia (ca. 200 km to Issaquah) which had a mean radiocarbon age of 16,059 $\pm$ 71 ^{14}C yr BP (Clague et al. 1988) with a reported IntCal93 calibrated age of 18,925 cal yr BP.

The difference in mean calibrated ages between Allison Pool and Issaquah delta gave an advance rate of 135 m/yr. With IntCal20 the mean of the Allison Pool radiocarbon dates calibrates to 19,175-19,550 cal yr BP (at 2 σ , rounding out to 5 years). Calculating the difference in the calibrated probability distributions using OxCal (Bronk Ramsey et al. 2009; 2017) gives 1939-1355 ¹⁴C years at 95% probability resulting in an advance rate of 103-148 m/yr. The estimated advance rate from Porter and Swanson (1998) of 135 m/yr falls well within that range.

Conclusions

For much of the Holocene, the IntCal20 curve will not have a large effect on the calibration of radiocarbon ages from single samples with the exception of potentially intercepting younger radiocarbon ages with the sharp radiocarbon declines resulting from the ¹⁴C events at AD 774-75-AD and AD 993-AD. For older periods, calibrated age ranges may shift by several hundred years either direction compared to IntCal13 and there may be additional interecepts-calibrated age ranges where the curve is comprised of higher resolution data. Despite increased detail in calibration curves over time, the overall shape of the IntCal20 curve back to a least 25,000 cal BP does not differ greatly from much older curves as seen by the relatively small change for the advance rate of the Cordilleran ice sheet into the Puget Lowland of Washington State as calibrated with IntCal93.

IntCal20 now extends to 55,000 cal-ka BP so that it is now possible to calibrate radiocarbon ages including two standard deviations up to ca. 50,000 ¹⁴C yr BP. The entire IntCal20 curve is available to download and access to the database can be found at <http://intcal.org>. The calibration programs CALIB <http://calib.org> and OxCal <https://c14.arch.ox.ac.uk/> have been updated to use the IntCal20 curve. It should be noted that IntCal20 is intended for the calibration of Northern Hemisphere atmospheric samples. SHCal20 should be used for the calibration of Southern Hemisphere atmospheric samples (Hogg et al. 2020) and Marine20 (with application of a local reservoir adjustment) for the calibration of marine samples (Heaton et al. 2020a).

Acknowledgements

I would like to thank the IntCal Working Group (<http://intcal.org>), focus group members and others who made the IntCal20 curve possible. I would also like to thank Ron Reimer for producing the figures for this paper. I am grateful to Julie Brigham-Grette and an anonymous reviewer for their constructive comments.

References

- Adolphi, F., Muscheler, R., Friedrich, M., Gütler, D., Wacker, L., Talamo, S., Kromer, B., 2017. Radiocarbon calibration uncertainties during the last deglaciation: Insights from new floating ~~tree-ring~~tree ring chronologies. *Quaternary Science Reviews* 170, 98-108.
- Bard, E., Ménot, G., Rostek, F., Licari, L., Böning, P., Edwards, R.L., Cheng, H., Wang, Y., Heaton, T.J., 2013. Radiocarbon calibration/comparison records based on marine sediments from the Pakistan and Iberian margins. *Radiocarbon* 55, 1999-2019.
- Beck, J.W., Richards, D.A., Edwards, R.L., Silverman, B.W., Smart, P.L., Donahue, D.J., Hererra-Osterheld, S., Burr, G.S., Calsoyas, L., Jull, A.J.T., Biddulph, D., 2001. Extremely Large Variations of Atmospheric ^{14}C Concentration During the Last Glacial Period. *Science* 292, 2453-2458.
- Bronk Ramsey, C., 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon* 51, 337-360.
- Bronk Ramsey, C., 2017. OxCal Program, Version 4.3, Oxford Radiocarbon Accelerator Unit: University of Oxford.
- Bronk Ramsey, C., Heaton, T.J., Schlolaut, G., Staff, R.A., Bryant, C.L., Brauer, A., Lamb, H.F., Marshall, M.H., Nakagawa, T., 2020. Reanalysis of the atmospheric radiocarbon calibration record from Lake Suigetsu. *Radiocarbon* 62, in press.
- Butzin M., Heaton, T.J., Köhler, P., Lohmann, G., 2020. A short note on marine reservoir age simulations used in IntCal20. *Radiocarbon* 62, in press.
- Capano, M., Miramont, C., Guibal, F., Kromer, B., Tuna, T., Fagault, Y., Bard, E., 2018. Wood ^{14}C Dating with AixMICADAS: Methods and Application to ~~Tree-Ring~~Tree ring Sequences from the Younger Dryas Event in the Southern French Alps. *Radiocarbon* 60, 51-74.
- Capano, M., Miramont, C., Shindo, L., Guibal, F., Marschal, C., Kromer, B., Tuna, T., Bard, E., 2020. Onset of the Younger Dryas recorded with ^{14}C at annual resolution in French subfossil trees. *Radiocarbon* 62, in press.
- Cheng, H., Edwards, R.L., Sinha, A., Spötl, C., Yi, L., Chen, S., Kelly, M., Kathayat, G., Wang, X., Li, X., 2016. The Asian monsoon over the past 640,000 years and ice age terminations. *Nature* 534, 640.

- Cheng, H., Edwards, R.L., Southon, J., Matsumoto, K., Feinberg, J.M., Sinha, A., Zhou, W., Li, H., Li, X., Xu, Y., 2018. Atmospheric $^{14}\text{C}/^{12}\text{C}$ changes during the last glacial period from Hulu Cave. *Science* 362, 1293-1297.
- Clague, J.J., Saunders, I.R., Roberts, M.C., 1988. Ice-free conditions in southwestern British Columbia at 16000 years BP. *Canadian Journal of Earth Sciences* 25, 938-941.
- Clark, R.M., 1975. A calibration curve for radiocarbon dates. *Antiquity* 49, 251-266.
- Dee, M.W., Brock, F., Harris, S.A., Ramsey, C.B., Shortland, A.J., Higham, T.F.G., and Rowland, J.M., 2010, Investigating the likelihood of a reservoir offset in the radiocarbon record for ancient Egypt. *Journal of Archaeological Science* 37, 687-693.
- Fahrni, SM, Southon, J, Fuller, BT, Park, J, Friedrich, M, Muscheler, R, Wacker, L, and Taylor, RE, 2020. Single-year German oak and Californian bristlecone pine ^{14}C data at the beginning of the Hallstatt plateau from 856 BC to 626 BC. *Radiocarbon* 62, in press.
- Friedrich, R, Kromer, B, Wacker, L, Olsen, J, Remmele, S, Lindauer, S, Land, A, Pearson, C. 2020. A new annual ^{14}C dataset for calibrating the Thera eruption. *Radiocarbon*, 62 in press.
- Friedrich, M., Remmele, S., Kromer, B., Hofmann, J., Spurk, M., Kaiser, K.F., Orcel, C., and Küppers, M., 2004. The 12,460-year Hohenheim oak and pine ~~tree-ring~~tree ring chronology from central Europe—a unique annual record for radiocarbon calibration and paleoenvironment reconstructions: *Radiocarbon* 46, 1111-1122.
- Heaton, T.J., Bard, E., Hughen, K.A., 2013. Elastic tie-pointing—transferring chronologies between records via a Gaussian process. *Radiocarbon* 55, 1975-1997.
- Heaton, TJ, Köhler, P, Butzin, M, Bard, E, Reimer, RW, Austin, WEN, Bronk Ramsey, C, Grootes, PM, Hughen, KA, Kromer, B, Reimer, PJ, Adkins, JF, Burke, A, Cook, MS, Olsen, J, Skinner, LC, 2020a. Marine20 --- the marine radiocarbon age calibration curve (0-55,000 cal BP). *Radiocarbon* 62, in press.
- Heaton, TJ, Blaauw, M, Blackwell, PG, Bronk Ramsey, C, Reimer, P, Scott, EM, 2020b. The IntCal20 approach to radiocarbon calibration curve construction: A new methodology using Bayesian splines and errors-in-variables. *Radiocarbon* 62, in press.

252 Hoffmann, D.L., Beck, J.W., Richards, D.A., Smart, P.L., Singarayer, J.S., Ketchmark, T.,
 253 and Hawkesworth, C.J., 2010. Towards radiocarbon calibration beyond 28 ka using
 254 speleothems from the Bahamas. *Earth and Planetary Science Letters* 289, 1-10.

255 Hogg, A.G., Hua, Q., Blackwell, P.G., Niu, M., Buck, C.E., Guilderson, T.P., Heaton, T.J.,
 256 Palmer, J.G., Reimer, P.J., Reimer, R.W., Turney, C.S.M., Zimmerman, S.R.H., 2013.
 257 SHCal13 Southern Hemisphere Calibration, 0–50,000 Years cal BP: *Radiocarbon* 55, 1889 -
 258 1903.

259 Hogg, A., Southon, J., Turney, C., Palmer, J., Ramsey, C.B., Fenwick, P., Boswijk, G.,
 260 Büntgen, U., Friedrich, M., Helle, G., 2016. Decadally resolved lateglacial radiocarbon
 261 evidence from New Zealand kauri. *Radiocarbon* 58, 709-733.

262 Hogg, A., Heaton, T.J., Hua, Q., Bayliss, A., Blackwell, P.G., Boswijk, G., Ramsey, C.B.,
 263 Palmer, J., Petchey, F., Reimer, P., Reimer, R.W., Southon, J., Turney, C., Wacker, L., 2020.
 264 SHCAL20 Southern Hemisphere calibration, 0-50,000 years cal BP. *Radiocarbon* [62](#), in
 265 press.

266 Huguen, K., Heaton, T.J., 2020. Updated Cariaco Basin ^{14}C Calibration Dataset from 0-60k
 267 BP. *Radiocarbon* [62](#), in press.

268

269 Jull A.T., Panyushkina I., Miyake F., Masuda K., Nakamura T., Mitsutani T., Lange T.E.,
 270 Cruz R.J., Baisan C., Janovics R., 2018. More Rapid ^{14}C Excursions in the Tree-Ring Record:
 271 A Record of Different Kind of Solar Activity at About 800 BC? *Radiocarbon* 60(4),1237-48.
 272

273 Kaiser, K.F., Friedrich, M., Miramont, C., Kromer, B., Sgier, M., Schaub, M., Boeren, I.,
 274 Remmele, S., Talamo, S., Guibal, F., Sivan, O., 2012. Challenging process to make the
 275 Lateglacial tree-ring chronologies from Europe absolute - an inventory. *Quaternary Science*
 276 *Reviews* 36, 78-90.

277 Klein, J., Lerman, J.C., Damon, P.E., Ralph, E.K., 1982. Calibration of Radiocarbon-Dates -
 278 Tables Based on the Consensus Data of the Workshop on Calibrating the Radiocarbon Time
 279 Scale: *Radiocarbon* 24, 103-150.

280 [Manning, S.W., Barbetti, M., Kromer, B., Kuniholm, P.I., Levin, I., Newton, M.W., and](#)
 281 [Reimer, P.J., 2002, No systematic early bias to Mediterranean C-14 ages: *Radiocarbon*](#)

[measurements from tree-ring and air samples provide tight limits to age offsets. Radiocarbon 44, 739-754](#)

Kuitens, M., Plicht, J.v.d., Jansma, E., 2020. Wood from the Netherlands around the time of the Santorini eruption dated by dendrochronology and Radiocarbon. Radiocarbon [62](#), in press.

Miyake, F., Nagaya, K., Masuda, K., Nakamura, T., 2012. A signature of cosmic-ray increase in AD 774-775 from tree rings in Japan. Nature 486, 240-242.

Miyake, F., Masuda, K., Nakamura, T., 2013. Another rapid event in the carbon-14 content of tree rings. Nat. Commun. 4.

Miyake, F., Jull, A.J.T., Panyushkina, I.P., Wacker, L., Salzer, M., Baisan, C.H., Lange, T., Cruz, R., Masuda, K., Nakamura, T., 2017a. Large ^{14}C excursion in 5480 BC indicates an abnormal sun in the mid-Holocene. Proceedings of the National Academy of Sciences 114, 881-884.

Miyake, F., Masuda, K., Nakamura, T., Kimura, K., Hakozaiki, M., Jull, A.J.T., Lange, T.E., Cruz, R., Panyushkina, I.P., Baisan, C., Salzer, M.W., 2017b. Search for Annual ^{14}C Excursions in the Past. Radiocarbon 59, 315-320.

Park, J., Southon, J., Fahrni, S., Creasman, P.P., Mewaldt, R., 2017. Relationship between solar activity and $\Delta^{14}\text{C}$ peaks in AD 775, AD 994, and 660 BC. Radiocarbon 59, 1147-1156.

Pearson, C.L., Brewer, P.W., Brown, D., Heaton, T.J., Hodgins, G.W., Jull, A.T., Lange, T., Salzer, M.W., 2018. Annual radiocarbon record indicates 16th century BCE date for the Thera eruption. Science advances 4, eaar8241.

Pearson, C.L., Wacker, L., Bayliss, A., Brown, D.M., Salzer, M., Brewer, P.W., Bollhalder, S., Boswijk, G., Hodgins, G.W.L., 2020. Annual variation in atmospheric ^{14}C between 1700 BC and 1480 BC. Radiocarbon [62](#), [doi:10.1017/RDC.2020.14](https://doi.org/10.1017/RDC.2020.14)in press.

Porter, S.C., Swanson, T.W., 1998. Radiocarbon Age Constraints on Rates of Advance and Retreat of the Puget Lobe of the Cordilleran Ice Sheet during the Last Glaciation: Quaternary Research 50, 205-213.

Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hafliðason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F.,

312 Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon,
 313 J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J., 2013a. IntCal13 and Marine13
 314 radiocarbon age calibration curves 0-50,000 years cal BP. Radiocarbon 55, 1869–1887.

315 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Brown, D.M.,
 316 Buck, C.E., Edwards, R.L., Friedrich, M., 2013b. Selection and treatment of data for
 317 radiocarbon calibration: an update to the International Calibration (IntCal) criteria.
 318 Radiocarbon 55, 1923-1945.

319 Reimer, PJ, Austin, WEN, Bard, E, Bayliss, A, Blackwell, PG, Ramsey, CB, Butzin, M,
 320 Cheng, H, Edwards, RL, Friedrich, M, Grootes, PM, Guilderson, TP, Hajdas, I, Heaton, TJ,
 321 Hogg, AG, Hughen, KA, Kromer, B, Manning, SW, Muscheler, R, Palmer, JG, Pearson, C,
 322 Plicht, Jvd, Reimer, RW, Richards, DA, Scott, EM, Southon, JR, Turney, CSM, Wacker, L,
 323 Adolphi, F, Büntgen, U, Capano, M, Fahrni, S, Fogtmann-Schulz, A, Friedrich, R, Kudsk, S,
 324 Miyake, F, Olsen, J, Reinig, F, Sakamoto, M, Sookdeo, A, Talamo, S, 2020. The IntCal20
 325 Northern Hemisphere radiocarbon calibration curve (0-55 cal ka BP). Radiocarbon [62](#), in
 326 press.

327 Reinig, F., Sookdeo, A., Esper, J., Friedrich, M., Guidobaldi, G., Helle, G., Kromer, B.,
 328 Nievergelt, D., Pauly, M., Tegel, W., Treydte, K., Lukas Wacker, Büntgen, U., 2020.
 329 Illuminating IntCal during the Younger Dryas. Radiocarbon [62](#), in press.

330 Schlolaut, G., Staff, R.A., Brauer, A., Lamb, H.F., Marshall, M.H., Ramsey, C.B., Nakagawa,
 331 T., 2018. An extended and revised Lake Suigetsu varve chronology from ~ 50 to ~ 10 ka BP
 332 based on detailed sediment micro-facies analyses. Quaternary Science Reviews 200, 351-366.

333 Scott, E.M., Naysmith, P., Cook, G.T., 2017. Why do we need 14C inter-comparisons?: The
 334 Glasgow-14C inter-comparison series, a reflection over 30 years. Quaternary Geochronology
 335 43, 72-82.

336 ~~Sookdeo, A., Kromer, B., Adolphi, F., Beer, J., Brehm, N., Büntgen, U., Christl, M.,~~
 337 ~~Eglinton, T.I., Friedrich, M., Guidobaldi, G., Helle, G., Nievergelt, D., Pauly, M., Reinig, F.,~~
 338 ~~Tegel, W., Treydte, K., Synal, H. A., Wacker, L., submitted. There Goes the Sun: 14C in~~
 339 ~~trees reveals reduced solar activity during the Younger Dryas. Nature Communications.~~
 340 ~~submitted.~~

341 Sookdeo A, Kromer B, Buentgen U, Friedrich M, Friedrich R, Helle G, Pauly M, Nievergelt
 342 D, Reinig F, Treydte K, Synal HA, Wacker, L. 2020b. Quality Dating: A well-defined

protocol implemented at ETH for high-precision ^{14}C dates tested on Late Glacial wood.
Radiocarbon [62](#), in press.

Stuiver, M., Suess, H.E., 1966. On the relationship between radiocarbon dates and true sample ages. Radiocarbon 8, 534-540.

Stuiver, M., Reimer, P.J., 1993. Extended C-14 Data-Base and Revised Calib 3.0 C-14 Age Calibration Program. Radiocarbon ~~v.~~35, ~~p.~~215-230.

[Telford, R.J., Heegaard, E., Birks, H.J.B., 2004, The intercept is a poor estimate of a calibrated radiocarbon age. The Holocene 14, 296-298.](#)

Turney, C.S.M., Fifield, L.K., Hogg, A.G., Palmer, J.G., Hughen, K., Baillie, M.G.L., Galbraith, R., Ogden, J., Lorrey, A., Tims, S.G., Jones, R.T., 2010. The potential of New Zealand kauri (*Agathis australis*) for testing the synchronicity of abrupt climate change during the Last Glacial Interval (60,000–11,700 years ago). Quaternary Science Reviews 29, 3677-3682.

Turney, C.S., Palmer, J., Ramsey, C.B., Adolphi, F., Muscheler, R., Hughen, K.A., Staff, R.A., Jones, R.T., Thomas, Z.A., Fogwill, C.J., 2016. High-precision dating and correlation of ice, marine and terrestrial sequences spanning Heinrich Event 3: Testing mechanisms of interhemispheric change using New Zealand ancient kauri (*Agathis australis*). Quaternary Science Reviews 137, 126-134.

Wang, Y.J., Cheng, H., Edwards, R.L., An, Z.S., Wu, J.Y., Shen, C.C., Dorale, J.A., 2001. A high-resolution absolute-dated Late Pleistocene monsoon record from Hulu Cave, China. Science 294, 2345-2348.

Figure captions:

Figure 1. Calibrated probability distributions are shown for hypothetical radiocarbon ages of 5000 ± 20 ^{14}C yr BP, $15,000 \pm 30$ ^{14}C yr BP, $30,000 \pm 50$ ^{14}C yr BP, and $40,000 \pm 200$ ^{14}C yr BP. 1 sigma age ranges are shown as thick lines and 2 sigma as thin lines at the base of each distribution. Calibrations were done using IntCal13 and CALIB 7.04 and using IntCal20 with CALIB 8.1 (Stuiver and Reimer 1993).

373 Figure 2. Map showing the extent of the Puget Lobe of the Cordilleran ice sheet in
374 Washington State during the last glaciation (shaded region) adapted from Porter and
375 Swanson1998. Dark grey arrows indicate inferred flow direction.