



**QUEEN'S
UNIVERSITY
BELFAST**

Mechanisms of virus-induced airway immunity dysfunction in the pathogenesis of COPD disease, progression, and exacerbation

Guo-Parke, H., Linden, D., Weldon, S., Kidney, J. C., & Taggart, C. C. (2020). Mechanisms of virus-induced airway immunity dysfunction in the pathogenesis of COPD disease, progression, and exacerbation. *Frontiers in Immunology*, 11, Article 1205. <https://doi.org/10.3389/fimmu.2020.01205>

Published in:
Frontiers in Immunology

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2020 The Authors.

This is an open access article published under a Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>



Mechanisms of Virus-Induced Airway Immunity Dysfunction in the Pathogenesis of COPD Disease, Progression, and Exacerbation

Hong Guo-Parke¹, Dermot Linden¹, Sinéad Weldon¹, Joseph C. Kidney^{2†} and Clifford C. Taggart^{1*†}

¹ Airway Innate Immunity Research Group, Wellcome Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry & Biomedical Sciences, Queens University Belfast, Belfast, United Kingdom, ² Department of Respiratory Medicine Mater Hospital Belfast, Belfast, United Kingdom

OPEN ACCESS

Edited by:

Christian Herr,
Saarland University Hospital, Germany

Reviewed by:

Giuseppe Lungarella,
University of Siena, Italy
Franz Puttur,
Imperial College London,
United Kingdom

*Correspondence:

Clifford C. Taggart
c.taggart@qub.ac.uk

[†]These authors share
senior authorship

Specialty section:

This article was submitted to
Mucosal Immunity,
a section of the journal
Frontiers in Immunology

Received: 02 December 2019

Accepted: 14 May 2020

Published: 16 June 2020

Citation:

Guo-Parke H, Linden D, Weldon S,
Kidney JC and Taggart CC (2020)
Mechanisms of Virus-Induced Airway
Immunity Dysfunction in the
Pathogenesis of COPD Disease,
Progression, and Exacerbation.
Front. Immunol. 11:1205.
doi: 10.3389/fimmu.2020.01205

Chronic obstructive pulmonary disease (COPD) is the integrated form of chronic obstructive bronchitis and pulmonary emphysema, characterized by persistent small airway inflammation and progressive irreversible airflow limitation. COPD is characterized by acute pulmonary exacerbations and associated accelerated lung function decline, hospitalization, readmission and an increased risk of mortality, leading to huge social-economic burdens. Recent evidence suggests ~50% of COPD acute exacerbations are connected with a range of respiratory viral infections. Nevertheless, respiratory viral infections have been linked to the severity and frequency of exacerbations and virus-induced secondary bacterial infections often result in a synergistic decline of lung function and longer hospitalization. Here, we review current advances in understanding the cellular and molecular mechanisms underlying the pathogenesis of COPD and the increased susceptibility to virus-induced exacerbations and associated immune dysfunction in patients with COPD. The multiple immune regulators and inflammatory signaling pathways known to be involved in host-virus responses are discussed. As respiratory viruses primarily target airway epithelial cells, virus-induced inflammatory responses in airway epithelium are of particular focus. Targeting virus-induced inflammatory pathways in airway epithelial cells such as Toll like receptors (TLRs), interferons, inflammasomes, or direct blockade of virus entry and replication may represent attractive future therapeutic targets with improved efficacy. Elucidation of the cellular and molecular mechanisms of virus infections in COPD pathogenesis will undoubtedly facilitate the development of these potential novel therapies that may attenuate the relentless progression of this heterogeneous and complex disease and reduce morbidity and mortality.

Keywords: chronic obstructive pulmonary disease, virus, inflammation, infection, lung damage, acute pulmonary exacerbation

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) is the umbrella term for chronic obstructive bronchitis and pulmonary emphysema, and is characterized by persistent small airway inflammation and progressive irreversible airflow limitation (1–5). COPD is associated with acute pulmonary exacerbations, accelerated lung function decline and increased risk of mortality (6, 7). As a common global epidemic, COPD affects 10% of the population and is the third leading cause of death worldwide (3). Viral and bacterial infections are key elements in the pathogenesis of exacerbations (5–9). Recent evidence suggests respiratory viral infections cause ~50% of COPD acute exacerbations (5, 6, 10). Secondary bacterial infections often ensue with pronounced illness (6).

However, the underlying mechanisms of how viruses subvert host immune defense systems in COPD exacerbations are not completely understood. Herein, we review current advances in understanding the cellular and molecular mechanisms associated with the increased susceptibility to virus infections. As respiratory viruses preferentially infect airway epithelial cells, we focus on virus-induced inflammatory responses in airway epithelium. Understanding these pathogenic pathways may facilitate the development of potential novel therapies to attenuate the relentless progression of the disease.

COPD PATHOGENESIS

Cigarette smoking is the predominant etiologic factor in the development of COPD (3–5). Other risk factors include host genetic factors, which is most evident in alpha-1 antitrypsin (AAT) deficiency (11–13). Recently, childhood respiratory viral infections have been postulated as an independent risk factor associated with COPD later in life (14). Other environmental factors such as pollutant and occupational exposure to dusts or fumes, particularly organic dusts are strongly associated with COPD (4, 13, 15, 16). Social deprivation is also a factor in the development of COPD (6, 17, 18).

Cigarette smoke and other inhaled noxious gases induce an abnormal inflammatory response, that is further amplified by protease and oxidative stress, which are central to COPD pathogenesis (8, 11). Persistent small airway inflammation and the resulting destruction of the lung architecture leads to emphysema and loss of lung elastic recoil, chronic bronchitis induced mucus hypersecretion and airflow obstruction, as well as peribronchial fibrosis (11, 19, 20). Excessive neutrophilic infiltration and associated proteolytic enzymes including neutrophil elastase are hallmark features of smoke-induced inflammation (19, 21–25). Consequently, the protease/antiprotease imbalance contributes to the pathogenesis of emphysema due to the increased breakdown of elastin and loss of elastic recoil in the lung parenchyma (19, 21–24). Diminished activity of protein phosphatase 2A (PP2A), a regulator of the inflammatory response in the airways, has been demonstrated in COPD and upregulation of PP2A activity can ameliorate inflammation in a cigarette smoke model of COPD by reducing activity of the cysteine protease, cathepsin S (26). Recent

research has proposed a role for formylated peptides and formyl peptide receptor (FPR) receptor signaling in the initiation and progression of lung disease in current and former smokers (27, 28). These peptides are present in tobacco leaves and are actively secreted by bacteria or passively released from dead and dying host cells after tissue injury (29). FPR1 and FPR2 activation may play a role in neutrophil migration, degranulation, reactive oxygen species (ROS) production, and phagocytosis (29, 30). A novel cross-talk mechanism was identified in neutrophils, by which signals generated by the purinergic receptor for ATP (P2Y₂) reactivate ligand-bound inactive FPRs, which resume signaling (31). Furthermore, a role for purinergic receptors in the pathophysiology of COPD has been demonstrated in human and experimental models (32–35), however, further work is needed to elucidate its role in the immune dysfunction associated with COPD (36). Excessive production of ROS results in an oxidant-antioxidant imbalance leading to oxidative stress and is a major predisposing feature in the development of the disease (37–41). Therefore, a vicious cycle is created in which inflammation drives a protease-antiprotease and oxidant-antioxidant imbalance, as well as multiple intracellular cell signaling mechanisms, which potentiate inflammation, goblet cell hyperplasia and mucus hypersecretion (8, 40).

Chronic low-grade respiratory syncytial virus (RSV) infection has also been implicated in COPD pathogenesis (42–45). However, the detection of RSV infection in stable COPD remains controversial (46, 47). Hogg and colleagues showed that the E1A region of the adenovirus may contribute to COPD pathogenesis by enhancing soluble ICAM-1 expression and inflammatory cells infiltration (48). In contrast, another study failed to demonstrate the persistent presence of adenovirus V or E1A (49). Polosukhin et al. detected Epstein Barr Virus (EBV) positive cells in COPD lung tissue sections by immunohistochemistry staining (50). Consistent with this finding, we have demonstrated that EBV DNA is frequently present in COPD sputum compared with unaffected smokers (51). Latent viral infections and cigarette smoke may synergistically contribute to the chronic inflammation in COPD (52). COPD is a heterogeneous disease with a complex etiology, however, acute and chronic lower respiratory tract infections occur with increased frequency in patients with COPD. Whatever the cause, it is clear that a defective host response plays an important role and improving our understanding of the mechanisms involved is essential to improving prevention and treatment strategies.

AIRWAY EPITHELIUM DYSFUNCTION IN COPD

Normal airway epithelial cells play a pivotal role in innate immune defense. They act as a barrier to pathogens and noxious stimuli and produce mediators and enzymes to orchestrate and maintain proper functioning of the innate and adaptive immune responses (24, 53, 54). As illustrated in **Figure 1**, the COPD airway epithelium responds to cigarette smoke by secreting inflammatory mediators and recruiting immune cells to the site of damage to orchestrate the inflammatory response. A robust

infiltration of macrophages and CD8⁺ T cells, and to a lesser extent CD4⁺ T cells, in the airway mucosa as well as elevated neutrophils in the airway lumen are the hallmark features of COPD inflammation, the degree of which correlates to disease severity (46, 55). Increased levels of epithelial-derived CXCL9 (MIG), CXCL10 (IP-10), and CXCL11 (I-TAC) and their receptor CXCR3 has been demonstrated to contribute, in part, to the mechanism of CD8⁺ cellular accumulation (40, 53, 54). CD8⁺ T cells release IP-10, TNF- α , IFN- γ , perforins, and granzyme, and have been associated with alveolar epithelial cell apoptosis (19, 37, 56, 57). As COPD progresses, elevated numbers of dendritic cells and B lymphocytes also appear in the airways and alveolar walls. CD8⁺ T cells and B cells organize into lymphoid follicles and may contribute to increased “immune surveillance” in COPD (19, 37, 39). The airway epithelium also releases a cascade of secondary mediators including cytokines, lipid mediators, growth factors, proteases, antiproteases and ROS to escalate COPD inflammation (53, 54, 58). Cigarette smoke and other irritants also activate epithelial cells and macrophages to release neutrophil and macrophage chemoattractants, such as LTB₄, IL-8, and related CXC chemokines (MCP-1, GRO- α and GM-CSF), which contribute to the development of emphysema (39, 46, 59, 60).

The mechanism of neutrophilic inflammation has been linked to CD11b/CD18 on neutrophils binding to ICAM-1 on bronchial epithelium, which is up-regulated in COPD (54, 61–63). Neutrophils migrate to the respiratory tract and release serine proteases, matrix metalloproteinases (MMPs) and oxidants (24, 40, 46). Neutrophil serine proteases are associated with emphysema, mucus hypersecretion, increased risk of exacerbation and accelerated forced expiratory volume in 1 s (FEV₁) decline (64–66). Subsequently, these proteases degrade extracellular matrix components leading to the destruction of the alveolar wall, epithelial barrier dysfunction, reduction in mucociliary clearance, mucus hypersecretion and goblet cell metaplasia through activation of the epidermal growth factor receptor (EGFR) (37, 59, 64). Moreover, alveolar epithelial cells also secrete transforming growth factor- β (TGF- β) which may contribute to small airway fibrosis and emphysema (67).

MOLECULAR MECHANISMS ASSOCIATED WITH VIRAL-INDUCED COPD EXACERBATIONS

Viral-Induced COPD Exacerbations

Acute exacerbations of COPD are characterized by a sudden decline in lung function, hospitalization and high mortality (7, 9, 46). The complicated interaction between the host and viral or bacterial infections or co-infection, as well as environmental factors, precipitate the onset of exacerbations. These factors amplify the inflammatory burden in the small airway, overpowering host anti-inflammatory mechanisms leading to profound airway obstruction in COPD (46, 68–70). Severe virus-associated exacerbations also induce elevated levels of CD8⁺ T cells, neutrophils, eosinophils, TNF- α and IL-6 in the sputum of COPD patients (68–70).

Exacerbations often occur seasonally accompanied by common cold-like symptoms implicating respiratory viral infections rather than hitherto suspected bacterial infection (43, 44). Respiratory virus infection, including human rhinovirus (HRV), influenza virus (IAV), coronavirus, RSV, human parainfluenza, metapneumovirus (hMPV) and adenovirus initiate nearly 50% of COPD exacerbations often with more severe symptoms (69–73). Viruses have developed a myriad of aversion strategies to subvert and manipulate host immune responses and these have been recently reviewed elsewhere (74, 75). Most respiratory viruses target airway epithelial cells leading to epithelial barrier destruction, microvascular dilatation, oedema and immune cell infiltration (58, 70–72). These viruses are associated with small airway secondary bacterial infection, thus magnifying the inflammatory response in COPD leading to a synergistic deterioration in lung function and prolonged hospitalization (42, 44, 71).

As detailed below, recent research has focused on immune regulators and inflammatory signaling pathways orchestrating the underlying mechanisms of increased susceptibility to virus-associated exacerbation and the exaggerated inflammatory response in COPD airways and potential therapeutic inventions.


T Cell Exhaustion

Although accumulated CD8⁺ T cells are present in greater numbers in severe COPD, a diminished CD8⁺ T cell antiviral response, worsened airflow limitation and respiratory symptoms have been reported in IAV and HRV-induced COPD exacerbations (68, 71, 76, 77). As a result, CD8⁺ cells potentially amplify airway epithelium destruction and promote tissue injury through mechanisms including direct cytotoxic effects, pro-inflammatory signaling and recruitment of other immune cells, leading to increased susceptibility to virus infections of airway epithelium (42–44, 69).

In COPD, prolonged receptor–ligand interaction during T cell activation may be linked to T cell exhaustion. McKendry and colleagues investigated increased CD8⁺ activation through the programmed cell death protein (PD)-1 exhaustion pathway as a potential mechanism of viral-induced COPD exacerbations (76). Dysregulation of T-cell cytotoxicity was associated with elevated levels of PD-1, which further increased following influenza infection in COPD patients (76). In contrast, infection-induced expression of the ligand PD-L1 on COPD macrophages was diminished, with a concomitant increase in IFN- γ release. These synergistic effects may cause excessive T-cell inflammation in response to virus infection.

The NF- κ B Pathway

The NF- κ B pathway is consistently activated in COPD macrophages and airway epithelium, in particular, during bacterial or viral infections (78). Upon pathogen stimulation, the canonical pathway is mainly triggered by Toll like receptors (TLRs) and pro-inflammatory cytokines such as TNF α and IL-1 leading to the activation of the RelA containing NF- κ B complexes. This initiates the translocation of RelA (p65)/p50 to the nucleus, where it induces the transcriptional response of pro-inflammatory and cell survival genes (78–80). The alternative



non-canonical NF- κ B pathway signals through a subset of receptors to activate the kinase NIK and IKK α complexes and downstream NF- κ B2 p100 leading to the p52/RelB nucleus translocation and lymphoid organogenesis and B cell activation (78, 79).

Persistent or prolonged activation of NF- κ B may contribute to COPD pathogenesis by switching on the transcriptional response of pro-inflammatory cytokines, chemokines, cell adhesion molecules (CAMs), proteases, and inhibitors of apoptosis to amplify inflammation. Therefore, strategies, which block

the activation of NF- κ B, offer attractive therapeutic options to regulate COPD inflammation. Several IKK- β inhibitors have been identified to inhibit p65 nuclear translocation and exert anti-inflammatory effects (81, 82). Lung-targeted overexpression of RelB has also been demonstrated to protect against cigarette smoke-induced inflammation by reducing inflammatory mediator production (83). In COPD airway epithelium, influenza virus infection increased microRNA-125a/b, which directly inhibits A20 and mitochondrial antiviral-signaling protein (MAVS) to promote inflammation and impair

antiviral responses in COPD (84). Thus, miR-125a/b may provide a potential therapeutic target for both inflammation and antiviral responses in COPD.

TLR Sensing and EGFR Signaling

Figure 2 illustrates key virus innate recognition signaling pathways in COPD airway epithelium. Briefly, ssRNAs of HRV, RSV, and IAV are recognized by TLR3 in the endosomes which consequently activate IRF-3 via the Toll/IL-1 receptor domain-containing adaptor (TRIF), leading to the induction of IFN- β and IFN- λ 1. Other endosomal TLRs (TLR7/8 and TLR7/9) recognize the dsRNAs of IAV and adenovirus through MyD88-dependent pathway to activate NF- κ B and IRF-7 to secrete pro-inflammatory mediators and IFNs, respectively. TLR4 expressed on the cell surface senses RSV and IAV, signaling through both the MyD88 and TRIF pathways to activate NF- κ B and IRF-7. The airway epithelium may recognize EBV by endosomal TLRs and TLR2 at the cell surface to activate downstream pathways (85, 86). As a risk factor for RSV-induced COPD exacerbations, TLR3 activation has been found to correlate with lung function deterioration during exacerbations highlighting TLR3 blockade as a therapeutic target (87). However, Silkoff et al. showed that TLR3 inhibition was inefficient in attenuating HRV-induced experimental asthma exacerbation (88).

Many TLRs recognize pathogen-associated molecular patterns (PAMPs) to activate airway epithelial EGFR signaling cascades. Aberrant EGFR signaling promotes progressive lung fibrosis and mucus hypersecretion; characteristic features of COPD, asthma and cystic fibrosis pathogenesis (24, 89). The EGFR cascade consists of multiple receptors and extracellular ligands that function via receptor auto-phosphorylation and cytoplasmic protein binding of four downstream complexes including the mitogen-activated protein kinases/extracellular signal-regulated kinases (MEK/ERK), phosphatidylinositol 3-kinases/protein kinase B (PKB) (PI3K/AKT), Just Another Kinase/signal transducer and activator of transcription (JAK/STAT) and mammalian target of rapamycin (mTOR) pathways (89). In a murine COPD model, EGFR activation through PI3K inhibited ciliated cell apoptosis and allowed IL-13 to stimulate the trans-differentiation of ciliated to goblet cell metaplasia (90). HRV infection induced the phosphorylation of PKD, a downstream kinase of PI3K. PKD inhibitors have been reported to effectively block HRV, poliovirus (PV) and foot-and-mouth disease virus (FMDV) replication at an early stage of infection, highlighting the potential of PKD inhibition in anti-HRV therapy in COPD (91). Chronic inflammation can also induce ICAM-1 and its ligand fibrinogen has been shown to promote EGFR-dependent mucin production in the airways of subjects with mucus hypersecretion (92).

EGF and the EGFR ligand, TGF- α , have been reported to directly enhance TNF- α -induced IL-8 secretion in airway inflammation (93). Ganesan et al. found that abnormal EGFR activation contributed to enhanced IL-8 expression in COPD airways via the NF- κ B regulator, FoxO3A (94). Interestingly, TLR3 also induced EGFR activation and EGFR ligands (TGF- α and amphiregulin), which in turn promote EGFR-ERK signaling and mucin production through an autocrine/paracrine loop (95).

Collectively, TLR antiviral defense mechanisms integrate with the EGFR mediated epithelial proliferation/repair pathways and may play an important role in viral-induced airway remodeling and airway disease exacerbations (93, 95, 96).

Viral infection *per se* also activates EGFR and EGFR signaling to ERK1/2, while STATs control the severity of HRV mediated airway inflammation. *In vitro*, HRV induced goblet cell hyperplasia was demonstrated to function through NF- κ B-dependent MMP-mediated TGF- α release, leading to EGFR activation and mucus secretion (97). Interestingly, virus-induced EGFR activation suppressed interferon regulatory factor 1 (IRF1)-dependent IFN- λ airway epithelial antiviral signaling (98, 99). Inhibiting virus-mediated EGFR signaling augmented IRF1, IFN- λ secretion and viral clearance, indicating EGFR pathways as potential therapeutic targets in viral-induced COPD exacerbations (99).

Cytoplasmic-Sensing Pathways

As shown in **Figure 2**, the airway epithelium also detects viral invasion through cytoplasmic pathogen recognition receptors. DNA and RNA viruses release their genomes into cytoplasm, which are detected by the host through cytoplasmic retinoic acid-inducible gene I/melanoma differentiation-associated protein 5-mitochondrial antiviral-signaling protein (RIG-I/MDA5-MAVS) RNA-sensing and the cyclic GMP-AMP synthase- signaling effector stimulator of interferon genes (cGAS-STING) DNA-sensing pathways, respectively (100). Upon ss/dsRNA binding, the RNA helicases, RIG-I and MDA5, interact with the adaptor protein MAVS on the mitochondrial outer membrane to activate the downstream signaling of type I interferon antiviral responses (100, 101). In contrast, the cGAS receptor senses retroviral replication products, dsDNA and RNA/DNA hybrids, to induce the synthesis of cGAMP which binds and activates STING (100). Interferon γ -inducible protein 16 (IFI16), a novel DNA sensor, has been found to recruit STING to activate type I IFN signaling through an unknown molecular mechanism (102). STING and MAVS also stimulate downstream multiple kinase signaling cascades resulting in IRF3 phosphorylation and NF- κ B nuclear translocation (101, 102).

The primary consequence of these virus-sensing pathways is the induction of type I/type III IFNs and IFN stimulated genes as well as the production of inflammatory cytokines and chemokines. Attenuation of the IFN response following virus infection could result in uncontrolled viral replication and an escalated inflammatory response, a potential mechanism of virus-induced exacerbations in COPD. IFN α / β deficiency has been demonstrated in bronchial biopsies of asthmatic patients with rhinovirus-induced exacerbations and smoking-induced COPD (103). Farazuddin et al. have demonstrated that quercetin, a potent antioxidant and anti-inflammatory agent with antiviral properties, effectively mitigates rhinovirus-induced COPD exacerbation in a mouse model (104). Elevated ICAM-1 expression on the surface of airway epithelium has been directly linked to the mechanism of increased susceptibility of HRV-induced acute exacerbation. As the receptor of the major group of HRV and a ligand of lymphocyte function-associated antigen 1 (LFA-1) on neutrophils, ICAM-1 over-expression has been




FIGURE 2 | Intracellular Viral Sensing Pathways. DNA and RNA viruses release their genomes in the cytoplasm, where host innate sensors for nucleic acids reside. Upon ss/dsRNA binding, RIG-I engages the adaptor protein MAVS on the mitochondrial outer membrane. The cGAS receptor recognizes dsDNA and the RNA:DNA hybrids generated during retroviral replication and catalyzes the synthesis of cGAMP, which is the primary agonist of the adaptor protein STING. Another sensor, IFI16 can recruit STING in response to cytoplasmic DNA through a molecular mechanism yet to be described. Both STING and MAVS stimulate downstream signaling cascades that involve multiple kinases and finally lead to IRF3 phosphorylation and nuclear translocation. The primary consequence of these virus sensing pathways is the induction of type I IFN and IFN stimulated genes. cGAS, cyclic GMP-AMP synthase; cGAMP, 2'3'guanosine-adenosine monophosphate; IFI16, interferon-g inducible protein 16; IKK, IκB kinase; IRF3, interferon regulatory factor 3; MAVS, mitochondrial antiviral-signaling protein; RIG-I, retinoic acid inducible gene-1; ss/dsRNA, single-stranded/double-stranded RNA; vRNA/DNA, viral RNA/DNA; STING, stimulator of interferon genes; TANK, TRAF-associated NF-κB activator; TBK1, TANK binding kinase 1.

shown on epithelial cells in smokers and patients with COPD (63, 105, 106). Blocking ICAM-1 may also represent as a potential therapeutic option in HRV-induced exacerbations.

Direct Targeting of Viral Binding, Entry, and Replication

Strategies that directly prevent virus binding, entry and replication may provide attractive alternatives in the treatment of COPD exacerbations (107). Capsid binders represent attractive potential inhibitors of HRV entry, however, they

are strain-specific and have shown no effect on improving lung function and exacerbation in clinical trials to date (106). Mousnier and colleagues demonstrated that a dual inhibitor of human N-myristoyltransferases NMT1 and NMT2 can inhibit host-cell N-myristoylation and completely prevent rhinoviral replication, highlighting the therapeutic potential of targeting myristoylation in blocking rhinovirus infection in COPD (108). Short palate, lung, nasal epithelium clone 1 (SPLUNC1), a multifunctional host defense protein, was demonstrated to inhibit IAV binding and entry into airway epithelial cells,

indicating an antiviral role for this protein in the airways (109). Therefore, in the COPD lung, SPLUNC1 degradation by proteases such as neutrophil elastase and/or inactivation by cigarette smoke may increase susceptibility to viral as well as bacterial infections, in addition to airway dehydration (110, 111). Recent research suggests that, in addition to modulating neutrophil chemotaxis, FPR2 signaling may be an important player in viral replication and IAV pathogenesis (30, 112, 113).

Inflammasome

The inflammasome is a multiprotein pro-inflammatory complex and serves as an important link between the innate and adaptive immune responses. Inflammasomes that are activated by IAV RNA, EBV and adenoviral DNA include the nucleotide binding and oligomerization domain (NOD)-like receptor family pyrin domain-containing 3 (NLRP3) protein, absent in melanoma 2 (AIM2) protein and IFI16 protein (114). The inflammasome complexes assemble after recognition of PAMPs or danger-associated molecular patterns (DAMPs) induced by virus-killed cells or tissue damage and interact with apoptosis-associated speck like protein containing a caspase recruitment domain (ASC) via caspase activation and recruitment domains (CARD)-CARD/caspase-1 pathway (115–117). Activation of the inflammasome complex results in the autocatalytic cleavage of caspase-1 and ultimately leads to the production of pro-inflammatory cytokines including IL-1 β , IL-18 and pro-IL-33 (116, 117). Upon maturation, these cytokines mediate inflammatory responses by activating lymphocytes and facilitating their infiltration to the site of primary infection

and by inducing IFNs and other pro-inflammatory cytokines secretions (116).

CONCLUDING REMARKS

COPD is a heterogeneous and complex disease resulting from the deregulation of multiple immune regulators and inflammatory signaling pathways. Significant progress has been made to elucidate the causative mechanism of COPD pathophysiology including viral infection in disease development, severity and exacerbations. Targeting virus-induced inflammatory pathways such as T cell exhaustion, NF- κ B, TLRs, EGFR, interferons and the inflammasome provide attractive future therapeutic options. Understanding the cellular and molecular mechanisms of virus-induced COPD pathogenesis could potentially limit pathogen-mediated disease exacerbations and minimize viral-associated inflammation, tissue destruction and pulmonary function deterioration.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

This work was funded by the Mater Hospital Young Philanthropist (YP) trustees (to JK), Pfizer UK and Chiesi Farmaceutici (to CT and JK). The funders were not involved in the study design, collection, analysis, interpretation of data, the writing of this article or the decision to submit it for publication.

REFERENCES

- Buist AS, McBurnie MA, Vollmer WM, Gillespie S, Burney P, Mannino DM, et al. International variation in the prevalence of COPD (the BOLD study): a population-based prevalence study. *Lancet*. (2007) 370:741–50. doi: 10.1016/S0140-6736(07)61377-4
- Mirza S, Clay RD, Koslow MA, Scanlon PD. COPD guidelines: a review of the 2018. GOLD Report. *Mayo Clin Proc*. (2018) 93:1488–502. doi: 10.1016/j.mayocp.2018.05.026
- Barnes PJ. Chronic obstructive pulmonary disease: a growing but neglected global epidemic. *PLoS Med*. (2007) 4:e112. doi: 10.1371/journal.pmed.0040112
- Salvi S. Tobacco smoking and environmental risk factors for chronic obstructive pulmonary disease. *Clin Chest Med*. (2014) 35:17–27. doi: 10.1016/j.ccm.2013.09.011
- Bauer CMT, Morissette MC, Stämpfli MR. The influence of cigarette smoking on viral infections. *Chest*. (2013) 143:196–206. doi: 10.1378/chest.12-0930
- Mohan A, Chandra S, Agarwal D, Guleria R, Broor S, Gaur B, et al. Prevalence of viral infection detected by PCR and RT-PCR in patients with acute exacerbation of COPD: a systematic review. *Respirology*. (2010) 15:536–42. doi: 10.1111/j.1440-1843.2010.01722.x
- Merinopoulou E, Raluy-Callado M, Ramagopalan S, MacLachlan S, Khalid JM. COPD exacerbations by disease severity in England. *Int J Chron Obstruct Pulmon Dis*. (2016) 11:697–709. doi: 10.2147/COPD.S100250
- Sethi S, Murphy TF. Infection in the pathogenesis and course of chronic obstructive pulmonary disease. *N Engl J Med*. (2008) 359:2355–65. doi: 10.1056/NEJMra0800353
- Ko FW, Chan KP, Hui DS, Goddard JR, Shaw JG, Reid DW, et al. Acute exacerbation of COPD. *Respirology*. (2016) 21:1152–65. doi: 10.1111/resp.12780
- Linden D, Guo-Parke H, Coyle P V., Fairley D, McAuley DE, Taggart CC, et al. Respiratory viral infection: a potential “missing link” in the pathogenesis of COPD. *Eur Respir Rev*. (2019) 28:180063. doi: 10.1183/16000617.0063-2018
- Calverley PMA, Walker P. Chronic obstructive pulmonary disease. *Lancet*. (2003) 362:1053–61. doi: 10.1016/S0140-6736(03)14416-9
- Pauwels RA, Rabe KF. Burden and clinical features of chronic obstructive pulmonary disease (COPD). *Lancet*. (2004) 364:613–20. doi: 10.1016/S0140-6736(04)16855-4
- Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet*. (2007) 370:765–73. doi: 10.1016/S0140-6736(07)61380-4
- Marsico S, Caccuri F, Mazzuca P, Apostoli P, Roversi S, Lorenzin G, et al. Human lung epithelial cells support human metapneumovirus persistence by overcoming apoptosis. *Pathog Dis*. (2018) 76:fty013. doi: 10.1093/femspd/fty013
- Marsh S, Aldington S, Shirtcliffe P, Weatherall M, Beasley R. Smoking and COPD: what really are the risks? *Eur Respir J*. (2006) 28:883–4. doi: 10.1183/09031936.06.00074806
- Lundbäck B, Lindberg A, Lindström M, Rönmark E, Jonsson AC, Jönsson E, et al. Obstructive lung disease in northern Sweden studies. Not 15 but 50% of smokers develop COPD?—Report from the obstructive lung disease in northern Sweden studies. *Respir Med*. (2003) 97:115–22. doi: 10.1053/rmed.2003.1446

17. GBD 2015 Chronic Respiratory Disease Collaborators. Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990-2015: a systematic analysis for the global burden of disease study 2015. *Lancet Respir Med.* (2017) 5:691-706. doi: 10.1016/S2213-2600(17)30293-X
18. Rycroft CE, Heyes A, Lanza L, Becker K. Epidemiology of chronic obstructive pulmonary disease: a literature review. *Int J Chron Obstruct Pulmon Dis.* (2012) 7:457-94. doi: 10.2147/COPD.S32330
19. Hogg JC, Chu F, Utokaparch S, Woods R, Elliott WM, Buzatu L, et al. The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med.* (2004) 350:2645-53. doi: 10.1056/NEJMoa032158
20. Kim V, Criner GJ. Chronic bronchitis and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* (2013) 187:228-37. doi: 10.1164/rccm.201210-1843CI
21. Navratilova Z, Kolek V, Petrek M. Matrix metalloproteinases and their inhibitors in chronic obstructive pulmonary disease. *Arch Immunol Ther Exp.* (2016) 64:177-93. doi: 10.1007/s00005-015-0375-5
22. Owen CA. Roles for proteinases in the pathogenesis of chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis.* (2008) 3:253-68. doi: 10.2147/COPD.S2089
23. Shapiro SD. Proteolysis in the lung. *Eur Respir J.* (2003) 22:30s-2. doi: 10.1183/09031936.03.00000903a
24. Bagdonas E, Raudoniute J, Bruzauskaite I, Aldonyte R. Novel aspects of pathogenesis and regeneration mechanisms in COPD. *Int J Chron Obstruct Pulmon Dis.* (2015) 10:995-1013. doi: 10.2147/COPD.S82518
25. Hoenderdos K, Condliffe A. The neutrophil in chronic obstructive pulmonary disease. Too Little, Too Late or Too Much, Too Soon? *Am J Respir Cell Mol Biol.* (2013) 48:531-9. doi: 10.1165/rcmb.2012-0492TR
26. Doherty DF, Nath S, Poon J, Foronjy RF, Ohlmeyer M, Dabo AJ, et al. Protein phosphatase 2a reduces cigarette smoke-induced cathepsin s and loss of lung function. *Am J Respir Crit Care Med.* (2019) 200:51-62. doi: 10.1164/rccm.201808-1518OC
27. Cardini S, Dalli J, Fineschi S, Perretti M, Lungarella G, Lucattelli M. Genetic ablation of the fpr1 gene confers protection from smoking-induced lung emphysema in mice. *Am J Respir Cell Mol Biol.* (2012) 47:332-9. doi: 10.1165/rcmb.2012-0036OC
28. De Cunto G, Bartalesi B, Cavarra E, Balzano E, Lungarella G, Lucattelli M. Ongoing lung inflammation and disease progression in mice after smoking cessation: beneficial effects of formyl-peptide receptor blockade. *Am J Pathol.* (2018) 188:2195-206. doi: 10.1016/j.ajpath.2018.06.010
29. Dorward DA, Lucas CD, Chapman GB, Haslett C, Dhaliwal K, Rossi AG. The role of formylated peptides and formyl peptide receptor 1 in governing neutrophil function during acute inflammation. *Am J Pathol.* (2015) 185:1172-84. doi: 10.1016/j.ajpath.2015.01.020
30. Bozinovski S, Anthony D, Anderson GP, Irving LB, Levy BD, Vlahos R. Treating neutrophilic inflammation in COPD by targeting ALX/FPR2 resolution pathways. *Pharmacol Ther.* (2013) 140:280-9. doi: 10.1016/j.pharmthera.2013.07.007
31. Önnheim K, Christenson K, Gabl M, Burbiel JC, Müller CE, Oprea TI, et al. A novel receptor cross-talk between the ATP receptor P2Y2 and formyl peptide receptors reactivates desensitized neutrophils to produce superoxide. *Exp Cell Res.* (2014) 323:209-17. doi: 10.1016/j.yexcr.2014.01.023
32. Lommatzsch M, Cicko S, Müller T, Lucattelli M, Bratke K, Stoll P, et al. Extracellular adenosine triphosphate and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* (2010) 181:928-34. doi: 10.1164/rccm.200910-1506OC
33. Lazar Z, Müllner N, Lucattelli M, Ayata CK, Cicko S, Yegutkin GG, et al. NTPDase1/CD39 and aberrant purinergic signalling in the pathogenesis of COPD. *Eur Respir J.* (2016) 47:254-63. doi: 10.1183/13993003.02144-2014
34. Cicko S, Lucattelli M, Müller T, Lommatzsch M, De Cunto G, Cardini S, et al. Purinergic receptor inhibition prevents the development of smoke-induced lung injury and emphysema. *J Immunol.* (2010) 185:688-97. doi: 10.4049/jimmunol.0904042
35. Lucattelli M, Cicko S, Müller T, Lommatzsch M, De Cunto G, Cardini S, et al. P2X7 receptor signaling in the pathogenesis of smoke-induced lung inflammation and emphysema. *Am J Respir Cell Mol Biol.* (2011) 44:423-9. doi: 10.1165/rcmb.2010-0038OC
36. Antonioli L, Blandizzi C, Pacher P, Haskó G. The purinergic system as a pharmacological target for the treatment of immune-mediated inflammatory diseases. *Pharmacol Rev.* (2019) 71:345-82. doi: 10.1124/pr.117.014878
37. Barnes PJ. Cellular and molecular mechanisms of asthma and COPD. *Clin Sci.* (2017) 131:1541-58. doi: 10.1042/CS20160487
38. Eapen MS, Myers S, Walters EH, Sohal SS. Airway inflammation in chronic obstructive pulmonary disease (COPD): a true paradox. *Expert Rev Respir Med.* (2017) 11:827-39. doi: 10.1080/17476348.2017.1360769
39. Barnes PJ, Shapiro SD, Pauwels RA. Chronic obstructive pulmonary disease: molecular and cellular mechanisms. *Eur Respir J.* (2003) 22:672-88. doi: 10.1183/09031936.03.00040703
40. MacNee W. Pathogenesis of chronic obstructive pulmonary disease. *Proc Am Thorac Soc.* (2005) 2:258-66. doi: 10.1513/pats.200504-045SR
41. McGuinness A, Sapey E. Oxidative stress in COPD: sources, markers, and potential mechanisms. *J Clin Med.* (2017) 6:21. doi: 10.3390/jcm6020021
42. Sikkel MB, Quint JK, Mallia P, Wedzicha JA, Johnston SL. Respiratory syncytial virus persistence in chronic obstructive pulmonary disease. *Pediatr Infect Dis J.* (2008) 27:S63-70. doi: 10.1097/INF.0b013e3181684d67
43. Seemungal T, Harper-Owen R, Bhowmik A, Moric I, Sanderson G, Message S, et al. Respiratory viruses, symptoms, and inflammatory markers in acute exacerbations and stable chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* (2001) 164:1618-23. doi: 10.1164/ajrccm.164.9.2105011
44. Singanayagam A, Joshi P V, Mallia P, Johnston SL. Viruses exacerbating chronic pulmonary disease: the role of immune modulation. *BMC Med.* (2012) 10:27. doi: 10.1186/1741-7015-10-27
45. Wilkinson TMA, Donaldson GC, Johnston SL, Openshaw PJM, Wedzicha JA. Respiratory syncytial virus, airway inflammation, and FEV1 decline in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* (2006) 173:871-6. doi: 10.1164/rccm.200509-1489OC
46. Papi A, Bellettato CM, Braccioni F, Romagnoli M, Casolari P, Caramori G, et al. Infections and airway inflammation in chronic obstructive pulmonary disease severe exacerbations. *Am J Respir Crit Care Med.* (2006) 173:1114-21. doi: 10.1164/rccm.200506-859OC
47. Falsey AR, Formica MA, Hennessey PA, Criddle MM, Sullender WM, Walsh EE. Detection of respiratory syncytial virus in adults with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* (2006) 173:639-43. doi: 10.1164/rccm.200510-1681OC
48. Matsuse T, Hayashi S, Kuwano K, Keuneeck H, Jefferies WA, Hogg JC. Latent adenoviral infection in the pathogenesis of chronic airways obstruction. *Am Rev Respir Dis.* (1992) 146:177-84. doi: 10.1164/ajrccm/146.1.177
49. McManus TE, Marley A-M, Baxter N, Christie SN, Elborn JS, Heaney LG, et al. Acute and latent adenovirus in COPD. *Respir Med.* (2007) 101:2084-90. doi: 10.1016/j.rmed.2007.05.015
50. Polosukhin VV, Cates JM, Lawson WE, Zaynagetdinov R, Milstone AP, Massion PP, et al. Bronchial secretory immunoglobulin a deficiency correlates with airway inflammation and progression of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* (2011) 184:317-27. doi: 10.1164/rccm.201010-1629OC
51. McManus TE, Marley A-M, Baxter N, Christie SN, Elborn JS, O'Neill HJ, et al. High levels of epstein-barr virus in COPD. *Eur Respir J.* (2008) 31:1221-6. doi: 10.1183/09031936.00107507
52. Foronjy RF, Dabo AJ, Taggart CC, Weldon S, Geraghty P. Respiratory syncytial virus infections enhance cigarette smoke induced COPD in Mice. *PLoS ONE.* (2014) 9:e90567. doi: 10.1371/journal.pone.0090567
53. Gao W, Li L, Wang Y, Zhang S, Adcock IM, Barnes PJ, et al. Bronchial epithelial cells: the key effector cells in the pathogenesis of chronic obstructive pulmonary disease? *Respirology.* (2015) 20:722-9. doi: 10.1111/resp.12542
54. Vareille M, Kieninger E, Edwards MR, Regamey N. The airway epithelium: soldier in the fight against respiratory viruses. *Clin Microbiol Rev.* (2011) 24:210-29. doi: 10.1128/CMR.00014-10
55. O'Shaughnessy TC, Ansari TW, Barnes NC, Jeffery PK. Inflammation in bronchial biopsies of subjects with chronic bronchitis: inverse relationship of CD8+ T lymphocytes with FEV1. *Am J Respir Crit Care Med.* (1997) 155:852-7. doi: 10.1164/ajrccm.155.3.9117016
56. Kim W-D, Chi H-S, Choe K-H, Oh Y-M, Lee S-D, Kim K-R, et al. A possible role for CD8 + and non-CD8 + cell granzyme B in early small

- airway wall remodelling in centrilobular emphysema. *Respirology*. (2013) 18:688–96. doi: 10.1111/resp.12069
57. Majo J, Ghezzi H, Cosio MG. Lymphocyte population and apoptosis in the lungs of smokers and their relation to emphysema. *Eur Respir J*. (2001) 17:946–53. doi: 10.1183/09031936.01.1750.9460
 58. Aghapour M, Raei P, Moghaddam SJ, Hiemstra PS, Heijink IH. Airway epithelial barrier dysfunction in chronic obstructive pulmonary disease: role of cigarette smoke exposure. *Am J Respir Cell Mol Biol*. (2018) 58:157–69. doi: 10.1165/rcmb.2017-0200TR
 59. Barnes PJ. Inflammatory mechanisms in patients with chronic obstructive pulmonary disease. *J Allergy Clin Immunol*. (2016) 138:16–27. doi: 10.1016/j.jaci.2016.05.011
 60. Wang Y, Xu J, Meng Y, Adcock IM, Yao X. Role of inflammatory cells in airway remodeling in COPD. *Int J Chron Obstruct Pulmon Dis*. (2018) 13:3341–8. doi: 10.2147/COPD.S176122
 61. Lopez-Campos JL, Calero C, Arellano-Orden E, Marquez-Martín E, Cejudo-Ramos P, Ortega Ruiz F, et al. Increased levels of soluble ICAM-1 in chronic obstructive pulmonary disease and resistant smokers are related to active smoking. *Biomark Med*. (2012) 6:805–11. doi: 10.2217/bmm.12.64
 62. Kidney JC, Proud D. Neutrophil transmigration across human airway epithelial monolayers: mechanisms and dependence on electrical resistance. *Am J Respir Cell Mol Biol*. (2000) 23:389–95. doi: 10.1165/ajrcmb.23.3.4068
 63. Shukla SD, Mahmood MQ, Weston S, Latham R, Muller HK, Sohal SS, et al. The main rhinovirus respiratory tract adhesion site (ICAM-1) is upregulated in smokers and patients with chronic airflow limitation (CAL). *Respir Res*. (2017) 18:6. doi: 10.1186/s12931-016-0483-8
 64. Dey T, Kalita J, Weldon S, Taggart CC. Proteases and their inhibitors in chronic obstructive pulmonary disease. *J Clin Med*. (2018) 7:244. doi: 10.3390/jcm7090244
 65. Sommerhoff CP, Nadel JA, Basbaum CB, Caughey GH. Neutrophil elastase and cathepsin G stimulate secretion from cultured bovine airway gland serous cells. *J Clin Invest*. (1990) 85:682–9. doi: 10.1172/JCI114492
 66. Weiss SJ. Tissue destruction by neutrophils. *N Engl J Med*. (1989) 320:365–76. doi: 10.1056/NEJM198902093200606
 67. Aschner Y, Downey GP. Transforming growth factor- β : master regulator of the respiratory system in health and disease. *Am J Respir Cell Mol Biol*. (2016) 54:647–55. doi: 10.1165/rcmb.2015-0391TR
 68. Kurai D, Saraya T, Ishii H, Takizawa H. Virus-induced exacerbations in asthma and COPD. *Front Microbiol*. (2013) 4:293. doi: 10.3389/fmicb.2013.00293
 69. Frickmann H, Jungblut S, Hirche TO, Groß U, Kuhns M, Zautner AE. The influence of virus infections on the course of COPD. *Eur J Microbiol Immunol*. (2012) 2:176–85. doi: 10.1556/EuJMI.2.2012.3.2
 70. Allie SR, Randall TD. Pulmonary immunity to viruses. *Clin Sci*. (2017) 131:1737–62. doi: 10.1042/CS20160259
 71. Schneider D, Ganesan S, Comstock AT, Meldrum CA, Mahidhara R, Goldsmith AM, et al. Increased cytokine response of rhinovirus-infected airway epithelial cells in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. (2010) 182:332–40. doi: 10.1164/rccm.200911-1673OC
 72. Potena A, Caramori G, Casolari P, Contoli M, Johnston SL, Papi A. Pathophysiology of viral-induced exacerbations of COPD. *Int J Chron Obstruct Pulmon Dis*. (2007) 2:477–83.
 73. Jafarinejad H, Moghoofei M, Mostafaei S, Salimian J, Azimzadeh Jamalkandi S, Ahmadi A. Worldwide prevalence of viral infection in AECOPD patients: a meta-analysis. *Microb Pathog*. (2017) 113:190–6. doi: 10.1016/j.micpath.2017.10.021
 74. Christiaansen A, Varga SM, Spencer J V. Viral manipulation of the host immune response. *Curr Opin Immunol*. (2015) 36:54–60. doi: 10.1016/j.coi.2015.06.012
 75. Moreno-Altamirano MMB, Kolstoe SE, Sánchez-García FJ. Virus control of cell metabolism for replication and evasion of host immune responses. *Front Cell Infect Microbiol*. (2019) 9:95. doi: 10.3389/fcimb.2019.0095
 76. McKendry RT, Spalluto CM, Burke H, Nicholas B, Cellura D, Al-Shamkhani A, et al. Dysregulation of antiviral function of cd8(+) t cells in the chronic obstructive pulmonary disease lung. Role of the PD-1-PD-L1 Axis. *Am J Respir Crit Care Med*. (2016) 193:642–51. doi: 10.1164/rccm.201504-0782OC
 77. Singanayagam A, Loo S-L, Calderazzo MA, Finney LJ, Trujillo Torralba M-B, Bakhsholiani E, et al. Anti-viral immunity is impaired in COPD patients with frequent exacerbations. *Am J Physiol Lung Cell Mol Physiol*. (2019) 317:L893–903. doi: 10.1101/632372
 78. Schuliga M. NF-kappaB signaling in chronic inflammatory airway disease. *Biomolecules*. (2015) 5:1266–83. doi: 10.3390/biom5031266
 79. Lawrence T. The nuclear factor NF- κ B pathway in inflammation. *Cold Spring Harb Perspect Biol*. (2009) 1:a001651. doi: 10.1101/cshperspect.a001651
 80. Zhou L, Liu Y, Chen X, Wang S, Liu H, Zhang T, et al. Over-expression of nuclear factor- κ B family genes and inflammatory molecules is related to chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis*. (2018) 13:2131–8. doi: 10.2147/COPD.S164151
 81. Gagliardo R, Chanez P, Profita M, Bonanno A, Albano GD, Montalbano AM, et al. κ B kinase-driven nuclear factor- κ B activation in patients with asthma and chronic obstructive pulmonary disease. *J Allergy Clin Immunol*. (2011) 128:635–45.e1-2. doi: 10.1016/j.jaci.2011.03.045
 82. Banerjee A, Koziol-White C, Panettieri R. p38 MAPK inhibitors, IKK2 inhibitors, and TNF α inhibitors in COPD. *Curr Opin Pharmacol*. (2012) 12:287–92. doi: 10.1016/j.coph.2012.01.016
 83. McMillan DH, Baglole CJ, Thatcher TH, Maggirwar S, Sime PJ, Phipps RP. Lung-targeted overexpression of the NF- κ B member RelB inhibits cigarette smoke-induced inflammation. *Am J Pathol*. (2011) 179:125–33. doi: 10.1016/j.ajpath.2011.03.030
 84. Hsu AC-Y, Dua K, Starkey MR, Haw T-J, Nair PM, Nichol K, et al. MicroRNA-125a and -b inhibit A20 and MAVS to promote inflammation and impair antiviral response in COPD. *JCI Insight*. (2017) 2:e90443. doi: 10.1172/jci.insight.90443
 85. Shehab M, Sherri N, Hussein H, Salloum N, Rahal EA. Endosomal toll-like receptors mediate enhancement of interleukin-17a production triggered by Epstein-Barr virus DNA in mice. *J Virol*. (2019) 93:e00987–19. doi: 10.1128/JVI.00987-19
 86. West JA, Gregory SM, Damania B. Toll-like receptor sensing of human herpesvirus infection. *Front Cell Infect Microbiol*. (2012) 2:122. doi: 10.3389/fcimb.2012.00122
 87. Liu D, Chen Q, Zhu H, Gong L, Huang Y, Li S, et al. Association of respiratory syncytial virus toll-like receptor 3-mediated immune response with COPD exacerbation frequency. *Inflammation*. (2018) 41:654–66. doi: 10.1007/s10753-017-0720-4
 88. Silkoff PE, Flavin S, Gordon R, Loza MJ, Sterk PJ, Lutter R, et al. Toll-like receptor 3 blockade in rhinovirus-induced experimental asthma exacerbations: a randomized controlled study. *J Allergy Clin Immunol*. (2018) 141:1220–30. doi: 10.1016/j.jaci.2017.06.027
 89. Vallath S, Hynds RE, Succony L, Janes SM, Giangreco A. Targeting EGFR signalling in chronic lung disease: therapeutic challenges and opportunities. *Eur Respir J*. (2014) 44:513–22. doi: 10.1183/09031936.00146413
 90. Tyner JW, Kim EY, Ide K, Pelletier MR, Roswit WT, Morton JD, et al. Blocking airway mucous cell metaplasia by inhibiting EGFR antiapoptosis and IL-13 transdifferentiation signals. *J Clin Invest*. (2006) 116:309–21. doi: 10.1172/JCI25167
 91. Guedán A, Swieboda D, Charles M, Toussaint M, Johnston SL, Asfor A, et al. Investigation of the role of protein kinase D in human rhinovirus replication. *J Virol*. (2017) 91:e00217–17. doi: 10.1128/JVI.00217-17
 92. Kim S, Nadel JA. Fibrinogen binding to ICAM-1 promotes EGFR-dependent mucin production in human airway epithelial cells. *Am J Physiol Lung Cell Mol Physiol*. (2009) 297:L174–83. doi: 10.1152/ajplung.00032.2009
 93. Subauste MC, Proud D. Effects of tumor necrosis factor- α , epidermal growth factor and transforming growth factor- α on interleukin-8 production by, and human rhinovirus replication in, bronchial epithelial cells. *Int Immunopharmacol*. (2001) 1:1229–34. doi: 10.1016/S1567-5769(01)00063-7
 94. Ganesan S, Unger BL, Comstock AT, Angel KA, Mancuso P, Martinez FJ, et al. Aberrantly activated EGFR contributes to enhanced IL-8 expression in COPD airways epithelial cells via regulation of nuclear FoxO3A. *Thorax*. (2013) 68:131–41. doi: 10.1136/thoraxjnl-2012-201719
 95. Zhu L, Lee P-K, Lee W-M, Zhao Y, Yu D, Chen Y. Rhinovirus-induced major airway mucin production involves a novel TLR3-EGFR-dependent pathway. *Am J Respir Cell Mol Biol*. (2009) 40:610–9. doi: 10.1165/rcmb.2008-0223OC
 96. Hewson CA, Haas JJ, Bartlett NW, Message SD, Laza-Stanca V, Kebabdzic T, et al. Rhinovirus induces MUC5AC in a human infection model and

- in vitro* via NF- κ B and EGFR pathways. *Eur Respir J.* (2010) 36:1425–35. doi: 10.1183/09031936.00026910
97. Stolarczyk M, Scholte BJ. The EGFR-ADAM17 axis in chronic obstructive pulmonary disease and cystic fibrosis lung pathology. *Mediators Inflamm.* (2018) 2018:1067134. doi: 10.1155/2018/1067134
 98. Ueki IF, Min-Oo G, Kalinowski A, Ballon-Landa E, Lanier LL, Nadel JA, et al. Respiratory virus-induced EGFR activation suppresses IRF1-dependent interferon λ and antiviral defense in airway epithelium. *J Exp Med.* (2013) 210:1929–36. doi: 10.1084/jem.20121401
 99. Kalinowski A, Galen BT, Ueki IF, Sun Y, Mulenon A, Osafo-Addo A, et al. Respiratory syncytial virus activates epidermal growth factor receptor to suppress interferon regulatory factor 1-dependent interferon-lambda and antiviral defense in airway epithelium. *Mucosal Immunol.* (2018) 11:958–67. doi: 10.1038/mi.2017.120
 100. Abe T, Marutani Y, Shoji I. Cytosolic DNA-sensing immune response and viral infection. *Microbiol Immunol.* (2019) 63:51–64. doi: 10.1111/1348-0421.12669
 101. Chan YK, Gack MU. Viral evasion of intracellular DNA and RNA sensing. *Nat Rev Microbiol.* (2016) 14:360–73. doi: 10.1038/nrmicro.2016.45
 102. Orzalli MH, Broekema NM, Diner BA, Hancks DC, Elde NC, Cristea IM, et al. cGAS-mediated stabilization of IFI16 promotes innate signaling during herpes simplex virus infection. *Proc Natl Acad Sci USA.* (2015) 112:E1773–81. doi: 10.1073/pnas.1424637112
 103. Zhu J, Message SD, Mallia P, Keadze T, Contoli M, Ward CK, et al. Bronchial mucosal IFN- α/β and pattern recognition receptor expression in patients with experimental rhinovirus-induced asthma exacerbations. *J Allergy Clin Immunol.* (2019) 143:114–25.e4. doi: 10.1016/j.jaci.2018.04.003
 104. Farazuddin M, Mishra R, Jing Y, Srivastava V, Comstock AT, Sajjan US. Quercetin prevents rhinovirus-induced progression of lung disease in mice with COPD phenotype. *PLoS ONE.* (2018) 13:e0199612. doi: 10.1371/journal.pone.0199612
 105. Traub S, Nikonova A, Carruthers A, Dunmore R, Vousden KA, Gogsadze L, et al. An anti-human ICAM-1 antibody inhibits rhinovirus-induced exacerbations of lung inflammation. *PLoS Pathog.* (2013) 9:e1003520. doi: 10.1371/journal.ppat.1003520
 106. Mirabelli C, Scheers E, Neyts J. Novel therapeutic approaches to simultaneously target rhinovirus infection and asthma/COPD pathogenesis. *F1000Res.* (2017) 6:1860. doi: 10.12688/f1000research.11978.1
 107. Beigel JH, Nam HH, Adams PL, Krafft A, Ince WL, El-Kamary SS, et al. Advances in respiratory virus therapeutics - a meeting report from the 6th isirv antiviral group conference. *Antiviral Res.* (2019) 167:45–67. doi: 10.1016/j.antiviral.2019.04.006
 108. Mousnier A, Bell AS, Swieboda DP, Morales-Sanfrutos J, Pérez-Dorado I, Brannigan JA, et al. Fragment-derived inhibitors of human N-mycristoyltransferase block capsid assembly and replication of the common cold virus. *Nat Chem.* (2018) 10:599–606. doi: 10.1038/s41557-018-0039-2
 109. Schaefer N, Li X, Seibold MA, Jarjour NN, Denlinger LC, Castro M, et al. The effect of BPIFA1/SPLUNC1 genetic variation on its expression and function in asthmatic airway epithelium. *JCI Insight.* (2019) 4:e127237. doi: 10.1172/jci.insight.127237
 110. Jiang D, Wenzel SE, Wu Q, Bowler RP, Schnell C, Chu HW. Human neutrophil elastase degrades SPLUNC1 and impairs airway epithelial defense against bacteria. *PLoS ONE.* (2013) 8:e64689. doi: 10.1371/journal.pone.0064689
 111. Seys LJM, Verhamme FM, Dupont LL, Desauter E, Duerr J, Seyhan Agircan A, et al. Airway surface dehydration aggravates cigarette smoke-induced hallmarks of COPD in mice. *PLoS ONE.* (2015) 10:e0129897. doi: 10.1371/journal.pone.0129897
 112. Tcherniuk S, Cenac N, Comte M, Frouard J, Errazuriz-Cerda E, Galabov A, et al. Formyl peptide receptor 2 plays a deleterious role during influenza A virus infections. *J Infect Dis.* (2016) 214:237–47. doi: 10.1093/infdis/jiw127
 113. Ampomah PB, Moraes LA, Lukman HM, Lim LHK. Formyl peptide receptor 2 is regulated by RNA mimics and viruses through an IFN- β -STAT3-dependent pathway. *FASEB J.* (2018) 32:1468–78. doi: 10.1096/fj.201700584RR
 114. Lupfer C, Malik A, Kanneganti T-D. Inflammasome control of viral infection. *Curr Opin Virol.* (2015) 12:38–46. doi: 10.1016/j.coviro.2015.02.007
 115. Chen I-Y, Ichinohe T. Response of host inflammasomes to viral infection. *Trends Microbiol.* (2015) 23:55–63. doi: 10.1016/j.tim.2014.09.007
 116. Colarusso C, Terlizzi M, Molino A, Pinto A, Sorrentino R. Role of the inflammasome in chronic obstructive pulmonary disease (COPD). *Oncotarget.* (2017) 8:81813–24. doi: 10.18632/oncotarget.17850
 117. Hikichi M, Mizumura K, Maruoka S, Gon Y. Pathogenesis of chronic obstructive pulmonary disease (COPD) induced by cigarette smoke. *J Thorac Dis.* (2019) 11:S2129–40. doi: 10.21037/jtd.2019.10.43

Conflict of Interest: JK was a non-executive director and shareholder of HiberGene Diagnostics Ltd.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Guo-Parke, Linden, Weldon, Kidney and Taggart. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.