



**QUEEN'S
UNIVERSITY
BELFAST**

Decreasing cerebral palsy prevalence in multiple births in the modern era: A population cohort study of European data

Perra, O., Rankin, J., Platt, M. J., Sellier, E., Arnaud, C., De La Cruz, J., Kraegeloh-Mann, I., Sweet, D., & Bjellmo, S. (2020). Decreasing cerebral palsy prevalence in multiple births in the modern era: A population cohort study of European data. *Archives of Disease in Childhood*. Advance online publication. <https://doi.org/10.1136/archdischild-2020-318950>

Published in:
Archives of Disease in Childhood

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2020 the authors.

This is an open access Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits use, distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>



OPEN ACCESS

Decreasing cerebral palsy prevalence in multiple births in the modern era: a population cohort study of European data

Oliver Perra,¹ Judith Rankin,² Mary Jane Platt,³ Elodie Sellier,^{4,5} Catherine Arnaud,^{6,7} Javier De La Cruz,⁸ Ingeborg Krägeloh-Mann,⁹ David G Sweet,¹⁰ Solveig Bjellmo^{11,12}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/archdischild-2020-318950>).

For numbered affiliations see end of article.

Correspondence to

Dr Oliver Perra, School of Nursing and Midwifery; Centre for Evidence and Social Innovation, Queen's University Belfast, Belfast BT9 7LB, UK; o.perra@qub.ac.uk

Received 31 January 2020
Revised 26 May 2020
Accepted 15 June 2020

ABSTRACT

Multiple births (twins or higher order multiples) are increasing in developed countries and may present higher risk for cerebral palsy (CP). However, few studies can reliably investigate trends over time because these outcomes are relatively rare.

Objective We pooled data from European CP registers to investigate CP birth prevalence and its trends among single and multiple births born between 1990 and 2008.

Design Population cohort study.

Setting 12 population-based registers from the Surveillance of Cerebral Palsy in Europe collaboration.

Participants 4 446 125 single and multiple live births, of whom 8416 (0.19%) had CP of prenatal or perinatal origin.

Main outcomes CP diagnosis ascertained in childhood using harmonised methods; CP subtype; Motor impairment severity among CP cases.

Results The rate of multiple births increased from 1990. Multiples displayed higher risk for CP (RR=4.27, 95% CI 4.00 to 4.57). For singletons and multiples alike, risk for CP was higher among births of lower gestational age (GA) or birth weight (BW). However, CP birth prevalence declined significantly among very preterm (<32 weeks) and very low BW (<1500 g) multiples. Singletons and multiples with CP displayed similar severity of motor impairment.

Conclusions Between 1990 and 2008, CP birth prevalence decreased steadily among multiples with low GA or BW. Furthermore, multiples with CP display similar profiles of severe motor impairment compared with CP singletons. Improvements in management of preterm birth since the 1990s may also have been responsible for providing better prospects for multiples.

INTRODUCTION

The rate of multiple births in developed countries^{1–6} has risen in the last four decades. Studies^{7–11} report that multiple births are associated with increased risk for cerebral palsy (CP), defined as lifetime physical disabilities due to non-progressive injuries to the immature or developing brain.^{12–14} Twins and higher order multiples are more likely to be born before term¹ and to present low birth weight (LBW);¹⁵ prematurity and LBW are, in turn, significant risk factors for CP;¹⁶ thus, increased rates of multiple births might expose more newborns to risk for CP.¹⁷ However, European-wide data¹⁸ suggest that the prevalence of CP has declined between birth years 1980 and 2003, particularly among

What is already known on this topic?

- The rate of multiple births is increasing, and multiple births are at higher risk for cerebral palsy (CP). However, changes in obstetric practice since the 1990s may have impacted on risk for CP among preterm and low birth weight multiples.
- Multiple births and CP are relatively rare events: their study necessitates pooling of data from different centres and countries.

What this study adds?

- Multiples remain at higher risk for CP, but this risk has been decreasing steadily since the 1990s among very preterm or low birth weight multiples.
- Singleton and multiple births with CP display similar severity of motor impairment.
- Advances in obstetric care may be responsible for these changes in CP risk among multiples.

very and moderate LBW infants. Improvements in neonatal care responsible for these trends may also have impacted on risk among multiple births. An analysis of recent trends in CP prevalence among multiple births is timely.

We investigated the birth prevalence of CP among single and multiple births born between 1990 and 2008 using data collected by the multi-centre Surveillance of Cerebral Palsy in Europe (SCPE): by harmonising definitions and methods, this collaboration provides a reliable source of data on CP prevalence. Topp and colleagues¹⁰ analysed data from SCPE cohorts until 1990: in our study, we analyse trends in births between 1990 and 2008 to investigate cohorts born in the antenatal corticosteroids¹⁹ and surfactant²⁰ era. A diagnosis of CP can only be confirmed reliably in childhood;²¹ data on children with ascertained CP were available up to birth year 2008 at the time of analyses.

Topp and colleagues¹⁰ reported a higher proportion of second-born twins among twins with CP and suggested that this may indicate risks for intrapartum asphyxia of the second-born twin during a vaginal delivery. A different study⁸ on USA and Australia populations did not confirm differences in



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Perra O, Rankin J, Platt MJ, et al. *Arch Dis Child Fetal Neonatal Ed* Epub ahead of print: [please include Day Month Year]. doi:10.1136/archdischild-2020-318950

risk for CP associated with twins' birth order. Investigating the order of birth of twins across different types of birth may help clarify this issue.

Our main aim was to investigate changes in risk for CP among multiple births. For this purpose, we analysed trends in CP prevalence by categories of gestational age (GA) and birth weight (BW). Our objectives were to:

1. Investigate prevalence of CP over time in multiples and singletons overall, and across GA and BW categories;
2. Compare CP type and severity of impairment between multiples and singletons with CP;
3. Investigate associations between mode of delivery and birth order of twins with CP.

METHODS

Study design

A population cohort study: population-based registers participating to SCPE provided information on children with CP, as well as denominators (total live births) by type of birth (single versus multiple) and by GA and BW categories in their catchment area. Some of these registers cover entire countries, while others cover defined areas within countries.

Participants

CP is a group of permanent disorders of movement and/or posture caused by non-progressive interferences/lesions to or abnormalities of the developing brain.^{12 14} SCPE registers use a range of sources for the notification and diagnostic ascertainment of eligible participants. Children with a postneonatal onset of CP were excluded from the CP population.²² Methods used by participating registers are detailed elsewhere.^{14 23}

Setting

Overall, 23 population-based registers provided data for births between 1 January 1990 and 31 December 2008. A diagnosis of CP is reliable in later childhood:²¹ SCPE registers confirm CP diagnoses after age 4 years, with the exception of deaths that take place before a suspected diagnosis is confirmed (age 2–4 years). The age when registers consider ascertainment of CP diagnoses complete varies across countries, but it takes place by the time children are of primary education age. At the time of analyses, data were available for children born until 2008.

In order to capture changes over time, we stipulated that centres had to provide information on CP and multiple and single births denominators by GA or BW categories for at least 3 years: data from 11 centres were excluded for not providing this information (see online supplementary appendix 1). Overall, 12 centres were included in the analyses: 2 provided only denominators on multiple births by BW and are thus excluded in prevalence analyses stratified by GA. The number of births by centre and birth type are reported in table 1.

Variables

CP subtypes were classified as unilateral spastic, bilateral spastic, dyskinetic and ataxic.²²

The Gross Motor Function Classification System (GMFCS) provided a reliable assessment of children's severity of motor impairment.^{24 25} However, some SCPE registers adopted this measure later than others. To avoid missing data and provide a meaningful measure of *Severity of Motor Impairment*, we categorised individuals with *moderate-severe impairment* to those with GMFCS level III to V or who were not able to walk without assistive devices.

Table 1 Number of children with CP, multiples with CP, live births, multiple live births and number of cohort birth years for which data were available by register

Register	N CP	Multiples CP	Live births	Multiple live births	Cohort birth years
Cork & Kerry, Ireland	127	26	82 019	2204	9
Gothenburg, Sweden	835	98	421 050	12 229	19
NW England, UK	1153	106	534 316	14 167	16
Oxford, UK	737	84	483 734	15 014	14
Denmark	2106	263	1 029 591	39 613	18
Viterbo, Italy	142	14	37 886	751	18
Norway	1257	158	638 888	22 556	10
Madrid, Spain	79	15	54 851	1180	9
Slovenia	309	45	142 311	4651	8
Portugal	1440	150	871 439	22 583	8
Latvia	93	4	61 210	1418	9
Hungary	138	15	88 830	2883	9

CP, cerebral palsy.

In analyses that involved only children with CP and their outcomes we controlled for GA and BW: to differentiate between low BW because of prematurity and small for gestation age, CP children were assigned a *Standardised BW z Score* to indicate deviation from the expected weight for their GA and sex. BW norms were different for singletons and multiples (see online supplementary appendix 2).

Statistical analysis

CP prevalence is reported per 1000 live births. CP prevalence across multiples and singletons is compared using risk ratios (RR).

We analysed trends over time using multilevel Negative Binomial models whereby observations were nested within registers. Multilevel models allow controlling for heterogeneity across registers (see online supplementary appendix 3). Models were adjusted for offset terms to account for differences in the number of live births across registers' catchment areas. Time trends described changes in prevalence year by year. We investigated linear and non-linear (quadratic) trends, and tested nested models using likelihood-ratio χ^2 tests with a threshold of $p < 0.005$. Coefficients reported are incidence rate ratios (IRR).

In analyses of outcomes among children with CP, we used multilevel logistic regressions to model associations between the outcome and birth type, successively adding covariates GA category and BW *z scores* (see online supplementary appendix 3). Whether the CP birth was single or multiple was unknown for 272 cases (3.23%), who were excluded from analyses: sensitivity analyses were run assigning unknown births to singletons and then to multiples, which replicated the patterns of results presented.

RESULTS

Multiple births and CP

Results indicated an increase of multiple births over time that followed a quadratic trend: 2.47% of total births were multiples in 1990, increasing to 3.34% in 2000, and a successive deceleration (eg, 2.76% in 2008): trends differed across centres (see online supplementary appendix 4).

The overall prevalence of CP cases per 1000 live births among singletons was 1.64 (95% CI 1.60 to 1.68), but was 7.02 CP

Table 2A Multilevel negative binomial regression coefficients of the total number of CP births on total live births by singletons and multiples

Model parameters	Singletons	95% CI	Multiples	95% CI
Intercept	0.002	0.001 to 0.003	0.010	0.001 to 0.013
Yearly rate of change	0.976	0.952 to 1.001	0.972	0.955 to 0.990
Overdispersion: Ln(Alpha)	-5.691	-7.332 to -4.049	-3.008	-4.039 to -1.977
Between-centre variation				
Initial status	0.239	0.092 to 0.621	0.042	0.006 to 0.288
Rate of change	0.002	0.001 to 0.004	NA	
Covariance	-0.017	-0.035 to 0.001	NA	
Model fit	Wald $\chi^2(1)=3.49, p=0.062$		Wald $\chi^2(1)=8.94, p=0.003$.	

Parameters in bold are significant at $p<0.005$. Comparisons of models using the likelihood ratio test indicated that estimating centre-level variance around the rate of change and the covariance between intercept and rate of change of Multiples did not improve model fit (at $p<0.005$). Therefore, these terms were not modelled.

cases per 1000 live births among multiples (95% CI 6.58 to 7.48): risk ratio (RR)=4.27, 95% CI 4.00 to 4.57. We report CP prevalence trends across singletons and multiples in [table 2A](#): these indicated a non-significant linear decrease in prevalence among singletons (IRR=0.98), while a linear decrease among multiples (IRR=0.97) was significant.

Prevalence of CP by multiple birth and GA

Results between birth years 1990–2008 did not indicate substantial differences across singletons and multiples born extremely or very preterm (<32 weeks), RR=0.97 (95% CI 0.86 to 1.08) or between singletons and multiples born moderately preterm (32–36 weeks), RR=0.96 (95% CI 0.82 to 1.11) (see [table 2B](#)). However, full-term multiples displayed higher risk for CP compared with full-term singletons, RR=1.89 (95% CI 1.56 to 2.30).

Significant decreases in the prevalence of CP were revealed for singletons and multiples born less than 32 weeks GA and among singletons born 32–36 weeks GA (see [figure 1](#) and [table 3](#)). Prevalence of CP remained stable for multiples born 32–36 weeks GA and did not change significantly for full-term singletons or full-term multiples (see also online supplementary appendix 5).

Prevalence of CP by multiple birth and BW

The prevalence of CP between 1990 and 2008 was similar for singletons and multiples born weighing less than 1500g, RR=0.95 (95% CI 0.85 to 1.05). The prevalence of CP was higher among singletons born weighing between 1500 and less than 2500g when compared with multiples of the same BW, RR=0.79 (95% CI 0.70 to 0.88). In contrast, multiples born




Figure 1 Prevalence of cerebral palsy across singletons and multiples by year and by categories of GA and BW. Raw trends over time are represented by smoothed curves, estimated using locally weighted scatterplot smoothing. BW, birth weight; GA, gestational age.

with normal BW (≥ 2500 g) displayed a higher risk for CP compared with normal BW singletons, RR=1.92, (95% CI 1.61 to 2.30) (see [table 2B](#)).

A linear decrease in CP prevalence was observed for both very LBW singletons and multiples and among moderately LBW singletons, but not among moderately LBW multiples (see [figure 1](#) and [table 3](#)). No significant changes across time were observed among normal BW singletons or normal BW multiples (see online supplementary file 1).

Table 2B CP prevalence by singletons and multiples and by categories of gestational age and birth weight

	Singletons	95% CI	Multiples	95% CI	RR	95% CI
Gestational age						
<32 weeks	47.35	44.50 to 50.34	45.84	41.43 to 50.60	0.97	0.86 to 1.08
32–36 weeks	5.47	5.09 to 5.87	5.24	4.56 to 5.99	0.96	0.82 to 1.11
Birth weight						
37+ weeks	1.07	1.03 to 1.11	2.02	1.65 to 2.45	1.89	1.56 to 2.30
<1500g	42.79	40.37 to 45.32	40.48	36.76 to 44.47	0.95	0.85 to 1.05
1500–2499g	8.37	7.92 to 8.83	6.58	5.90 to 7.31	0.79	0.70 to 0.88
2500+ g	1.07	1.03 to 1.10	2.05	1.71 to 2.44	1.92	1.61 to 2.30

RRs represent the CP risk of multiples compared with singletons. Parameters in bold are significant at $p<0.005$

CP, cerebral palsy; RR, risk ratio.

Table 3 IRRs indicating yearly changes in the total number of CP cases on the total number of live births, by GA or BW categories and by singletons or multiples

	Singletons	IRR	95% CI	Model Wald χ^2	Cross-centre variability in slope?	Multiples	IRR	95% CI	Model Wald χ^2	Cross-centre variability in slope?
GA										
<32 weeks	Linear yearly decrease	0.96*	0.95 to 0.98	$\chi^2(1)=21.38^*$	No	Linear yearly decrease	0.95*	0.93 to 0.98	$\chi^2(1)=14.62^*$	No
32–36 weeks	Linear yearly decrease	0.96*	0.94 to 0.98	$\chi^2(1)=17.50^*$	No	No yearly change	0.99	0.96 to 1.03	$\chi^2(1)=0.10$	No
37+ weeks	No yearly change	0.99	0.96 to 1.02	$\chi^2(1)=0.29$	Yes	No yearly change	0.98	0.94 to 1.02	$\chi^2(1)=0.77$	No
BW										
<1500 g	Linear yearly decrease	0.96*	0.95 to 0.98	$\chi^2(1)=20.87^*$	No	Linear yearly decrease	0.95*	0.93 to 0.97	$\chi^2(1)=19.87^*$	No
1500–<2500 g	Linear yearly decrease	0.97*	0.96 to 0.98	$\chi^2(1)=18.88^*$	No	No yearly change	0.99	0.96 to 1.01	$\chi^2(1)=0.73$	No
≥2500 g	No yearly change	0.98	0.95 to 1.01	$\chi^2(1)=1.08$	Yes	No yearly change	0.96	0.92 to 1.00	$\chi^2(1)=4.01$	No

We also reported whether the models indicated cross-centre variability in the rate of change (slope) of CP prevalence. IRRs were estimated using multilevel negative binomial regressions. Parameters of the models are introduced in online supplementary appendix 4 and full parameters reported in online supplementary appendices 5 and 6. Cross-centre variability in slope indicated that different centres displayed varying rates of change in the CP prevalence for that category of birth and GA or BW. Conversely, if no cross-centre variability was observed, the rate of change of CP prevalence was estimated to be consistent across centres. The lack of significant cross-centre variability in the slopes of CP prevalence for singletons and multiples in lower GA and BW categories indicated that the significant decrease of CP cases among these at-risk categories was consistent across centres.

*P<0.005.

BW, birth weight; CP, cerebral palsy; GA, gestation age; IRR, incidence rate ratios.

Differences in CP type and motor impairment

A higher percentage of multiples met criteria for spastic bilateral CP (see table 4) and displayed higher odds of presenting this subtype rather than other subtypes, adjusted OR=1.39 (95% CI 1.21 to 1.61); ORs were adjusted for birth year and centre-level effects. However, these differences were non-significant once we controlled for variations in GA and BW z scores: adjusted OR=0.87 for multiples compared with singletons (95% CI 0.74 to 1.01). Compared with full-term birth, birth before 32 weeks was associated with a threefold increase in the odds of spastic bilateral CP, OR=3.16 (95% CI 2.80 to 3.56). Higher BW for GA was also associated with reduced odds of spastic bilateral CP, OR=0.95 (95% CI 0.93 to 0.98).

The percentage of singletons and multiples with moderate-severe motor impairment was 43% in both cases: (n=2975/6880) and (n=402/941), respectively. Models (see online supplementary appendix 7) indicated a linear decrease in the odds of moderate-severe impairment (OR=0.98, 95% CI 0.97 to 0.99), but an interaction term suggested this trend did not differ between singletons and multiples, LR test $\chi^2(1)=0.26$, p=0.61.

CP risk and birth order

Twin children with CP did not display different birth order: 50.39% (n=384/762) were second-born (binomial test assuming equal probability of being first-born or second-born: p=0.86). A higher proportion of twins with CP born by vaginal delivery was first-born rather than second-born: 55.39% (n=113/204). The proportion of first-born and second-born CP twins did not differ

significantly among emergency caesarean deliveries, 47.57% being first born (n=98/206).

DISCUSSION

We investigated the prevalence of CP in multiples and singletons using population-based data collated from a European collaboration. By harmonising methods and definitions, and by adopting robust procedures for ascertainment, this collaboration provides reliable data on the condition over time. The results indicate that multiple births had been increasing since 1990, and they were associated with increased risk for CP. This increased risk was related to the higher prevalence of preterm or LBW births among multiples, which was also underlined by the finding that multiples with CP displayed the bilateral spastic CP-subtype typical of lower GA and BW groups. Despite multiples being at increased risk for CP, CP prevalence decreased linearly among multiples in at-risk categories because of prematurity and LBW.

In particular, while CP prevalence among multiples born less than 32 weeks was higher for cohorts born in the early 1990s, this declined significantly, becoming comparable to the CP prevalence among singletons of the same GA (see figure 1). A significant decline in CP prevalence was also observed among multiples weighing less than 1500 g (LBW). Results based on SCPE data indicated an analogous decline in CP prevalence among LBW European children:¹⁸ here, we show that this decline involves both multiples and singletons. Furthermore, the decline in CP prevalence among the lowest GA and BW categories of singletons and multiples was consistent across participating registers (see table 3): this has potential clinical significance (see online supplementary file 1).

The higher risk for CP among singletons in the Moderate LBW category (1500–2499 g) is consistent with findings from another large population study:⁸ however, our findings also show that this difference has been narrowing over time.

In contrast, CP prevalence among multiples born at term did not change significantly between 1990 and 2008. Overall, 2.02 per 1000 full-term multiples births were diagnosed with CP, which is substantially higher than the prevalence of 1.07 per 1000 births among full-term singletons. A specific pathogenic mechanism may be suspected for CP in full-term multiples. Some research suggests a role for intrapartum risks, for example, twin-twin transfusion syndrome.⁸ Other studies provided evidence of intrauterine growth divergence being associated with increasing risk for CP:^{26,27} since growth divergence is a common occurrence

Table 4 Number and percentages of singletons and multiples by CP type

	Singletons	Multiples	Total	
Spastic bilateral	N	3610	570	4180
	%	50.38	58.28	51.33
Spastic unilateral	N	2471	309	2780
	%	34.48	31.60	34.14
Other CP type	N	929	80	1009
	%	12.96	8.18	12.39
CP type uncertain	N	156	19	175
	%	2.18	1.94	2.15

Pearson $\chi^2(3)=28.70$, p<0.001.

CP, cerebral palsy.

in multiple pregnancies,⁸ this may be a key factor in determining differential patterns of risk between singletons and multiples, particularly when the pregnancy reaches term.

Among children with CP, multiples and singletons displayed similar profiles of motor impairment severity. Furthermore, we report a significant linear decrease in the proportion of the CP population with moderate-severe motor impairment that did not vary significantly between singletons and multiples. Improvements in care that have contributed to an overall decrease of moderate-severe CP across European countries¹⁸ may have also reduced risk for severe impairment among multiples.

Results did not indicate a higher proportion of second-born among twins with CP, and when considering those twins born by vaginal delivery, second-born twins were proportionally fewer (45%). If intrapartum asphyxia could explain CP risk of second-born twins in cohorts before the 1990s,²⁸ our results suggest that changes in obstetric care since 1990 have countered this pathogenic mechanism.

Our findings were limited by our inability to investigate trends in birth years after 2008: this is due to the need for reliable CP ascertainment by registers. Furthermore, many variables that could inform about potential causal mechanisms linking multiple birth and CP were not reported. For example, a causal pathway may involve increased risk for placental abruption in multiple births,²⁹ which in turn may contribute to preterm birth³⁰ and CP.^{11 31} Advanced maternal age³² and assisted reproductive technologies (ARTs)³³ may also confound the association between multiple birth and CP.³⁴

CONCLUSION

Our study is unique in using reliable observations of multiple births and CP collected over a significant period, thus providing statistical power to detect key trends. Results support reports of an increased risk for CP in multiples.

Studies linked increased rates of multiples to availability of ARTs and higher maternal age:^{35–37} our findings of increased CP risk in multiples highlight the importance of limiting the multiple gestation risk of ARTs.^{33 38} However, multiples have profited equally from improvements in management of premature births since the 1990s, as indicated by the decrease in CP prevalence among multiples of lower GA or BW.

The continuously higher prevalence of CP in multiples born at term is, however, of concern. Other studies indicate multiples' increased risk for mortality and severe disability after reaching 38 weeks' gestation^{8 15} and recommendations acknowledge these risks: for example, NICE guidelines recommend informing women with uncomplicated dichorionic diamniotic and monochorionic diamniotic twin pregnancies about the risks of continuing the pregnancy beyond 37⁺⁶ and 36⁺⁶ weeks, respectively.³⁹ More research using population-based studies is needed to investigate the causal mechanisms that link multiple births to increased risk for CP.³⁴

Author affiliations

¹School of Nursing and Midwifery, Centre for Evidence and Social Innovation, Queen's University Belfast, Belfast, UK

²Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK

³Norwich Medical School, University of East Anglia, Norwich, UK

⁴University Grenoble Alpes, CNRS, Grenoble INP, CHU Grenoble Alpes, TIMC-IMAG, Grenoble, France

⁵RHEOP, Grenoble, France

⁶Public Health Department; UMR 1027 SPHERE, Université Toulouse III Paul Sabatier, Toulouse, France

⁷Clinical Epidemiology Unit, University Hospital Centre Toulouse, Toulouse, France

⁸Health Research Institute, University Hospital of Madrid, Madrid, Spain

⁹Department of Paediatric Neurology and Developmental Medicine, University Children's Hospital, Eberhard Karls Universität Tübingen, Tübingen, Germany

¹⁰Regional Neonatal Unit, Royal Maternity Hospital, Belfast Health and Social Care Trust, Belfast, UK

¹¹Department of Obstetrics and Gynecology, Helse More og Romsdal HF, Alesund, Norway

¹²NTNU, Norwegian University of Science and Technology, Trondheim, Norway

Twitter Oliver Perra @oliver_perra and David G Sweet @dsweetbelfast

Acknowledgements We wish to acknowledge the contribution of all SCPE centres and collaborators: the RHEOP, France; RHE31, France; Lavanagh Centre, Cork and Kerry counties, Ireland; CP Register of Western Sweden, Sweden; Eastern Ireland Area CP study, Dublin, Ireland; NPEU, Oxford, UK; The Danish Cerebral Palsy Register, Denmark; Central Italy Cerebral Palsy Register, Italy; The Cerebral Palsy Registry of Norway, Norway; ASL, Bologna, Italy; G Gaffney, Galway, Ireland; Madrid Cerebral Palsy Register, Spain; The Northern Ireland Cerebral Palsy Register, Northern Ireland, UK; Slovenian Register of Cerebral Palsy, Slovenia; Programa Vigilancia Nacional da Paralisia Cerebral aos 5 anos, Portugal; Riga Association Rehabilitation Center, Latvia; Cerebral Palsy Register of South-West Hungary, Hungary; Nicosia, Cyprus; Icelandic Cerebral Palsy Register, Iceland; Register of Cerebral Palsy in Tyrol, Austria; Belgian Cerebral Palsy Registry, Belgium; Croatian Cerebral Palsy Register, Croatia; CP register –St Gallen Canton, Switzerland; The Cerebral Palsy Register of Malta, Malta; The Cerebral Palsy Register of Attica-Greece, Greece; Sunderland, Washington, Coalfields and North Easington Cerebral Palsies Register, UK; ISDSHS, Edinburgh, UK; University of Newcastle, Newcastle, UK. This project was completed on behalf of the Surveillance of Cerebral Palsy Europe Network group: A Ego, C Tronc, C Arnaud, M Delobel, V Mc Manus, A Lyons, K Himmelmann M Pahlman, V Dowding, K McConnell, O Perra, J Kurinczuk, G Surman, P Uldall, G Rackauskaite, MG Torrioli, M Marcelli, G L Andersen, S Julsen Hollung, M Bottos, G Gaffney, J de la Cruz, C Pallas, D Neubauer, Anja Radsel, D Virella, M Andrada, A Greitane, K Hollody, I Dakovic, S Sigurdardottir, I Einarsson, K Rostasy, F Zeiner, E Ortibus, I Franki, V Mejaski-Bosnjak, C Kuenzle, A Tschertner, S Attard, A Papavasiliou, K Horridge, MJ Platt, I Krägeloh-Mann, V Horber. We also thank the generous contribution and support provided by the Joint Research Centre of the European Commission, and particularly Simona Martin, Agnieszka Kinsner-Ovaskainen and Monica Lanzoni.

Contributors OP designed and conducted the analyses reported in this study and wrote the manuscript. JR contributed significantly to the design of these analyses. JR, MJP, ES, CA, JDL, IK-M, DGS and SB all contributed significantly to the final draft of this manuscript by discussing results and their interpretations and by revising earlier drafts.

Funding Data reported are collected by the Surveillance of Cerebral Palsy in Europe (SCPE) collaboration. SCPE work was supported by grants from the European Commission (SCPE-NET 2008 13 07/FIB_H120_FY2014 OG).

Disclaimer The European Commission had no role in study design, data collection, data analysis, data interpretation, or the writing of the report.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The SCPE network has no specific ethical approval as it only gathers anonymised data. Each register had its own ethical approval that follows the legislative rules of its country.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. Data have been obtained under request to the Joint Research Centre of the European Commission, which manages and provides them on behalf of the Surveillance of Cerebral Palsy in Europe.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Blondel B, Macfarlane A, Gissler M, *et al*. Preterm birth and multiple pregnancy in European countries participating in the PERISTAT project. *BJOG* 2006;113:528–35.
- Luke B, Martin JA. The rise in multiple births in the United States: who, what, when, where, and why. *Clin Obstet Gynecol* 2004;47:118–33.
- Ghosh K. Birth characteristics in England and Wales: 2017. *Stat Bull Off Natl Stat* 2019:1–14.
- Martin JA, Hamilton BE DP, *et al*. National Vital Statistics Reports Births : Final Data for 2013. *Statistics* 2015;64:1–104.

- 5 Choi SH, Park YS, Shim KS, *et al.* Recent trends in the incidence of multiple births and its consequences on perinatal problems in Korea. *J Korean Med Sci* 2010;25:1191–6.
- 6 Fell DB, Joseph K. Temporal trends in the frequency of twins and higher-order multiple births in Canada and the United States. *BMC Pregnancy Childbirth* 2012;12:103.
- 7 Williams K, Hennessy E, Alberman E. Cerebral palsy: effects of twinning, birthweight, and gestational age. *Arch Dis Child Fetal Neonatal Ed* 1996;75:F178–82.
- 8 Scher AI, Petterson B, Blair E, *et al.* The risk of mortality or cerebral palsy in twins: a collaborative population-based study. *Pediatr Res* 2002;52:671–81.
- 9 Pharoah PO, Cooke T. Cerebral palsy and multiple births. *Arch Dis Child Fetal Neonatal Ed* 1996;75:F174–7.
- 10 Topp M, Huusom LD, Langhoff-Roos J, *et al.* Multiple birth and cerebral palsy in Europe: a multicenter study. *Acta Obstet Gynecol Scand* 2004;83:548–53.
- 11 Trønnes H, Wilcox AJ, Lie RT, *et al.* Risk of cerebral palsy in relation to pregnancy disorders and preterm birth: a national cohort study. *Dev Med Child Neurol* 2014;56:779–85.
- 12 Rosenbaum P, Paneth N, Leviton A, *et al.* A report: the definition and classification of cerebral palsy April 2006. *Dev Med Child Neurol Suppl* 2007;109:8–14.
- 13 Kinsner-Ovaskainen A, Lanzoni M, Delobel M. *Surveillance of cerebral palsy in Europe development of the JRC-SCPE central database and public health indicators.* Luxembourg: Publications Office of the European Union, 2017.
- 14 Cans C *et al.* Surveillance of cerebral palsy in Europe: a collaboration of cerebral palsy surveys and registers. *Dev Med Child Neurol* 2000;42:816–24.
- 15 Luke B, Keith LG. The contribution of singletons, twins and triplets to low birth weight, infant mortality and handicap in the United States. *J Reprod Med* 1992;37:661–6.
- 16 Colver A, Fairhurst C, Pharoah POD. Cerebral palsy. *Lancet* 2014;383:1240–9.
- 17 Kiely JL, Kiely M, Blickstein I. Contribution of the rise in multiple births to a potential increase in cerebral palsy. *Pediatr Res* 2000;47:314A.
- 18 Sellier E, Platt MJ, Andersen GL, *et al.* Decreasing prevalence in cerebral palsy: a multi-site European population-based study, 1980 to 2003. *Dev Med Child Neurol* 2016;58:85–92.
- 19 WHO. WHO recommendations on interventions to improve preterm birth outcomes, 2019. Available: https://www.who.int/reproductivehealth/publications/maternal_perinatal_health/preterm-birth-guideline/en/ [Accessed 25 Sep 2019].
- 20 Engle WA, American Academy of Pediatrics Committee on Fetus and Newborn. Surfactant-replacement therapy for respiratory distress in the preterm and term neonate. *Pediatrics* 2008;121:419–32.
- 21 Nelson KB, Ellenberg JH. Children who "outgrew" cerebral palsy. *Pediatrics* 1982;69:529–36.
- 22 Cans C, Dolk H, Platt MJ, *et al.* Recommendations from the SCPE Collaborative group for defining and classifying cerebral palsy. *Dev Med Child Neurol* 2007;49:35–8.
- 23 Cans C, Guillem P, Arnaud C, *et al.* Prevalence and characteristics of children with cerebral palsy in Europe. *Dev Med Child Neurol* 2002;44:633–40.
- 24 Palisano RJ, Gorter JW, Morris C, *et al.* The creation and purpose of a classification system of children's abilities: Gross Motor Function Classification System (GMFCS). *J Intellect Disabil Res* 2004;48:346.
- 25 Wood EP, Rosenbaum P. The gross motor function classification system for cerebral palsy: a reliability. *Ann Neurol* 1998;44:560.
- 26 Jarvis S, Glinianaia SV, Arnaud C, *et al.* Case gender and severity in cerebral palsy varies with intrauterine growth. *Arch Dis Child* 2005;90:474–9.
- 27 Glinianaia SV, Jarvis S, Topp M, *et al.* Intrauterine growth and cerebral palsy in twins: a European multicenter study. *Twin Res Hum Genet* 2006;9:460–6.
- 28 Topp M, Langhoff-Roos J, Uldall P. Preterm birth and cerebral palsy. predictive value of pregnancy complications, mode of delivery, and Apgar scores. *Acta Obstet Gynecol Scand* 1997;76:843–8.
- 29 Lindqvist PG, Hapbach C. Risk and risk estimation of placental abruption. *Eur J Obstet Gynecol Reprod Biol* 2006;126:160–4.
- 30 Ananth CV, Berkowitz GS, Savitz DA, *et al.* Placental abruption and adverse perinatal outcomes. *JAMA* 1999;282:1646.
- 31 Pariente G, Wiznitzer A, Sergienko R, *et al.* Placental abruption: critical analysis of risk factors and perinatal outcomes. *J Matern Fetal Neonatal Med* 2011;24:698–702.
- 32 Schneider RE, Ng P, Zhang X, *et al.* The association between maternal age and cerebral palsy risk factors. *Pediatr Neurol* 2018;82:25–8.
- 33 Hvidtjørn D, Grove J, Schendel D, *et al.* Multiplicity and early gestational age contribute to an increased risk of cerebral palsy from assisted conception: a population-based cohort study. *Hum Reprod* 2010;25:2115–23.
- 34 Ananth CV, Schisterman EF. Confounding, causality, and confusion: the role of intermediate variables in interpreting observational studies in obstetrics. *Am J Obstet Gynecol* 2017;217:167–75.
- 35 Fauser BCJM, Devroey P, Macklon NS. Multiple birth resulting from ovarian stimulation for subfertility treatment. *Lancet* 2005;365:1807–16.
- 36 Blondel B, Kaminski M. Trends in the occurrence, determinants, and consequences of multiple births. *Semin Perinatol* 2002;26:239–49.
- 37 Reynolds MA, Schieve LA, Martin JA, *et al.* Trends in multiple births conceived using assisted reproductive technology, United States, 1997–2000. *Pediatrics* 2003;111:1159–62.
- 38 Strömberg B, Dahlquist G, Ericson A, *et al.* Neurological sequelae in children born after in-vitro fertilisation: a population-based study. *Lancet* 2002;359:461–5.
- 39 NICE. Overview | Twin and triplet pregnancy | Guidance | NICE, 2019. Available: <https://www.nice.org.uk/guidance/ng137/chapter/Recommendations#timing-of-birth> [Accessed 3 Dec 2019].