**QUEEN'S
UNIVERSITY
BELFAST**

Multivariate Approach for the Authentication of Vanilla Using Infrared and Raman Spectroscopy

Jiménez, A. M., Tonolini, M., McAleer, O., Granato, D., Cuadros-Rodríguez, L., & Koidis, T. (2021). Multivariate Approach for the Authentication of Vanilla Using Infrared and Raman Spectroscopy. *Food Research International*. <https://doi.org/10.1016/j.foodres.2021.110196>

Published in:
Food Research International

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2021 Elsevier.

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License

(<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

MULTIVARIATE APPROACH FOR THE AUTHENTICATION OF VANILLA USING INFRARED AND RAMAN SPECTROSCOPY

Ana M. Jiménez-Carvelo ¹, Margherita Tonolini ², Orla McAleer ², Luis Cuadros-Rodríguez ¹,
Daniel Granato ³, Anastasios Koidis ²

¹ Department of Analytical Chemistry, University of Granada, c/ Fuentenueva, s.n. E-18071 Granada, Spain.

² Institute for Global Food Security, Queen's University, 18-30 Malone Road, Belfast BT9 5BN, Northern Ireland, United Kingdom.

³ Food Processing and Quality, Natural Resources Institute Finland, Latokartanonkaari 9,
00790 Helsinki, Finland.

✉ Corresponding author (e-mail: t.koidis@gub.ac.uk , tel. +44 28 9097 5569)

ABSTRACT

Many different versions of vanilla extracts exist in the market in a variety of origins, purity levels and composition with little effective regulation. In this study, vanilla is authenticated both in terms of purity and geographical origin applying a multivariate approach using near infrared (NIR), mid infrared (MIR) and Raman spectroscopy following a complex experimental design. Partial least squares-discriminant analysis (PLS-DA) was applied to the spectral data to produce qualitative models. The prediction accuracy of the models was externally validated from the specific success/error contingencies. The results showed that MIR and Raman are reliable for authenticating vanilla in terms of purity, obtaining sensitivity, specificity, precision, and efficiency values equal to 1.00, and Raman is especially suitable for indicating the geographical origin of vanilla extracts, achieving performance metrics around 0.9.

Keywords:; authenticity, vanilla; fingerprinting, multivariate analysis; infrared spectroscopy; Raman spectroscopy; discriminant analysis.

1. INTRODUCTION

Vanilla is the second-most expensive spice worldwide, as well as the world's most popular flavouring material in the food industry. The spice derives from the orchid pods of the genus *Vanilla*, a native plant of Mexico. The majority of the world's vanilla (80% of global production) belongs to *Vanilla planifolia* species, commonly known as "Bourbon" vanilla, grown in Madagascar and neighbouring islands with 1000-1200 tons of cured vanilla beans exported every year (**Bomgardner, 2017**). The second type is Tahiti Vanilla (*V. tahitensis*) cultivated in French Polynesia and New Papua Guinea. Mexico, India, and Indonesia produce *V. planifolia*. Vanilla can be marketed in different forms, including extract, beans, and powder. The vanilla beans, which can be found inside the pods, can greatly vary in price according to their particular type/species, geographical origin and overall quality (**Ranadive, 1992**).

Real vanilla pods have a pure delicate and complex spicy flavour, which can be mainly attributed to the phenolic aldehyde vanillin ($C_8H_8O_3$), the latter accounts for up to 2% of the actual vanilla bean. Apart from vanillin, *p*-hydroxybenzaldehyde, guaiacol (and in the case of Tahiti vanilla, also anise alcohol) complement the main aroma profile of real vanilla (**Westcott et al., 1994; Bettazzi et al., 2006; Sharma et al., 2006**).

Consequently, vanilla beans are in limited global supply, and command a premium price (\$4,000/kg; **Rao & Ravishankar, 2000**). This has led to development of numerous synthetic or semi synthetic products in order to minimise the cost and satisfy the demand. These products contain synthetically produced vanillin as the base ingredient. Synthetic vanillin is synthesised from ferulic acid or guaiacol (synthetic pathway) and to a lesser extend from eugenol or lignin (semi-synthetic). Most of these products also contain ethyl vanillin, coumarin or piperonal used to mimic the flavour of natural vanilla but do not naturally occur in the authentic vanilla (**Sinha et al., 2008a; Huesgen, 2011a**).

Both EU and USA legislation state that if a product is labelled as 'vanilla extract', it has to derive from authentic vanilla beans and should not contain any synthetic vanillin. Vanillin synthesis and the production of synthetic vanilla extracts, however, might involve the use of

hazardous ingredients, or processes. Biotechnologically produced vanillin from the microbial enzymatic oxidation of naturally occurring 4-hydroxy-3-methoxycinnamic acid (found in rice bran), will have the same biogenic signature as the one deriving from the natural vanilla beans, so even if it is illegal to substitute with it, it is very hard to discriminate using analytical methods (Greule, Mosandl, Hamilton & Keppler, 2015). One can understand how the complexity of the legislation around 'vanilla extracts' or 'vanilla flavouring' and the fine line between a legit and a non-legit product fuels the motivation for fraud either in the form of blending or substitution adulteration (Poole *et al.*, 1995). The most common type of vanilla adulteration involves partly or wholly substituting vanilla flavour extracted from vanilla beans with cheaper synthetic vanillin, while labelling the result flavouring as "pure" and coming only from beans. Other types of falsification include the substitution of tonka bean extract (lower cost plant extract with a strong flavour) for some of the vanilla extract or labelling with the incorrect geographical origin of growth.

The cured vanilla bean has a complex composition, and its composition is highly affected by different climatic conditions, processing, and origins. Thus, detecting any adulteration is not an easy task. Many analytical methods have been developed for the determination of the authenticity and traceability of vanilla. Sophisticated targeted methods for traceability mainly focused on stable isotopes analysis and on elemental analysis. Gas chromatography-isotope ratio mass spectrometry (GC-IR/MS) was employed to determine $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values of the vanillin from the vanilla extracts for determination the geographical origin of vanilla pods (Hansen *et al.*, 2014). On the other hand, X-ray fluorescence spectroscopy and separately, inductive coupled plasma-mass spectroscopy (ICP-MS) were used to study the elemental composition and classify vanilla from different origins (Hondrogiannis, Rotta & Zapf, 2013a; Hondrogiannis, Ehrlinger, Poplaski & Lisle, 2013b). The aforementioned studies have demonstrated some success especially in species differentiation (*V. planifolia* vs *V. tahitensis*) and in the broad geographic discriminations which is intrinsically linked to the cultivated species. More specific studies also focused on the determining the source of vanillin (natural

vs biotechnologically produced) using site-specific natural isotope fractionation–nuclear magnetic resonance (SNIF-NMR) (Remaud, Martin, Martin, & Martin, 1997) and specific methoxy group targeted $\delta^{13}\text{C}$ and $\delta^2\text{H}$ stable isotope analysis (Greule, Mosandl, Hamilto, & Keppler, 2009; Greule *et al.*, 2010).

Although it is true that authenticity is better correlated with stable isotopes and minerals/trace elements, rapid analysis is needed in the field and along the supply chain (Geibler *et al.*, 2017; Galvin-King, Haughey, & Elliot, 2018). Recently, the use of molecular fingerprinting techniques such as vibrational spectroscopy (infrared or Raman) is rising due to the increased sensitivity reduced cost and ease of use. In the case of herbs and spices these techniques demonstrated their potential (Galvin-King, Haughey, & Elliot, 2018). Sharp *et al.* (2012) attempted to use mid infrared-attenuated total reflectance (MIR-ATR) coupled with soft independent modelling of class analogy (SIMCA) to cluster the vanilla extracts from the same origins and species. This study showed some separation between clusters of molecular signatures (representing different vanilla origins) and although concluded that MIR less powerful for discrimination due to its non-specificity, the potential of this rapid method was demonstrated. There are no studies concerning the use of other types of vibrational spectroscopy such as Near Infrared (NIR) and Raman. Based on the ease of use, speed of analysis, portability of the instrument, instrument prevalence in many industries including the food industry, as well as current demonstrated applications of all types of vibrational spectroscopy in the area of food authentication (Danezis *et al.*, 2016), a comprehensive study evaluating their potential in vanilla authentication is missing. The aim of this study, therefore, was to develop a screening protocol for rapid evaluation of authenticity and geographical origin detection of commercial vanilla samples using IR and Raman spectroscopy. According to the design, the authenticity is evaluated based on the detection of synthetic vanillin or tonka bean extract mixed in different ratios in the vanilla sample and the origin is evaluated based on whether a vanilla sample is coming from Madagascar or not.

2. MATERIALS AND METHODS

2.1 Vanilla samples

A total of 125 vanilla samples were used in this study. From them, 65 were pure vanilla from different origin geographic and 60 were mixtures of pure extract with synthetic extract vanilla prepared in the laboratory (the design and preparation of these mixtures is described in the 2.3 section). Table 1 includes details of the samples.

Table 1

The majority of the pure vanilla was acquired from trusted sourced including local authorities using overseas contacts and others were purchased either from online retailers or standard retail both in France and in the UK during the period 2018-19. The retail price range was 160-882 £/kg for the samples claimed to originate from Madagascar and 0.8-924 £/kg for the rest of the samples. Lastly, two types of vanillin were purchased, a synthetically produced (99.9% pure, Sigma-Aldrich, Dorset, UK) and a biotechnologically produced (fermentation of natural eugenol sold by Solvay, Zurich, Switzerland) respectively. Pure tonka beans were also purchased from Sigma-Aldrich.

2.2 Extract preparation of vanilla pods

The vanilla pod samples were opened carefully to collect the beans, labelled and frozen (–25°C) upon arrival in individual bags. After at least 24 h, frozen beans were transferred to a freeze dryer and dried over 48 h. The beans were removed from the freeze dryer and immediately air sealed. The dry bean samples were then grounded individually with a small laboratory grinder to a fine powder (5 mm pore sleeve was used) and transferred into airtight plastic containers. For the solid-liquid solvent extraction, 4.000 g of each powdered sample

were placed in a 50 ml tube and 25 ml of 50:50 ethanol:water (v/v) were added. After 1 min of vigorous agitation, the tubes were left to macerate for 72 h in slow speed. The solids were separated from the extracted liquid using a centrifuge (20 min at room temperature, 2000 min⁻¹ × g). The extracts were then transferred to a pre-weighted glass tubes and dried in batches using a Turbovap at 50°C under a nitrogen flow. The solids were reconstituted at 3% w/v using the appropriate amount of 35% ethanol: water. The pure vanilla extracts were transferred to glass vials and frozen (–25 °C) until further analysis.

The exact same extraction procedure was used to prepare the 3% Tonka bean extracts starting from raw Tonka beans. This extract was also infused with caramel colouring (3% w/v final concentration) at a later stage to make "spiked vanilla extract". Both these ingredients were added to simulate the composition and colour of real vanilla extracts.

2.3 Experimental design and admixtures preparation

The purity and the geographic origin of the vanilla samples were tested using two scenarios. In the first scenario (focusing on authenticity of the vanilla) the main research question was whether the sample tested was 100% pure vanilla or it has some adulterants, namely synthetic vanillin or Tonka bean extract, often added to lower the cost and mimic the flavour of pure vanilla. For this, 107 vanilla extracts were considered, of those: a) 59 were the pure extracts (PE group, PE = M + NM + MX, where M is the Madagascar vanilla samples, NM is the non-Madagascar samples and MX are the admixtures of these two sample categories) originating from the extraction of the vanilla beans (as per the procedure described in 2.2) - regardless of their exact geographical origin, b) 45 were vanillin-spiked vanilla extracts and c) 15 were Tonka-spiked vanilla extracts. The latter two (spiked) groups were admixtures prepared in house. More specifically, the 45 vanillin-spiked vanilla samples contained pure extracts (35-85%) mixed with synthetic vanillin (0.03-0.13%) and caramel flavour (0-3.5%) using 3 different vanillin final concentration (0.2%, 0.3%, 0.35%). On the other hand, the 15 Tonka-spiked

vanilla samples (CODE) contained pure vanilla extracts (50-90%) mixed with various amounts of caramel-infused 3% Tonka bean extract. To prepare this set, the 3% w/v pure Tonka bean extract (prepared as in 2.2) was infused with caramel flavour at a 95:5 w/w ratio (see Supplementary Information for more details). In total the adulterated vanilla samples were 55 (AD group).

In the second scenario (focusing on the origin), the main research question was whether the vanilla sample under investigation is coming from Madagascar or not. For this, the obtained pure vanilla extracts (see 2.2) were used to form two district groups: the M group (n=21 samples) strictly deriving from Madagascar, the NM group (Non-Madagascar, n=26 samples from elsewhere) as well as certain in-house admixtures of both groups, to simulate potential adulteration of Madagascar with vanilla from other locations. More specifically, to create this admixture group (MX), 6 Madagascar and 6 Non-Madagascar extracts were randomly selected to prepare 12 simple admixtures in ratios ranging from 5-95%, using each N and each NM vanilla extract only once. This occurred to create some variability in the dataset. More specifically, the admixtures were: 5% 17M+ 95% 35NM; 5% 11M+ 95% 36NM; 5% M12 + 95% 38NM; 10% M9 + 90% 23NM; 10% M12 + 90% 21NM; 10% M14 + 90% 36NM 15% M22 + 85% 17NM; 15% M14 + 85% 38NM; 20% M17 + 80% 20% M11 + 80% 21NM; 23NM; 5% M9 + 95% 17NM; 10% M22 + 90% 35NM (see Supplementary Information for more details).

2.4 MIR, NIR and Raman spectroscopy acquisition

All samples were analysed directly as extracts in their liquid form. Measurements were performed in triplicate using all 3 types of benchtop vibrational spectroscopy in benchtop instruments using typical acquisition parameters. The mid infrared spectra were acquired using the Thermo Nicolet iS5 spectrometer (Thermo Scientific, Dublin, Ireland) coupled with ATR. The resolution was set at 4 cm⁻¹, the frequency range at 4000 to 700 cm⁻¹ and number of scans to 64 with 0.482 cm⁻¹ intervals resulting in 6846 wavenumbers collected. The

background noise was acquired before each measurement and was appropriately subtracted from the signal. The Near Infrared spectrum was acquired using the Antaris II FT-NIR (Thermo Fisher Scientific, Dublin, Ireland) in reflectance mode with a spinning module. All spectra were computed at 8 cm^{-1} resolution in 4 mm pathlength across the spectral range $3999\text{--}11998\text{ cm}^{-1}$ with 1.5 cm^{-1} intervals resulting in 3022 wavenumbers collected. The spectra were recorded at ambient temperature and a total of 64 scans were acquired for each spectrum after appropriate background measurement. A DeltaNu Advantage Raman Spectrometer equipped with a 1064 nm laser was used to acquire the Raman spectra. The instrument was calibrated before every use with a blank vial followed by a random vial from the sample set to be tested. The laser power was set to 1000 mW, integration time was 10 sec and the number of spectra for each acquisition was set to 32 collecting signals from 1867 wavenumbers. In all cases the three NIR or MIR or Raman spectra were averaged in one spectrum using the respective acquisition software.

2.5 Chemometric analysis of spectral data

Spectra were obtained in the instrument-specific format, averaged and exported as CSV, then converted to MATLAB format (version R2013b). Before performing the multivariate analysis to any chemometric analysis, the spectra were submitted to a data processing stage of autoscaling and a variable selection was performed.

Principal component analysis (PCA), soft independent modelling by class analogy (SIMCA), partial least squares-discriminant analysis (PLS-DA) and support vector machine-classification mode (SVM-C) were built using PLS Toolbox (version 8.6.1, Eigenvector Research, Wenatchee, WA). For validation purpose of classification methods, the original data set was split into training and external validation set using the CADEX algorithm, developed by Kennard and Stone (**Kennard & Stone, 1969**), and at least 30% of the samples from each class were set to define the external validation set while the remaining samples constitutes the training set. Two authentication scenarios have been achieved attending to: (i)

discrimination of pure vanilla/non-pure vanilla, and (ii) classification according to their geographical origin (Madagascar / Non-Madagascar). Table 2 details the samples used for each model according to the two scenarios.

Table 2

3. RESULTS AND DISCUSSION

3.1 Interpretation of Spectra

The MIR spectral features corroborate with similar studies. The broad peak at 3200-3550 cm^{-1} is attributed to O-H stretching vibrations mainly due to ethanol present in the extract; The sharp peak at 2980 cm^{-1} is due to C-H, (such as those in phenols and fatty acids) whereas the shoulder peak at 2905 cm^{-1} is aliphatic C-H stretching (methyl, $-\text{CH}_3$, and methylene, $-\text{CH}_2-$) of organic compounds abundant in the extracts (**Moreno-Ley et al. 2019**). The sharp peak at 1645 cm^{-1} is due to C=O and C=C aromatic stretching vibrations (aldehydes, ketones, esters of vanilla as well specific pyrones (**Brunschwig et al., 2009**). Peaks in the fingerprint region such as the 1088 and 1044 cm^{-1} are less specific and are attributed to stretching vibrations of the C-H, C-O bonds also abundant in all plant extracts (**Aljaff et al., 2013**).

In general, the harmonic vibrations and overtone absorption spectra of NIR spectroscopy provide less specific information on fundamental molecular vibrations (raw data shown in Fig 3b). Absorption peaks were observed in the raw spectra at approximately 8400-8200 cm^{-1} , (second and third C-H overtones) and the regions 6800-7250, 6200-6500 and 5700-6000 cm^{-1} overlap with the first C-H overtone. The two peaks at 5200-4500 cm^{-1} fall within the region associated with C-H and O-H combination bands coarsely attributed to the solvent and the other organic constituents of the extracts. The spectral features broadly agree with those reported in the literature (**Wongsheree et al., 2014**).

On the other hand, the most intense Raman bands appear in four wavenumbers or Raman shifts: 877 cm^{-1} (low frequency carbon-carbon vibrations), 1040-1185 cm^{-1} double peak (linked to aromatic rings), 1275 cm^{-1} , 1456 cm^{-1} ($-\text{CH}_3$ and $-\text{CH}_2-$ deformations). There are no other studies that reported Raman spectral features of vanilla extracts or vanilla powder although the observed vibration patterns of the spectra are in line with Raman spectra of similar herbs and spices (Galvin-King et al., 2018). The MIR and Raman spectral features of the vanilla extracts are shown in Figure 1.

Figure 1

3.2 Selection of variables by means on principal component analysis

When working with instrumental fingerprints it is common to have a data set, characterised by thousands of variables, that is too large so long computational times are required to build the classification model. For this reason, a selection of variables to reduce the dimensionality of the data is commonly applied. Usually two ways of selecting the region of interest are based on: a) selecting a specific number of raw variables by implementing variable selection methods, such as interval partial least-squares regression (iPLS), genetic algorithms (GA) or variable importance in projection (VIP), or b) generating new variables, which are usually a combination of the raw variables, for example when hierarchical cluster analysis (HCA) or PCA are employed as tool to reduce the dimensionality of data and not as techniques of exploratory analysis.

In this work PCA has been employed as technique of exploratory analysis and, in addition variable selection was performed examining the PCA loading plot for each data set acquired by NIR, MIR and Raman. The region of interest selected was a selection of the relevant raw

variables of the spectra. For that purpose, the regions of the spectra where the intensity of the loading was high were selected.

Initially, a PCA model from NIR data was developed with four principal components (PCs) which explain 97.51% of the variance (see Figure 2). In this case, PCA allowed to visualize the two groups of samples which are correlated with adulterated vanilla (positive scores in PC2) and pure extract of vanilla (negative scores in PC2). Then, the distribution of the loadings was examined to understand the groupings observed in the scores plot, to select the region of interest (ROI) of the NIR spectra. Figure 3 shows both the PC2 loading plot, aligned on the NIR spectra from two samples of pure and non-pure vanilla. The profile from the PC2 loading plot shows a sharp and intense boosting and busting which corresponds to the central peak of the spectrum ($5926\text{-}6890\text{ cm}^{-1}$) and is defining of the ROI from which the PC2 which sets the spacing between the two groups is principally composed.

Figure 2

Figure 3

For the PCA model from Raman dataset four principal components were selected which explain 99.92% of the variance (see figure 4). In this case no groupings were observed. The PCA loadings shown the same shape than the Raman spectra, thus no region of interest was selected (see figure 5). Although the initial region of the Raman spectrum ($200\text{-}777\text{ cm}^{-1}$) shows a high loading value, it was not finally selected since this part corresponds with fluorescence interference and it was not considered. Therefore, only the $778\text{-}1549\text{ cm}^{-1}$ range was employed for the development of the classification methods.

Figure 4

293

Figure 5

294

295 In the case of the data from MIR, PCA model was built with 7 PCs that explained a 95.54% of
296 the variance. As can be observed in Figure 6 there is a grouping pattern of the pure vanilla
297 samples with negative scores for PC1 and positive scores for PC2 whereas that the non-pure
298 samples are distributed with scores positives for PC1 and positive and negative scores for
299 PC2. Then, when PCA loading plots were examined was not possible to select a specific
300 region of interest since the entire loading signal shown similar importance for the grouping
301 pattern of both sample dataset. Therefore, for the development of the multivariate
302 classification models the entire MIR signal was used.

303

Figure 6

304

305 **3.3 Scenario 1: Purity (discrimination between pure and non-pure vanilla)**

306 The classification rule was based on setting the boundaries for the different classes defined
307 ('Pure' and 'Non-pure') by the corresponding training sample dataset. Three classification
308 methods, SIMCA, PLS-DA and SVM-C, were compared. The classification threshold for each
309 class was established by the software.

310 For each classification method, three models were built from NIR, MIR and Raman dataset.
311 The 'Pure' class was defined by value equal to 0, while the 'Non-pure' class was defined by a
312 value of 1. In all cases, the PLS-DA method provided the best results. Table 3 shows the
313 numbers of LVs chosen for each model, and Table 4 shows the quality performance metrics
314 for the three binary classification methods, calculated according to the external validation set
315 using the success/errors contingency for each class and for the three techniques (see
316 supplementary Information for the classification and loading plots).

317

Table 3

318

Table 4

319

320 MIR and Raman models correctly classified all samples. All the vanilla samples from 'Pure'
321 class were well classified (probability = 0) and the samples from the 'Non-pure' class were
322 also classified correctly (probability = 1). Thus, all the quality metrics of the models were equal
323 to 1.00 what involves this method could be considered reliable for authenticating the purity of
324 vanilla samples.

325 Conversely, the results of the NIR model were remarkably worse than the last ones, obtaining
326 values of sensitivity, specificity, and precision of 0.82, 0.72 and 0.76 for the 'Pure' class and
327 0.72, 0.89 and 0.87 for the 'Non-pure' class, respectively. Seven vanilla samples were
328 misclassified. Two samples of the class 'Pure' were classified as 'Non-pure', and the rest of
329 samples from 'Non-pure' class were incorrectly classified as belonging to 'Pure' class.

330

331 **3.4 Scenario 2: Geographical origin (classification according to Madagascar/Non-**
332 **Madagascar origin)**

333 The same classification methodology was used to build the different PLS-DA multivariate
334 classification models. The 'Madagascar' class was defined by value equal to 0, while the 'Non-
335 Madagascar' class was defined by a value of 1. Table 5 shows the numbers of LVs chosen
336 for each model, and Table 6 shows the quality performance metrics for the three binary
337 classification methods (see supplementary Information the loading plots).

338

Table 5

339

Table 6

340

341 In this scenario NIR and MIR were not useful to discriminate vanilla samples from Madagascar
342 origin. NIR and MIR achieved high classification errors of the 56% and 38% respectively,
343 which is reflected in the performance classification metrics values shown in the Table 6 for
344 both techniques. By contrast, Raman classified correctly almost all samples except one
345 sample from 'Madagascar' considered as 'Non-Madagascar'. The sensitivity, specificity and
346 precision of the model were equal to 0.86, 1.00 and 1.00 for the 'Madagascar' class and 1.00,
347 0.86 and 0.89 for 'Non-Madagascar' class, respectively.

348 The results indicate that the main chemical composition of vanilla is not strongly dependent
349 on the geographical origin and therefore the vibrational analytical signals are not enough
350 different to be able to make a right effective discrimination. Possibly, the differences could be
351 focused on some minority compounds whose presence or concentration does not greatly
352 influences the spectra.

353

354 4. CONCLUSIONS

355 This work proposes a useful methodology to rapidly authenticate the purity of vanilla extracts
356 applying a comprehensive multivariate analytical approach considering the different
357 compositions of fraudulent vanilla extracts. To this end, different modes of vibrational
358 spectroscopy (NIR, MIR and Raman) have been explored. The PLS-DA classification method
359 is the one which produces the best results. It performs well when the composition of the
360 fraudulent samples is different from the original extract and is not sensitive enough to
361 determine, for example, differences between biotechnologically produced vanillin and the
362 native variant present in the vanilla pod. It can however distinguish the "gross offenders" such
363 as spiking with tonka beans, caramel, and other additions because the difference on spectral

signal can be discriminative between different samples. It is therefore useful as a screening method to be used by the spice industry and retailers because access to equipment and the measurement is low cost and generally accessible.

In addition, the same multivariate analytical approach maybe be also able to identify the geographical origin of a vanilla extract, namely if it is from Madagascar or not. Geographical origin is more complex analytical problem with many factors including the vanilla species also playing a major role, which was not explored in this study. There is room for other more sophisticated methods such as DNA fingerprinting. Further chemometric classification model based on larger datasets and diversity in the samples used, representing global vanilla supply and multi-year sampling, would improve model robustness and general accuracy of the analytical method.

- Bettazzi, F., Palchetti, I., Sisalli S., Mascini M. (2006). A disposable electrochemical sensor for vanillin detection. *Analytica Chimica Acta*, 55: 134–138. <https://doi.org/10.1016/j.aca.2005.08.069>
- Bomgardner, M. (2017). The problem with vanilla | September 12, 2016 Issue - Vol. 94 Issue 36 | Chemical & Engineering News. [online] Cen.acs.org. Available at: <http://cen.acs.org/articles/94/i36/problem-vanilla.html> [Accessed 6/9/2020].
- Danezis, G.P., Tsagkaris, A.S., Camin, F., Brusica, V., & Georgiou, C.A. (2016). Food authentication: Techniques, trends & emerging approaches. *Trends in Analytical Chemistry*, 85, 123-132. <https://doi.org/10.1016/j.trac.2016.02.026>
- Food and Drug Administration - FDA (2019). Code of Federal Regulations Title 21, Vol. 3. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=189&showFR=1> [Accessed 6/9/2020]
- Galvin-King, P., Haughey, S.A., & Elliott, C.T. (2018). Herb and spice fraud; the drivers, challenges and detection. *Food Control*, 88, 85-97. <https://doi.org/10.1016/j.foodcont.2017.12.031>
- Geibler, K., Greule, M., Schafer, U., Hans, J., Geibler, T., Meier, L., Keppler, F., & Krammer, G. (2017). Vanilla authenticity control by DNA barcoding and isotope data aggregation. *Flavour and Fragrance Journal*, 32 (4), 228-237. <https://doi.org/10.1002/ffj.3379>
- Greule, M., Mosandl, A., Hamilton, J.T.G., & Keppler, F. (2009). A simple rapid method to precisely determine $^{13}\text{C}/^{12}\text{C}$ ratios of plant methoxyl groups. *Rapid Communications in Mass Spectrometry*, 23, 1710-1714. <https://doi.org/10.1002/rcm.4057>
- Greule, M., Mosandl, A., Hamilton, J.T.G., & Keppler, F. (2015). Comment on authenticity and traceability of vanilla flavors by analysis of stable isotopes of carbon and hydrogen. *Journal of Agricultural and Food Chemistry*, 63, 5305-5306. <https://doi.org/10.1021/jf506172q>
- Greule, M., Tumino, L.D., Kronewald, T., Hener, U., Schleucher, J., Mosandl, A., & Keppler, F. (2010). Improved rapid authentication of vanillin using $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values. *European Food Research and Technology*, 231, 933-941. <https://doi.org/10.1007/s00217-010-1346-z>

- Hansen, A.M.S., Fromberg, A., & Frandsen, H.L. (2014). Authenticity and traceability of vanilla flavors by analysis of stable isotopes of carbon and hydrogen. *Journal of Agricultural and Food Chemistry*, 62(42), 10326-10331.
<https://doi.org/10.1021/jf503055k>
- Hondrogiannis, E., Rotta, K., & Zapf, C.M. (2013). The use of wavelength dispersive X-ray fluorescence in the identification of the elemental composition of vanilla samples and the determination of the geographic origin by discriminant function analysis. *Journal of Food Science*, 78(3), C395-C401. <https://doi.org/10.1111/1750-3841.12050>
- Hondrogiannis, E.M., Ehrlinger, E., Poplaski, A., & Lisle, M. (2013). Use of laser ablation–inductively coupled plasma–time of flight–mass spectrometry to identify the elemental composition of vanilla and determine the geographic origin by discriminant function analysis. *Journal of Agricultural and Food Chemistry*, 61(47), 11332-11337.
<https://doi.org/10.1021/jf4025536>
- Huesgen, A.G. (2011). Analysis of Natural And Artificial Vanilla Preparations. Application Notes. 1st ed. Germany: Agilent Technologies, Inc. Web. 17 Feb. 2017. [Accessed 6/9/2020]
- Kennard, R.W., & Stone, L.A. (1969). Computer aided design of experiments. *Technometrics*, 11, 137–148. <http://doi.org/10.1080/00401706.1969.10490666>
- Poole, S.K., Kiridena, W., Miller, K.G., & Poole, C.F. (1995). Planar chromatographic methods for determination of the quality of spices and flavours as exemplified by cinnamon and vanilla. *Journal of Planar Chromatography*, 8, 257-268.
- Ranadive, A.S. (2005). Vanilla cultivation. *In: Vanilla: The first international congress*. Princeton: NJ: Allured, Carol Stream. p 25–32.
- Rao, S.R., & Ravishankar, G. A. (2000). Vanilla flavour: Production by conventional and biotechnological routes. *Journal of the Science of Food and Agriculture*, 80: 289–304.
[https://doi.org/10.1002/1097-0010\(200002\)80:3<289::AID-JSFA543>3.0.CO;2-2](https://doi.org/10.1002/1097-0010(200002)80:3<289::AID-JSFA543>3.0.CO;2-2)
- Remaud, G.S., Martin, Y.L., Martin, G.G. & Martin, G.J. (1997). Detection of sophisticated adulterations of natural vanilla flavors and extracts: application of the SNIF-NMR method to vanillin and p-hydroxybenzaldehyde. *Journal of Agricultural and Food Chemistry*, 45, 859-866. <https://doi.org/10.1021/jf960518f>
- Sharma, A., Verma, S.C., Saxena N., Chadda, N., Singh, N.P., & Sinha A.K. (2006). Microwave and ultrasound-assisted extraction of vanillin and its quantification by high-

- performance liquid chromatography in *Vanilla planifolia*. *Journal of Separation Science*, 29: 613-619. <https://doi.org/10.1002/jssc.200500339>
- Sharp, M.D., Kocaoglu-Vurma, N.A., Langford, V., Rodriguez-Saona, L.E., & Harper, W.J. (2012). Rapid discrimination and characterization of vanilla bean extracts by attenuated total reflection infrared spectroscopy and selected ion flow tube mass spectrometry. *Journal of Food Science*, 77(3), C284-C292. <https://doi.org/10.1111/j.1750-3841.2011.02544.x>
- Sinha, A.K., Sharma, U.K., & Sharma, N. (2008). A comprehensive review on vanilla flavor: Extraction, isolation and quantification of vanillin and other constituents. *International Journal of Food Sciences and Nutrition* 59(4): 299-326. <https://doi.org/10.1080/09687630701539350>
- Westcott, R.J., Cheetham, P.S.J., & Arraclough, A.J.B. (1994). Use of organized viable vanilla plant aerial roots for the production of natural vanillin. *Phytochemistry*, 35: 135-138. [https://doi.org/10.1016/S0031-9422\(00\)90521-1](https://doi.org/10.1016/S0031-9422(00)90521-1).
- Aljaff, P., Rasheed, B. O., & Omer, T. A. (2013). A comparison between natural and synthetic food flavoring extracts using infrared spectra and optical activity. *Journal of Applied Physics*, 5(3), 1-6.
- Brunschwig, C., Collard, F. X., Bianchini, J. P., & Raharivelomanana, P. (2009). Evaluation of chemical variability of cured vanilla beans (*Vanilla tahitensis* and *Vanilla planifolia*). *Natural product communications*, 4(10), 1934578X0900401016.
- Moreno-Ley, C. M., Hernández-Martínez, D. M., Osorio-Revilla, G., Tapia-Ochoategui, A. P., Dávila-Ortiz, G., & Gallardo-Velázquez, T. (2019). Prediction of coumarin and ethyl vanillin in pure vanilla extracts using MID-FTIR spectroscopy and chemometrics. *Talanta*, 197, 264-269.
- Wongsheree, T., Rittiron, R., Wongs-Aree, C. and Thongtheing, T. (2016). Vanilla content evaluation in cured bean by near-infrared technique. *Acta Horticulture* 1119, 297-304 DOI: 10.17660/ActaHortic.2016.1119.41

TABLES

Table 1. Types of vanilla samples analysed

Type of vanilla	Nº	Origin
Pure Extract (<i>V. planifolia</i> and <i>V. tahitensis</i>)	53	Madagascar (19) Non-Madagascar (34)
Blends of pure extract ¹	12	<i>Not considered</i>
Mixture of pure extract with synthetic extract	60	<i>Not considered</i>

¹ *Blends of pure extract from Madagascar and non-Madagascar*

Table 2. Distribution of the samples used in the different classification models.

Dataset	Scenario 1: Purity		Scenario 2: Origin	
	Pure	Non-Pure	Madagascar	Non-Madagascar
Training set	44	44	12	27
Validation set	19	18	7	7

Table 3. Characteristics of the PLS-DA models for the Scenario 1: Purity

NIR		MIR		Raman	
LVs	% var	LVs	% var	LVs	% var
4	98.23	7	92.95	3	97.75

Table 4. Classification performance metrics of the PLS-DA model for the Scenario 1: Purity

Performance metrics	Pure	Non-Pure
<i>NIR</i> (5926 – 6890 cm⁻¹)		
Sensitivity (or Recall)	0.89	0.72
Specificity	0.72	0.89
Positive predictive value (Precision)	0.76	0.87
Negative predictive value	0.87	1.00
Youden index	0.61	0.61
Positive likelihood rate	3.20	6.50
Negative likelihood rate	0.15	0.31
F-measure	0.82	0.79
Discriminant power	0.73	0.73
Efficiency (or Accuracy)	0.81	0.81
AUC (Correctly classified rate)	0.81	0.81
Matthews correlation coefficient	0.62	0.62
Kappa coefficient	0.61	0.61
<i>MIR</i> (700 – 4000 cm⁻¹)		
Sensitivity (or Recall)	1.00	1.00
Specificity	1.00	1.00
Positive predictive value (or Precision)	1.00	1.00
Negative predictive value	1.00	1.00
Youden index	1.00	1.00
Positive likelihood rate	-	-
Negative likelihood rate	0.00	0.00
F-measure	1.00	1.00
Discriminant power	-	-
Efficiency (or Accuracy)	1.00	1.00
AUC (Correctly classified rate)	1.00	1.00
Matthews correlation coefficient	1.00	1.00
Kappa coefficient	1.00	1.00

394 *Continue Table 4*

395

Performance metrics	Pure	Non-Pure
<i>Raman</i> (778 – 1549 cm⁻¹)		
Sensitivity (or Recall)	1.00	1.00
Specificity	1.00	1.00
Positive predictive value (or Precision)	1.00	1.00
Negative predictive value	1.00	1.00
Youden index	1.00	1.00
Positive likelihood rate	-	-
Negative likelihood rate	0.00	0.00
F-measure	1.00	1.00
Discriminant power	-	-
Efficiency (or Accuracy)	1.00	1.00
AUC (Correctly classified rate)	1.00	1.00
Matthews correlation coefficient	1.00	1.00
Kappa coefficient	1.00	1.00

The hyphen "-" is signifying that the performance feature cannot be determined.

396

Table 5. Characteristics of the PLS-DA models for the Scenario 2: Origin

NIR		MIR		Raman	
LVs	% var	LVs	% var	LVs	% var
3	97.07	9	96.73	2	99.72

Table 6. Classification performance metrics of the PLS-DA model for the Scenario 2: Origin

Performance metrics	Madagascar	Non-Madagascar
<i>NIR</i> (3999 – 11998 cm⁻¹)		
Sensitivity (or Recall)	0.29	0.50
Specificity	0.50	0.29
Positive predictive value (or Precision)	0.33	0.44
Negative predictive value	0.44	1.00
Youden index	-0.21	-0.21
Positive likelihood rate	0.57	0.70
Negative likelihood rate	1.43	1.75
F-measure	0.31	0.47
Discriminant power	-0.22	-0.22
Efficiency (or Accuracy)	0.40	0.40
AUC (Correctly classified rate)	0.39	0.39
Matthews correlation coefficient	-0.22	-0.22
Kappa coefficient	-0.22	-0.22
<i>MIR</i> (700 – 4000 cm⁻¹)		
Sensitivity (or Recall)	0.57	0.63
Specificity	0.63	0.57
Positive predictive value (or Precision)	0.57	0.63
Negative predictive value	0.63	1.00
Youden index	0.20	0.20
Positive likelihood rate	1.52	1.46
Negative likelihood rate	0.69	0.66
F-measure	0.57	0.63
Discriminant power	0.19	0.19
Efficiency (or Accuracy)	0.60	0.60
AUC (Correctly classified rate)	0.60	0.60
Matthews correlation coefficient	0.20	0.20
Kappa coefficient	0.20	0.20

Continue table 6

Performance metrics	Madagascar	Non-Madagascar
<i>Raman</i> (778 – 1549 cm ⁻¹)		
Sensitivity (or Recall)	0.86	1.00
Specificity	1.00	0.86
Positive predictive value (Precision)	1.00	0.89
Negative predictive value	0.89	1.00
Youden index	0.86	0.86
Positive likelihood rate	–	7.00
Negative likelihood rate	0.14	0.00
F-measure	0.92	0.94
Discriminant power	–	–
Efficiency (or Accuracy)	0.93	0.93
AUC (Correctly classified rate)	0.93	0.93
Matthews correlation coefficient	0.87	0.87
Kappa coefficient	0.86	0.86

The hyphen "–" is signifying that the performance feature cannot be determined.

FIGURE CAPTIONS

Figure 1. Raw spectra of all vanilla extracts by (a) MIR and (b) Raman.

Figure 2. PCA scores plot in PC1-PC2 plane for pure and non-pure vanilla samples from NIR data.

Figure 3. (a) PC2 loading plot of all vanilla samples and (b) superposed NIR spectra from two vanilla samples (one pure and other non-pure) zooming the region of interest selected.

Figure 4. PCA scores plot in PC1-PC2 plane for pure and non-pure vanilla samples from Raman data.

Figure 5. (a) Loading plot of all vanilla samples and (b) Raman spectrum of a pure vanilla sample.

Figure 6. PCA scores plot in PC1-PC2 plane for pure and non-pure vanilla samples from MIR data.

<Figure 1>

(a)

(b)

SMCA 15 - 25/06/2020

429 <Figure 2>

430

431

432

433 <Figure 3>

434

435

436

<Figure 4>

441 <Figure 5>

442

443

444

445 <Figure 6>

446

447

448

SUPPLEMENTARY INFORMATION

Concentrations of in-house mixtures

The following tables show the varying ratios the in-house mixtures were made to.

The admixtures were prepared by selecting 6 Madagascar and 6 Non-Madagascar extracts;

Sample name	% Madagascar extract	% Non-Madagascar extract
V-MX1	5	95
V-MX2	10	90
V-MX3	15	85
V-MX4	20	80
V-MX5	5	95
V-MX6	10	90
V-MX7	5	95
V-MX8	10	90
V-MX9	15	85
V-MX10	20	80
V-MX11	5	95
V-MX12	10	90

Another 45 samples were also prepared enriched with synthetic vanillin using 3 different vanillin final concentrations:

➤ TOTAL VANILLIN CONCENTRATION 0.2%

Sample N	Pure extract Number	Pure extract	Solvent	Caramel	Vanillin
V-AD1-Val	1	85.69%	14.28%	0.00%	0.03%
V-AD2-Val	1	49.95%	49.95%	0.00%	0.10%
V-AD3-Val	1	35.67%	64.20%	0.00%	0.13%
V-AD4-Val	2	85.69%	14.28%	0.00%	0.03%
V-AD5-Val	2	49.95%	49.95%	0.00%	0.10%
V-AD6-Val	2	35.67%	64.20%	0.00%	0.13%
V-AD7-Val	3	89.53%	9.70%	0.75%	0.03%
V-AD8-Val	3	51.23%	46.11%	2.56%	0.10%
V-AD9-Val	3	36.32%	60.28%	3.27%	0.13%
V-AD10-Val	4	89.53%	9.70%	0.75%	0.03%
V-AD11-Val	4	51.23%	46.11%	2.56%	0.10%
V-AD12-Val	4	36.32%	60.28%	3.27%	0.13%
V-AD13-Val	5	89.53%	9.70%	0.75%	0.03%
V-AD14-Val	5	51.23%	46.11%	2.56%	0.10%
V-AD15-Val	5	36.32%	60.28%	3.27%	0.13%

458 ➤ TOTAL VANILLIN CONCENTRATION 0.3%

Sample N	Pure extract Number	Pure extract	Solvent	Caramel	Vanillin
V-AD16-Val	1	89.43%	9.69%	0.75%	0.13%
V-AD17-Val	1	51.18%	46.06%	2.56%	0.20%
V-AD18-Val	1	36.28%	60.22%	3.27%	0.23%
V-AD19-Val	2	89.43%	9.69%	0.75%	0.13%
V-AD20-Val	2	51.18%	46.06%	2.56%	0.20%
V-AD21-Val	2	36.28%	60.22%	3.27%	0.23%
V-AD22-Val	3	89.43%	9.69%	0.75%	0.13%
V-AD23-Val	3	51.18%	46.06%	2.56%	0.20%
V-AD24-Val	3	36.28%	60.22%	3.27%	0.23%
V-AD25-Val	4	85.60%	14.27%	0.00%	0.13%
V-AD26-Val	4	49.90%	49.90%	0.00%	0.20%
V-AD27-Val	4	35.63%	64.14%	0.00%	0.23%
V-AD28-Val	5	85.60%	14.27%	0.00%	0.13%
V-AD29-Val	5	49.90%	49.90%	0.00%	0.20%
V-AD30-Val	5	35.63%	64.14%	0.00%	0.23%

459

460

461 ➤ TOTAL VANILLIN CONCENTRATION 0.35%

Sample N	Pure extract Number	Pure extract	Solvent	Caramel	Vanillin
V-AD31-Val	1	85.56%	14.26%	0.00%	0.18%
V-AD32-Val	1	49.88%	49.88%	0.00%	0.25%
V-AD33-Val	1	35.62%	64.11%	0.00%	0.28%
V-AD34-Val	2	89.39%	9.68%	0.74%	0.19%
V-AD35-Val	2	51.15%	46.04%	2.56%	0.26%
V-AD36-Val	2	36.26%	60.19%	3.26%	0.28%
V-AD37-Val	3	85.56%	14.26%	0.00%	0.18%
V-AD38-Val	3	49.88%	49.88%	0.00%	0.25%
V-AD39-Val	3	35.62%	64.11%	0.00%	0.28%
V-AD40-Val	4	89.39%	9.68%	0.74%	0.19%
V-AD41-Val	4	51.15%	46.04%	2.56%	0.26%
V-AD42-Val	4	36.26%	60.19%	3.26%	0.28%
V-AD43-Val	5	85.56%	14.26%	0.00%	0.18%
V-AD44-Val	5	49.88%	49.88%	0.00%	0.25%
V-AD45-Val	5	35.62%	64.11%	0.00%	0.28%

462

463

464

465 Finally, the “fake” vanilla extract was used to make 15 adulterated samples as follows:

466

Sample N	Pure extract Number	PVE	FVE
V-AD46-TB	1	90%	10%
V-AD47-TB	2	90%	10%
V-AD48-TB	3	90%	10%
V-AD49-TB	4	90%	10%
V-AD50-TB	5	90%	10%
V-AD51-TB	1	50%	50%
V-AD52-TB	2	50%	50%
V-AD53-TB	3	50%	50%
V-AD54-TB	4	50%	50%
V-AD55-TB	5	50%	50%
V-AD56-TB	1	80%	20%
V-AD57-TB	2	80%	20%
V-AD58-TB	3	80%	20%
V-AD59-TB	4	80%	20%
V-AD60-TB	5	80%	20%

467 PVE: pure vanilla extract; FVE: "fake" vanilla extract

468 In total, 60 adulterated vanilla in house admixtures were prepared.

469

470

Loading and classification plots of the different classification models developed

Supplementary Figure 1. Loading plot from PLS-DA models for the scenario 1: purity: (a) NIR, (b) MIR and (c) Raman

Supplementary Figure 2. Classification plot from PLS-DA models for the scenario 1: purity: (a) NIR, (b) MIR and (c) Raman

Supplementary Figure 3. Loading plot from PLS-DA models for the scenario 2: origin: (a) NIR, (b) MIR and (c) Raman