



**QUEEN'S
UNIVERSITY
BELFAST**

Divergent responses of permafrost peatlands to recent climate change

Sim, T. G., Swindles, G. T., Morris, P. J., Baird, A. J., Cooper, C. L., Gallego-Sala, A. V., Charman, D. J., Roland, T. P., Borken, W., Mullan, D. J., Aquino-López, M. A., & Gaika, M. (2021). Divergent responses of permafrost peatlands to recent climate change. *Environmental Research Letters*, 16(3), Article 034001. <https://doi.org/10.1088/1748-9326/abe00b>

Published in:

Environmental Research Letters

Document Version:

Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2021 the authors.

This is an open access article published under a Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

LETTER • OPEN ACCESS

Divergent responses of permafrost peatlands to recent climate change

To cite this article: Thomas G Sim *et al* 2021 *Environ. Res. Lett.* **16** 034001

View the [article online](#) for updates and enhancements.

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

Divergent responses of permafrost peatlands to recent climate change

OPEN ACCESS

RECEIVED
22 July 2020REVISED
22 January 2021ACCEPTED FOR PUBLICATION
26 January 2021PUBLISHED
11 February 2021

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Thomas G Sim¹ , Graeme T Swindles^{2,3}, Paul J Morris¹, Andy J Baird¹, Claire L Cooper⁴, Angela V Gallego-Sala⁵ , Dan J Charman⁵ , Thomas P Roland⁵, Werner Borken⁶, Donal J Mullan², Marco A Aquino-López⁷ and Mariusz Gałka⁸

¹ School of Geography, University of Leeds, Leeds, United Kingdom

² Geography, School of Natural and Built Environment, Queen's University Belfast, Belfast, United Kingdom

³ Ottawa-Carleton Geoscience Centre and Department of Earth Sciences, Carleton University, Ottawa, Ontario, Canada

⁴ School of Geography, University of Lincoln, Lincoln, United Kingdom

⁵ Geography, College of Life and Environmental Sciences, University of Exeter, Exeter, United Kingdom

⁶ Soil Ecology, University of Bayreuth, Bayreuth Center of Ecology and Environmental Research (BayCEER), Dr.-Hans-Frisch-Str. 1-3, 95448, Bayreuth, Germany

⁷ Centro de Investigacion en Matemáticas, Guanajuato, Mexico

⁸ Faculty of Biology and Environmental Protection, Department of Biogeography, Paleoecology and Nature Protection, University of Lodz, 12/16 Banacha Str., Lodz, Poland

E-mail: gy12tgs@leeds.ac.uk

Keywords: permafrost, peatlands, climate change, hydrology, carbon

Supplementary material for this article is available [online](#)

Abstract

Permafrost peatlands are found in high-latitude regions and store globally-important amounts of soil organic carbon. These regions are warming at over twice the global average rate, causing permafrost thaw, and exposing previously inert carbon to decomposition and emission to the atmosphere as greenhouse gases. However, it is unclear how peatland hydrological behaviour, vegetation structure and carbon balance, and the linkages between them, will respond to permafrost thaw in a warming climate. Here we show that permafrost peatlands follow divergent ecohydrological trajectories in response to recent climate change within the same rapidly warming region (northern Sweden). Whether a site becomes wetter or drier depends on local factors and the autogenic response of individual peatlands. We find that bryophyte-dominated vegetation demonstrates resistance, and in some cases resilience, to climatic and hydrological shifts. Drying at four sites is clearly associated with reduced carbon sequestration, while no clear relationship at wetting sites is observed. We highlight the complex dynamics of permafrost peatlands and warn against an overly-simple approach when considering their ecohydrological trajectories and role as C sinks under a warming climate.

1. Introduction

Permafrost peatlands have developed in cold regions during the Holocene and store a disproportionate amount of organic carbon (C) for their extent, estimated to total ~277 Gt C (Tarnocai *et al* 2009)—making up around a fifth of all permafrost soil C (Hugelius *et al* 2014). These ecosystems experience a short growing season where a seasonal active layer thaws (French 2017), and C accumulates when the addition of plant litter exceeds losses from decomposition (Yu *et al* 2011). The maintenance of a near-surface water table by seasonal active layer

thaw, snow melt and summer precipitation (Woo and Winter 1993) limits C losses from microbial decomposition (Heffernan *et al* 2020) and encourages the growth of decomposition-resistant plants such as *Sphagnum* mosses (Rydin *et al* 2006).

High-latitude regions of the Northern Hemisphere are now experiencing warming at a rate two to three times the global average (Masson-Delmotte *et al* 2018). Permafrost extent is shifting northwards with warming, evidenced by the thawing of peatlands in the discontinuous permafrost zones of North America (Camill 2005) and Eurasia (Åkerman and Johansson 2008, Payette *et al* 2004). Deeper thaw

increases the amount of soil organic matter vulnerable to decomposition, while rising temperatures simultaneously increase the rate of microbial decomposition; both contribute to increased greenhouse gas (GHG) emissions and a positive feedback with climate (Jeong *et al* 2018). In addition, climate-driven drying may expose peat to increased aerobic decomposition, leading to increases in carbon dioxide (CO₂) emissions (Ise *et al* 2008), while thaw-induced wetting has been associated with elevated methane (CH₄) emissions (Christensen *et al* 2004). However, these C losses may be partially offset or even reversed by improved plant productivity during longer growing seasons (Gallego-Sala *et al* 2018, Taylor *et al* 2019, Heffernan *et al* 2020).

Although studies of degrading permafrost peatlands have established a relationship between GHG fluxes, permafrost thaw and hydrological conditions, intensive monitoring in most areas began no earlier than the 1990s (Johansson *et al* 2006). Therefore, a palaeoecological approach using proxies—such as testate amoebae (Lamarre *et al* 2012, Jones *et al* 2013, Swindles *et al* 2015a, Pelletier *et al* 2017, Taylor *et al* 2019) and plant macrofossils (Fritz *et al* 2016, Gałka *et al* 2017a)—provides a valuable longer-term perspective. A recent palaeohydrological study of European peatlands in mainly temperate latitudes showed widespread 20th century drying (Swindles *et al* 2019); however, permafrost peatlands are subject to processes that are unique to cold regions. Gradual permafrost thaw increases active layer thickness and can lead to evaporation-driven drying (van Bellen *et al* 2018, Zhang *et al* 2018a). If a threshold point in permafrost thaw is reached structural collapse and wetting can occur, often linked to deep C losses (O'Donnell *et al* 2012, Jones *et al* 2017, Turetsky *et al* 2020), but there is evidence that in some instances abrupt permafrost thaw can lead to increased post-thaw C accumulation that partially or completely offsets deep C losses (Jones *et al* 2013, Swindles *et al* 2015b, Heffernan *et al* 2020). This non-linear response suggests ecological thresholds and autogenic feedbacks may be important, but these are not yet fully understood.

The uncertainty over the future of permafrost peatland C stocks mirrors that of the entire permafrost zone, arising from limited understanding of feedbacks between changes in hydrological regime, vegetation shifts, and permafrost thaw (Abbott *et al* 2016). In this study we investigate the ecohydrological and carbon dynamics response of permafrost peatlands in a rapidly warming region of subarctic Sweden. We conduct multiproxy palaeoecological reconstructions from ten peat profiles across eight sites that are reliably dated at high-resolution using ²¹⁰Pb, ¹⁴C and tephrochronology. More specifically we aim to (a) reconstruct changes in peatland vegetation, moisture conditions and carbon dynamics over at least

the last 300 years, (b) determine the relationship between any important changes and measured climatic variables and (c) better quantify autogenic ecohydrological feedbacks operating in permafrost peatlands.

2. Methods


2.1. Study region and sampling

Our study region near Abisko in northern Sweden (figure 1) is in the discontinuous permafrost zone and is characterised by extensive palsas, peat plateaus, bogs, and fens, many of which are currently experiencing permafrost degradation (Åkerman and Johansson 2008) and are no longer in equilibrium with climate (Olvmo *et al* 2020). Therefore, our study region may indicate how peatland areas currently with more extensive permafrost may respond to future warming. In total, we sampled ten peat profiles across eight sites within a ~60 km radius of each other. All sites were underlain by permafrost and were sampled to the base of the active layer, excluding Maunuvuoma fen where no permafrost was present. Monoliths were cut out from the peat at all sites, with the exception of Stordalen palsa and Maunuvuoma fen, that were sampled using a Russian corer (De Vleeschouwer *et al* 2010). Refer to supplementary section 4 (available online at stacks.iop.org/ERL/16/034001/mmedia) for imagery of sampling site locations.

The peatlands in our study region formed as early as ~9500 yr BP following the retreat of the Fennoscandian Ice Sheet (Sannel *et al* 2018), with peatland initiation linked to warming growing seasons and potentially increased precipitation (Morris *et al* 2018). Yet, permafrost may only have started forming in Fennoscandia from ~1500 yr BP (Treat and Jones 2018), perhaps as late as the Little Ice Age. Abisko has warmed by 1.59 °C in the last century (figure 1), far above the mean global increase of 0.91 °C (Lenssen *et al* 2019, GISTEMP Team 2021) (averages for 1913–1922 against 2003–2012), with annual average temperatures in some areas of our study region now above 0 °C—a key threshold for permafrost and ecological dynamics (Callaghan *et al* 2010).

2.2. Age modelling

The chronology of these peat profiles was determined from ²¹⁰Pb, ¹⁴C and tephra layers. ¹⁴C dates were calibrated using the IntCal13 calibration curve (Reimer *et al* 2013) and the northern hemisphere zone 1 post-bomb calibration curve (Hua *et al* 2013). Annually-resolved tephra layers found in Stordalen palsa (Hekla 1158 at 23 cm; Hekla 1104 at 30 cm)—see Cooper *et al* (2019)—were included in the age-depth model. Age-depth profiles (supplementary section 3) were constructed using PLUM (Aquino-López *et al* 2018), which is a Bayesian modelling approach, for a more robust integration of ²¹⁰Pb, ¹⁴C and tephra dates.



2.3. Peat properties

Peat was stored at 4 °C before bulk density and loss-on-ignition (LOI; %) analyses were completed in the laboratory following Chambers *et al* (2011). Bulk density was measured for contiguous 0.5 cm thick layers, while LOI was determined for 1 cm thick layers. Bulk density (g cm^{-3}) was calculated by dividing dry mass of peat (g; dried overnight at 105 °C) by the total sample volume (cm^3). LOI was calculated by subtracting ash mass (g; after 8 h in 550 °C furnace) from dry mass (g), dividing this by dry mass and multiplying the product by 100. C and nitrogen (N) content was measured for 0.5 cm layers in each peat profile on a Thermo Scientific Flash (2000) Series CHNS/O analyser.

2.4. Carbon accumulation


Apparent C accumulation rate (ACAR; $\text{g C m}^{-2} \text{yr}^{-1}$) was calculated for each peat profile by multiplying the accumulation rate of peat ($\text{g m}^{-2} \text{yr}^{-1}$), determined from respective age-depth models, by the proportion of C in each sample. Peat decay modelling (Clymo 1984) has been successfully used in other studies to calculate C accumulation under constant conditions—i.e. partially accounting for incomplete decomposition of surface peats—and to interpret the influence of allogenic (external) forcing (Zhang *et al* 2018b, 2020). However, here subjectivity in determining the transition of oxic to anoxic peat in these

permafrost peatland sites, confounded by uncertainties associated with fitting exponential curves to data (Belyea and Baird 2006), made use of such an approach inappropriate. Therefore, we have used a case-by-case discursive approach to interpret the relationship between ACAR and ecohydrological or climatic variables.

2.5. Water-table reconstructions

Testate amoebae were prepared and analysed following a modified version of Booth *et al* (2010); peat samples of 2 cm^3 were boiled in water for 10 min and stirred with a glass rod. This solution was rinsed through a 300 μm sieve, back-sieved through a 15 μm mesh and left to settle. Slides were made up for microscopy and a minimum of 100 individual testate amoebae were counted per sample at 200–400 \times magnification. In 10.5% (24 out of 228) of the samples a minimum of 50 individuals were counted owing to low abundance (supplementary section 5). Testate amoeba species identification was aided by reference to relevant literature (Charman *et al* 2000, Siemensma 2021). Refer to Swindles *et al* (2015b) for information on counting of testate amoebae in Stordalen palsa and Maunuvuoma fen (an additional 54 samples).

Taxonomies were harmonised to that of the Amesbury *et al* (2016) European-wide transfer function which was then applied to testate amoeba



abundance data to reconstruct past water-table depth (WTD). This transfer function uses a weighted average tolerance-down weighted model with inverse de-shrinking. Errors are based on 999 bootstrap cycles. Full testate amoeba abundance data can be found in supplementary section 1. Standardised z-scores of WTD reconstructions were calculated following Swindles *et al* (2015c) for the periods 1500 CE to present (see figure 3) and 1913 CE to present (see figure 4).

Testate amoeba species were grouped by k-means clustering into three groups (wet, intermediate and dry indicators—see figure 2) based upon their water-table optima values in the transfer function (Amesbury *et al* 2016). The testate amoeba indicator

percentages allow for an assessment of the homogeneity of WTD optima values within each sample—high heterogeneity could in theory indicate seasonal variation in hydrological regime or that wet and dry periods are captured within a single sample.

2.6. Plant macrofossils

Plant macrofossils were analysed for contiguous 1 cm thick layers in all peat profiles. Samples of 5 cm³ were washed under a warm-water spray and sieved using a 0.2 mm mesh. Initially, the entire sample was examined with a stereomicroscope to obtain volume percentages of individual subfossils of vascular plants and mosses. The subfossil car-pological remains and vegetative fragments (leaves,





Figure 2. (Continued.)

rootlets, epidermis) were identified using identification keys (Hadenäs 2003, Smith 2004, Mauquoy and van Geel 2007). Identification of *Sphagnum* to species level was carried out separately based upon analyses of stem leaves using specialist keys (Hölzer 2010, Laine *et al* 2011). See Gałka *et al* (2017b) for a more detailed methodology for plant macrofossil analysis. Entire plant macrofossil records can be found in supplementary section 2, along with the species groupings used for presentation in figure 2. Refer to Gałka *et al* (2017a) regarding

the plant macrofossil analysis of Stordalen palsa and Maunuvuoma fen.

2.7. Climatic data

Temperature and precipitation data for regional weather stations were downloaded from the Swedish Meteorological and Hydrological Institute (www.smhi.se/en). Climate data were also provided by Abisko Scientific Research Station. Growing degree days (GDD_0) were calculated annually by summing daily temperature values above 0°C .




For the Abisko station Penman–Monteith potential evapotranspiration (mm d^{-1}) was calculated for the period July 1984 CE to December 2002 CE, where data of sufficient climatic variables were available. For this period, gaps made up 6.04% of the data; therefore, gaps (up to 20 d) were filled using linear interpolation—following this process gaps made up only 0.34% of the data. This interpolation was based upon the average of 5 d prior to and after a gap. If there were no data for 5 d prior to or after the gap, then data from 15 d before or after respectively was extrapolated to fill the gap. For the Penman–Monteith calculations, roughness length (z_0) values measured at a boreal Swedish peatland (Alekseychik *et al* 2017) and albedo values from a study at Stordalen (Stiegler *et al* 2016) were used.

The period for which there is Penman–Monteith evapotranspiration data is limited in timespan, although temperature, GDD_0 and sunlight hours

data—related to evapotranspiration rates—are available from 1913 to present. Therefore multiple linear regression models using these variables were fitted to the Penman–Monteith evapotranspiration data at a monthly resolution for the period July 1984 to December 2002 in R v.3.6.1 (R Core Team 2019). The relative quality of these models was assessed by comparing corrected Akaike information criterion (AICc) values in the R package ‘MuMIn’ v.43.6 (Barton 2019). The best performing model with a zero intercept had GDD_0 , temperature and sunlight hours as explanatory variables and had an adjusted R-squared of 0.87. This multiple linear regression model was then used to model monthly evapotranspiration at Abisko from 1913 to 2017 CE.

2.8. Theil-sen regression

The relationship between climate variables (GDD_0 , evapotranspiration and precipitation) at Abisko and



Standardised WTD by core

Theil-sen beta slope			SP	SB	RB	BB	RP	GP1	GP2	MF	VB	OP
1	GDD ₀ (°C days)	Coefficient	0.002	0.004	0.003	0.006	0.002	-0.005	-0.005	-0.006	-0.004	0.000
		Beta	0.29	0.55	0.42	0.69	0.27	-0.68	-0.85	-0.78	-0.45	-0.01
0.5	ET (mm)	Coefficient	0.005	0.028	0.027	0.048	0.017	-0.029	-0.042	-0.025	-0.018	-0.012
		Beta	0.10	0.52	0.45	0.85	0.26	-0.52	-0.87	-0.45	-0.35	-0.17
0	Precipitation (mm)	Coefficient	0.001	0.006	0.009	0.010	0.002	-0.011	-0.005	-0.011	-0.013	-0.012
		Beta	0.04	0.26	0.40	0.48	0.06	-0.44	-0.27	-0.59	-0.63	-0.44
-0.5	P-E (mm)	Coefficient	0.002	0.001	0.005	0.011	-0.007	-0.013	0.001	-0.012	-0.011	-0.011
		Beta	0.08	0.04	0.25	0.51	-0.21	-0.53	0.06	-0.66	-0.63	-0.35
-1												

Figure 4. Relationship between climatic variables and water-table depth (WTD) across all sites. Annual (a) growing degree days above 0 °C (GDD₀; °C days), (b) evapotranspiration (ET; mm) and (c) precipitation (P; mm); (d) precipitation minus evapotranspiration (P-E; mm) for Abisko, 1913 CE to present. Standardised WTD data from (e) drying and (f) wetting or fluctuating peat profiles since 1913 CE, and (g) Theil-sen regression matrix (coefficients and betas) of standardised WTD by peat profile against each climatic variable. SP = Stordalen palsa, SB = Stordalen bog, RB = Ribasvuomuš bog, BB = Bergfors bog, RP = Rensjön palsa, GP1 = Gurasáhpí palsa 1, GP2 = Gurasáhpí palsa 2, MF = Maunuvuoma fen, VB = Veigi bog and OP = Orusjohka palsa.

WTD from 1913 to 2012 CE was explored using Theil-Sen robust regression in the R package 'deming' v.1.4 (Therneau 2018). WTD data were standardised following Swindles *et al* (2015c) for the time period 1913–2012 CE. Climatic data were averaged for the time period each WTD sample represented in the age-depth model to make data comparable. The testate amoeba subfossil assemblage has accumulated over the time period represented by each sample slice (typically 1 cm) and therefore provides a WTD reconstruction averaged to this timeframe. Theil-sen regression coefficients were then standardised to produce beta coefficients.

3. Results and discussion

3.1. Response of permafrost peatlands to recent warming

Our palaeoecological reconstructions span at least the last 500 years and show a varied ecohydrological response to recent warming (figure 2). In terms of hydrological regime, local regression (loess) models highlight a divergent response to recent warming (figure 3). Stordalen palsa, Stordalen bog, Ribasvuomuš bog, Bergfors bog and Rensjön palsa have all undergone clear drying since the mid-20th century. In contrast, Gurasáhpi palsa 1 and 2 and Maunuvuoma fen have experienced a period of 20th century drying, followed by rapid wetting from ~1990 CE. Meanwhile, Veigi bog and Orusjohka palsa in Paittasjärvi valley to the south record asynchronously fluctuating wet and dry periods in recent decades. Individual sites have undergone wet or dry shifts earlier in the record (e.g. Stordalen palsa and Stordalen bog), but it is only within the last century that high-magnitude shifts in hydrological regime have occurred across all sites.

Vegetation across all sites demonstrates a degree of resistance (the ability of a population to avoid displacement during a period of stress; Harrison 1979), and in some cases resilience (the ability of a population to recover from change or disturbance when a period of stress has subsided; Harrison 1979) to recent hydrological shifts and climate change (figure 2). In Rensjön palsa, Maunuvuoma fen and Veigi bog, *Sphagnum fuscum* populations remain stable during the last century, demonstrating resistance to warming and changing hydrological regimes, probably because of its broad hydrological tolerance (Rydin and McDonald 1985). In Gurasáhpi palsa 1, *S. fuscum* demonstrates resilience with populations recovering following a period of intense drying—inferred from WTD reconstructions and high presence of unidentified organic matter (UOM)—in the mid-20th century. The resilience and resistance of *S. fuscum* may have been aided by it forming dense carpets and its increased growth rate with warming; however, a reduced bulk density with faster growth could hamper moisture

retention capabilities in the long-term and limit future productivity (Dorrepaal *et al* 2004). Both Ribasvuomuš bog and Bergfors bog have been dominated by *Dicranum* spp.—predominantly the hummock species *Dicranum elongatum* (supplementary section 2)—since ~1600 CE, with a period of Ericaceae dominance in the late 20th century. Initially, drying may have allowed Ericaceae to outcompete *D. elongatum*, but further drying appears to have favoured *D. elongatum*—which can tolerate extreme dry habitats (Sonesson *et al* 2002)—allowing the species to re-emerge as dominant. Stordalen palsa and Stordalen bog experience a shift from ~1950 CE to more ombrotrophic conditions, from sedge (Cyperaceae) and herbs to bryophyte dominance. From ~1800 CE to ~2000 CE, Stordalen palsa experiences a stable WTD of ~6 cm, while exhibiting shifts in plant communities (figure 2(a)). From ~1800 CE, brown moss (*Drepanocladus* sp., see supplementary section 4) dominates, from ~1900 CE there is a period of *Sphagnum lindbergii* and sedge and herb dominance before a transition in recent decades to *Sphagnum balticum*. In the absence of clear hydrological change during this period, a reduction in available nutrients could be driving the shift from *S. lindbergii* to *S. balticum* (Gunnarsson *et al* 2004). *S. balticum* is a species typical of moderately-wet conditions (Johansson and Linder 1980), yet demonstrates resistance to considerable drying since ~2000 CE. Similarly, in Stordalen bog the typically wet hollow species *Sphagnum fallax* demonstrates resistance to drying in recent decades, likely owing to its tolerance to desiccation (Wagner and Titus 1984).

All sample locations—excluding Maunuvuoma fen—are currently underlain by permafrost. A transition from Cyperaceae and herb and UOM dominated assemblages to *Sphagnum* spp. or *Dicranum* spp. dominance may represent the aggradation of permafrost in Maunuvuoma fen ~2150 cal. BP, in Orusjohka palsa ~400 cal. BP and ~200 cal. BP in Ribasvuomuš bog (figure 2). In general permafrost is thought to have begun aggrading in Fennoscandia ~450 cal. BP (Treat and Jones 2018), with evidence that permafrost aggradation may have been occurred in Stordalen mire as early as 2650 cal BP and late as 120 cal BP (Kokfelt *et al* 2010). Greater certainty in the reconstruction of permafrost dynamics would likely be obtained through the study of subfossil oribatid mite communities (Markkula and Kuhry 2020).

Any increases in C accumulation owing to improved productivity with warming or hydrological changes are difficult to distinguish from the artefact of incomplete decomposition in recent peats (Young *et al* 2019)—characterised by an uptick in recent ACAR (see figure 2). Nonetheless, it is clear there is a decrease in C accumulation in Stordalen palsa, Rensjön palsa, Ribasvuomuš bog and Gurasáhpi palsa 1 during drying periods in the last century. A decrease in ACAR from the late-20th century in Stordalen

palsa (figure 2(a)) indicates a reduction in C accumulation great enough to overcome any artefact of incomplete decomposition, likely responsible for the increase in ACAR during the mid-20th century. We suggest this decreased C accumulation in the surface peat is probably because of high levels of aerobic decomposition associated with drying from ~2000 CE. This drying appears to be to such an extent as to have caused decomposition further down the profile (secondary decomposition) and consequently a reduced ACAR in the late-20th century (Frolking *et al* 2014, Morris *et al* 2015a). In Rensjön palsa, there is a decrease in ACAR coincident with drying prior to the recent uptick in ACAR (figure 2(g)). Similarly, increased UOM—associated with high levels of decomposition—is observed alongside drying at the surface of Ribasvuomuš bog (figure 2(c)) and at ~1950 CE in Gurasáhpi palsa 1 (figure 2(d)). The UOM layer in both of these sites is stratigraphically above older, better preserved peat, suggesting an increase in the rate of decomposition that led to poorer preservation of more recent plant material. The subsequent replacement of UOM by a dominance of *S. fuscum* from ~1990 CE alongside wetting in Gurasáhpi palsa 1 suggests lower decomposition due to wetting and may indicate an increase in C sequestration. A complete lack of macrocharcoal in all our peat profiles shows fire has not been important in affecting C dynamics, in contrast to the importance of fire in some North American permafrost peatlands (Robinson and Moore 2000, Camill *et al* 2009, Jones *et al* 2013, Gibson *et al* 2018).

3.2. Climatic and autogenic drivers of ecosystem change

Abisko instrumental records show an increase in temperature, GDD₀, evapotranspiration and precipitation during the early- and late-20th century, while precipitation minus evapotranspiration (P-E) experiences a slight increase since the late 20th century (figures 1 and 4). Precipitation observed at Abisko is lower than that of other local stations (figure 1). Sites experiencing drying show a positive relationship between WTD and climatic variables, while sites experiencing wetting or fluctuating WTD trends exhibit a negative relationship (figure 4). This suggests that autogenic processes and site-specific factors play a key role in the hydrological response to climatic forcing. Under a steady-state climate, models of cyclic palsa formation and degradation associated with vegetation and snow accumulation feedbacks have been proposed, with drying during permafrost aggradation and wetting during degradation (Zuidhoff and Kolstrup 2005). However, recent and continued warming in our study region makes further permafrost aggradation and palsa formation all but impossible.

Greater winter snow depth, combined with rising summer temperatures, are thought to be the cause of

thaw in the permafrost peatlands of northern Sweden (Sannel *et al* 2016). Therefore, permafrost thaw, combined with increased growing season moisture from precipitation and snow melt, provide a plausible explanation for sites experiencing recent wetting (figure 4). Wetting from ~1990 CE in Gurasáhpi palsa 1 and 2 and Maunuvuoma fen was preceded by a period of drying (figure 3). Here, a threshold point in drying and permafrost thaw may have been reached causing surface collapse and associated rewetting (Swindles *et al* 2015b), suggesting the state of localised permafrost degradation is an important control on hydrological regime. Additionally, increases in decomposition with drying—as evidenced by a high mid-20th century concentration of UOM (~60%) in Gurasáhpi palsa 1 (figure 2(d))—may have reduced peatland surface hydraulic conductivity and drainage enough to facilitate recent wetting (Morris *et al* 2015b).

Recent drying in Stordalen palsa, Stordalen bog, Ribasvuomuš bog, Bergfors bog, Rensjön palsa (and drying prior to wetting in Gurasáhpi palsa 1 and 2 and Maunuvuoma fen; figure 3) is likely driven by increases in evapotranspiration and potentially productivity (i.e. GDD₀; figure 4). Evaporation-driven drying with recent warming has also been observed in other Fennoscandian and northwest Russian permafrost peatlands (Zhang *et al* 2018a). The lower precipitation experienced at Stordalen palsa and Stordalen bog (figure 1) has perhaps made these sites more susceptible to drying. The slight recent increase in P-E appears to be of lesser importance to WTD in drying sites, but shows a stronger relationship with WTD at wetting sites (figure 4(g)). A potential explanation for this is differences in permafrost structure affecting runoff rate and drainage. For example, localised permafrost collapse could create a topographic depression more favourable for retention of surplus precipitation. In other studies (Sonesson *et al* 2002), greater *S. fuscum* or *D. elongatum* growth has been associated with increases in temperature and precipitation, and this increased productivity may be the cause of a deepening of the water-table as the peat surface grows rapidly upwards (van Bellen *et al* 2018). However, this explanation could only apply where there is no clear decrease in C accumulation associated with drying, e.g. Bergfors bog with recent *D. elongatum* growth (figure 2). The insulating properties of dry bryophyte layers are likely to have reduced the amplitude of soil temperatures and conversely may have acted to slow the rate of permafrost thaw (Soudzilovskaia *et al* 2013). Nevertheless, increased connectivity of drainage pathways with permafrost thaw could be increasing runoff, leading to drying (Haynes *et al* 2018)—here local topographic setting is likely to be important. Comparison of imagery of our study sites from 1959 to 1960 CE and 2012 to 2019 CE provides some evidence of changes in permafrost peatland structure and drainage (supplementary section 4). These

structural changes associated with permafrost thaw have the potential to cause both wetting (e.g. surface collapse) and drying (e.g. drainage of surface water features).

Both Orusjohka palsa and Veigi bog experience fluctuations in wetness since the mid-20th century that are asynchronous to each other and the other sites (figure 3). This fluctuating WTD may represent a flickering of ecosystem state (Wang *et al* 2012), with negative autogenic feedbacks moderating the response to recent climate forcing (Swindles *et al* 2012, Waddington *et al* 2015, Morris *et al* 2015a). Despite comparable growing season conditions, colder annual temperatures recorded near these sites at Nikkaloukta station (figure 1) may be slowing rates of localised permafrost thaw, allowing autogenic feedbacks to dominate. For example, increases in bryophyte productivity with higher GDD_0 may facilitate rapid vertical growth and an increase in WTD below the peatland surface. Higher aerobic decomposition with drying—evidenced in Orusjohka palsa ~1985 CE by a high UOM concentration (60%) and fungal remains count (figure 2(j); supplementary section 2)—may cause a decrease in hydraulic conductivity of upper peats that then reduces water loss, inducing wetting. These more favourable (wetter) hydrological conditions may then again allow for greater bryophyte productivity as GDD_0 rise—continuing to drive fluctuations in hydrological conditions.

3.3. Implications for ecohydrological and carbon dynamics in global permafrost peatlands

Here we show a divergent response of permafrost peatland ecohydrological regimes to climate change over the last century (figures 2–4) and highlight the importance of internal autogenic and site-specific factors in these ecosystems. We link climate-driven drying with reduced C accumulation at four sites, but observe no clear relationship between wetting sites and C accumulation (figure 2). Our data provide multi-proxy high-resolution evidence to substantiate previous suggestions of a heterogeneous response to recent warming in permafrost peatlands from longer Holocene archives in Fennoscandia (Zhang *et al* 2018b) and late Holocene reconstructions in High Arctic Canada (Sim *et al* 2019). Our findings illustrate how localised climatic variations between sites and autogenic processes linked to permafrost thaw, peatland structure, productivity and decomposition can combine in many ways to determine the future trajectory of permafrost peatlands, exemplified by the potential to cause both wetting and drying. Consequently, models of catastrophic C loss associated with drying (e.g. Ise *et al* 2008) do not capture the complex ecological and hydrological dynamics of permafrost peatlands. Models incorporating permafrost, vegetation and hydrological dynamics (e.g. Chaudhary *et al* 2020) are likely to provide a truer representation of reality and here we provide

empirical data to inform and test such models. Future research may be better able to disentangle the relative importance of autogenic and site-specific factors by: (a) collating existing palaeoecological records coupled with climatic data from a variety of permafrost peatlands; and (b) combining a palaeoecological approach with models of peatland ecosystem development (e.g. Morris *et al* 2015a), specifically adapted for permafrost conditions.

We show that bryophyte populations have demonstrated resistance and in some cases resilience to climatic and hydrological changes over the last century (figure 2). Bryophytes are abundant across high-latitude ecosystems, with *Sphagnum* spp.—such as *S. fuscum*—dominating boreal and tundra permafrost peatlands (Treat *et al* 2016). This bryophyte resistance and resilience may allow for widespread increases in productivity with warming (Charman *et al* 2013), perhaps to an extent where post-thaw surface C accumulation offsets C losses from deeper peat (Heffernan *et al* 2020). However, bryophyte populations may be vulnerable to replacement by vascular plants with a greater availability of near-surface nitrogen (N) (Berendse *et al* 2001) associated with permafrost thaw (Keuper *et al* 2012) and increasing atmospheric N deposition (Galloway *et al* 2004). Nonetheless, despite levels of N deposition being greater in Fennoscandia than other permafrost regions in recent centuries (Galloway *et al* 2004), our data show stable bryophyte populations. This resistance and resilience suggests a degree of long-term bryophyte sustainability in global permafrost peatlands and through their insulating properties (Soudzilovskaia *et al* 2013) they may help slow rates of inevitable permafrost thaw.

The divergent response to recent warming seen here in peatland ecosystems on the limits of discontinuous permafrost extent, may act as an indication for the future trajectory of more northerly or continental ecosystems currently exhibiting continuous permafrost (Brown *et al* 2002). These areas of extensive permafrost peatlands—such as northern Canada, Alaska and northern Russia—are likely to experience greater increases in temperature and precipitation than Fennoscandia in the 21st century (Christensen *et al* 2013). Furthermore, there is likely to be regional variation in the importance of certain ecological processes, such as fire frequency—for which our findings suggest has a limited role, currently, in northern Sweden. Therefore, we can likely expect a degree of heterogeneity between Arctic regions in the future trajectory of permafrost peatlands.

4. Conclusions

Our findings suggest that: (a) permafrost peatlands have the potential to exhibit both wetting and drying under future climate change, owing to autogenic processes linked to permafrost thaw, peatland structure, productivity and decomposition; (b) although

ACAR should be interpreted cautiously, hydrological conditions appear to be an important control on C dynamics; (c) models of catastrophic C loss associated with drying do not capture the complex ecological and hydrological dynamics of permafrost peatlands; and (d) bryophyte populations—specifically *S. fuscum* and *Dicranum* spp.—demonstrate resistance and in some cases resilience to recent climatic and hydrological changes. Our study provides a detailed insight into the recent response of permafrost peatlands to climate change in Fennoscandia and warns against an overly-simple approach to considering their future ecohydrological dynamics and role in the global C cycle.

Data availability

All data that support the findings of this study are included within the article (and any supplementary files).

Acknowledgments

T G S is in receipt of a UK Natural Environment Research Council Training Grant (NE/L002574/1). G T acknowledges the support of a Worldwide University Network (WUN) Grant which helped fund the fieldwork (Project: Arctic Environments, Vulnerabilities and Opportunities). C L C acknowledges a Leeds Anniversary Research Scholarship at the University of Leeds, in addition to support from the Climate Research Bursary Fund awarded by the Priestley International Centre for Climate. A V G-S and D J C were funded by the Natural Environment Research Council (NERC standard Grant Numbers NE/I012915/1). Staff from the Abisko Scientific Research Station assisted with field logistics and provided climatic data. Kallax Flyg AB provided helicopter support. We thank Matt Amesbury for useful discussions on testate amoeba transfer functions.

Author contributions

T G S, G T S, P J M and A J B designed the study. T G S performed testate amoebae analysis, M G conducted plant macrofossil analysis and C L C undertook tephra analysis. A V G-S and T G S carried out bulk density determinations, A V G-S analysed C and N content and D J C and T P R carried out ²¹⁰Pb dating. WB carried out a number of ¹⁴C dates. T G S, P J M and D J M, analysed climatic data. G T S carried out peat profile sampling in the field. M A A-L carried out modelling of the age with depth profiles. T G S wrote the manuscript, under supervision of G T S, P J M and A J B, with input from all authors.

Conflict of interest

The authors declare no competing interests.

ORCID iDs

Thomas G Sim  <https://orcid.org/0000-0001-8604-9996>

Angela V Gallego-Sala  <https://orcid.org/0000-0002-7483-7773>

Dan J Charman  <https://orcid.org/0000-0003-3464-4536>

References

- Abbott B W *et al* 2016 Biomass offsets little or none of permafrost carbon release from soils, streams, and wildfire: an expert assessment *Environ. Res. Lett.* **11** 034014
- Åkerman H J and Johansson M 2008 Thawing permafrost and thicker active layers in sub-arctic Sweden *Permafrost. Periglacial Process.* **19** 279–92
- Alekseychik P K, Korrensalo A, Mammarella I, Vesala T and Tuittila E S 2017 Relationship between aerodynamic roughness length and bulk sedge leaf area index in a mixed-species boreal mire complex *Geophys. Res. Lett.* **44** 5836–43
- Amesbury M J *et al* 2016 Development of a new pan-European testate amoeba transfer function for reconstructing peatland palaeohydrology *Quat. Sci. Rev.* **152** 132–51
- Aquino-López M A, Blaauw M, Christen J A and Sanderson N K 2018 Bayesian analysis of 210 Pb dating *J. Agric. Biol. Environ. Stat.* **23** 317–33
- Bartoń K 2019 MuMIn: Multi-Model Inference, Version 1.43.6 R *Packag*
- Belyea L R and Baird A J 2006 Beyond ‘The Limits to Peat Bog Growth’: cross-scale feedback in peatland development *Ecol. Monogr.* **76** 299–322
- Berendse F *et al* 2001 Raised atmospheric CO₂ levels and increased N deposition cause shifts in plant species composition and production in Sphagnum bogs *Glob. Change Biol.* **7** 591–8
- Booth R K, Lamentowicz M and Charman D J 2010 Preparation and analysis of testate amoebae in peatland palaeoenvironmental studies *Mires Peat* **7** 1–7
- Brown J, Ferrians O, Heginbottom J A and Melnikov E 2002 Circum-Arctic Map of Permafrost and Ground-Ice Conditions, Version 2 (<https://doi.org/10.7265/skbg-kf16>)
- Callaghan T V, Bergholm F, Christensen T R, Jonasson C, Kokfelt U and Johansson M 2010 A new climate era in the sub-Arctic: accelerating climate changes and multiple impacts *Geophys. Res. Lett.* **37** L14705
- Camill P 2005 Permafrost thaw accelerates in boreal peatlands during late-20th century climate warming *Clim. Change* **68** 135–52
- Camill P, Barry A, Williams E, Andreassi C, Limmer J and Solick D 2009 Climate-vegetation-fire interactions and their impact on long-term carbon dynamics in a boreal peatland landscape in northern Manitoba, Canada *J. Geophys. Res.* **114** G04017
- Chambers F M, Beilman D W and Yu Z 2011 Methods for determining peat humification and for quantifying peat bulk density, organic matter and carbon content for palaeostudies of climate and peatland carbon dynamics *Mires Peat* **7** 1–10 (<http://mires-and-peat.net/pages/volumes/map07/map0707.php>)
- Charman D J *et al* 2013 Climate-related changes in peatland carbon accumulation during the last millennium *Biogeosciences* **10** 929–44
- Charman D J, Hendon D and Woodland W A 2000 *The Identification of Testate Amoebae (Protozoa: Rhizopoda) in Peats: QRA Technical Guide No. 9* (London: Quaternary Research Association)
- Chaudhary N, Westermann S, Lamba S, Shurpali N, Sannel A B K, Schurgers G, Miller P A and Smith B 2020 Modelling past

- and future peatland carbon dynamics across the pan-Arctic *Glob. Change Biol.* **26** 4119–33
- Christensen J H *et al* 2013 Climate phenomena and their relevance for future regional climate change *The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker, D Qin, G-K Plattner, M Tignor, S K Allen, J Boschung, A Nauels, Y Xia, V Bex and P M Midgley (Cambridge: Cambridge University Press) pp 1217–310
- Christensen T R, Johansson T, Åkerman H J, Mastepanov M, Malmer N, Friberg T, Crill P and Svensson B H 2004 Thawing sub-arctic permafrost: effects on vegetation and methane emissions *Geophys. Res. Lett.* **31** L04501
- Clymo R S 1984 The limits to peat bog growth *Philos. Trans. R. Soc. B* **303** 605–54
- Cooper C L, Swindles G T, Watson E J, Savov I P, Gałka M, Gallego-Sala A and Borken W 2019 Evaluating tephrochronology in the permafrost peatlands of northern Sweden *Quat. Geochronol.* **50** 16–28
- De Vleeschouwer F, Chambers F M and Swindles G T 2010 Coring and sub-sampling of peatlands for palaeoenvironmental research *Mires Peat* **7** 1–10 (<http://mires-and-peat.net/pages/volumes/map07/map0701.php>)
- Dorrepaal E, Aerts R, Cornelissen J H C, Callaghan T V and Van Logtestijn R S P 2004 Summer warming and increased winter snow cover affect *Sphagnum fuscum* growth, structure and production in a sub-arctic bog *Glob. Change Biol.* **10** 93–104
- French H M 2017 *The Periglacial Environment* (New York: Wiley)
- Fritz M, Wolter J, Rudaya N, Palagushkina O, Nazarova L, Obu J, Rethemeyer J, Lantuit H and Wetterich S 2016 Holocene ice-wedge polygon development in northern Yukon permafrost peatlands (Canada) *Quat. Sci. Rev.* **147** 279–97
- Frolking S, Talbot J and Subin Z M 2014 Exploring the relationship between peatland net carbon balance and apparent carbon accumulation rate at century to millennial time scales *The Holocene* **24** 1167–73
- Gałka M, Szal M, Watson E J, Gallego-Sala A, Amesbury M J, Charman D J, Roland T P, Edward Turner T and Swindles G T 2017a Vegetation succession, carbon accumulation and hydrological change in subarctic Peatlands, Abisko, Northern Sweden. *Permafr. Periglac. Process.* **28** 589–604
- Gałka M, Tobolski K, Lamentowicz Ł, Ersek V, Jassey V E J, van der Knaap W O and Lamentowicz M 2017b Unveiling exceptional Baltic bog ecohydrology, autogenic succession and climate change during the last 2000 years in CE Europe using replicate cores, multi-proxy data and functional traits of testate amoebae *Quat. Sci. Rev.* **156** 90–106
- Gallego-Sala A V *et al* 2018 Latitudinal limits to the predicted increase of the peatland carbon sink with warming *Nat. Clim. Chang* **8** 907–13
- Galloway J N *et al* 2004 Nitrogen cycles: past, present, and future *Biogeochemistry* **70** 153–226
- Gibson C M, Chasmer L E, Thompson D K, Quinton W L, Flannigan M D and Olefeldt D 2018 Wildfire as a major driver of recent permafrost thaw in boreal peatlands *Nat. Commun.* **9** 3041
- GISTEMP Team 2021 GISS Surface Temperature Analysis (GISTEMP), version 4 NASA *Goddard Institute for Space Studies* (<https://data.giss.nasa.gov/gistemp/>)
- Gunnarsson U, Granberg G and Nilsson M 2004 Growth, production and interspecific competition in Sphagnum: effects of temperature, nitrogen and sulphur treatments on a boreal mire *New Phytol.* **163** 349–59
- Hadenäs L 2003 The European species of the *Calliargon–Scorpidium–Drepanocladus* complex, including some related or similar species *Meylania* **28** 1–116
- Harrison G W 1979 Stability under environmental stress: resistance, resilience, persistence, and variability *Am. Nat.* **113** 659–69
- Haynes K M, Cannon R F and Quinton W L 2018 Permafrost thaw induced drying of wetlands at Scotty Creek, NWT, Canada *Environ. Res. Lett.* **13** 114001
- Heffernan L, Estop-Aragonés C, Knorr K, Talbot J and Olefeldt D 2020 Long-term impacts of permafrost thaw on carbon storage in peatlands: deep losses offset by surficial accumulation *J. Geophys. Res. Biogeosciences* **125** e2019JG005501
- Hölzer A 2010 *Die Torfmoose Südwestdeutschlands Und Der Nachbargebiete* (Jena: Weissdorn-Verlag)
- Hua Q, Barbetti M and Rakowski A Z 2013 Atmospheric radiocarbon for the period 1950–2010 *Radiocarbon* **55** 2059–72
- Hugelius G *et al* 2014 Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps *Biogeosciences* **11** 6573–93
- Ise T, Dunn A L, Wofsy S C and Moorcroft P R 2008 High sensitivity of peat decomposition to climate change through water-table feedback *Nat. Geosci.* **1** 763–6
- Jeong S-J *et al* 2018 Accelerating rates of Arctic carbon cycling revealed by long-term atmospheric CO₂ measurements *Sci. Adv.* **4** eaao1167
- Johansson L G and Linder S 1980 Photosynthesis of Sphagnum in different microhabitats on a subarctic mire *Ecol. Bull.* **30** 181–90
- Johansson M, Christensen T R, Åkerman H J and Callaghan T V 2006 What determines the current presence or absence of permafrost in the Torneträsk region, a sub-arctic landscape in Northern Sweden? *Ambio* **35** 190–7
- Jones M C, Booth R K, Yu Z and Ferry P 2013 A 2200-Year Record of Permafrost Dynamics and Carbon Cycling in a Collapse-Scar Bog, Interior Alaska *Ecosystems* **16** 1–19
- Jones M C, Harden J, O'Donnell J, Manies K, Jorgenson T, Treat C and Ewing S 2017 Rapid carbon loss and slow recovery following permafrost thaw in boreal peatlands *Glob. Change Biol.* **23** 1109–27
- Keuper F, van Bodegom P M, Dorrepaal E, Weedon J T, van Hal J, van Logtestijn R S P and Aerts R 2012 A frozen feast: thawing permafrost increases plant-available nitrogen in subarctic peatlands *Glob. Change Biol.* **18** 1998–2007
- Kokfelt U, Reuss N, Struyf E, Sonesson M, Rundgren M, Skog G, Rosén P and Hammarlund D 2010 Wetland development, permafrost history and nutrient cycling inferred from late Holocene peat and lake sediment records in subarctic Sweden *J. Paleolimnol.* **44** 327–42
- Laine J, Harju P, Timonen T, Laine A, Tuittila E-S, Minkkinen K and Vasander H 2011 *The Intricate Beauty of Sphagnum Mosses: A Finnish Guide to Identification* (Helsinki: Department of Forest Sciences, University of Helsinki)
- Lamarre A, Garneau M and Asnong H 2012 Holocene paleohydrological reconstruction and carbon accumulation of a permafrost peatland using testate amoeba and macrofossil analyses, Kuujjuarapik, subarctic Québec, Canada *Rev. Palaeobot. Palynol.* **186** 131–41
- Lenssen N J, Schmidt G A, Hansen J E, Menne M J, Persin A, Ruedy R and Zys D 2019 Improvements in the GISTEMP uncertainty model *J. Geophys. Res. Atmos.* **124** 6307–26
- Markkula I and Kuhry P 2020 Subfossil oribatid mite communities indicate Holocene permafrost dynamics in Canadian mires *Boreas* **49** 730–38
- Masson-Delmotte V *et al* 2018 Intergovernmental Panel on Climate Change, 2018: summary for Policymakers *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change* pp 3–24
- Mauquoy D and van Geel B 2007 Mire and peat macros *Encyclopedia of Quaternary Science* ed S A Elias (Amsterdam: Elsevier) pp 2315–36
- Morris P J, Baird A J and Belyea L R 2015a Bridging the gap between models and measurements of peat hydraulic conductivity *Water Resour. Res.* **51** 5353–64
- Morris P J, Baird A J, Young D M and Swindles G T 2015b Untangling climate signals from autogenic changes in

- long-term peatland development *Geophys. Res. Lett.* **42** 10788–97
- Morris P J, Swindles G T, Valdes P J, Ivanovic R F, Gregoire L J, Smith M W, Tarasov L, Haywood A M and Bacon K L 2018 Global peatland initiation driven by regionally asynchronous warming *Proc. Natl Acad. Sci.* **115** 4851–6
- O'Donnell J A, Jorgenson M T, Harden J W, McGuire A D, Kanevskiy M Z and Wickland K P 2012 The effects of permafrost thaw on soil hydrologic, thermal, and carbon dynamics in an alaskan peatland *Ecosystems* **15** 213–29
- Olvmo M, Holmer B, Thorsson S, Reese H and Lindberg F 2020 Sub-arctic palsa degradation and the role of climatic drivers in the largest coherent palsa mire complex in Sweden (Vissávuopmi), 1955–2016 *Sci. Rep.* **10** 1–10
- Payette S, Delwaide A, Caccianiga M and Beauchemin M 2004 Accelerated thawing of subarctic peatland permafrost over the last 50 years *Geophys. Res. Lett.* **31** L18208
- Pelletier N, Talbot J, Olefeldt D, Turetsky M, Blodau C, Sonnentag O and Quinton W L 2017 Influence of Holocene permafrost aggradation and thaw on the paleoecology and carbon storage of a peatland complex in northwestern Canada *The Holocene* **27** 1391–405
- R Core Team 2019 R: a language and environment for statistical computing (www.r-project.org)
- Reimer P J et al 2013 IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP *Radiocarbon* **55** 1869–87
- Robinson S D and Moore T R 2000 The influence of permafrost and fire upon carbon accumulation in high boreal peatlands, northwest territories, Canada *Arctic, Antarct. Alp. Res.* **32** 155–66
- Rydin H, Gunnarsson U and Sundberg S 2006 The role of Sphagnum in peatland development and persistence *Boreal Peatland Ecosystems* ed R K Wieder and D H Vitt (Berlin: Springer) pp 47–65
- Rydin H and McDonald A J S 1985 Tolerance of Sphagnum to water level *J. Bryol.* **13** 571–8
- Sannel A B K, Hempel L, Kessler A and Prėskienis V 2018 Holocene development and permafrost history in sub-arctic peatlands in Tavvavuoma, northern Sweden *Boreas* **47** 454–68
- Sannel A B K, Hugelius G, Jansson P and Kuhry P 2016 Permafrost warming in a subarctic peatland—which meteorological controls are most important? *Permafr. Periglac. Process.* **27** 177–88
- Siemasma F J 2021 Microworld, world of amoeboid organisms *World-wide Electron. Publ.* (www.arcella.nl)
- Sim T G, Swindles G T, Morris P J, Gałka M, Mullan D and Galloway J M 2019 Pathways for ecological change in Canadian high arctic wetlands under rapid twentieth century warming *Geophys. Res. Lett.* **46** 4726–37
- Smith A J E 2004 *The Moss Flora of Britain and Ireland* (Cambridge: Cambridge University Press)
- Sonesson M, Carlsson B Å, Callaghan T V, Halling S, Björn L O, Bertgren M and Johanson U 2002 Growth of two peat-forming mosses in subarctic mires: species interactions and effects of simulated climate change *Oikos* **99** 151–60
- Soudzilovskaia N A, van Bodegom P M and Cornelissen J H C 2013 Dominant bryophyte control over high-latitude soil temperature fluctuations predicted by heat transfer traits, field moisture regime and laws of thermal insulation ed J Schweitzer *Funct. Ecol.* **27** 1442–54
- Stiegler C, Johansson M, Christensen T R, Mastepanov M and Lindroth A 2016 Tundra permafrost thaw causes significant shifts in energy partitioning *Tellus B Chem. Phys. Meteorol.* **68** 30467
- Swindles G T et al 2015a Evaluating the use of testate amoebae for palaeohydrological reconstruction in permafrost peatlands *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **424** 111–22
- Swindles G T et al 2015c The long-term fate of permafrost peatlands under rapid climate warming *Sci. Rep.* **5** 17951
- Swindles G T et al 2019 Widespread drying of European peatlands in recent centuries *Nat. Geosci.* **12** 922–8
- Swindles G T, Holden J, Raby C L, Turner T E, Blundell A, Charman D J, Menberu M W and Kløve B 2015b Testing peatland water-table depth transfer functions using high-resolution hydrological monitoring data *Quat. Sci. Rev.* **120** 107–17
- Swindles G T, Morris P J, Baird A J, Blaauw M and Plunkett G 2012 Ecohydrological feedbacks confound peat-based climate reconstructions *Geophys. Res. Lett.* **39** L11401
- Tarnocai C, Canadell J G, Schuur E A G, Kuhry P, Mazhitova G and Zimov S 2009 Soil organic carbon pools in the northern circumpolar permafrost region *Glob. Biogeochem. Cycles* **23** GB2023
- Taylor L S, Swindles G T, Morris P J, Gałka M and Green S M 2019 Evidence for ecosystem state shifts in Alaskan continuous permafrost peatlands in response to recent warming *Quat. Sci. Rev.* **207** 134–44
- Therneau T 2018 Deming, Version 1.4. R Package (<https://cran.r-project.org/web/packages/deming/deming.pdf>)
- Treat C C et al 2016 Effects of permafrost aggradation on peat properties as determined from a pan-Arctic synthesis of plant macrofossils *J. Geophys. Res. Biogeosciences* **121** 78–94
- Treat C C and Jones M C 2018 Near-surface permafrost aggradation in Northern Hemisphere peatlands shows regional and global trends during the past 6000 years *The Holocene* **28** 998–1010
- Turetsky M R et al 2020 Carbon release through abrupt permafrost thaw *Nat. Geosci.* **13** 138–43
- van Bellen S, Magnan G, Davies L, Froese D, Mullan-Boudreau G, Zaccone C, Garneau M and Shoty W 2018 Testate amoeba records indicate regional 20th-century lowering of water tables in ombrotrophic peatlands in central-northern Alberta, Canada *Glob. Change Biol.* **24** 2758–74
- Waddington J M, Morris P J, Kettridge N, Granath G, Thompson D K and Moore P A 2015 Hydrological feedbacks in northern peatlands *Ecohydrology* **8** 113–27
- Wagner D J and Titus J E 1984 Comparative desiccation tolerance of two Sphagnum mosses *Oecologia* **62** 182–7
- Wang R, Dearing J A, Langdon P G, Zhang E, Yang X, Dakos V and Scheffer M 2012 Flickering gives early warning signals of a critical transition to a eutrophic lake state *Nature* **492** 419–22
- Woo M-K and Winter T C 1993 The role of permafrost and seasonal frost in the hydrology of northern wetlands in North America *J. Hydrol.* **141** 5–31
- Young D M, Baird A J, Charman D J, Evans C D, Gallego-Sala A V, Gill P J, Hughes P D M, Morris P J and Swindles G T 2019 Misinterpreting carbon accumulation rates in records from near-surface peat *Sci. Rep.* **9** 1–8
- Yu Z, Beilman D W, Frolking S, MacDonald G M, Roulet N T, Camill P and Charman D J 2011 Peatlands and their role in the global carbon cycle *Eos, Trans. Am. Geophys. Union* **92** 97–8
- Zhang H, Gallego-Sala A V, Amesbury M J, Charman D J, Piilo S R and Väliranta M M 2018a Inconsistent response of Arctic permafrost peatland carbon accumulation to warm climate phases *Global Biogeochem. Cycles* **32** 1605–20
- Zhang H, Piilo S R, Amesbury M J, Charman D J, Gallego-Sala A V and Väliranta M M 2018b The role of climate change in regulating Arctic permafrost peatland hydrological and vegetation change over the last millennium *Quat. Sci. Rev.* **182** 121–30
- Zhang H, Väliranta M, Piilo S, Amesbury M J, Aquino-López M A, Roland T P, Salminen-Paatero S, Paatero J, Lohila A and Tuittila E 2020 Decreased carbon accumulation feedback driven by climate-induced drying of two southern boreal bogs over recent centuries *Glob. Change Biol.* **26** 2435–48
- Zuidhoff F S and Kolstrup E 2005 Palsa development and associated vegetation in Northern Sweden palsa development and associated vegetation in Northern Sweden *Arctic Antarct. Alp. Res.* **37** 49–60