



**QUEEN'S  
UNIVERSITY  
BELFAST**

## Combination of Natural Compounds With Novel Non-thermal Technologies for Poultry Products: A Review

Barroug, S., Chaple, S., & Bourke, P. (2021). Combination of Natural Compounds With Novel Non-thermal Technologies for Poultry Products: A Review. *Frontiers in Nutrition*, 8, Article 628723. Advance online publication. <https://doi.org/10.3389/fnut.2021.628723>

**Published in:**  
Frontiers in Nutrition

**Document Version:**  
Publisher's PDF, also known as Version of record

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

### **Publisher rights**

Copyright 2021 the authors.

This is an open access article published under a Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

### **General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

### **Open Access**

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>



# Combination of Natural Compounds With Novel Non-thermal Technologies for Poultry Products: A Review

Soukaina Barroug<sup>1†</sup>, Sonal Chaple<sup>1†</sup> and Paula Bourke<sup>1,2\*</sup>

<sup>1</sup> School of Biosystems and Food Engineering, University College Dublin, Dublin, Ireland, <sup>2</sup> School of Biological Sciences, Institute Global Food Security, The Queens University Belfast, Belfast, United Kingdom

## OPEN ACCESS

### Edited by:

Maria João Fraqueza,  
University of Lisbon, Portugal

### Reviewed by:

Rosario Ramirez Bernabé,  
Center for Scientific and Technological  
Research of Extremadura  
(CICYTEX), Spain  
Isabel Odrizola-Serrano,  
Universitat de Lleida, Spain

### \*Correspondence:

Paula Bourke  
paula.bourke@ucd.ie

<sup>†</sup>These authors have contributed  
equally to this work

### Specialty section:

This article was submitted to  
Nutrition and Food Science  
Technology,  
a section of the journal  
Frontiers in Nutrition

**Received:** 12 November 2020

**Accepted:** 18 March 2021

**Published:** 08 June 2021

### Citation:


Barroug S, Chaple S and Bourke P  
(2021) Combination of Natural  
Compounds With Novel Non-thermal  
Technologies for Poultry Products: A  
Review. *Front. Nutr.* 8:628723.  
doi: 10.3389/fnut.2021.628723

Ensuring safe, fresh, and healthy food across the shelf life of a commodity is an ongoing challenge, with the driver to minimize chemical additives and their residues in the food processing chain. High-value fresh protein products such as poultry meat are very susceptible to spoilage due to oxidation and bacterial contamination. The combination of non-thermal processing interventions with nature-based alternatives is emerging as a useful tool for potential adoption for safe poultry meat products. Natural compounds are produced by living organisms that are extracted from nature and can be used as antioxidant, antimicrobial, and bioactive agents and are often employed for other existing purposes in food systems. Non-thermal technology interventions such as high-pressure processing, pulsed electric field, ultrasound, irradiation, and cold plasma technology are gaining increasing importance due to the advantages of retaining low temperatures, nutrition profiles, and short treatment times. The non-thermal unit process can act as an initial obstacle promoting the reduction of microflora, while natural compounds can provide an active obstacle either in addition to processing or during storage time to maintain quality and inhibit and control growth of residual contaminants. This review presents the application of natural compounds along with emerging non-thermal technologies to address risks in fresh poultry meat.

**Keywords:** poultry, non-thermal processing, natural compounds, *Campylobacter*, essential oils

## INTRODUCTION

Fresh poultry meat and poultry products are highly perishable products but also have high potential as sources of human infection due to the presence and persistence of key pathogens in the poultry process chain. Outbreaks of foodborne illnesses in association with poultry products are one of the primary causes of outbreaks in the US and the EU. Among the reported numbers and notification rates of confirmed zoonoses in the EU in 2018, the top 5 are *campylobacteriosis* (246,158), salmonellosis (91,662), yersiniosis (6,823), and Shiga toxin-producing *Escherichia coli* (STEC) infections (6,073) (1, 2). Also, according to the Centers for Disease Control and Prevention (3), around 11, 2, and 1% of foodborne outbreaks are associated with chicken, turkey, and other poultry products, respectively, while turkey (609 illnesses) had the most outbreak-associated illnesses followed by chicken (487 illnesses). The pathogen *Campylobacter* caused up to 1.5 million illnesses each year in the US (4); thus, a focus on comprehensive and emerging methods for safety control in poultry processing is warranted.




In order to provide safer poultry products, the food industry has developed and implemented preventive measures based on the Hazard Analysis and Critical Control Points (HACCP) and food safety management systems in combination with technological interventions, such as sanitization processes, refrigeration, and modified atmosphere packaging that can control identified potential microbial hazards during food processing and storage. However, according to the policy and consumer demand requiring sustainable, safe, and high-quality minimally processed foods, the food industry seeks alternative approaches to extend safe shelf life; such as irradiation, high-pressure processing (HPP), and natural green chemicals including bio-preservation or intelligent packaging. **Figure 1** (6) illustrates the consumers' perception of 10 different meat decontamination processes based on how natural and invasive the process is considered.

The intensity of non-thermal processing treatment is vital for product safety and inactivation of pathogenic microbes. Han et al. (7) reported reactivation of sublethally injured microorganisms in a favorable environment during storage in meat and meat products. Spore-forming organisms such as *Clostridium botulinum* are resistant to HPP (8). The increase in a non-thermal process intensity to deal with recalcitrant microbiological issues may adversely affect foods' sensory properties (9). HPP can also alter the structure of polysaccharides and proteins, leading to textural changes in terms of hardness (10). Meat tenderness is the essential attribute that drives its consumer acceptability (11). Post slaughter, meat tenderization results from protease

proteolysis of myofibrillar and cytoskeletal proteins as well as from the degradation of connective tissue substances, in particular collagen (12). Applying power ultrasound can produce free radicals, which imparts effective microbiological control, but which can also impact the product quality of high-fat foods due to oxidation (13). Similarly, irradiation can also cause undesirable organoleptic changes to high-fat foods and can induce color, odor, and taste effect on fresh meat products (14).

Thus, to overcome these shortcomings, there is potential to optimize non-thermal technology effects in combination with incorporation of natural compounds. Careful consideration of the mechanisms of action of individual non-thermal approaches may reveal what combinations can be successful across a range of food systems. For example, pulsed electric field (PEF) can cause cell membrane damage of microorganisms, enhancing sensitivity to antimicrobial agents like nisin (15). Similarly, application of naturally occurring antioxidant compounds such as rosemary extract, blueberry, or ascorbic acid can be used as an alternative for synthetic preservatives like butylated hydroxyanisole, butylated hydroxytoluene, and tertiary butylhydroquinone (16). Synergistic inactivation of microorganisms using non-thermal technology with natural compounds can be a promising way to increase the safety of poultry products while diminishing undesirable effects on some food characteristics. This review summarizes the key risks and published findings where non-thermal technologies have been combined with natural compounds and comments on the effects on the microbiological and physicochemical characteristics.



## CURRENT INTERVENTION PROCESS TECHNOLOGIES ENSURING POULTRY MICROBIOLOGICAL SAFETY

There are many approaches applied in poultry processing to maintain fresh poultry meat safety, which are generally classed as biological, physical, or chemical interventions (6). Electrolyzed water, hot water combined with rapid cooling, chilling and freezing (cold air and ice water), activated oxygen, and organic acids (lactic acid, oxidizing acids, and peroxyacetic acid) are examples of these decontamination processes currently used in poultry processing plants (17). The overview of the poultry processing plant is given in **Figure 2**. The proposed application of antimicrobial interventions in poultry processing can be implemented at carcass washing, scalding, defeathering, chilling/cooling, or packaging. The US Department of Agriculture (USDA) recommends the use of hot water above 74°C for sanitizing effect on carcasses.

High pressure, steam, and steam vacuum, as well as hot and cold water are some physical treatments commonly applied to meat carcass surface decontamination. Physical decontamination approaches currently used for chicken are steam and immersion in hot water (18). Due to the high temperature (100°C), microorganisms including natural microflora should be inactivated on the surface of the product within limited exposure times. Some limitations with steam usage are the deterioration of sensory characteristics attributed to changes of the color, and

samples that look partially cooked, with shrunken skin (19). In steam vacuum processes, steam or hot water is sprayed on carcasses followed by vacuum treatment. This process is an effective method for spot decontamination at the slaughtering unit before the final chilling. The solutions of organic acids are frequently used in the chemical rinse to decontaminate the entire surface of carcasses. The most commonly used organic acids are acetic and lactic acids (20).

Peracetic acid, disodium phosphate, hexadecylpyridinium chloride, and sodium hypochlorite are the most utilized sanitizers during poultry processing (scalding and pre/post chilling) in poultry plants. Treatments with these antimicrobials can be online or off-line for reprocessing at different stages and temperatures and for different treatment times (21). Peracetic acid is an artificial disinfectant retaining good efficacy against poultry meat-related pathogens: *Salmonella* and *Campylobacter* (22). Sodium hypochlorite is commonly applied in water used for chilling/cleaning by spraying and/or immersion to reduce microbial load; however, the efficiency drops down significantly because of the interaction between organic matter in the meat and chlorine (23). Chlorine usage is prohibited in some countries including Germany, Denmark, and Belgium because of the potential interactions with organic matter within poultry carcasses, which can generate harmful chlorinated compounds (halo acetic acids, trihalomethanes, and chloramines) reported to be mutagenic and carcinogenic (23).

However, chlorine dioxide has also been used for sterilization, sanitization, and as disinfectant depending on its form (liquid or gas). Acidified sodium chlorite is an oxidative antimicrobial agent with a large activity spectrum (yeast, fungi, protozoa, viruses, pathogens, and molds). It is authorized by Food and Drug Administration (FDA) and Environmental Protection Agency (EPA) to be used in poultry (24). One of the best alternatives to classical sodium hypochlorite is electrolyzed water due to its low cost, high sterilization effect, and non-harmful effect (23). Specific organic acids can also be effective in terms of microbial inactivation and stability in the presence of organic material, namely, citric acid, lactic acid, succinic acid, and acetic acid. Nevertheless, some limitations are associated with the usage of such organic acids such as off-colors, odors, and flavors in addition to material corrosion (24).

## MICROBIOLOGICAL CONTAMINANTS OF CONCERN IN THE POULTRY SECTOR

### Microbial Spoilage

Owing to microbial spoilage, millions of pounds per annum of fresh poultry meat products are lost (25, 26). Poultry meat spoilage has dramatic effects by limiting shelf life (27) and negatively affecting the economy (28). The deterioration may result from quality and sensory damage to change in texture, odor, color, taste (29), and slime formation (27, 30). These changes are induced by enzymatic reactions, lipid oxidation (30), and action of the natural microflora within the poultry meat (28). Odor quality is affected by the production of volatile catabolites, while the deterioration in color happens throughout storage that

is frequently related to biochemical reactions (between meat pigments, oxygen, and volatile microbial catabolites) and higher meat pH (30).

Several bacteria may be involved in poultry meat spoilage including coliforms, Enterobacteriaceae, *Brochothrix thermosphacta*, *Pseudomonas*, *Aeromonas* sp., *Serratia*, lactic acid bacteria as *Lactobacillus oligofermentans*, *Leuconostoc gelidum* subsp. *gasicomitatum* (29), *Lactococcus*, *Vagococcus*, and *Carnobacterium* (27). The dominant spoilage bacteria is *Pseudomonas* spp. (26) due to its ability to assimilate, penetrate, and metabolize many meat compounds that other bacteria cannot use (30). When the total viable count achieves or exceeds 7 Log CFU/g, spoilage is deemed to ensue (29, 30).

## Pathogens

*Campylobacter*, *Salmonella*, *Listeria monocytogenes*, and *Staphylococcus aureus* are some of the primary pathogens naturally present or contaminants of poultry meat products (31, 32). *Clostridium perfringens*, *Listeria innocua*, and *Aeromonas* spp. were also identified in poultry meat (29). Recently, *Helicobacter pullorum* and *Acrobacter* gained considerable attention as relevant agents of poultry meat infections (33). *Staphylococcus saprophyticus* is another relevant agent of poultry product contamination, which varies from *S. aureus* because of its virulence factors and genetic profile (34). *C. perfringens* is widespread in the tract of birds (35), is responsible for enteric diseases in poultry (36), and is known for potential to generate extracellular enzymes and large numbers of toxins (37). At a certain microbial load, toxin production is upregulated (38). The enterotoxins delivered are associated with human gastrointestinal illnesses—enterotoxemia where toxins induce organ damage upon entering the circulation (36, 38).

## *Campylobacter* spp.

Poultry is the natural host/reservoir of *Campylobacter* spp. (39), which is present in the intestinal tract of birds, skin, and feathers (40). *Campylobacter jejuni* and *Campylobacter coli* are the main zoonotic enteropathogen causative serovars of human campylobacteriosis (41) and with high prevalence (42). *C. jejuni* is dominant by comparison with *C. coli* (39). At each point of the production chain, the proportions of these two isolates may be reversed by passing from one stage to the other. This could be attributable to the resistance or the susceptibility to a particular isolation technique introduced during the test of the collected samples and/or the feed withdrawal (41). They are microaerophilic and thermophilic (43) and require specific environmental conditions to develop (44).

It was reported that for 25.7% of chicken broiler carcasses tested, both liver (surface and internal tissue) and ceca were *Campylobacter* positive. However, for 83% (58/70) of carcasses tested, *Campylobacter* was isolated at least once in one of these compartments (42), and it is of note that the study pointed out that dissimilar subtypes of *Campylobacter* could simultaneously contaminate the same broiler carcasses.

## *Salmonella* spp.

*Salmonella* spp. is a facultative anaerobic Gram-negative genus (44), belonging to Enterobacteriaceae family. It is motile (except for *Salmonella enterica* Gallinarum and Pullorum) (45). It grows at optimum environmental conditions of pH 6.5–7 and temperature around 37°C (45). It is ubiquitous and able to survive in water for several months and in a dry environment for up to 2 weeks (46).

*Salmonella* is one of the main causative agents of foodborne disease globally (47). Animal-based foods such as beef, poultry, and pork are the major sources of salmonellosis (48), where human salmonellosis is mostly due to consumption of poultry products (44). *Salmonella* can persist throughout the processing chain from the farm to the fork (47). Enteritidis, Newport, and Typhimurium are serotypes commonly identified (49). The main genes encoding for the virulence are in both virulence-associated plasmid and pathogenicity islands. These genes are involved in internalization, epithelial cell invasion, survival, and replication, have a significant role in systemic infection (50), and may present a target for intervention technologies design.

## NATURAL COMPOUND-BASED INTERVENTIONS SIGNIFICANT TO POULTRY SECTOR

### Spices and Herbs

These natural compounds are used in foods for flavoring and preservation and as additives but also for medicinal and therapeutic goals (anti-oxidative, immune modulators, anti-inflammatory, and antimutagenic). Their utilization in foods can have beneficial effects for shelf life extension as well as improvement of the organoleptic characteristics (51). The antioxidant potential of herbs or spices can prevent or decrease lipid oxidation (52) attributed to the action of phenolic compounds (53).

Clove and rosemary are two aromatic spices known for their antimicrobial and antioxidant potential. Antimicrobial activities of rosemary and clove extracts were tested separately or in combination against pathogenic and spoilage bacteria related to meat, namely, *L. monocytogenes*, *E. coli*, *Pseudomonas fluorescens*, and *Lactobacillus sake*, as well as against native microflora in poultry meat samples, where the combination of both provided enhanced microbial inactivation potential. The combination of the herb and spice also maintained or improved the sensory characteristics of fresh meat, reduced lipid oxidation, and extended shelf life up to 15 days (53).

Curcumin is a US FDA-approved safe plant pigment used for cooking, reputed for its health benefits. It acts as antioxidant, anti-inflammatory, and antiproliferative, but drawbacks for use in foods are related to the color, taste, and quality alterations (54). Corrêa et al. (54) analyzed the antimicrobial effect of two photonic approaches, ultraviolet (UV)-C and curcumin-mediated photodynamic inactivation (PDI), for control of *E. coli* and *S. aureus* inoculated on chicken breast cubes. Limitations were associated with these treatments, as the UV-C light or the curcumin-mediated PDI treatment with emission at 450 nm were



not absorbed in all areas and did not significantly penetrate the chicken meat surface, with 1–2 log<sub>10</sub> CFU/ml reductions observed (54).

## Essential Oils

Essential oils (EOs) are aromatic secondary metabolites and concentrated plant extracts. They can be obtained by steam distillation, expression or supercritical extraction with carbon dioxide from different parts of the plant, for example, bark, flower, fruits, seeds, leaves, or roots. Many EOs have found application in poultry feed as an alternative to antibiotics due to antioxidant, antiseptic, and insect repellent properties as well as immune-modulatory effects (55). EOs are chemically diverse compounds; hence, their antimicrobial activity varies from compound to compound. However, due to their hydrophobic nature, they are likely to enter cell membranes of microbes or eukaryotes (56). The main limitations of using EOs in food products are the strong flavor imparted on foods (56), the heat-labile nature, and volatile characteristics (57).

Rosemary extract is well-known for its antimicrobial activity, which is related to its phenolic composition (e.g., rosmarinic and carnistic acids). Inactivation of cellular enzymes was seen to result from the effect of phenolic compounds (16). Treatment of *Salmonella typhimurium* with thyme EO caused a rise in the electrical conductivity, which appears to result from the destruction of the cell membrane as well as electrolyte leakage. Additionally, quantitative analysis reported a significant drop in the protein contents, DNA, and ATP by 55.42, 54.03, and 52.64%, respectively, when compared to the control (57). Likewise, EOs (lemon oil, lemon grass oil, lime oil, garlic oil, onion oil, pimento berry oil, oregano oil, thyme oil, and rosemary oil) had higher antimicrobial potential against four *Campylobacter* strains when compared to organic acid (ascorbic acid, citric acid, and lactic acid). Oregano EOs specifically displayed the higher inactivation potential against *C. jejuni*, where the minimal inhibitory concentration was equivalent to 62.5 ppm (18).

The application of 0.1% oregano EO was effective for extending the shelf life up to 5–6 days for fresh chicken breast meat before packaging (58). The authors pipetted 0.1% of oregano EO in the low-density polyethylene (LDPE)/polyamide (PA)/LDPE barrier pouches, which was later subjected to either air or modified atmosphere packaging (MAP). The lipid peroxidation and deterioration of sarcoplasmic proteins were controlled to extend the shelf life of chicken breast up to 2 weeks at 4°C with the application of 0.5% of both thyme and *Melissa officinalis* balm EOs. These EOs were applied on the chicken breast slices by dipping method for 15 min. The results highlighted that thyme (0.5%) was more effective in inhibiting the growth of *E. coli*, whereas balm (0.5%) was more effective on the *Salmonella* spp. (59). The combined effect of ethylenediaminetetraacetic acid (EDTA) (1.5% w/w) lysozyme (1.5% w/w), rosemary oil (0.2% v/w), and oregano oil (0.2% v/w) was effective on extending the shelf life of vacuum-packed semi-cooked coated chicken filets stored at 4°C (60). EDTA and lysozyme were applied by spraying technique on the surface of the chicken surface, while rosemary and oregano oil

were pipetted in the LDPE/PA/LDPE pouch barriers containing chicken samples.

## Organic Acids

Organic acids are naturally occurring compounds present in many foods and can be produced during the fermentation process. They are added in foods as acidulants, preservatives, or flavorants. The commonly used organic acids are lactic, acetic, malic, and ascorbic acid, etc. The mechanism of inactivation of these acids is through lowering of pH, pKa value along with penetration of undissociated compounds through the cell membrane and its dissociation inside the cell, thus affecting the bacterial membrane (61, 62). In poultry products, the salts of organic acids such as potassium or sodium lactate and sodium diacetate are used to inactivate *L. monocytogenes*, and buffered citrate is used to enhance flavor (63). The maximum level for potassium and sodium lactate is 4.8% by weight of total formulation in various meat and poultry products. For sodium diacetate, the maximum permitted level is 0.25% by weight of total formulation when used as either antimicrobial agent or flavoring agent (64).

## NON-THERMAL TECHNOLOGIES AND THEIR COMBINATIONS WITH NATURAL COMPOUNDS IN POULTRY PROCESSING


Non-thermal technologies, such as HPP, PEF, ultrasound, UV, irradiation, and cold atmospheric plasma (CAP) can retain nutritional as well as sensory properties of food in shorter treatment times and low operational temperatures (65). Extensive research on the application of various non-thermal technologies to poultry meat has been conducted in recent years. The effect of non-thermal technologies on microbial and physicochemical properties will be discussed in this section. The incorporation of natural compounds with non-thermal technologies can give an additional hurdle enhancing antimicrobial efficacy. Additionally, this will help adjust the processing conditions at lower intensity, giving improved physicochemical properties (66). Few studies have focused on combining natural compounds with the non-thermal treatment as summarized in **Table 1** illustrating its impact on both physicochemical and microbial properties. A pictorial representation of governing processing parameters, mechanism for microbial decontamination, and key physicochemical parameters to take into consideration while application of the non-thermal technology in the poultry products is depicted in **Figure 3**, and it will be further discussed in coming sections.

## High-Pressure Processing

HPP is a non-thermal technology for the sterilization and preservation of food products in which the product is subjected to high pressure (300–600 MPa) with or without the combination of heat. With the application of HPP, covalent bonds in food matrices are not broken and the effect on the food characteristics is minimal. HPP is based on Le Chatelier's principle and the

**TABLE 1** | Summary of combination trials of different non-thermal technologies with natural compounds on poultry products.

Non-thermal technology	Natural compounds	Poultry products	Chemical observations	Microbiological observations	References
CAP	Rosemary extract	Poultry ground meats	NA	<ul style="list-style-type: none"> <li>- Reduction of the bacterial functional diversity</li> <li>- The lowest Maximum Population Size (54.65, 95% confidence interval [CI95%] ranges, 54.03–55.16) and slowest growth rate (hour) (0.03258, CI95% ranges, 0.0179–0.04726) in day 0</li> <li>- At day 5 of storage at 4°C, the maximum population sizes of treated samples were statistically not significant comparing to day 0</li> </ul>	(67)
	Rosemary extract	Ground chicken patties	<ul style="list-style-type: none"> <li>- Lower pH values for rosemary samples</li> <li>- a* value was significantly affected by rosemary addition</li> <li>- Addition of rosemary extract prevented lipid oxidation for CAP</li> </ul>	Rosemary extract significantly reduced the total plate counts with and without cold plasma treatment	(16)
	Thyme oil (TO)/ Silk fibroin (SF) nanofiber	Chicken and duck meat	<ul style="list-style-type: none"> <li>- Thyme oil release was enhanced due to surface modification of SF by plasma treatment</li> <li>- Higher overall acceptability of chicken meat treated with plasma treatment and combination of TO/SF nanofiber</li> </ul>	- The population of <i>Salmonella</i> Typhimurium on treated chicken meat reached 1.15 and 1.96 log CFU/g when stored at respectively 4 and 25°C for 7 days after been wrapped with plasma-Thymol oil-Silk fibroin nanofibers. Identical effects seen with the duck meat treated with the same process	(57)
	Essential oils: <i>Crocus sativus</i> L., <i>Allium sativum</i> L., and <i>Zataria multiflora</i> Boiss	Breast chicken fillets	Overall acceptability and no undesirable impacts on both flavour and odour	<ul style="list-style-type: none"> <li>- Associating CP and essential oils treatments of breast chicken fillet infected by <i>S. aureus</i> and <i>E. coli</i> lead to significant microbial reductions by at least 3–4 logCFU/g.</li> <li>- A synergetic effect due to the combination of three different EOs (<i>Crocus sativus</i> L., <i>Allium sativum</i> L., and <i>Zataria multiflora</i> Boiss.) and CP treatment reaching microbial reductions to great extent.</li> <li>- After 14 days storage, 2–2.7 logCFU/g microbial inactivation reported comparing to 4.9 logCFU/g of samples treated with only EOs</li> </ul>	(68)
HPP	Articoat-DLP (lactic acid, acetic acid and sodium diacetate-active compounds)	Chicken breast fillets	<ul style="list-style-type: none"> <li>- Significant increase in L-value</li> <li>- TBARS value remained same during storage</li> <li>- Increase in pH due to HPP</li> </ul>	<ul style="list-style-type: none"> <li>- <i>Pseudomonas</i> spp., <i>B. thermosphacta</i>, <i>coliforms</i>, <i>E. coli</i> inactivated below detection limit</li> <li>- LAB reformed after 7 days storage time</li> </ul>	(69)
	Carvacrol	Turkey breast ham	<ul style="list-style-type: none"> <li>- Higher TBARS value for pressurised samples</li> <li>- Carvacrol addition decreased TBARS value of samples</li> </ul>	<ul style="list-style-type: none"> <li>- Carvacrol+HPP extend the lag phase for <i>Listeria</i></li> <li>- Reduced the growth rate of LAB spoilage groups</li> </ul>	(70)
	Thymol	Ground chicken	NA	- addition of thymol impacted the HPP sensitivity for iPEC O157:H7 and UPEC	(71)
PEF	Oregano essential oils	Raw chicken	NA	<ul style="list-style-type: none"> <li>- No significant inhibition of <i>C. jejuni</i> if only treatment with PEF (0.25–1 kV/cm) applied.</li> <li>- Sequential treatment of PEF with immersion for 20 min in oregano essential oil (15.625 ppm) were effective against <i>C. jejuni</i> 1146 DF with maximum reduction of 1.5 log CFU/g</li> </ul>	(18)
Ultrasound	Lactic acid	Broiler drumstick skin	NA	Ultra-sonication alone and with 1% lactic acid did not significantly affect aerobic plate count	(72)
	Lactic acid	Poultry skin	NA	<ul style="list-style-type: none"> <li>- <i>Pseudomonas</i> was most sensitive to lactic acid than other gram-negative bacteria</li> <li>- Degree of reduction of gram-negative bacteria was dependent on treatment time and liquid medium (water or lactic acid)</li> </ul>	(73)
	Oregano essential oil	Chicken breast	NA	0.3% oregano oil and ultrasound showed better inactivation of lactic acid bacteria, mesophiles and anaerobic bacteria at day 0 and during 21 days of storage	(74)



isostatic principle. Le Chatelier's principle states that "if a change in conditions is applied on a system in equilibrium, then the system will try to counteract that change and restore the equilibrium." The isostatic principle states that food products are compressed by uniform pressure from every direction and then returned to their original shape when pressure is released. HPP is currently used for liquid and high-moisture solid products (75). The first commercialized HPP meat products available are sliced cooked ham and precooked meals containing poultry, pork, chorizo, and different sausages in Spanish market (76, 77). Further details about commercialized meat product of HPP are detailed elsewhere (78).

### Effect of High-Pressure Processing on Microbial Decontamination of Poultry Products

Tracz et al. (2015) investigated the potential of HPP to destroy a mixed culture of three stains of *C. jejuni* inoculated in chicken breast under different pressure conditions (200, 300, and 400 MPa) and treatment times (5, 10, and 15 min), where D values were lowest at the highest pressure applied (79). According to the pressure applied, the temperature varies from 0 to 10°C. When the lowest pressure (200 MPa) was applied, *C. jejuni* exposed resistance and no significant reduction was achieved regardless of the duration of HPP treatment. Gram-negative bacteria are generally more susceptible to pressure compared



to Gram-positive bacteria (79). Sheen et al. (2015) reported also that the inactivation of *Salmonella* spp. in ground chicken was dependent on both treatment time and pressure level applied (80). It was highlighted that even at high pressure (550 MPa), while the highest temperature reached was 28°C, *Salmonella* recovered and resuscitated over storage at 10°C to achieve ~6 Log CFU/g at day 9 of storage. Therefore, despite the mechanisms noted of surface structure damage or disintegration, internal cell compound disappearance, and appearance of internal voids in the cells, some cells survived the HPP 15-min treatment (80), pointing to a need for combination approaches. Argyri et al. (2018) HPP treatment at 500 MPa for 10 min at 18–20°C resulted in a significant reduction, below detection limit, of both the native microbiota of chicken and a cocktail culture of three different strains of *Salmonella* (31). Furthermore, *Salmonella enteritidis* inoculated on chicken at different initial concentration levels stayed below or just at detection limits during the storage at 4°C over 18 days (30). Working with a cocktail of *Listeria monocytogenes*, also Argyri et al. (2019) perceived the capability of HPP in maintaining safety and extending the shelf life of chicken (30, 32). Xu et al. (2020) reported D<sub>10</sub> values for multi-isolated cocktails of extraintestinal pathogenic *E. coli* (ExPEC) to HPP (400 MPa, 0–25 min) on ground chicken, where 3.26 min was the average and the highest temperature value reached during the treatment was 25°C (81). Increasing the pressure to 600 MPa provided more than 6 log reduction within 3 min with no bacterial recovery after 4 min (81). The inactivation effect of HPP on two different strains of *E. coli* on ground chicken was assessed while the temperature remained under 40°C, where a significant resistance of uropathogenic *E. coli* (UPEC) by comparison with intestinal pathogenic *E. coli* (iPEC) O157:H7 at 450 and 500 MPa was reported (71). Liu et al. (2012) stated that *C. jejuni* HCCJ2316 exhibited high resistance to pressure (2.8 Log CFU/g reduction), whereas the others were more susceptible to treatment and achieved 5 Log CFU/g reduction (82). However, the microbial recovery upon pressure treatment of *C. jejuni* is iron-dependent (82).

### Effect of High-Pressure Processing on Physicochemical Properties of Poultry Products

Lipid oxidation is one of the major causes of deterioration of meat during storage. The chicken meat contains a higher amount of unsaturated fatty acids compared to other animal meats, which makes it more susceptible to lipid oxidation. The most common method to determine the lipid oxidation is Thiobarbituric acid reactive substances (TBARS) analysis (24). The lipid oxidation was particularly affected by working pressure; for low pressure (400 MPa and less), significantly less change in TBARS value was reported, while high pressure (500 MPa) have a higher impact on TBARS value (83). Similar effect was observed when chicken breast filets was treated with 450 and 600 MPa, while no significant change in TBARS value was observed at 300 MPa (84). The pressure of 800 MPa has the most detrimental on TBARS value (85).

### Combined Effect of High-Pressure Processing and Natural Compounds on Poultry Products

A synergistic effect was recorded using nisin (200 ppm) and HPP (450 MPa) at 20°C, enhancing the microbial reduction of mechanically recovered poultry meat, specifically, the inactivation of both aerobic mesophile and psychrotroph populations was greater by comparison with HPP treatment on its own (86). Other researchers outlined the strong synergistic effect of combining hydrostatic pressure treatment (250 MPa for 30 min at 25°C) and 1% food additive (citric acid, nisin, and wasabi extract) in completely reducing the microbial concentration of *S. enteritidis* to undetectable levels (87). Combining HPP (300–400 MPa) with thymol (100–200 ppm) provided a large inactivation effect on separate cocktails of iPEC O157:H7 (0.94–5.16 Log CFU/g) and UPEC (0.41–4.66 Log CFU/g) in ground chicken samples (71).

### Pulsed Electric Field

PEF uses short pulses of high voltage (5–80 kV) for microbial inactivation. The food is placed between two electrodes, and an external electric field is applied, which induces the movement of ions along the direction of lines of force of the applied electric field inside as well as outside the cells. This causes the accumulation of ions on the membranes, causing polarization of the cell, which results in thickness reduction of the membranes due to the forces of attraction between oppositely charged ions on either side of the membrane (88). Because of the potential for cell membrane permeabilization, PEF is a promising technology to modify several qualities of meat, such as color, texture, and water-holding capacity, and enhance mass transfer during curing and brining. However, applications to date can be limited in solid products due to conductivity requirements (89).

### Effect of Pulsed Electric Field on Microbial Decontamination of Poultry Products

The cell membrane is commonly referred to as the only target of PEF contributing to bacterial cell death (90). Treatments with PEF display reversible or irreversible damages on the cell membrane by disorganizing the structure, which yields the breakdown of the semipermeable barrier due to formation of pores in the membrane (18), leading to irreversible electro-permeabilization of the cell membrane; however, recovery can occur in optimal conditions (91). Process parameters of pulse frequency and strength of the electric field of PEF have been demonstrated to affect the microbial inactivation (92).

Reduction in population densities of *S. enteritidis* and *S. typhimurium* strains suspended in citrate-phosphate buffer was greater with increasing both treatment time and electric field above 9 kV/cm (93). It was demonstrated by Clemente et al. (18) that the treatment of chicken thighs with PEF did not result in any significant reduction of *C. jejuni*. PEF was not sufficient to reduce cell concentration of *S. enteritidis*, *E. coli*, and *C. jejuni* on raw chicken (92). However, this non-thermal technology is suggested to be suitable for treating process waters used in poultry processing as well as for poultry scald (92).

## Effect of Pulsed Electric Field on Physicochemical Properties of Poultry Products

Several studies have reported the ability of PEF treatment to modify sensory characteristics, texture, and water-holding property of the meat products, further improving the mass transfer properties (94–96). Meat is considered an excellent source of minerals, such as zinc, iron, phosphorus, and calcium (97). PEF induces irreversible electroporation in meat and affects cellular permeability and mass transfer. Studies suggest that PEF-applied products have changes in mineral content when compared to control. Khan et al. (98) examined the effect of low (2.5 kV, 200 Hz) and high PEF (10 kV, 200 Hz) on four nutritionally important minerals (P, K, Fe, and Zn) of raw and cooked chicken breast. For raw chicken, non-significant changes in mineral content were noted; however, with cooking, a decrease in P, K, and Zn was observed and the concentration of Fe was not affected by treatment or cooking. In another study conducted by Khan et al. (99), the authors found higher concentrations of Ni and Cu for both low (2.5 kV, 200 Hz) and high PEF (10 kV, 200 Hz) than control. Thus, it is vital to study the migration of minerals into meat products due to PEF treatment and should be checked under regulatory limits.

## Combined Effect of Pulsed Electric Field and Natural Compounds on Poultry Products

Recent work revealed that chicken oyster thigh artificially contaminated by *C. jejuni* 1,146 DF (final concentration  $4.41 \pm 0.20 \log_{10}$  CFU/g) and treated with only PEF (0.25–1 kV/cm) did not demonstrate any significant inhibition potential. However, sequential treatment of PEF (1 kV/cm) and immersion in buffer with oregano EO (15.625 ppm) for 20 min resulted in a significant reduction close to  $1.5 \log_{10}$  CFU/g (18).

## Ultraviolet

UV light is electromagnetic radiation with wavelength from 10 to 400 nm. UV lights fall in the range between visible light and X-rays. To control surface contaminations on food products, UV-C light has received US FDA approval (Approval-2010). High-intensity pulsed UV light has been approved by FDA up to  $12 \text{ J/cm}^2$  (100). UV-C light can be used in Europe; however, in Germany, the use is limited to water, fruit, vegetables, and stored hard cheese (101). UV-C has a wavelength range of 220–300 nm (102) and is known for its antimicrobial effect (103), where the specific mechanisms of action include targeting of the nucleic acids (DNA, RNA) within the bacterial cell and generation of pyrimidine dimers (104). This latter results in the bonding of two adjacent pyrimidine bases, provoking obstruction of transcription and translation, respectively, and suspending vital cellular functions (102, 105, 106).

## Effect of UV on Microbial Decontamination of Poultry Products

There are some limitations of using UV in poultry processing: UV-C light is not absorbed and cannot penetrate the chicken meat surface, which may affect microbial reduction. The antimicrobial efficacy of pulsed UV and UV-C has limitations also in terms of product density and treatment time (104,

107). The bactericidal effect of UV-C irradiation against *C. jejuni*, *L. monocytogenes*, and *S. typhimurium* on chicken breast was dose-dependent, where treatment at  $5 \text{ kJ/m}^2$  reduced *L. monocytogenes*, *C. jejuni*, and *S. typhimurium*, respectively by 1.29, 1.26, and 1.19 log cycles (102). Haughton et al. (105) examined UV effects against *S. enteritidis*, *E. coli*, and *Campylobacter* (*C. jejuni* and *C. coli*) when inoculated in liquid matrix, chicken skin and skinless chicken breast, food contact surfaces, as well as packaging materials. Treatment at a high dose equivalent to  $0.192 \text{ J/cm}^2$  provided complete microbial inactivation of *Campylobacter* strains suspended in a liquid matrix. By contrast, *Salmonella* and *E. coli* were more resistant to the similar UV dose (105). Food surface topography can shield microorganisms and limit UV treatment efficacy (104, 107). Isohanni and Lyhs (103) highlighted that although UV treatment was effective in reducing *C. jejuni* on surface medium by 6.3 log cycles per square centimeter. However, only 0.8 and 0.7 log cycles reduction were achieved on broiler skin and on broiler meat, respectively, with a dose of 32.9 mW/s per square centimeter. UV light seems to work well on smooth surfaces (108). Bacterial multilayer overloading as well as overlapping, and in the presence of cell, organic compounds protect to target bacteria from UV irradiation (104, 107).

## Effect of UV on Physicochemical Properties of Poultry Products

UV light can form off-flavors due to the photochemical effect on the lipid fractions of product or due to absorption of ozone and oxides of nitrogen (101). This leads to the development of lipid peroxidation causing off-flavor. The hexanal aldehyde is a volatile secondary lipid oxidation product, and it is indicative of fatty aldehydes by headspace/gas chromatography–mass spectrometry (GC-MS). McLeod et al. (104) detected an increase in hexanal content of the raw chicken filets treated with  $10.8 \text{ J/cm}^2$  in air, which was noted by the sensory panel as a “sunburnt flavor” giving a low sensory score. Interestingly, when the same chicken samples were cooked, the sensory panel was unable to identify the difference, and it scored fairly with the untreated sample (104).

## Combined Effect of UV and Natural Compounds on Poultry Products

The combination treatment of UV-C light and clove EO was assessed against poultry-related pathogen *S. typhimurium* biofilms, generated on stainless steel coupon surfaces. Treatment with 1.2 mg/ml of clove EO followed with UV-C ( $76.41 \text{ mJ/cm}^2$ ) induced a synergistic effect and resulted in no surviving cells ( $6.8 \log \text{ CFU/cm}^2$ ) embedded within the biofilms. It was demonstrated that the contact with clove EO made the cell easily accessible by UV-C due to the morphological damage occurring: flatter structure (109).

## Ultrasound

Ultrasound as a non-thermal approach applies sound waves with higher frequency (above 20 kHz) than the normal human hearing. The ultrasound frequencies used in the food industry are classified into three categories based on the frequency-power ultrasound: low frequency, high power range (20–100 kHz) and

large-amplitude waves where typical applications are within altering physicochemical properties or structure of foods. For low-intensity ultrasound, in the range of 100 kHz to 1 MHz, chemical reactions are activated, and free radicals can form like hydroxyl ions that can have antimicrobial properties. High-frequency ultrasound is usually used in the food processing and food safety industry. When the cavitation bubble breaks, it forms hydroxyl ions, which can have antimicrobial properties (110).

### Effect of Ultrasound on Microbial Decontamination of Poultry Products

The mechanisms of action are connected to cavitation generation, which eventually disturbs the cell permeability as well as causing thinning (111) and damage on the bacterial membrane (112) and “localized heating” (73) that yields cell inactivation. The cell metabolism is disturbed due to ion penetration of the cell cytoplasm upon permeability disruption caused by pressure gradients of ultrasound. These mechanisms are the result of the collapse of cavitation bubbles during the acoustic cavitation (73). A further utility of ultrasound treatments is the “de-agglomeration of bacterial clusters” (73).

Apparent characteristics of target microorganism type, physiological state, and morphology determine the efficacy of ultrasound. The efficacy is also dependent on the surface of food matrix and temperature (111). Moreover, other parameters interfere with efficacy, such as frequency and sonication treatment time (73). The peptidoglycan in the cell membrane of Gram-positive could be a reason behind the resistance to ultrasound by these bacteria compared to Gram-negative (112), and the susceptibility to ultrasound treatment may vary between strains from the same type. The resistance of cells on plates during *in vitro* experiments to sonication by ultrasound was higher in contrast to the susceptibility of both *Campylobacter* and *Enterobacteriaceae* in raw poultry (112).

It was highlighted that chicken breast subjected to high-intensity ultrasound promoted the growth of mesophilic, psychrophilic, and lactic acid bacteria compared to untreated samples, possibly resulting from the release of nutrients (113). However, it was pointed out that the presence of *E. coli* was lower for samples subjected to longer treatment (30–50 min) compared to non-treated, whereas for *S. aureus*, it significantly decreased after 50 min. The microbial reduction in previously contaminated chicken wings depends on both treatment time (3–6 min) and sonication environment solution where the treatment was in (1% solution of lactic acid or sterile distilled water). The combination of lactic acid and sonication (40 kHz, 2.5 W/cm<sup>2</sup>) had a bactericidal effect on all the bacteria tested and was considered suitable for poultry carcass skin decontamination (73). However, combining ultrasound treatment (37 kHz, 380 W, 5 min) with 70% ethanol induced the highest microbial reduction from chicken skin for three types of attachment by *S. typhimurium* loosely, intermediately, and tightly attached by respectively 2.86, 2.49, and 1.63 log CFU/g (114).

### Effect of Ultrasound on Physicochemical Properties of Poultry Products

Marinating is mostly used to increase meat tenderness, enhance flavor profile, reduce cooking time, and increase the shelf life of

meat. Ultrasound (40 kHz, 22 W/cm<sup>2</sup>) increased the marination efficiency of chicken breast when treated with 15 and 20 min (115). A similar increase in marination efficiency, cooking yield, and tenderization was reported when broiler chicken was treated for 20 min and 18 h marination (91% water, 6% NaCl, 3% sodium tripolyphosphate) (116). A positive influence of ultrasound frequency (25, 45, and 130 kHz) and treatment time (1, 3, 6, 16, and 24 h) on marination efficiency was also reported for chicken breast, giving higher uptake of sodium chloride (117). Ultrasound improved the marination properties of meat by breaking the integrity of muscle cell or by enhancing the enzymatic reactions in cell (11, 111). Thus, ultrasound can be used as an alternative to standard marination techniques used in the industry.

### Combined Effect of Ultrasound and Natural Compounds on Poultry Products

The exposure of broiler drumstick skin to ultrasonic energy in water and submerged in 1% lactic acid did not show consistent effect in terms of reducing aerobic plate counts. The irregular characteristics of broiler skin surface were proposed as the reason behind the lack of microbial reduction by protecting bacteria in the skin crevices and avoidance of the cavitation (72). In contrast, other work showed the decontamination efficacy of sonication (40 kHz) of chicken wing skin in 1% lactic acid aqueous solution, where the reduction of *E. coli*, *Proteus* sp., *Salmonella anatum*, and *P. fluorescens* inoculated on the surface of the chicken skin significantly increased with treatment time rising from 3 to 6 min. Except for *E. coli*, the microbial reduction was higher when sonication was performed in an aqueous solution of lactic acid instead of water. This was explained by the presence of ions penetrating the cytoplasm due to the action of gradient pressure yielding from ultrasound and the presence of free radicals received in sonochemical reactions (73). Combining high-intensity ultrasound with 0.3% oregano EO treatment was the most appropriate combination to achieve the best reduction of lactic acid bacteria (2.30 log<sub>10</sub> CFU ml<sup>-1</sup>), mesophilic populations (3.36 log<sub>10</sub> CFU ml<sup>-1</sup>), and anaerobic bacteria (3.11 log<sub>10</sub> CFU ml<sup>-1</sup>) present in chicken breasts at day 0 of refrigeration. However, the treatment with ultrasound alone was ineffective to control microbial growth during chilled storage, where the release of nutrient was suggested as a reason permitting microbial growth (74).

### Cold Atmospheric Plasma

Plasma is a quasi-neutral ionized gas composed of ions, free electrons, atoms, and molecules in their ground as well as the excited state. Plasma can be generated using any kind of energy, which can ionize the gas, and mostly electric or electromagnetic source are used for generation of plasma species. Plasma can be classified as a thermal plasma or non-thermal plasma. Thermal or non-thermal plasma processes can be designed to be delivered in a format that is cold or near room temperature at the point of application, which is of value for retaining quality and nutritional characteristics while providing efficient bio-decontamination resulting from reactive oxygen or nitrogen species, charged particles, electric field, UV as components of diverse mechanisms of action (118).



## Effect of Cold Plasma on Microbial Decontamination of Poultry Products

It is well-documented that cold plasma (CP) has a large potential for controlling microbial quality, extending shelf life, and avoiding post-processing contamination (25). It not only induces bacterial decontamination but also inactivates a broad spectrum of microorganisms including fungi, viruses, and spores (119). Various plasma compounds have critical interventions in the microbial decontamination process like NO<sub>2</sub>, NO, O, O<sub>3</sub>, OH, H<sub>2</sub>O<sub>2</sub>, UV photons, charged particles, and electric fields (120). Bacterial cell etching, erosion, morphological alteration, nucleic acid damaging, protein oxidation, and loss of cell viability are the mechanisms of CP to retain microbial safety in foods (121). However, different parameters and mechanisms interfere with the gravity of damage occurring. The antimicrobial effects of CP are a function of process: duration of treatment, gas mixture, mode of exposure (direct or indirect), power source intensity, as well as intrinsic characteristics of product: surface topology, nature of samples treated (liquid, solid, or semisolid), and characteristics of the target cell (122).

In-package CP treatment configuration enhanced the microbial safety, avoided postprocess contamination, and retained toxicological safety of ready-to-eat chicken products (123). Using the *Salmonella* mutagenicity assay, no genotoxicity was seen in plasma-treated chicken breast (120). Tulane virus (1.08 ± 0.15 log CFU/cube), indigenous mesophilic bacteria (0.70 ± 0.12 log CFU/cube), and *Salmonella* (1.45 ± 0.05 log CFU/cube) from chicken samples were significantly reduced upon CP treatment (24 kV for 3 min), where increasing voltage (from 22 to 24 kV) and treatment time had a positive impact on the microbiological quality (123). However, the majority of cells showed morphological changes (cell flattened and other distortions) at 2,000 Hz, whereas at 1,000 Hz, only cell clumps appeared, and other cells were hollowed out (124). *S. typhimurium*, *E. coli* O157: H7, and *L. monocytogenes* populations on chicken breast reduced from 5.48, 5.84, and 5.88 log CFU/g, respectively, at 0 min to 2.77, 3.11, and 3.74 log CFU/g at 10-min plasma exposure (120). Other studies highlight the potential of in-package dielectric barrier discharge (DBD) (70 kV) in controlling poultry-related pathogens, namely, *Salmonella* and *Campylobacter*, and inhibiting the growth of spoiling bacteria (psychrophiles) from chicken breast treated and stored (5 days/4°C) (23), where increasing CP treatment time to beyond 60 s improved microbial reduction of psychrophiles, while no significant effect was seen against foodborne pathogens. The *in situ* decontamination potential of plasma-activated water (PAW) against *P. fluorescens* ATCC13525 previously inoculated on chicken skin pieces was associated with the plasma process parameters of plasma discharge frequency and treatment time (124), where the concentration of plasma-generated reactive species of nitrite, nitrate, peroxide, hydroxyl, and ozone increased with the discharge frequency (124).

## Effect of Cold Plasma on Physicochemical Properties of Poultry Products

The reaction of myoglobin with hydrogen peroxide may produce choleglobin-inducing discoloration of meat. An increase in both

L\* and b\* value was observed when chicken breast was treated with flexible thin-layer DBD (123); in contrast, application of DBD applied on chicken breast (110 kV, 60 kHz) showed a decrease in L\* value mainly due to slime formation after 9 days of storage at 4°C (28). However, Zhuang et al. (25) did not report significant changes in L\*, a\*, and b value when chicken breast was treated with 70 kV. The reactive species generated from the plasma can induce lipid peroxidation. Zhuang et al. (2019), Lee et al. (2019), and Moutiq et al. (2020) reported no changes in lipid profile of chicken breast after CP treatment, attributed to plasma reactive species being less damaging on chicken breast as compared to red meat due to variation in fat content (25, 28, 120, 123).

## Combined Effect of Cold Plasma and Natural Compounds on Poultry Products

Yeh et al. (2019) considered the combined treatment effect of rosemary (*Rosmarinus officinalis*) extract (1%) and in-package DBD-CP on poultry ground meat (Table 1) (67). The treatment induced a reduction of the diversity in bacterial communities, and by day 5 of storage at 4°C, the maximum population densities of treated samples were similar to those at day 0 (67). Rosemary extract had a significant effect in reducing the total microbial counts not only of previously plasma-treated chicken patties but also of non-plasma-treated chicken patties (16). Thyme oil/silk fibroin nanofibers treated with CP were proposed as an active packaging approach with antimicrobial potential against *S. typhimurium* inoculated on poultry meat (chicken meat and duck meat). The inhibition potential of thyme EO/silk fibroin nanofibers treated with CP was reported as higher compared to thyme EO/silk fibroin nanofibers, with an increase in the rate of thyme oil released (23.5–25%) upon plasma treatment clearly noted (57). Sahebkar et al. (2020) found that associating CP (10 min at 32 kHz) and EO (in marinade solutions) treatments of breast chicken filet inoculated with *S. aureus* and *E. coli* challenge populations leads to significant microbial reductions by up to 3–4 log CFU/g (68). A synergetic effect was identified by combining three different EOs (*Crocus sativus* L., *Allium sativum* L., and *Zataria multiflora* Boiss.) and CP treatment, where the advantage of combining the EOs with CP was retained after 14 days of storage (68).

## FUTURE CHALLENGES FOR THE APPLICATION OF NON-THERMAL TECHNOLOGIES TO POULTRY PROCESSING

In recent years, there has been intensive research and development of non-thermal process technologies for application in fresh food processing. EU regulations refer to “fresh meat” as meat that has not undergone any preserving process other than chilling, freezing, or quick-freezing, including meat that is vacuum-wrapped or wrapped in a controlled atmosphere (125). The application of a novel non-thermal technology will be subject to these rules, which may on the one hand lead to consumer skepticism toward acceptance of these technologies while on the

other hand limit adoption of processes that can enhance safe and sustainable processing of food resources. To date, HPP and UV (with limitations in some countries) are approved. Thus, combining these approaches with other generally recognized safe approaches including approved natural compounds provides technical options to enhance poultry processing outcomes. The morphological characteristics of poultry products and the skin make the application of non-thermal technology very critical. For example, from a microbial safety perspective, the successful application of UV in poultry processing must consider density, and effects may be limited due to non-absorption and non-penetration of light in the chicken meat surface (104, 107).

While many plant EOs are considered generally safe by FDA, with increasing use, the daily dose intake remains a safety question (126). The adoption of EOs is controlled by a nexus of dosage level, antimicrobial efficacy (127), and the effect of organoleptic characteristics on consumer acceptability (128). The stability, strong smell, volatility, and limited solubility are technical issues to be considered from an efficacy perspective. These should also be considered in tandem with other processing features as the process or environment may stimulate the degradability of these compounds (129).

## REFERENCES

- Center for Disease Control. (CDC) 2015. Surveillance for Foodborne Disease Outbreaks — United States, 2009 – 2015. *MMWR Morb Mortal Wkly Rep.* (2018) 67:2009–15. doi: 10.15585/mmwr.ss6710a1
- Efsa and EcDc. The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2017. *Efsa J.* (2018) 16:5500. doi: 10.2903/j.efsa.2018.5500
- CDC. Surveillance for foodborne disease outbreaks - United States, 2009–2010. *Ann Emerg Med.* (2013) 62:91–3. doi: 10.1016/j.annemergmed.2013.04.001
- CDC. *Campylobacter (Campylobacteriosis)*. Atlanta, GA: Center for Disease Control (CDC) (2019).
- Safefood. *Consumer Preferences of Poultry Decontamination Methods on the Island of Ireland*. Dublin: Safefood (2017).
- Buncic S, Sofos J. Interventions to control *Salmonella* contamination during poultry, cattle and pig slaughter. *Food Res Int.* (2012) 45:641–55. doi: 10.1016/j.foodres.2011.10.018
- Han Y, Xu X, Jiang Y, Zhou G, Sun X, Xu B. Inactivation of food spoilage bacteria by high pressure processing: evaluation with conventional media and PCR–DGGE analysis. *Food Res Int.* (2010) 43:1719–24. doi: 10.1016/j.foodres.2010.05.012
- Black EP, Setlow P, Hocking AD, Stewart CM, Kelly AL, Hoover DG. Response of spores to high-pressure processing. *Compr Rev Food Sci Food Saf.* (2007) 6:103–19. doi: 10.1111/j.1541-4337.2007.00021.x
- Voigt DD, Kelly AL, Huppertz T. High-pressure processing of milk and dairy products. *Emerg Dairy Process Technol Oppor Dairy Ind.* (2015) 3:71–92. doi: 10.1002/9781118560471.ch3
- Marušić Radović N, Ježek D, Markov K, Frece J, Ćurić D, Medić H. The effect of high pressure treatment on the quality of chicken breast meat. *Hrvat časopis za prehrambenu Tehnol Biotehnol i Nutr.* (2020) 14:76–81. doi: 10.31895/hcptbn.14.3-4.6
- Xiong G, Fu X, Pan D, Qi J, Xu X, Jiang X. Influence of ultrasound-assisted sodium bicarbonate marination on the curing efficiency of chicken breast meat. *Ultrason Sonochem.* (2020) 60:104808. doi: 10.1016/j.ultsonch.2019
- Lawrie RA, Ledward D. *Lawrie's Meat Science*. Cambridge: Woodhead Publishing (2014).

## CONCLUSION

Effective interventions, based on the combinations of emerging process technologies with the well-understood efficacies of nature-based compounds, can be designed to enhance safety and quality and minimize food loss in poultry processing. Studies to date have demonstrated that the order or sequence of application can be variable to address the key risks at different process stages, providing great flexibility if being considered as effective replacements for thermal or conventional chemical strategies.

## AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

## ACKNOWLEDGMENTS

The authors would like to acknowledge the Department of Agriculture, Food and Marine under grant number 17/F/275 and Science Foundation Ireland (SFI) under grant number SFI/16/BBSRC/3391 along with the BBSRC under grant reference BB/P008496/1.

- Al-Hilphy AR, Al-Temimi AB, Al Rubaiy HHM, Anand U, Delgado-Pando G, Lakhssassi N. Ultrasound applications in poultry meat processing: a systematic review. *J Food Sci.* (2020) 85:1386–96. doi: 10.1111/1750-3841.15135
- Mostafavi HA, Fathollahi SMMH. *The Potential of Food Irradiation: Benefits and Limitations*. Tehran: Trends in Vital Food and Control Engineering (2012).
- Calderón-Miranda ML, Barbosa-Cánovas GV, Swanson BG. Inactivation of *Listeria innocua* in liquid whole egg by pulsed electric fields and nisin. *Int J Food Microbiol.* (1999) 51:7–17. doi: 10.1016/S0168-1605(99)00070-7
- Gao Y, Zhuang H, Yeh HY, Bowker B, Zhang J. Effect of rosemary extract on microbial growth, pH, color, and lipid oxidation in cold plasma-processed ground chicken patties. *Innov Food Sci Emerg Technol.* (2019) 57:102168. doi: 10.1016/j.ifset.2019.05.007
- Huffman RD. Current and future technologies for the decontamination of carcasses and fresh meat. *Meat Sci.* (2002) 62:285–94. doi: 10.1016/S0309-1740(02)00120-1
- Clemente I, Condón-Abanto S, Pedrós-Garrido S, Whyte P, Lyng JG. Efficacy of pulsed electric fields and antimicrobial compounds used alone and in combination for the inactivation of *Campylobacter jejuni* in liquids and raw chicken. *Food Control.* (2020) 107:106491. doi: 10.1016/j.foodcont.2019.01.017
- Kure CF, Axelsson L, Carlehög M, Måge I, Jensen MR, Holck A. The effects of a pilot-scale steam decontamination system on the hygiene and sensory quality of chicken carcasses. *Food Control.* (2020) 109:106948. doi: 10.1016/j.foodcont.2019.106948
- Silano V, Barat Baviera JM, Bolognesi C, Brüscheweiler BJ, Chesson A, Cocconcelli PS, et al. Evaluation of the safety and efficacy of the organic acids lactic and acetic acids to reduce microbiological surface contamination on pork carcasses and pork cuts. *Efsa J.* (2018) 16:5482. doi: 10.2903/j.efsa.2018.5482
- Cadena M, Kelman T, Marco ML, Pitesky M. Understanding antimicrobial resistance (AMR) profiles of *Salmonella* biofilm and planktonic bacteria challenged with disinfectants commonly used during poultry processing. *Foods.* (2019) 8:275. doi: 10.3390/foods8070275



22. Micciche AC, Feye KM, Rubinelli PM, Lee JA, Knueven CJ, Ricke SC. Comparison of acid sanitizers on *Salmonella* typhimurium inoculated commercial poultry processing reuse water. *Front Sustain Food Syst.* (2019) 2:90. doi: 10.3389/fsufs.2018.00090
23. Wang H, Qi J, Duan D, Dong Y, Xu X, Zhou G. Combination of a novel designed spray cabinet and electrolyzed water to reduce microorganisms on chicken carcasses. *Food Control.* (2018) 86:200–6. doi: 10.1016/j.foodcont.2017.11.027
24. Keener KM, Bashor MP, Curtis PA, Sheldon BW, Kathariou S. Comprehensive review of campylobacter and poultry processing. *Compr Rev Food Sci Food Saf.* (2004) 3:105–16. doi: 10.1111/j.1541-4337.2004.tb00060.x
25. Zhuang H, Rothrock MJ, Hiatt KL, Lawrence KC, Gamble GR, Bowker BC, et al. In-package air cold plasma treatment of chicken breast meat: treatment time effect. *J Food Qual.* (2019) 2019:1837351. doi: 10.1155/2019/1837351
26. Wang J, Zhuang H, Hinton A, Zhang J. Influence of in-package cold plasma treatment on microbiological shelf life and appearance of fresh chicken breast fillets. *Food Microbiol.* (2016) 60:142–6. doi: 10.1016/j.fm.2016.07.007
27. Vester C, Kjeldgaard J, Ingmer H, Bisgaard M, Christensen H. International Journal of Food Microbiology Microbiota encompassing putative spoilage bacteria in retail packaged broiler meat and commercial broiler abattoir. *Int J Food Microbiol.* (2019) 300:14–21. doi: 10.1016/j.ijfoodmicro.2019.04.003
28. Moutiq R, Misra NN, Mendonça A, Keener K. In-package decontamination of chicken breast using cold plasma technology: microbial, quality and storage studies. *Meat Sci.* (2020) 159:107942. doi: 10.1016/j.meatsci.2019.107942
29. Tresse O. *Bacterial Contaminants of Poultry Meat : Sources, Species, and Dynamics.* Nantes: Microorganisms (2017).
30. Katiyo W, Kock HL De, Coorey R, Buys EM. LWT - Food Science and Technology Sensory implications of chicken meat spoilage in relation to microbial and physicochemical characteristics during refrigerated storage. *LWT Food Sci Technol.* (2020) 128:109468. doi: 10.1016/j.lwt.2020.109468
31. Argyri AA, Papadopoulou OS, Nisiotiou A, Tassou CC, Chorianopoulos N. Effect of high pressure processing on the survival of *Salmonella* Enteritidis and shelf-life of chicken fillets. *Food Microbiol.* (2018) 70:55–64. doi: 10.1016/j.fm.2017.08.019
32. Argyri AA, Papadopoulou OS, Sourri P, Chorianopoulos N, Tassou CC. Quality and safety of fresh chicken fillets after high pressure processing : survival of indigenous *Brochothrix thermosphacta* and inoculated *Listeria monocytogenes*. *Microorganisms.* (2019) 7:520. doi: 10.3390/microorganisms7110520
33. Pasquali F, De Cesare A, Meunier M, Guyard M, Rivoal K, Chemaly M, et al. *Current Challenges in Poultry Meat Safety.* Duxford: Elsevier Ltd. (2017). doi: 10.1016/B978-0-08-100763-1.00007-6
34. Sommers C, Sheen S, Scullen OJ, Mackay W. Inactivation of *Staphylococcus saprophyticus* in chicken meat and purge using thermal processing, high pressure processing, gamma radiation, and ultraviolet light (254 nm). *Food Control.* (2017) 75:78–82. doi: 10.1016/j.foodcont.2016.12.020
35. Udhayavel S, Thippichettyalayam Ramasamy G, Gowthaman V, Malmarugan S, Senthilvel K. Occurrence of *Clostridium perfringens* contamination in poultry feed ingredients: isolation, identification and its antibiotic sensitivity pattern. *Anim Nutr.* (2017) 3:309–12. doi: 10.1016/j.aninu.2017.05.006
36. Golić N, Veljović K, Popović N, Djokić J, Strahinić I, Mrvaljević I, et al. *In vitro* and *in vivo* antagonistic activity of new probiotic culture against *Clostridium difficile* and *Clostridium perfringens*. *BMC Microbiol.* (2017) 17:1–9. doi: 10.1186/s12866-017-1015-5
37. Guo S, Liu D, Zhang B, Li Z, Li Y, Ding B, et al. Two *Lactobacillus* species inhibit the growth and  $\alpha$ -toxin production of *Clostridium perfringens* and induced proinflammatory factors in chicken intestinal epithelial cells *in vitro*. *Front Microbiol.* (2017) 8:1–12. doi: 10.3389/fmicb.2017.02081
38. Hamad GM, Abdelmotilib NM, Darwish AMG, Zeitoun AM. Commercial probiotic cell-free supernatants for inhibition of *Clostridium perfringens* poultry meat infection in Egypt. *Anaerobe.* (2020) 62:102181. doi: 10.1016/j.anaerobe.2020.102181
39. Bolton DJ. *Campylobacter* virulence and survival factors. *Food Microbiol.* (2015) 48:99–108. doi: 10.1016/j.fm.2014.11.017
40. Perez-arnedo I, Gonzalez-fandos E. Prevalence of *Campylobacter* spp. in poultry in three Spanish farms, a slaughterhouse and a further processing plant. *Foods.* (2019) 8:111. doi: 10.3390/foods8030111
41. Rossler E, Olivero C, Soto LP, Frizzo LS, Zimmermann J, Rosmini MR, et al. Prevalence, genotypic diversity and detection of virulence genes in thermotolerant *Campylobacter* at different stages of the poultry meat supply chain. *Int J Food Microbiol.* (2020) 326:108641. doi: 10.1016/j.ijfoodmicro.2020.108641
42. Berrang ME, Meinersmann RJ, Cox NA. *Campylobacter* subtypes detected in broiler ceca and livers collected at slaughter. *Poult Sci.* (2019) 98:5908–12. doi: 10.3382/ps/pez340
43. Gharajalar N, Sahar, Hassanzadeh P, Hosseinali Nejad N. Molecular detection of *Campylobacter* species and Cytolethal distending toxin isolated from chicken livers in Tabriz. *Comp Immunol Microbiol Infect Dis.* (2020) 71:101474. doi: 10.1016/j.cimid.2020.101474
44. Lamas A, Regal P, Vázquez B, Miranda JM, Cepeda A, Franco CM. *Salmonella* and *Campylobacter* biofilm formation : a comparative assessment from farm to fork. *J Sci Food Agric.* (2018) 98:4014–32. doi: 10.1002/jsfa.8945
45. Chlebicz A, Slizewska K. *Campylobacteriosis, Salmonellosis, Yersiniosis, and Listeriosis* as zoonotic foodborne diseases: a review. *Int J Environ Res Public Health.* (2018) 15:1–28. doi: 10.3390/ijerph15050863
46. WHO. *Salmonella (non-typhoidal).* (2018). Available online at: [https://www.who.int/news-room/fact-sheets/detail/salmonella-\(non-typhoidal\)](https://www.who.int/news-room/fact-sheets/detail/salmonella-(non-typhoidal)) (accessed June 9, 2020).
47. Jajere SM. A review of *Salmonella enterica* with particular focus on the pathogenicity and virulence factors, host specificity and antimicrobial resistance including multidrug resistance. *Vet World.* (2019) 12:504–21. doi: 10.14202/vetworld.2019.504-521
48. Bouchrif B, Paglietti B, Murgia M, Piana A, Cohen N, Ennaji MM, et al. Prevalence and antibiotic-resistance of *Salmonella* isolated from food in Morocco. *J Infect Dev Ctries.* (2009) 3:35–40. doi: 10.3855/jidc.103
49. Tack DM, Ray L, Griffin PM, Cieslak PR, Dunn J, Rissman T, et al. Preliminary incidence and trends of infections with pathogens transmitted commonly through food - foodborne diseases active surveillance network, 10 U.S. Sites, 2016–2019. *MMWR Morb Mortal Wkly Rep.* (2020) 69:509–14. doi: 10.15585/mmwr.mm6917a1
50. Dantas STA, Camargo CH, Tiba-casas MR, Vivian RC, Pinto JPAN, Pantoja JCF, et al. Environmental persistence and virulence of *Salmonella* spp. Isolated from a poultry slaughterhouse. *Food Res Int.* (2020) 129:108835. doi: 10.1016/j.foodres.2019.108835
51. El-Sayed SM, Youssef AM. Potential application of herbs and spices and their effects in functional dairy products. *Heliyon.* (2019) 5:e01989. doi: 10.1016/j.heliyon.2019.e01989
52. Shah MA, Bosco SJD, Mir SA. Plant extracts as natural antioxidants in meat and meat products. *Meat Sci.* (2014) 98:21–33. doi: 10.1016/j.meatsci.2014.03.020
53. Zhang H, Wu J, Guo X. Effects of antimicrobial and antioxidant activities of spice extracts on raw chicken meat quality. *Food Sci Hum Wellness.* (2016) 5:39–48. doi: 10.1016/j.fshw.2015.11.003
54. Corrêa TQ, Blanco KC, Garcia EB, Perez SML, Chianfrone DJ, Moraes VS, et al. Effects of ultraviolet light and curcumin-mediated photodynamic inactivation on microbiological food safety: a study in meat and fruit. *Photodiagnosis Photodyn Ther.* (2020) 30:101678. doi: 10.1016/j.pdpdt.2020.101678
55. Amerah AM, Ouwehand AC. *Use of Essential Oils in Poultry Production.* London: Elsevier Inc. (2016). doi: 10.1016/B978-0-12-416641-7.0010-9
56. Burt S. Essential oils: their antibacterial properties and potential applications in foods—a review. *Int J Food Microbiol.* (2004) 94:223–53. doi: 10.1016/j.ijfoodmicro.2004.03.022
57. Lin L, Liao X, Cui H. Cold plasma treated thyme essential oil/silk fibroin nanofibers against *Salmonella Typhimurium* in poultry meat. *Food Packag Shelf Life.* (2019) 21:100337. doi: 10.1016/j.fpsl.2019.100337
58. Chouliara E, Karatapanis A, Savvaides IN, Kontominas MG. Combined effect of oregano essential oil and modified atmosphere packaging on shelf-life extension of fresh chicken breast meat, stored at 4°C. *Food Microbiol.* (2007) 24:607–17. doi: 10.1016/j.fm.2006.12.005

59. Fratianni F, De Martino L, Melone A, De Feo V, Coppola R, Nazzaro F. Preservation of chicken breast meat treated with thyme and balm essential oils. *J Food Sci.* (2010) 75:M528–35. doi: 10.1111/j.1750-3841.2010.01791.x
60. Ntzimani A, Giatrakou V, Savvaidis I. Combined natural antimicrobial treatments (EDTA, lysozyme, rosemary and oregano oil) on semi cooked coated chicken meat stored in vacuum packages at 4°C: microbiological and sensory evaluation. *Innov food Sci Emerg Technol.* (2010) 11:187–96. doi: 10.1016/j.ifset.2009.09.004
61. Gurtler J, Mai TL. Traditional preservatives - organic acids. *Encycl Food Microbiol.* (2014) 3:119–30. doi: 10.1016/B978-0-12-384730-0.00260-3
62. Anyasi T, Jideani A, Edokpayi J, Anokwuru C. *Application of Organic Acids in Food Preservation.* New York, NY: Nova Publishers (2017). p. 45
63. Theron MM, Lues JFR. Organic acids and meat preservation : a review organic acids and meat preservation : a review. *Food Rev Int.* (2007) 23:141–58. doi: 10.1080/87559120701224964
64. FDA. Food additives for use in meat and poultry products: sodium diacetate, sodium acetate, sodium lactate and potassium. *Fed Regist.* (2000) 65:3121–3.
65. Zhang Z-H, Wang L-H, Zeng X-A, Han Z, Brennan CS. Non-thermal technologies and its current and future application in the food industry: a review. *Int J Food Sci Technol.* (2019) 54:1–13. doi: 10.1111/ijfs.13903
66. Wang L, Xia Q, Li Y. Synergistic effects of high pressure processing and slightly acidic electrolysed water on the inactivation of *Bacillus cereus* spores. *Int J Food Sci Technol.* (2017) 52:2429–35. doi: 10.1111/ijfs.13527
67. Yeh HY, Line JE, Hinton A Jr, Gao Y, Zhuang H. The effect of rosemary Extract and cold plasma treatments on bacterial community diversity in poultry ground meats. *Heliyon.* (2019) 5:e02719. doi: 10.1016/j.heliyon.2019.e02719
68. Sahebkar A, Hosseini M, Sharifan A. Plasma-assisted preservation of breast chicken fillets in essential oils-containing marinades. *LWT.* (2020) 131:109759. doi: 10.1016/j.lwt.2020.109759
69. Rodríguez-Calleja JM, Cruz-Romero MC, O'Sullivan MG, García-López ML, Kerry JP. High-pressure-based hurdle strategy to extend the shelf-life of fresh chicken breast fillets. *Food Control.* (2012) 25:516–24. doi: 10.1016/j.foodcont.2011.11.014
70. de Oliveira TLC, de Castro Leite BR, Ramos ALS, Ramos EM, Piccoli RH, Cristianini M. Phenolic carvacrol as a natural additive to improve the preservative effects of high pressure processing of low-sodium sliced vacuum-packed turkey breast ham. *LWT Food Sci Technol.* (2015) 64:1297–308. doi: 10.1016/j.lwt.2015.06.011
71. Chien S-Y, Sheen S, Sommers CH, Sheen L-Y. Modeling the inactivation of intestinal pathogenic *Escherichia coli* O157:H7 and uropathogenic *E. coli* in ground chicken by high pressure processing and thymol. *Front Microbiol.* (2016) 7:1–11. doi: 10.3389/fmicb.2016.00920
72. Sams AR, Ferial R. Microbial effects of ultrasonication of broiler drumstick skin. *J Food Sci.* (1991) 56:247–8. doi: 10.1111/j.1365-2621.1991.tb08020.x
73. Kordowska-Wiater M, Stasiak DM. Effect of ultrasound on survival of gram-negative bacteria on chicken skin surface. *Bull Vet Inst Pulawy.* (2011) 55:207–10.
74. Piñon MI, Alarcon-Rojo AD, Renteria AL, Mendez G, Janacua-Vidales H. Reduction of microorganisms in marinated poultry breast using oregano essential oil and power ultrasound. *Acta Aliment.* (2015) 44:527–33. doi: 10.1556/066.2015.44.0024
75. Yordanov DG, Angelova GV. *High Pressure Processing for Foods Preserving.* Plodiv: Biotechnology & Biotechnological Equipment (2014). p. 2818.
76. Toldrá F. *Safety of Meat and Processed Meat.* New York, NY: Springer (2009). doi: 10.1007/978-0-387-89026-5
77. Gassiot M, Masoliver P. *Commercial High Pressure Processing of Ham and Other Sliced Meat Products at Esteban España, S.A.* Oxford; Cambridge; Philadelphia, PA; New Delhi: Woodhead Publishing Limited (2010). doi: 10.1533/9780857090713.1.21
78. Bajovic B, Bolumar T, Heinz V. Quality considerations with high pressure processing of fresh and value added meat products. *MESCI.* (2012) 92:280–9. doi: 10.1016/j.meatsci.2012.04.024
79. Tracz M. Effects of high hydrostatic pressure on *Campylobacter jejuni* in poultry meat. *Pol J Vet Sci.* (2015) 18:261–6. doi: 10.1515/pjvs-2015-0034
80. Sheen S, Cassidy J, Scullen B, Uknalis J, Sommers C. Inactivation of *Salmonella* spp. in ground chicken using high pressure processing. *Food Control.* (2015) 57:41–7. doi: 10.1016/j.foodcont.2015.04.005
81. Xu A, Scullen OJ, Sheen S, Liu Y, Johnson JR, Sommers CH. Inactivation of extraintestinal pathogenic *E. coli* suspended in ground chicken meat by high pressure processing and identification of virulence factors which may affect resistance to high pressure. *Food Control.* (2020) 111:107070. doi: 10.1016/j.foodcont.2019.107070
82. Liu Y, Betti M, Gänzle MG. High pressure inactivation of *Escherichia coli*, *Campylobacter jejuni*, and spoilage microbiota on poultry meat. *J Food Prot.* (2012) 75:497–503. doi: 10.4315/0362-028X.JFP-11-316
83. Dissing J, Bruun-Jensen L, Skibsted LH. Effect of high-pressure treatment on lipid oxidation in turkey thigh muscle during chill storage. *Eur Food Res Technol.* (1997) 205:11–3. doi: 10.1007/s002170050115
84. Kruk ZA, Yun H, Rutley DL, Jung E, Ji Y, Jo C. The effect of high pressure on microbial population, meat quality and sensory characteristics of chicken breast fillet. *Food Control.* (2011) 22:6–12. doi: 10.1016/j.foodcont.2010.06.003
85. Orlien V, Hansen E, Skibsted LH. Lipid oxidation in high-pressure processed chicken breast muscle during chill storage: critical working pressure in relation to oxidation mechanism. *Eur Food Res Technol.* (2000) 211:99–104. doi: 10.1007/s002179900118
86. Yuste J, Pla R, Capellas M, Mor-Mur M. Application of high-pressure processing and nisin to mechanically recovered poultry meat for microbial decontamination. *Food Control.* (2002) 13:451–5. doi: 10.1016/S0956-7135(01)00071-8
87. Ogihara H, Yatuzuka M, Horie N, Furukawa S, Yamasaki M. Synergistic effect of high hydrostatic pressure treatment and food additives on the inactivation of *Salmonella enteritidis*. *Food Control.* (2009) 20:963–6. doi: 10.1016/j.foodcont.2008.11.004
88. Demirci A, Ngadi MO. *Microbial Decontamination in the Food Industry.* Pennsylvania: Woodhead Publishing (2012). doi: 10.1533/9780857095756
89. Bhat ZF, Morton JD, Mason SL, Bekhit AEDA. Current and future prospects for the use of pulsed electric field in the meat industry. *Crit Rev Food Sci Nutr.* (2019) 59:1660–74. doi: 10.1080/10408398.2018.1425825
90. Pillet F, Formosa-Dague C, Baaziz H, Dague E, Rols MP. Cell wall as a target for bacteria inactivation by pulsed electric fields. *Sci Rep.* (2016) 6:1–8. doi: 10.1038/srep19778
91. Espina L, Monfort S, Álvarez I, García-Gonzalo D, Pagán R. Combination of pulsed electric fields, mild heat and essential oils as an alternative to the ultrapasteurization of liquid whole egg. *Int J Food Microbiol.* (2014) 189:119–25. doi: 10.1016/j.ijfoodmicro.2014.08.002
92. Haughton PN, Lyng JG, Cronin DA, Morgan DJ, Fanning S, Whyte P. Efficacy of pulsed electric fields for the inactivation of indicator microorganisms and foodborne pathogens in liquids and raw chicken. *Food Control.* (2012) 25:131–5. doi: 10.1016/j.foodcont.2011.10.030
93. Álvarez I, Mañas P, Condón S, Raso J. Resistance variation of *Salmonella enterica* serovars to pulsed electric fields treatments. *J Food Sci.* (2003) 68:2316–20. doi: 10.1111/j.1365-2621.2003.tb05765.x
94. McDonnell C, Allen P, Chardonnerau F, Arimi J, Lyng J. The use of pulsed electric fields for accelerating the salting of pork. *Leb und-Technologie.* (2014) 59:1054–60. doi: 10.1016/j.lwt.2014.05.053
95. Arroyo C, Lascorz D, O'Dowd L, Noci F, Arimi J, Lyng JG. Effect of Pulsed Electric Field treatments at various stages during conditioning on quality attributes of beef longissimus thoracis et lumborum muscle. *Meat Sci.* (2015) 99:52–9. doi: 10.1016/j.meatsci.2014.08.004
96. Bekhit AE-DA, van de Ven R, Fahri F, Hopkins DL. Effect of pulsed electric field treatment on cold-boned muscles of different potential tenderness. *Food Bioprocess Technol.* (2014) 7:3136–46. doi: 10.1007/s11947-014-1324-8
97. Ahmad RS, Imran A, Hussain MB. Nutritional composition of meat. *Meat Sci Nutr.* (2018) 61:77045. doi: 10.5772/intechopen.77045
98. Khan AA, Randhawa MA, Carne A, Mohamed Ahmed IA, Barr D, Reid M, et al. Quality and nutritional minerals in chicken breast muscle treated with low and high pulsed electric fields. *Food Bioprocess Technol.* (2018) 11:122–31. doi: 10.1007/s11947-017-1997-x
99. Khan AA, Randhawa MA, Carne A, Mohamed Ahmed IA, Al-Juhaimi FY, Barr D, et al. Effect of low and high pulsed electric field processing on macro

- and micro minerals in beef and chicken. *Innov Food Sci Emerg Technol*. (2018) 45:273–9. doi: 10.1016/j.ifset.2017.11.012
100. Mandal R, Mohammadi X, Wiktor A, Singh A, Pratap Singh A. Applications of pulsed light decontamination technology in food processing: an overview. *Appl Sci*. (2020) 10:3606. doi: 10.3390/app10103606
  101. Bintsis T, Litopoulou-Tzanetaki E, Robinson RK. Existing and potential applications of ultraviolet light in the food industry - a critical review. *J Sci Food Agric*. (2000) 80:637–45. doi: 10.1002/(SICI)1097-0010(20000501)80:6<637::AID-JSFA603>3.0.CO;2-1
  102. Chun HH, Kim JY, Lee BD, Yu DJ, Song KB. Effect of UV-C irradiation on the inactivation of inoculated pathogens and quality of chicken breasts during storage. *Food Control*. (2010) 21:276–80. doi: 10.1016/j.foodcont.2009.06.006
  103. Isohanni PMI, Lyhs U. Use of ultraviolet irradiation to reduce *Campylobacter jejuni* on broiler meat. *Poult Sci*. (2009) 88:661–8. doi: 10.3382/ps.2008-00259
  104. McLeod A, Hovde Liland K, Haugen JE, Sørheim O, Myhrer KS, Holck AL. Chicken fillets subjected to UV-C and pulsed UV light: reduction of pathogenic and spoilage bacteria, and changes in sensory quality. *J Food Saf*. (2018) 38:12421. doi: 10.1111/jfs.12421
  105. Haughton PN, Lyng JG, Cronin DA, Morgan DJ, Fanning S, Whyte P. Efficacy of UV light treatment for the microbiological decontamination of chicken, associated packaging, and contact surfaces. *J Food Prot*. (2011) 74:565–72. doi: 10.4315/0362-028X.JFP-10-356
  106. Yang S, Sadekuzzaman M, Ha SD. Reduction of *Listeria monocytogenes* on chicken breasts by combined treatment with UV-C light and bacteriophage ListShield. *LWT Food Sci Technol*. (2017) 86:193–200. doi: 10.1016/j.lwt.2017.07.060
  107. Holck AL, Liland KH, Drømtorp SM, Carlehö GM, McLeod A. Comparison of UV-C and pulsed UV light treatments for reduction of salmonella, listeria monocytogenes, and enterohemorrhagic *Escherichia coli* on eggs. *J Food Prot*. (2018) 81:6–16. doi: 10.4315/0362-028X.JFP-17-128
  108. Begum M, Hocking AD, Miskelly D. Inactivation of food spoilage fungi by ultra violet (UVC) irradiation. *Int J Food Microbiol*. (2009) 129:74–7. doi: 10.1016/j.ijfoodmicro.2008.11.020
  109. Silva-Espinoza BA, Palomares-Navarro JJ, Tapia-Rodriguez MR, Cruz-Valenzuela MR, González-Aguilar GA, Silva-Campa E, et al. Combination of ultraviolet light-C and clove essential oil to inactivate *Salmonella Typhimurium* biofilms on stainless steel. *J Food Saf*. (2020) 40:12788. doi: 10.1111/jfs.12788
  110. Feng H, Barbosa-Cánovas GV, Weiss J. *Ultrasound Technologies for Food and Bioprocessing*. New York, NY: Springer New York (2011). doi: 10.1007/978-1-4419-7472-3
  111. Turantaş F, Kiliç GB, Kiliç B. Ultrasound in the meat industry: general applications and decontamination efficiency. *Int J Food Microbiol*. (2015) 198:59–69. doi: 10.1016/j.ijfoodmicro.2014.12.026
  112. Haughton PN, Lyng JG, Morgan DJ, Cronin DA, Noci F, Fanning S, et al. An evaluation of the potential of high-intensity ultrasound for improving the microbial safety of poultry. *Food Bioprocess Technol*. (2012) 5:992–8. doi: 10.1007/s11947-010-0372-y
  113. Piñon MI, Alarcon-Rojo AD, Renteria AL, Carrillo-Lopez LM. Microbiological properties of poultry breast meat treated with high-intensity ultrasound. *Ultrasonics*. (2020) 102:1. doi: 10.1016/j.ultras.2018.01.001
  114. Seo MK, Jeong HL, Han SH, Kang I, Ha SD. Impact of ethanol and ultrasound treatment on mesophilic aerobic bacteria, coliforms, and *Salmonella Typhimurium* on chicken skin. *Poult Sci*. (2019) 98:6954–63. doi: 10.3382/ps/pez486
  115. Leal-Ramos MY, Alarcon-Rojo AD, Mason TJ, Paniwnyk L, Alarjah M. Ultrasound-enhanced mass transfer in Halal compared with non-Halal chicken. *J Sci Food Agric*. (2011) 91:130–3. doi: 10.1002/jsfa.4162
  116. Smith DP. Effect of ultrasonic marination on broiler breast meat quality and *Salmonella* contamination. *Int J Poult Sci*. (2011) 10:757–9. doi: 10.3923/ijps.2011.757.759
  117. Inguglia ES, Burgess CM, Kerry JP, Tiwari BK. Ultrasound-assisted marination: role of frequencies and treatment time on the quality of sodium-reduced poultry meat. *Foods*. (2019) 8:473. doi: 10.3390/foods8100473
  118. Thirumdas R, Sarangapani C, Annapure US. Cold plasma: a novel non-thermal technology for food processing. *Food Biophys*. (2014) 10:1–11. doi: 10.1007/s11483-014-9382-z
  119. Fernández A, Thompson A. The inactivation of *Salmonella* by cold atmospheric plasma treatment. *Food Res Int*. (2012) 45:678–84. doi: 10.1016/j.foodres.2011.04.009
  120. Lee H, Yong HI, Kim HJ, Choe W, Yoo SJ, Jang EJ, et al. Evaluation of the microbiological safety, quality changes, and genotoxicity of chicken breast treated with flexible thin-layer dielectric barrier discharge plasma. *Food Sci Biotechnol*. (2016) 25:1189–95. doi: 10.1007/s10068-016-0189-1
  121. Ulbin-Figlewicz N, Jarmoluk A, Marycz K. Antimicrobial activity of low-pressure plasma treatment against selected foodborne bacteria and meat microbiota. *Ann Microbiol*. (2015) 65:1537–46. doi: 10.1007/s13213-014-0992-y
  122. Lu H, Patil S, Keener KM, Cullen PJ, Bourke P. Bacterial inactivation by high-voltage atmospheric cold plasma: influence of process parameters and effects on cell leakage and DNA. *J Appl Microbiol*. (2014) 116:784–94. doi: 10.1111/jam.12426
  123. Lee ES, Cheigh CI, Kang JH, Lee SY, Min SC. Evaluation of in-package atmospheric dielectric barrier discharge cold plasma treatment as an intervention technology for decontaminating bulk ready-to-eat chicken breast cubes in plastic containers. *Appl Sci*. (2020) 10:1–21. doi: 10.3390/app10186301
  124. Mai-Prochnow A, Alam D, Zhou R, Zhang T, Ostrikov K, Cullen PJ. The effect of rosemary E and cold plasma treatments on bacterial community diversity in poultry ground meats. Microbial decontamination of chicken using atmospheric plasma bubbles. *Plasma Process Polym*. (2020) 2020:52. doi: 10.1002/ppap.202000052
  125. The European Parliament and the council of European union. Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29 April 2004 laying down specific hygiene rules for on the hygiene of foodstuffs. *Off J Eur Union*. (2004) 139:1–155.
  126. Llana-Ruiz-Cabello M, Pichardo S, Maisanaba S, Puerto M, Prieto AI, Gutiérrez-Praena D, et al. *In vitro* toxicological evaluation of essential oils and their main compounds used in active food packaging: a review. *Food Chem Toxicol*. (2015) 81:9–27. doi: 10.1016/j.fct.2015.03.030
  127. Jayasena DD, Jo C. Essential oils as potential antimicrobial agents in meat and meat products: a review. *Trends Food Sci Technol*. (2013) 34:96–108. doi: 10.1016/j.tifs.2013.09.002
  128. Hyldgaard M, Mygind T, Meyer RL. Essential oils in food preservation: mode of action, synergies, and interactions with food matrix components. *Front Microbiol*. (2012) 3:1–24. doi: 10.3389/fmicb.2012.00012
  129. Fernández-López J, Viuda-Martos M. Introduction to the special issue: application of essential oils in food systems. *Foods*. (2018) 7:56. doi: 10.3390/foods7040056

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Barroug, Chaple and Bourke. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.