

Small Non-coding RNA Expression Following Respiratory Syncytial Virus or Measles Virus Infection of Neuronal Cells

Bakre, A. A., Duffy, C., Abdullah, H., Cosby, S. L., & Tripp, R. A. (2021). Small Non-coding RNA Expression Following Respiratory Syncytial Virus or Measles Virus Infection of Neuronal Cells. *Frontiers in Microbiology*, *12*, Article 671852. https://doi.org/10.3389/fmicb.2021.671852

Published in:

Frontiers in Microbiology

Document Version: Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:

Link to publication record in Queen's University Belfast Research Portal

Publisher rights

Copyright 2021 the authors.

This is an open access article published under a Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: http://go.qub.ac.uk/oa-feedback





Small Non-coding RNA Expression Following Respiratory Syncytial Virus or Measles Virus Infection of Neuronal Cells

Abhijeet A. Bakre^{1*}, Catherine Duffy², Hani'ah Abdullah³, S. Louise Cosby^{2,3} and Ralph A. Tripp^{1*}

OPEN ACCESS

Edited by:

Jianrong Li, The Ohio State University, United States

Reviewed by:

Qiyun Zhu, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences (CAS), China Mark Peeples, The Ohio State University, United States

*Correspondence:

Abhijeet A. Bakre bakre@uga.edu Ralph A. Tripp ratripp@uga.edu

Specialty section:

This article was submitted to Virology, a section of the journal Frontiers in Microbiology

Received: 24 February 2021 Accepted: 03 August 2021 Published: 03 September 2021

Citation:

Bakre AA, Duffy C, Abdullah H, Cosby SL and Tripp RA (2021) Small Non-coding RNA Expression Following Respiratory Syncytial Virus or Measles Virus Infection of Neuronal Cells. Front. Microbiol. 12:671852. doi: 10.3389/fmicb.2021.671852 ¹ Department of Infectious Diseases, University of Georgia, Athens, GA, United States, ² Virology Branch, Veterinary Sciences Division, Agri-Food and Biosciences Institute, Belfast, United Kingdom, ³ Wellcome Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Belfast, United Kingdom

Respiratory syncytial virus (RSV) or measles virus (MeV) infection modifies host responses through small non-coding RNA (sncRNA) expression. We show that RSV or MeV infection of neuronal cells induces sncRNAs including various microRNAs and transfer RNA fragments (tRFs). We show that these tRFs originate from select tRNAs (GCC and CAC for glycine, CTT and AAC for Valine, and CCC and TTT for Lysine). Some of the tRNAs are rarely used by RSV or MeV as indicated by relative synonymous codon usage indices suggesting selective cleavage of the tRNAs occurs in infected neuronal cells. The data implies that differentially expressed sncRNAs may regulate host gene expression *via* multiple mechanisms in neuronal cells.

Keywords: respiratory syncytial virus, measles virus, neuronal cells, microRNAs, piwi-associated RNAs, transfer RNAs

INTRODUCTION

Small non-coding RNAs (sncRNAs) are <200 nucleotides and include microRNAs (miRs), PIWIinteracting RNAs (piRs), and transfer RNA fragments (tRNA)-derived RNA fragments (tRFs) (Petrov et al., 2017). The expression of sncRNAs differs in their biogenesis and functional activity (Bartel, 2009; Bakre et al., 2019). Most sncRNAs induced by viral infection involve regulatory mechanisms affecting virus replication, persistence, or latency (Grassmann and Jeang, 2008; Roizman et al., 2011), and immune evasion (Cullen, 2013; Bernier and Sagan, 2018). RSV typically infects ciliated respiratory epithelial cells, while MeV may infect dendritic cells and subsequently infect the epithelia for virus release from the respiratory tract (Lay et al., 2016; Lin and Richardson, 2016; Singh et al., 2016; Taniguchi et al., 2019). These viruses cause a spectrum of diseases and may infect a spectrum of people, i.e., the very young to the elderly (Griffin, 2018; Mazur et al., 2018). RSV is a major cause of serious lower respiratory tract infections in childhood (Mirra et al., 2018; Paes, 2018), but also affects the elderly and immune-compromised (Haber, 2018). MeV causes acute viral infection with respiratory involvement leading to symptoms of rash and more serious complications, and in some cases to mortality (Bohmwald et al., 2018; Leung et al., 2018). Both RSV and MeV are paramyxoviruses with non-segmented negative-sense singlestranded RNA genomes (Russell et al., 2018). Unfortunately, there are no approved RSV vaccines and anti-viral treatments are limited. Although there is an effective MeV vaccine, incidences of MeV infection are increasing worldwide due to vaccine complacency.

Respiratory syncytial virus and MeV can infect neuronal cells, and MeV may cause persistent infection (Rota et al., 2017; Griffin et al., 2018). Compromised immunity may facilitate neuronal infection (Rima and Duprex, 2005; Omar et al., 2017; Ganesan et al., 2018; Ferren et al., 2019). MeV infection of neuronal cells may lead to subacute sclerosing panencephalitis (SSPE), and measles inclusion body encephalitis (MIBE) in immunocompromised individuals. In MIBE, MeV replicates in the central nervous system usually because of an inadequate immune response (McQuaid et al., 1998; Hardie et al., 2013; Rota et al., 2017). RSV infection in mice has been associated with exaggerated neurogenic inflammation in the airways, and studies have shown that the neuropeptide, substance P (SP) is a mediator of neurogenic inflammation (King et al., 2001; Tan et al., 2008).

Expression of the RSV G protein is associated with increased pulmonary expression of SP, and lung neurons express CX3CR1 (Tripp et al., 2000, 2002, 2003; Haynes et al., 2003). CX3CR1 is the receptor for fractalkine (aka neurotactin) and CX3CR1 has been shown to bind RSV G protein (Tripp et al., 2001; Harcourt et al., 2004; Choi et al., 2012; Bakre et al., 2017; Bergeron et al., 2021).

Studies in mice suggest that RSV may cause lasting infection (Dakhama et al., 1997; Li et al., 2006). RSV has been shown to infect mouse primary cortical neuronal cells as shown by co-localization of the RSV N protein and neuronal markers (Li et al., 2006). These findings support how RSV may cause neurological irregularities in patients (Eisenhut, 2006; Li et al., 2006; Morichi et al., 2017). Several studies have described an association between lower respiratory tract RSV infection in infancy and the subsequent development of persistent wheezing in children (Zhou N. et al., 2017; Zhong et al., 2018). In mice, RSV induces long-term airway disease characterized by chronic airway inflammation and airway hyperreactivity (Long et al., 2016; Zhou N. et al., 2017), and RSV antigens have been detected in the lungs >100 days (Schwarze et al., 2004). However, RSV persistence does not appear in the bronchial epithelium (the primary site of viral replication) but in deeper lung structures (Long et al., 2016; Zhou N. et al., 2017).

Respiratory syncytial virus and MeV infection modulate sncRNA expression inducing mRNA degradation or translation inhibition and have a role in determining the level of protein expression of host cells (Munday et al., 2012). It has been shown that tRNAs are cleaved during cellular stress, and that cleavage results in tRFs that contain the 5' end (5'tRFs) or the 3' end (3'tRFs) of the parent tRNA molecule (Sofos et al., 2015; Advani and Ivanov, 2019). Although the specific activity of tRFs is not well-understood, tRFs can behave as small interfering RNA leading to the degradation of transcripts (Ivanov et al., 2011) which can regulate ribosomal loading and protein chain elongation (Sobala and Hutvagner, 2013).

Accumulating data have shown that both coding and noncoding transcriptomes are modified during RSV infection. We have shown that RSV infection of human A549 lung cells modifies sncRNA expression and that the RSV G protein and NS1/NS2 proteins modulate miR expression (Bakre et al., 2012, 2015, 2017). Other studies have shown that the pattern of miR expression is modified following RSV infection of nasal epithelial cells (Inchley et al., 2015; Hasegawa et al., 2018), and PBMCs from children (Wang et al., 2017). Deregulated miR expression occurs in MeV-infected human neuroblastoma cells and PBMCs (Inchley et al., 2015; Yis et al., 2015; Naaman et al., 2017; Hasegawa et al., 2018), as well as in PBMCs from RSV-infected children (Wang et al., 2017). It is known that MeV persistence in neuroblastoma cells is assisted by the downregulation of CDK6, a component of cell cycle progression regulated by miRNA-124 (Riddell et al., 2007; Naaman et al., 2017; Griffin et al., 2018).

In this study, we examined sncRNA responses in neuronal SHSY5Y (SHS) cells infected with RSV or MeV and show that infection with either of these viruses modifies miRs and tRF expression in a temporal and virus-specific manner. Antimicrobial peptides (AMPs) are generated in the response to pathogens, and arginine and glycine are especially abundant in AMPs (Mishra and Wang, 2012). In this study, we show that RSV or MeV infection modifies the extent of transfer RNA fragments (tRFs), and the finding of decreased full-length tRNAs after infection indicates decreased translation rates possibly indicative of a lessened anti-viral state. Together, these findings strengthen the notion that sncRNAs have a key role in regulating the host cell response to infection and viral replication.

MATERIALS AND METHODS

Cells and Viruses

Human SHSY5Y (SHS) neuroblastoma cells (ATCC CRL 2266) were grown in DMEM (Sigma, St. Louis, MO, United States) containing 10% FBS (Atlanta Biologicals, Atlanta, GA, United States). SHS cells were maintained at 37°C, 5% CO₂. RSV strain A2 (ATCC CCL 81) was propagated in Vero cells (Oshansky et al., 2009; Dahiya and Atreya, 2014). Briefly, 80% confluent Vero cells were infected (MOI = 1.0) for 1 h at 37° C, 5% CO₂. After infection, 2% FBS was added to the DMEM, and the cells were incubated for 3 days until syncytia were evident. On the day of harvest, the cells were removed and centrifuged at 500 \times g for 15 min at 4°C. RSV strain A2 was used for the infection in SHS cells and rgRSV224 GFP (Hallak et al., 2000, 2007; Techaarpornkul et al., 2002) (a kind gift from Dr. Michael Teng, University of South Florida) for microscopy studies. The WT strain Dublin-3267 of MeV was grown in Vero cells expressing human CD150 (SLAM), a high-affinity receptor for MeV (Tatsuo et al., 2000).

RNA Isolation and Library Construction

Total RNA from mock-treated or virus-infected SHS cells was isolated using RNAzol RT (MRC, Cincinnati, OH, United States)

TABLE 1 | Summary of read numbers.

Sample	Original # of	% Trimmed	% Uniquely mapped reads	
Campie	reads	reads		
12 hpi–RSV-1	5,953	98%	5%	
12 hpi–RSV-2	18,599	92%	42%	
12 hpi–RSV-3	7,546	94%	23%	
12 hpi–MeV-1	101,182	78%	50%	
12 hpi–MeV-2	27,880	74%	41%	
12 hpi–MeV-3	12,088	41%	11%	
24 hpi–RSV-1	46,761	72%	10%	
24 hpi–RSV-2	239,242	74%	9%	
24 hpi–RSV-3	74,687	74%	12%	
24 hpi–MeV-1	27,320	66%	7%	
24 hpi–MeV-2	26,797	62%	6%	
24 hpi–MeV-3	92,746	66%	9%	
SHS-1	57,494	79%	51%	
SHS-2	77,356	79%	44%	

Numbers indicate raw read counts, percent of reads with adaptor trimming [59], and those mapping to unique loci on the genome.



per the manufacturer's instructions and divided into small and large RNA fractions and quantified using a Qubit fluorimeter-based RNA assay (Invitrogen Life Tech, Carlsbad, CA, United States). Illumina next-generation sequencing (NGS) libraries were constructed using a TruSeq[®] small RNA library (Illumina, San Diego, CA, United States). Briefly, small RNA from mock-treated SHS cells (n = 2/group), or RSV- or MeV-infected SHS cells (n = 3/group) was isolated at 12

TABLE 2 Normalized read counts between mock,	RSV- and MeV-infected samples as	analyzed by OASIS 2.0 (Rahman et al.	, 2018a).
			, ,

Mature	log ₂ fold-change	padj	Mock Mock	12 h		24 h	
				RSV	MeV	RSV	MeV
miR-10b-5p	-2	0.00	1,416, 1,636	94, 144, 78	369, 262, 128	165, 248, 158	166, 173, 243
miR-181c-5p	-2	0.03	25, 26	12, 3, 6	6, 10, 12	9, 3, 9	10, 5, 6
miR-125b-5p	-2	0.02	50, 34	12, 12, 10	9, 9, 27	12, 10, 10	7, 21, 12
miR-375	2	0.01	56, 49	82, 225, 273	137, 165, 155	168, 132, 193	128, 163, 136
p-miR-247	2	0.02	128, 147	222, 336, 458	131, 298, 530	560, 688, 694	416, 446, 387
miR-21-5p	2	0.03	11, 11	211, 19, 29	43, 23, 15	32, 65, 27	57, 26, 62
miR-877-5p	2	0.02	1, 3	12, 28, 45	5, 10, 19	5, 6, 8	14, 21, 4
miR-7704	3	0.03	1, 1	0, 1, 6	1, 2, 4	25, 16, 17	14, 10, 18
miR-100-5p	3	0.05	0, 1	35, 1, 3	3, 2, 4	2, 1, 4	10, 10, 3
p-miR-330	3	0.01	1, 2	23, 6, 10	3, 4, 15	17, 13, 6	17, 52, 16
miR-182-5p	3	0.00	3, 3	12, 22, 16	34, 30, 23	18, 37, 24	20, 47, 39
miR-4532	4	0.02	0, 0	0, 0, 0	0, 0, 0	11, 29, 10	7, 26, 10
miR-1273g-3p	5	0.00	0, 0	0, 1, 3	0, 1, 0	15, 20, 18	24, 21, 22
miR-483-3p	4	0.00	0, 0	12, 11, 6	3, 10, 12	3, 4, 9	14, 0, 3
miR-3141	5	0.05	0, 0	0, 0, 0	0, 0, 0	29, 32, 14	61, 47, 17
p-miR-338	5	0.01	0, 0	0, 0, 0	0, 0, 0	23, 32, 10	24, 47, 11
miR-1246	5	0.00	0, 1	0, 3, 3	0, 1, 0	32, 42, 22	44, 110, 32
p-miR-317	6	0.00	0, 0	0, 0, 0	0, 0, 0	42, 44, 36	47, 68, 41
miR-3196	6	0.02	0, 0	0, 0, 0	0, 0, 0	29, 52, 14	54, 68, 24
miR-4516	6	0.01	0, 0	0, 0, 3	0, 0, 0	20, 20, 6	41, 26, 11
p-miR-106	6	0.01	0, 0	0, 0, 0	0, 0, 0	22, 18, 12	27, 52, 7
miR-4488	6	0.00	0, 0	0, 0, 0	0, 0, 0	25, 22, 11	34, 63, 15
p-miR-113	6	0.00	0, 0	0, 0, 0	0, 0, 0	37, 61, 35	44, 68, 50
p-miR-256	6	0.00	0, 0	12, 0, 3	0, 1, 0	12, 11, 10	10, 10, 3
p-miR-232	6	0.00	0, 0	0, 0, 0	0, 0, 0	8, 16, 6	20, 26, 9
p-miR-103	7	0.00	0, 0	0, 1, 0	0, 0, 0	22, 29, 21	17, 5, 34

Negative log₂ fold-change values indicate downregulation.

"p," predicted miR; RSV, respiratory syncytial virus; MeV, measles virus; padj, Adjusted p-value.



analyzed for post-transcriptional modifications using Chimira (Vitsios and Enright, 2015). Total counts of internal, 3' arm and 5' arm modifications per sample type are shown. Differences in the number of modifications were analyzed by 2-way ANOVA with *post hoc* Dunnet's test. *p*-values are indicated by ** <0.01; ns, not significant. Error bars represent \pm SD from three replicate measurements. 3p ARM, 3' end of miRNA; 5P ARM, 5' arm of miRNA.

or 24 hpi and ligated with 3' and 5' RNA adaptors using T4 RNA ligase 2 (Epicenter Biotechnologies, Madison, WI, United States) at 28° C for 60 min. Ligated RNA was reverse

transcribed using proprietary reverse primers and the firststrand cDNA was amplified with a combination of a proprietary forward primer and reverse primers containing unique index bar codes per the manufacturer's instructions. Amplicons were run on a Tapestation 2200 (Agilent, Santa Clara, CA, United States) high-sensitivity DNA chip to visualize products. Indexed samples were pooled per the manufacturer's protocol and electrophoresed on a 6% TBE-urea PAGE gel. Bands corresponding to 147 and 157 bp corresponding to adaptorligated mature miRNA and other non-coding RNAs were cut out and cDNAs were precipitated. Libraries were validated on a Tapestation 2200 (Agilent) using a high-sensitivity DNA chip. Libraries were denatured for 5 min using 5 M NaOH and loaded onto a MiSeq (Agilent) for sequencing by synthesis. Reads were trimmed to remove adaptor sequences and then analyzed for miRNA differential expression using Chimira (Vitsios and Enright, 2015) and OASIS 2.0 (Rahman et al., 2018a). Read counts were normalized to a total number of reads per sample and then differential expression was calculated using DESeq2 (Love et al., 2014). All reads used for analysis had quality scores ($Q_{-} > 30$. This represents less than or equal to 1 error per 1,000 nts corresponding to a 99.99% accuracy.









miR Analysis

The host genes and pathways predicted to be regulated by miRs were determined by DIANA miRPath analysis (Vlachos et al., 2015). miRs were examined using the DIANA-miRPath server and targets extracted from the Tarbase database (Karagkouni et al., 2018). Enriched KEGG pathway analysis was used to determine intersections (Kanehisa et al., 2016, 2017). We applied a *p*-value threshold of \leq 0.05 along with a false-discovery rate correction.

Virus Quantitation

QRT-PCR of RSV M, or MeV M, and N gene primer-probes were examined for each replicate to confirm infection of SHS cells. Sequences of the primers/probes used were RSV M forward 5'- ACACCCAAGGGACCTTCACTAAGA, RSV M reverse 5'-GCCTTGATTTCACAGGGTGTGGTT-3', RSV M probe/56-FAM/AAG TGC AGT GCT AGC ACA AAT GCC CA/3BHQ_1. For MeV, primer sequences were forward primer MeVEdNPF-5'-CAAGACCCTGAGGGATTCAA-3' and reverse primer MeVEdNPR-5'-CTCTCGCATCACTTGCTCTG-3. For both viruses, 100 ng of RNA was reverse transcribed using Lunascript RT Supermix (New England Biolabs, Ipswitch, MA, United States) using the following conditions: primer annealing at 25°C for 2 min, reverse transcription at 55°C for 30 min followed by heat-inactivation of reverse transcriptase at 95°C for 1 min. Undiluted cDNA was used for PCR. PCR reactions contained cDNA, 0.4 µM of forward and reverse oligomers, 0.8 μM of the probe, 2X Luna Universal probe master mix (New England Biolabs) and were amplified as follows: initial denaturation at 95°C for 60 s, 40 cycles of denaturation at 95°C for 10 s, and combined annealing and extension at 60°C for 30 s. For MeV NP amplification, PCR reactions contained cDNA, 10 µM of forward and



reverse primers, Q5 2X High-Fidelity master mix (New England Biolabs), and was amplified using the conditions: initial denaturation at 98° C for 30 s, 25 cycles of 98° C for 10 s, annealing at 65° C for 30 s and extension at 72° C for 2 min, and a final extension at 72° C for 2 min. MeV NP amplicons were resolved on a 1% agarose gel in low osmolality buffer at 5 V/cm and imaged using a Protein Simple imager (Invitrogen).

RESULTS

sncRNAs and Virus-Infected SHS Cells

Previous studies from our laboratory have shown that RSV infection of A549 cells modifies miR expression (Bakre et al., 2012, 2015, 2017) however, information regarding miR expression in RSV- or MeV-infected neuronal SHS cells is inadequate (Naaman et al., 2017; Cakmak Genc et al., 2018;

Haralambieva et al., 2018). Neuronal cells are not the primary cell type infected by RSV, but they can be infected and may act as a reservoir of infection. Mechanisms that contribute to this are poorly understood. Thus, we examined SHS cells that were RSV- or MeV-infected (MOI = 1.0), or mock-treated and sncRNA expression determined at 12 and 24 hpi by Illumina NGS. We confirmed RSV infection of SHS neuronal cells using RSV-GFP. SHS cells were infected with RSV-GFP at a MOI = 0.1 (**Supplementary Figure 1**). At 48 h and 72 hpi, virus replication was evident using combined dark field and fluorescence microscopy (UV Eclipse Ti2000-U microscope, Nikon).

It is important to note that production of infectious viral particles is restricted in the neuronal environment and viral spread is mediated by fusogenic events leading to syncytia formation that are common to both MeV and RSV infection (Sato et al., 2018). Total reads expressed in a text-based format (fastq) were used and the corresponding reads passing quality scores (Q > 30) are listed in Table 1. Raw fastq reads were mapped to the human genome, and miRs using Chimira (Vitsios and Enright, 2015) and Oasis 2.0 (Rahman et al., 2018a) were normalized to the total read counts and used to calculate differential expression using DESeq 2.0 (Love et al., 2014) relative to mock-treated cells. We used principal component analysis (PCA) to determine sample clustering based on similarity for miR expression profiling. PCA is a method of taking combinations of measured variables and lowering the dimensionality of the data to find the principal source of variance. As the human genome encodes > 2,300miRs and we examined RSV and MeV and two time points, the dataset had > 9,200 variables. PCA identified two major component dimensions-PC1 consists of the time component in addition to virus and miRNA expression, and PC2 is the infection state. PC1 and PC2 are principal components and organized by the percent of variability they explain. Being derived from combinations of variables, PCs do not have labels/units. Samples 1-3 (12 h MeV), 4-6 (12 h RSV), 7-9 (24 h MeV), and 10-12 (24 h RSV) clustered together and away from mock infected cells (samples 13-14) suggesting that the sncRNA response to MeV or RSV infection is distinct from mock infected cells. In addition, clustering of 12 and 24 h MeV and RSV datasets suggests that infections elicit temporal responses. The PCA results indicate that the time post infection is a major component driving differential miRNA expression as shown on PC1 (x-axis with 54% variance) compared to pathogen (PC2 y-axis 13% variance). This is evident in the clustering among mock-treated vs. virus-infected cells (Figure 1). Similarly, miR expression profiles in RSV- or MeV-infected SHS cells at 12 and 24 hpi revealed distinct clustering. Most RSV samples clustered separately from MeV at each timepoint (Figure 1).

The data were filtered to contain only read numbers > 5 (**Table 1**), and those having *p*-values ≤ 0.05 , and fold-change ≤ 1.5 or ≥ 1.5 were used to help identify downregulated vs. upregulated miRs. RSV or MeV infection was confirmed by qPCR. The chief deregulated miRs expressed in SHS cells following RSV- or MeV-infection are shown in **Table 2**.

|--|

Codon	RSV_NS1	RSV_NS2	RSV_NP	RSV_P	RSV_M	RSV_SH	RSV_G	RSV_F	RSV_M2	RSV_L
GUU (V)	0.8	0	1.2	0.9	0.7	0	0	2.0	1.3	1.6
GUG (V)	1.2	0	1.0	2.2	1.4	0	0.5	0.4	0.8	0.6
GGC (G)	1.14	0	0.8	0	0.7	0	0.7	0.7	0	0.5
GGG (G)	0.57	2.0	0.8	1.7	0.7	4.0	0.7	0.3	0.4	0.5
Codon	MeV_N	MeV_P	MeV_M	MeV_F	MeV_H	MeV_L				
GUU (V)	2.0	0.9	0.6	0.9	0.6	1.9				
GUG (V)	0.2	2.0	2.3	0.65	1.6	0.7				
GGC (G)	0.7	0.6	1.1	0.8	0.5	1.0				
GGG (G)	0.5	1.3	0.4	1.4	1.8	0.8				

RSCU scores for individual RSV and MeV genes were determined using MEGA X (Kumar et al., 2018).

RSV_NS1, RSV Non-structural protein 1; RSV_NS2, RSV Non-structural gene 2; RSV_NP, RSV Nucleoprotein; RSV_P, RSV Phosphoprotein; RSV_M, RSV Matrix protein; RSV_SH, RSV Small hydrophobic protein; RSV_G, RSV glycoprotein; RSV_F, RSV Fusion protein; RSV_M2, RSV Matrix ORF2 protein; RSV_L, RSV Large polymerase protein; MeV_N, Measles virus Nucleocapsid protein; MeV_P, Measles virus Phosphoprotein; MeV_M, Measles virus Matrix protein; MeV_F, Measles virus polymerase.

Compared to mock-infection, expression of miR-10b-5p, miR-181c-5p, and miR-125b-5p were strikingly downregulated in SHS cells at 12 or 24 hpi following RSV or MeV infection. In contrast, expression of p-miR-330, p-miR-338, p-miR-317, p-miR-106, p-miR-113, p-miR-256, p-miR-232, p-miR-103 and miR-182-5p, miR-4532, miR-1273g-3p, miR-483-3p, miR-3141, miR-1246 miR-3196, miR-4516, and miR-4488 were substantially upregulated in SHS cells.

Studies suggest that modifications involving Adenosine to Inosine (A- > I) or Cytosine to Uracil (C- > U) within miRs, N6 methylation, and oxidation affect miR stability, maturation and target specificity (Alarcon et al., 2015; Wang et al., 2015; Correia De Sousa et al., 2019). Posttranscriptional modifications to miRNAs have been shown to alter their binding and function (Pirouz et al., 2019; Briand et al., 2020; Cheray et al., 2020). We therefore examined epitranscriptional modifications of the deregulated miRs using Chimira (**Figure 2**; Vitsios and Enright, 2015). Most modifications were located on the 3p arms of the deregulated miRs. RSV infection in SHS cells decreased the abundance of 3p modifications at 12 hpi (**Figure 2** and **Supplementary Table 1**).

RSV or MeV Infection Induces tRNA Fragments

Cellular stress triggers the degradation of tRNA to tRFs (Shen et al., 2018). tRFs may encompass the 5', 3', or internal anti-codon loop and are classified as 5'-, 3-, or class I tRFs (Shen et al., 2018). Functions of tRFs are poorly understood, but tRFs can perform as siRNAs degrading target transcripts (Maute et al., 2013), and regulate ribosomal loading and protein chain elongation (Sobala and Hutvagner, 2013). The biosynthesis of tRFs involves degradation of pre-tRNA molecules *via* the nuclear TRAMP pathway (Anderson, 2005; Maraia and Lamichhane, 2011) or cytosolic degradation of mature tRNAs *via* the rapid tRNA decay pathway. Analysis of the small RNA sequencing dataset in this

study showed a differing abundance of tRFs following RSV or MeV infection. Analysis of tRFs showed that among the most abundant were 5'tRFs derived from tRNAs for glycine and valine (Figure 3), suggesting that tRNAs for these two amino acids were targeted during RSV and MeV infection. Levels of 5'tRFs for glycine and valine were substantially increased at 24 hpi following RSV or MeV infection. Frequencies of glycine and valine 5'tRFs increased with time, i.e., from 12 to 24 hpi in RSV- and MeV-infected SHS cells. Analysis of the anti-codon sequence for these tRFs showed that GCC, AAC, and CAC anticodons are enriched in 5'tRFs in SHS cells following RSV or MeV infection (Figure 4). A previous report showed that 5'tRF-Glu-CTC produced during RSV infection of A549 cells interacted with apolipoprotein E receptor 2 (APOER2) mRNA to inhibit translation of the gene, favoring RSV replication (Deng et al., 2015).

We postulated that tRFs had a direct effect on translation and compared tRF profiles in RSV- or MeV-infected SHS cells with synonymous codon usage. Our analysis compared how frequently each codon in the RSV or MeV genome is utilized relative to the standard genetic code frequencies that are given as a codon adaptation index (CAI). CAI ranges from 0 to 1.0, with 0 representing least used, and 1.0 representing most common codons. A comparison of RSCU profiles for RSV and MeV genomes using Mega X (Kumar et al., 2018) showed the codon usage to be considerably different than standard human codon usage frequencies (Figures 5A,B). The assessment showed that of the four possible codons that encode glycine (GGU, GGC, GGA, GGG), three codons (GGC, GGU, GGG) are under-utilized for RSV genes, and valine, RSV uses the GUC codon less frequently. For MeV, two of the four codons for glycine (GGU, GGC) and GUA for valine are underutilized. The consequence of these findings is unknown but suggests that the tRNAs used during RSV or MeV infection are selectively processed to produce these tRNAs potentially for two reasons-to reduce host translation (Kim et al., 2017, 2019; Mesitov et al., 2017), and/or to function as siRNAs and

silence host genes (Emara et al., 2010; Wang et al., 2013; Kuscu et al., 2018; Jehn et al., 2020). For valine, and the increased abundance of valine 5'tRFs (AAC and CAC), the corresponding codon usage was not low suggesting that valine tRFs might have functions other than reducing translation rates. RSCU analysis of individual RSV and MeV genes showed that glycine and valine tRNAs were under-utilized following RSV or MeV infection in SHS cells (**Table 3**). Together these findings suggest that tRNA cleavage is linked with codon usage. The sncRNA results show that the expression profiles for small RNAs differ in kinetics of expression for RSV and MeV infection of SHS cells.

DISCUSSION

In this study, we examined differentially expressed sncRNAs following RSV or MeV infection of SHS cells at 12 and 24 hpi using NGS. NGS showed unique miR expression profiles following RSV or MeV infection (Figure 1 and Table 2). We identified that (1) miR-10b-5p was expressed in mock-treated cells but showed reduced expression in RSV- or MeV-infected cells, (2) p-miR-247 and miR-375 had robust expression in RSV- or MeV-infected cells, (3) miRs had no/low expression in mock-treated cells, but were increased by RSV or MeV infection, and (4) miRs were not expressed in mock-infected cells at 12 hpi but showed robust copy numbers at 24 hpi. These findings show that RSV or MeV infection induces distinct miR expression in SHS cells that is virus-type and time-point specific. While the sequencing data from our study has identified these deregulated miRNAs, the mechanisms contributing to their biogenesis remains unknown and is out of the scope of this manuscript.

Evidence suggests that epitranscriptomics (or posttranscriptional modification of cellular RNAs) especially 3' end uridylation and adenylation of miRs can alter both miRNA stability and target repertoire (Vitsios and Enright, 2015). We explored the nature of epitranscriptional miR modifications following RSV or MeV infection. The findings suggest that RSV and MeV induce sncRNA modifications affecting target cell specificity. The results showed extensive 3' modifications (mostly uridylation and adenylation) in the miRs following RSV or MeV infection that may alter host mRNA or miRNA transcript stability. Further studies are needed to elucidate the function of these epitranscriptional modifications on miRNA function.

Post-transcriptional regulatory mechanisms control translation of >60% of the human transcriptome (Friedman et al., 2009). The identification of cleaved tRNAs during RSV or MeV infection of SHS cells is novel. The tRFs identified contained exclusively of 5'tRFs during later stages of RSV or MeV infection, and of these tRNAs, glycine, and valine were the primary targets for 5'tRF formation. Other studies showed that RSV infection can induce angiogenin (ANG)-mediated cleavage of tRNAs to produce tRFs (Deng et al., 2015; Zhou J. et al., 2017). These studies identified a pro-viral role for 5'tRF-GlyCCC, 5'tRF-LysCTT, and 5'tRF-GluCTC by depression of APOER2 mRNA. We postulate that tRNA cleavage to tRFs

may have a broader anti-viral role. Degradation of host tRNAs may reduce the anti-viral protein response, and recent findings from proteomic studies in RSV-infected A549 cells have shown that the host nuclear proteins are considerably reduced following RSV infection (Munday et al., 2012, 2015). Similar results have also been shown for MeV, an attribute linked to the N protein (Sato et al., 2007; Okonski and Samuel, 2013). It is important to investigate why RSV or MeV infection induces selective cleavage of glycine and valine tRNAs as glycine, and valine tRNAs are selectively under-utilized by most RSV and MeV genes (**Table 3**). Transfer RNA fragments operate in mucosal immunity (Chen and Shen, 2021) regulating retrotransposon expression (Schorn et al., 2017) and viral adaptation to the host (Lauring et al., 2012; Pavon-Eternod et al., 2013; Rahman et al., 2018b; Le Nouën et al., 2020; Nunes et al., 2020).

The degradation of tRNAs may be a host stress response to viral infection, however, the increases in tRNA for glycine and valine hint that MeV and RSV may potentially target these tRNAs. The mechanisms that cause are unknown and are difficult to resolve in the context of this study. Currently, we do not understand the mechanisms involved in sncRNA deregulation, nor the functional outcome, thus further studies are necessary. Translation of mRNAs linked to the anti-viral response is key to control of virus infection and clearance. The findings show that RSV or MeV infection induces several classes of sncRNAs that regulate post-transcriptional gene expression and emphasizes the need to better understand the host-virus interface.

DATA AVAILABILITY STATEMENT

The data presented in the study are deposited in Gene Expression Omnibus (GEO) with accession numbers GSE167724 and GSM5112024–GSM-5112037.

AUTHOR CONTRIBUTIONS

AB contributed to conception, execution, and data analysis, and wrote manuscript drafts and revisions. CD contributed to execution. SC and RT contributed to study design, funding, data interpretation, manuscript writing, and review. HA contributed to validating infection using dark field fluorescence microscopy. All authors have seen the draft of the manuscript.

FUNDING

The studies were supported by a grant awarded to SC and RT (IE140652) to study "Paramyxovirus miRNA biogenesis and implications for host and viral gene expression" and the Georgia Research Alliance Foundation for RT.

ACKNOWLEDGMENTS

The authors would like to acknowledge the kind gift of rgRSV224GFP from Michael Teng at University of South Florida (USF).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fmicb. 2021.671852/full#supplementary-material

Supplementary Figure 1 | RSV infects SHS neuronal cells. SHS cells were infected with rgRSV-GFP at MOI = 0.1 for indicated time points. Viral replication

REFERENCES

- Advani, V. M., and Ivanov, P. (2019). Translational Control under Stress: Reshaping the Translatome. *Bioessays* 41:e1900009.
- Alarcon, C. R., Lee, H., Goodarzi, H., Halberg, N., and Tavazoie, S. F. (2015). N6-methyladenosine marks primary microRNAs for processing. *Nature* 519, 482–485. doi: 10.1038/nature14281
- Anderson, J. T. (2005). RNA turnover: unexpected consequences of being tailed. *Curr. Biol.* 15, R635–R638.
- Bakre, A. A., Harcourt, J. L., Haynes, L. M., Anderson, L. J., and Tripp, R. A. (2017). The Central Conserved Region (CCR) of Respiratory Syncytial Virus (RSV) G Protein Modulates Host miRNA Expression and Alters the Cellular Response to Infection. *Vaccines* 5:16. doi: 10.3390/vaccines503 0016
- Bakre, A. A., Maleki, A., and Tripp, R. A. (2019). MicroRNA and Nonsense Transcripts as Putative Viral Evasion Mechanisms. *Front. Cell Infect. Microbiol.* 9:152. doi: 10.3389/fcimb.2019.00152
- Bakre, A., Mitchell, P., Coleman, J. K., Jones, L. P., Saavedra, G., Teng, M., et al. (2012). Respiratory syncytial virus modifies microRNAs regulating host genes that affect virus replication. J. Gen. Virol. 93, 2346–2356. doi: 10.1099/vir.0. 044255-0
- Bakre, A., Wu, W., Hiscox, J., Spann, K., Teng, M. N., and Tripp, R. A. (2015). Human respiratory syncytial virus non-structural protein NS1 modifies miR-24 expression via transforming growth factor-beta. J. Gen. Virol. 96, 3179–3191. doi: 10.1099/jgv.0.000261
- Bartel, D. P. (2009). MicroRNAs: Target Recognition and Regulatory Functions. Cell 136, 215–233. doi: 10.1016/j.cell.2009.01.002
- Bergeron, H. C., Murray, J., Nunez Castrejon, A. M., Dubois, R. M., and Tripp, R. A. (2021). Respiratory Syncytial Virus (RSV) G Protein Vaccines With Central Conserved Domain Mutations Induce CX3C-CX3CR1 Blocking Antibodies. *Viruses* 13:352. doi: 10.3390/v13020352
- Bernier, A., and Sagan, S. M. (2018). The Diverse Roles of microRNAs at the Host(-)Virus Interface. Viruses 10:440. doi: 10.3390/v10080440
- Bohmwald, K., Galvez, N. M. S., Rios, M., and Kalergis, A. M. (2018). Neurologic Alterations Due to Respiratory Virus Infections. *Front. Cell Neurosci.* 12:386. doi: 10.3389/fncel.2018.00386
- Briand, J., Sérandour, A. A., Nadaradjane, A., Bougras-Cartron, G., Heymann, D., Ory, B., et al. (2020). N6-Adenosine Methylation of miRNA-200b-3p Influences Its Functionality and Is a Theranostic Tool. *Mol. Ther. Nucleic Acids* 22, 72–83. doi: 10.1016/j.omtn.2020.08.010
- Cakmak Genc, G., Dursun, A., Karakas Celik, S., Calik, M., Kokturk, F., and Piskin, I. E. (2018). IL28B, IL29 and micro-RNA 548 in subacute sclerosing panencephalitis as a rare disease. *Gene* 678, 73–78. doi: 10.1016/j.gene.2018.07.062
- Chen, Y., and Shen, J. (2021). Mucosal immunity and tRNA, tRF, and tiRNA. J. Mol. Med. 99, 47–56. doi: 10.1007/s00109-020-02008-4
- Cheray, M., Etcheverry, A., Jacques, C., Pacaud, R., Bougras-Cartron, G., Aubry, M., et al. (2020). Cytosine methylation of mature microRNAs inhibits their functions and is associated with poor prognosis in glioblastoma multiforme. *Mol. Cancer* 19:36.
- Choi, Y., Mason, C. S., Jones, L. P., Crabtree, J., Jorquera, P. A., and Tripp, R. A. (2012). Antibodies to the central conserved region of respiratory syncytial virus (RSV) G protein block RSV G protein CX3C-CX3CR1 binding and cross-neutralize RSV A and B strains. *Viral Immunol.* 25, 193–203.
- Correia De Sousa, M., Gjorgjieva, M., Dolicka, D., Sobolewski, C., and Foti, M. (2019). Deciphering miRNAs' Action through miRNA Editing. *Int. J. Mol. Sci.* 20:6249. doi: 10.3390/ijms20246249

was visualized by dark field fluorescence microscopy on a Nikon UV Eclipse TE2000-U microscope at 10X magnification.

Supplementary Table 1 | MiRNA modification data patterns in RSV and MeV infected neuronal cells. Table lists samples, microRNAs, type of modifications, modification site, pattern, modification position, internal modification type, internal modification position, repeat modifications and counts of modifications for all deregulated miRNAs identified in the study.

- Cullen, B. R. (2013). MicroRNAs as mediators of viral evasion of the immune system. Nat. Immunol. 14, 205–210. doi: 10.1038/ni.2537
- Dahiya, N., and Atreya, C. D. (2014). MicroRNAs and Major Bloodborne Infectious Viral Diseases. *Microrna* 2, 212–218. doi: 10.2174/ 2211536602666131118224225
- Dakhama, A., Vitalis, T. Z., and Hegele, R. G. (1997). Persistence of respiratory syncytial virus (RSV) infection and development of RSV-specific IgG1 response in a guinea-pig model of acute bronchiolitis. *Eur. Respir. J.* 10, 20–26. doi: 10.1183/09031936.97.10010020
- Deng, J., Ptashkin, R. N., Chen, Y., Cheng, Z., Liu, G., Phan, T., et al. (2015). Respiratory Syncytial Virus Utilizes a tRNA Fragment to Suppress Antiviral Responses Through a Novel Targeting Mechanism. *Mol. Ther.* 23, 1622–1629. doi: 10.1038/mt.2015.124
- Eisenhut, M. (2006). Extrapulmonary manifestations of severe respiratory syncytial virus infection–a systematic review. Crit. Care 10:R107.
- Emara, M. M., Ivanov, P., Hickman, T., Dawra, N., Tisdale, S., Kedersha, N., et al. (2010). Angiogenin-induced tRNA-derived stress-induced RNAs promote stress-induced stress granule assembly. *J. Biol. Chem.* 285, 10959–10968. doi: 10.1074/jbc.m109.077560
- Ferren, M., Horvat, B., and Mathieu, C. (2019). Measles Encephalitis: Towards New Therapeutics. Viruses 11:1017. doi: 10.3390/v11111017
- Friedman, R. C., Farh, K. K., Burge, C. B., and Bartel, D. P. (2009). Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* 19, 92–105. doi: 10.1101/gr.082701.108
- Ganesan, P., Chandwani, M. N., Creisher, P. S., Bohn, L., and O'donnell, L. A. (2018). The neonatal anti-viral response fails to control measles virus spread in neurons despite interferon-gamma expression and a Th1-like cytokine profile. *J. Neuroimmunol.* 316, 80–97. doi: 10.1016/j.jneuroim.2017.12.018
- Grassmann, R., and Jeang, K. T. (2008). The roles of microRNAs in mammalian virus infection. *Biochim. Biophys. Acta* 1779, 706–711. doi: 10.1016/j.bbagrm. 2008.05.005
- Griffin, D. E. (2018). Measles Vaccine. Viral Immunol. 31, 86-95.
- Griffin, D. E., Lin, W. W., and Nelson, A. N. (2018). Understanding the causes and consequences of measles virus persistence. *F1000Res* 7:237. doi: 10.12688/ f1000research.12094.1
- Haber, N. (2018). Respiratory syncytial virus infection in elderly adults. Med. Mal. Infect. 48, 377–382. doi: 10.1016/j.medmal.2018.01.008
- Hallak, L. K., Kwilas, S. A., and Peeples, M. E. (2007). Interaction between respiratory syncytial virus and glycosaminoglycans, including heparan sulfate. *Methods Mol. Biol.* 379, 15–34. doi: 10.1385/1-59745-393-5:15
- Hallak, L. K., Spillmann, D., Collins, P. L., and Peeples, M. E. (2000). Glycosaminoglycan sulfation requirements for respiratory syncytial virus infection. J. Virol. 74, 10508–10513. doi: 10.1128/jvi.74.22.10508-10513. 2000
- Haralambieva, I. H., Kennedy, R. B., Simon, W. L., Goergen, K. M., Grill, D. E., Ovsyannikova, I. G., et al. (2018). Differential miRNA expression in B cells is associated with inter-individual differences in humoral immune response to measles vaccination. *PLoS One* 13:e0191812. doi: 10.1371/journal.pone.0191812
- Harcourt, J. L., Karron, R. A., and Tripp, R. A. (2004). Anti-G protein antibody responses to respiratory syncytial virus infection or vaccination are associated with inhibition of G protein CX3C-CX3CR1 binding and leukocyte chemotaxis. *J. Infect. Dis.* 190, 1936–1940. doi: 10.1086/425516
- Hardie, D. R., Albertyn, C., Heckmann, J. M., and Smuts, H. E. (2013). Molecular characterisation of virus in the brains of patients with measles inclusion body encephalitis (MIBE). *Virol J.* 10:283.
- Hasegawa, K., Perez-Losada, M., Hoptay, C. E., Epstein, S., Mansbach, J. M., Teach, S. J., et al. (2018). RSV vs. rhinovirus bronchiolitis: difference in nasal

airway microRNA profiles and NFkappaB signaling. *Pediatr. Res.* 83, 606–614. doi: 10.1038/pr.2017.309

- Haynes, L. M., Jones, L. P., Barskey, A., Anderson, L. J., and Tripp, R. A. (2003). Enhanced disease and pulmonary eosinophilia associated with formalininactivated respiratory syncytial virus vaccination are linked to G glycoprotein CX3C-CX3CR1 interaction and expression of substance P. J. Virol. 77, 9831– 9844. doi: 10.1128/jvi.77.18.9831-9844.2003
- Inchley, C. S., Sonerud, T., Fjaerli, H. O., and Nakstad, B. (2015). Nasal mucosal microRNA expression in children with respiratory syncytial virus infection. *BMC Infect. Dis.* 15:150. doi: 10.1186/s12879-015-0878-z
- Ivanov, P., Emara, M. M., Villen, J., Gygi, S. P., and Anderson, P. (2011). Angiogenin-induced tRNA fragments inhibit translation initiation. *Mol. Cell* 43, 613–623. doi: 10.1016/j.molcel.2011.06.022
- Jehn, J., Treml, J., Wulsch, S., Ottum, B., Erb, V., Hewel, C., et al. (2020). 5' tRNA halves are highly expressed in the primate hippocampus and might sequence-specifically regulate gene expression. RNA 26, 694–707. doi: 10.1261/ rna.073395.119
- Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y., and Morishima, K. (2017). KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res.* 45, D353–D361.
- Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., and Tanabe, M. (2016). KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res.* 44, D457–D462.
- Karagkouni, D., Paraskevopoulou, M. D., Chatzopoulos, S., Vlachos, I. S., Tastsoglou, S., Kanellos, I., et al. (2018). DIANA-TarBase v8: a decade-long collection of experimentally supported miRNA-gene interactions. *Nucleic Acids Res.* 46, D239–D245.
- Kim, H. K., Fuchs, G., Wang, S., Wei, W., Zhang, Y., Park, H., et al. (2017). A transfer-RNA-derived small RNA regulates ribosome biogenesis. *Nature* 552, 57–62. doi: 10.1038/nature25005
- Kim, H. K., Xu, J., Chu, K., Park, H., Jang, H., Li, P., et al. (2019). A tRNA-Derived Small RNA Regulates Ribosomal Protein S28 Protein Levels after Translation Initiation in Humans and Mice. *Cell Rep.* 29, 3816.e–3824.e.
- King, K. A., Hu, C., Rodriguez, M. M., Romaguera, R., Jiang, X., and Piedimonte, G. (2001). Exaggerated neurogenic inflammation and substance P receptor upregulation in RSV-infected weanling rats. *Am. J. Respir. Cell Mol. Biol.* 24, 101–107. doi: 10.1165/ajrcmb.24.2.4264
- Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* 35, 1547–1549. doi: 10.1093/molbev/msy096
- Kuscu, C., Kumar, P., Kiran, M., Su, Z., Malik, A., and Dutta, A. (2018). tRNA fragments (tRFs) guide Ago to regulate gene expression post-transcriptionally in a Dicer-independent manner. *RNA* 24, 1093–1105. doi: 10.1261/rna.066126. 118
- Lauring, A. S., Acevedo, A., Cooper, S. B., and Andino, R. (2012). Codon usage determines the mutational robustness, evolutionary capacity, and virulence of an RNA virus. *Cell Host Microbe* 12, 623–632. doi: 10.1016/j.chom.2012.10.008
- Lay, M. K., Bueno, S. M., Galvez, N., Riedel, C. A., and Kalergis, A. M. (2016). New insights on the viral and host factors contributing to the airway pathogenesis caused by the respiratory syncytial virus. *Crit. Rev. Microbiol.* 42, 800–812.
- Le Nouën, C., Luongo, C. L., Yang, L., Mueller, S., Wimmer, E., Dinapoli, J. M., et al. (2020). Optimization of the Codon Pair Usage of Human Respiratory Syncytial Virus Paradoxically Resulted in Reduced Viral Replication In Vivo and Reduced Immunogenicity. J. Virol. 94, e1296–e1219.
- Leung, A. K., Hon, K. L., Leong, K. F., and Sergi, C. M. (2018). Measles: a disease often forgotten but not gone. *Hong Kong Med. J.* 24, 512–520.
- Li, X. Q., Fu, Z. F., Alvarez, R., Henderson, C., and Tripp, R. A. (2006). Respiratory syncytial virus (RSV) infects neuronal cells and processes that innervate the lung by a process involving RSV G protein. *J. Virol.* 80, 537–540. doi: 10.1128/ jvi.80.1.537-540.2006
- Lin, L. T., and Richardson, C. D. (2016). The Host Cell Receptors for Measles Virus and Their Interaction with the Viral Hemagglutinin (H) Protein. *Viruses* 8:250. doi: 10.3390/v8090250
- Long, X., Xie, J., Zhao, K., Li, W., Tang, W., Chen, S., et al. (2016). NK cells contribute to persistent airway inflammation and AHR during the later stage of RSV infection in mice. *Med. Microbiol. Immunol.* 205, 459–470. doi: 10.1007/ s00430-016-0459-9

- Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:550.
- Maraia, R. J., and Lamichhane, T. N. (2011). 3' processing of eukaryotic precursor tRNAs. *Wiley Interdiscip. Rev. RNA* 2, 362–375. doi: 10.1002/wrna.64
- Maute, R. L., Schneider, C., Sumazin, P., Holmes, A., Califano, A., Basso, K., et al. (2013). tRNA-derived microRNA modulates proliferation and the DNA damage response and is down-regulated in B cell lymphoma. *Proc. Natl. Acad. Sci. U S A*. 110, 1404–1409. doi: 10.1073/pnas.1206761110
- Mazur, N. I., Higgins, D., Nunes, M. C., Melero, J. A., Langedijk, A. C., Horsley, N., et al. (2018). The respiratory syncytial virus vaccine landscape: lessons from the graveyard and promising candidates. *Lancet Infect. Dis.* 18, e295–e311.
- McQuaid, S., Cosby, S. L., Koffi, K., Honde, M., Kirk, J., and Lucas, S. B. (1998). Distribution of measles virus in the central nervous system of HIV-seropositive children. Acta Neuropathol. 96, 637–642. doi: 10.1007/s004010050945
- Mesitov, M. V., Soldatov, R. A., Zaichenko, D. M., Malakho, S. G., Klementyeva, T. S., Sokolovskaya, A. A., et al. (2017). Differential processing of small RNAs during endoplasmic reticulum stress. *Sci. Rep.* 7:46080.
- Mirra, V., Ullmann, N., Cherchi, C., Onofri, A., Paglietti, M. G., and Cutrera, R. (2018). Respiratory syncytial virus prophylaxis and the "special population". *Minerva Pediatr.* 70, 589–599.
- Mishra, B., and Wang, G. (2012). The Importance of Amino Acid Composition in Natural AMPs: An Evolutional, Structural, and Functional Perspective. *Front. Immunol.* 3:221. doi: 10.3389/fimmu.2012.00221
- Morichi, S., Morishita, N., Ishida, Y., Oana, S., Yamanaka, G., Kashiwagi, Y., et al. (2017). Examination of neurological prognostic markers in patients with respiratory syncytial virus-associated encephalopathy. *Int. J. Neurosci.* 127, 44–50. doi: 10.3109/00207454.2016.1138951
- Munday, D. C., Howell, G., Barr, J. N., and Hiscox, J. A. (2015). Proteomic analysis of mitochondria in respiratory epithelial cells infected with human respiratory syncytial virus and functional implications for virus and cell biology. *J. Pharm. Pharmacol.* 67, 300–318.
- Munday, D. C., Surtees, R., Emmott, E., Dove, B. K., Digard, P., Barr, J. N., et al. (2012). Using SILAC and quantitative proteomics to investigate the interactions between viral and host proteomes. *Proteomics* 12, 666–672. doi: 10.1002/pmic. 201100488
- Naaman, H., Rall, G., Matullo, C., Veksler-Lublinsky, I., Shemer-Avni, Y., and Gopas, J. (2017). MiRNA-124 is a link between measles virus persistent infection and cell division of human neuroblastoma cells. *PLoS One* 12:e0187077. doi: 10.1371/journal.pone.0187077
- Nunes, A., Ribeiro, D. R., Marques, M., Santos, M. A. S., Ribeiro, D., and Soares, A. R. (2020). Emerging Roles of tRNAs in RNA Virus Infections. *Trends Biochem. Sci.* 45, 794–805. doi: 10.1016/j.tibs.2020.05.007
- Okonski, K. M., and Samuel, C. E. (2013). Stress granule formation induced by measles virus is protein kinase PKR dependent and impaired by RNA adenosine deaminase ADAR1. J. Virol. 87, 756–766. doi: 10.1128/jvi.02270-12
- Omar, S., Clarke, R., Abdullah, H., Brady, C., Corry, J., Winter, H., et al. (2017). Respiratory virus infection up-regulates TRPV1, TRPA1 and ASICS3 receptors on airway cells. *PLoS One* 12:e0171681. doi: 10.1371/journal.pone.0171681
- Oshansky, C. M., Krunkosky, T. M., Barber, J., Jones, L. P., and Tripp, R. A. (2009). Respiratory syncytial virus proteins modulate suppressors of cytokine signaling 1 and 3 and the type I interferon response to infection by a toll-like receptor pathway. *Viral Immunol.* 22, 147–161. doi: 10.1089/vim.2008.0098
- Paes, B. (2018). Respiratory Syncytial Virus in Otherwise Healthy Prematurely Born Infants: A Forgotten Majority. Am. J. Perinatol. 35, 541–544. doi: 10.1055/ s-0038-1637762
- Pavon-Eternod, M., David, A., Dittmar, K., Berglund, P., Pan, T., Bennink, J. R., et al. (2013). Vaccinia and influenza A viruses select rather than adjust tRNAs to optimize translation. *Nucleic Acids Res.* 41, 1914–1921. doi: 10.1093/nar/gks986
- Petrov, A. I., Kay, S. J. E., Kalvari, I., Howe, K. L., Gray, K. A., Bruford, E. A., et al. (2017). RNAcentral: a comprehensive database of non-coding RNA sequences. *Nucleic Acids Res.* 45, D128–D134.
- Pirouz, M., Ebrahimi, A. G., and Gregory, R. I. (2019). Unraveling 3'-end RNA uridylation at nucleotide resolution. *Methods* 155, 10–19. doi: 10.1016/j.ymeth. 2018.10.024
- Rahman, R.-U., Gautam, A., Bethune, J., Sattar, A., Fiosins, M., Magruder, D. S., et al. (2018a). Oasis 2: improved online analysis of small RNA-seq data. *BMC Bioinformatics* 19:54. doi: 10.1186/s12859-018-2047-z

- Rahman, S. U., Yao, X., Li, X., Chen, D., and Tao, S. (2018b). Analysis of codon usage bias of Crimean-Congo hemorrhagic fever virus and its adaptation to hosts. *Infect. Genet. Evol.* 58, 1–16. doi: 10.1016/j.meegid.2017.11.027
- Riddell, M. A., Moss, W. J., Hauer, D., Monze, M., and Griffin, D. E. (2007). Slow clearance of measles virus RNA after acute infection. *J. Clin. Virol.* 39, 312–317. doi: 10.1016/j.jcv.2007.05.006
- Rima, B. K., and Duprex, W. P. (2005). Molecular mechanisms of measles virus persistence. Virus Res. 111, 132–147. doi: 10.1016/j.virusres.2005.04.005
- Roizman, B., Zhou, G., and Du, T. (2011). Checkpoints in productive and latent infections with herpes simplex virus 1: conceptualization of the issues. *J. Neurovirol.* 17, 512–517. doi: 10.1007/s13365-011-0058-x
- Rota, P. A., Rota, J. S., and Goodson, J. L. (2017). Subacute Sclerosing Panencephalitis. *Clin. Infect. Dis.* 65, 233–234.
- Russell, C. J., Simoes, E. A. F., and Hurwitz, J. L. (2018). Vaccines for the Paramyxoviruses and Pneumoviruses: Successes, Candidates, and Hurdles. *Viral Immunol.* 31, 133–141. doi: 10.1089/vim.2017.0137
- Sato, H., Masuda, M., Kanai, M., Tsukiyama-Kohara, K., Yoneda, M., and Kai, C. (2007). Measles virus N protein inhibits host translation by binding to eIF3-p40. *J. Virol.* 81, 11569–11576. doi: 10.1128/jvi.00570-07
- Sato, Y., Watanabe, S., Fukuda, Y., Hashiguchi, T., Yanagi, Y., and Ohno, S. (2018). Cell-to-Cell Measles Virus Spread between Human Neurons Is Dependent on Hemagglutinin and Hyperfusogenic Fusion Protein. J. Virol. 92, e2166–e2117.
- Schorn, A. J., Gutbrod, M. J., Leblanc, C., and Martienssen, R. (2017). LTR-Retrotransposon Control by tRNA-Derived Small RNAs. *Cell* 170, 61.e–71.e.
- Schwarze, J., O'donnell, D. R., Rohwedder, A., and Openshaw, P. J. (2004). Latency and persistence of respiratory syncytial virus despite T cell immunity. *Am. J. Respir. Crit. Care Med.* 169, 801–805. doi: 10.1164/rccm.200308-12030c
- Shen, Y., Yu, X., Zhu, L., Li, T., Yan, Z., and Guo, J. (2018). Transfer RNA-derived fragments and tRNA halves: biogenesis, biological functions and their roles in diseases. J. Mol. Med. 96, 1167–1176. doi: 10.1007/s00109-018-1693-y
- Singh, B. K., Li, N., Mark, A. C., Mateo, M., Cattaneo, R., and Sinn, P. L. (2016). Cell-to-Cell Contact and Nectin-4 Govern Spread of Measles Virus from Primary Human Myeloid Cells to Primary Human Airway Epithelial Cells. *J. Virol.* 90, 6808–6817.
- Sobala, A., and Hutvagner, G. (2013). Small RNAs derived from the 5' end of tRNA can inhibit protein translation in human cells. *RNA Biol.* 10, 553–563. doi: 10.4161/rna.24285
- Sofos, N., Xu, K., Dedic, E., and Brodersen, D. E. (2015). Cut to the chase-Regulating translation through RNA cleavage. *Biochimie* 114, 10–17. doi: 10. 1016/j.biochi.2015.01.009
- Tan, Y. R., Yang, T., Liu, S. P., Xiang, Y., Qu, F., Liu, H. J., et al. (2008). Pulmonary peptidergic innervation remodeling and development of airway hyperresponsiveness induced by RSV persistent infection. *Peptides* 29, 47–56. doi: 10.1016/j.peptides.2007.10.020
- Taniguchi, M., Yanagi, Y., and Ohno, S. (2019). Both type I and type III interferons are required to restrict measles virus growth in lung epithelial cells. Arch. Virol. 164, 439–446. doi: 10.1007/s00705-018-4087-0
- Tatsuo, H., Ono, N., Tanaka, K., and Yanagi, Y. (2000). SLAM (CDw150) is a cellular receptor for measles virus. *Nature* 406, 893–897. doi: 10.1038/35022579
- Techaarpornkul, S., Collins, P. L., and Peeples, M. E. (2002). Respiratory syncytial virus with the fusion protein as its only viral glycoprotein is less dependent on cellular glycosaminoglycans for attachment than complete virus. *Virology* 294, 296–304. doi: 10.1006/viro.2001.1340
- Tripp, R. A., Barskey, A., Goss, L., and Anderson, L. J. (2002). Substance P receptor expression on lymphocytes is associated with the immune response to respiratory syncytial virus infection. *J. Neuroimmunol.* 129, 141–153. doi: 10.1016/s0165-5728(02)00169-8
- Tripp, R. A., Dakhama, A., Jones, L. P., Barskey, A., Gelfand, E. W., and Anderson, L. J. (2003). The G glycoprotein of respiratory syncytial virus

depresses respiratory rates through the CX3C motif and substance P. J. Virol. 77, 6580–6584. doi: 10.1128/jvi.77.11.6580-6584.2003

- Tripp, R. A., Jones, L. P., Haynes, L. M., Zheng, H., Murphy, P. M., and Anderson, L. J. (2001). CX3C chemokine mimicry by respiratory syncytial virus G glycoprotein. *Nat. Immunol.* 2, 732–738. doi: 10.1038/90675
- Tripp, R. A., Moore, D., Winter, J., and Anderson, L. J. (2000). Respiratory syncytial virus infection and G and/or SH protein expression contribute to substance P, which mediates inflammation and enhanced pulmonary disease in BALB/c mice. J. Virol. 74, 1614–1622. doi: 10.1128/jvi.74.4.1614-1622.2000
- Vitsios, D. M., and Enright, A. J. (2015). Chimira: analysis of small RNA sequencing data and microRNA modifications. *Bioinformatics* 31, 3365–3367. doi: 10.1093/ bioinformatics/btv380
- Vlachos, I. S., Zagganas, K., Paraskevopoulou, M. D., Georgakilas, G., Karagkouni, D., Vergoulis, T., et al. (2015). DIANA-miRPath v3.0: deciphering microRNA function with experimental support. *Nucleic Acids Res.* 43, W460–W466.
- Wang, J. X., Gao, J., Ding, S. L., Wang, K., Jiao, J. Q., Wang, Y., et al. (2015). Oxidative Modification of miR-184 Enables It to Target Bcl-xL and Bcl-w. *Mol. Cell* 59, 50–61. doi: 10.1016/j.molcel.2015.05.003
- Wang, Q., Lee, I., Ren, J., Ajay, S. S., Lee, Y. S., and Bao, X. (2013). Identification and functional characterization of tRNA-derived RNA fragments (tRFs) in respiratory syncytial virus infection. *Mol. Ther.* 21, 368–379. doi: 10.1038/mt. 2012.237
- Wang, S., Liu, P., Yang, P., Zheng, J., and Zhao, D. (2017). Peripheral blood microRNAs expression is associated with infant respiratory syncytial virus infection. *Oncotarget* 8, 96627–96635. doi: 10.18632/oncotarget.1 9364
- Yis, U., Tufekci, U. K., Genc, S., Carman, K. B., Bayram, E., Topcu, Y., et al. (2015). Expression patterns of micro-RNAs 146a, 181a, and 155 in subacute sclerosing panencephalitis. *J. Child Neurol.* 30, 69–74. doi: 10.1177/088307381453 1329
- Zhong, Q., Feng, H., Lu, Q., Liu, X., Zhao, Q., Du, Y., et al. (2018). Recurrent wheezing in neonatal pneumonia is associated with combined infection with Respiratory Syncytial Virus and Staphylococcus aureus or *Klebsiella pneumoniae*. Sci. Rep. 8:995.
- Zhou, J., Liu, S., Chen, Y., Fu, Y., Silver, A. J., Hill, M. S., et al. (2017). Identification of two novel functional tRNA-derived fragments induced in response to respiratory syncytial virus infection. J. Gen. Virol. 98, 1600–1610. doi: 10.1099/ jgv.0.000852
- Zhou, N., Li, W., Ren, L., Xie, X., and Liu, E. (2017). An Interaction of LPS and RSV Infection in Augmenting the AHR and Airway Inflammation in Mice. *Inflammation* 40, 1643–1653. doi: 10.1007/s10753-017-0604-7

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Bakre, Duffy, Abdullah, Cosby and Tripp. This is an openaccess article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.