



**QUEEN'S
UNIVERSITY
BELFAST**

Nanoparticles Isolated From Porcine Bone Soup Ameliorated Dextran Sulfate Sodium-Induced Colitis and Regulated Gut Microbiota in Mice

Wang, H., Huang, J., Ding, Y., Zhou, J., Gao, G., Han, H., Zhou, J., Ke, L., Rao, P., Chen, T., & Zhang, L. (2022). Nanoparticles Isolated From Porcine Bone Soup Ameliorated Dextran Sulfate Sodium-Induced Colitis and Regulated Gut Microbiota in Mice. *Frontiers in Nutrition*, 9, Article 821404. <https://doi.org/10.3389/fnut.2022.821404>

Published in:
Frontiers in Nutrition

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2022 the authors.

This is an open access article published under a Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>



Nanoparticles Isolated From Porcine Bone Soup Ameliorated Dextran Sulfate Sodium-Induced Colitis and Regulated Gut Microbiota in Mice

Huiqin Wang¹, Jin Huang¹, Yanan Ding¹, Jianwu Zhou¹, Guanzhen Gao^{1*}, Huan Han¹, Jingru Zhou¹, Lijing Ke¹, Pingfan Rao¹, Tianbao Chen² and Longxin Zhang³

¹ Food Nutrition Science Centre, School of Food Science and Biotechnology, Zhejiang Gongshang University, Hangzhou, China, ² School of Pharmacy, Queen's University Belfast, Belfast, United Kingdom, ³ Fujian Provincial Maternity and Children's Hospital, Affiliated Hospital of Fujian Medical University, Fuzhou, China

OPEN ACCESS

Edited by:

Xin Wang,
Zhejiang Academy of Agricultural
Sciences, China

Reviewed by:

Shusong Wu,
Hunan Agricultural University, China
Zhengpeng Li,
Chinese PLA General Hospital, China

*Correspondence:

Guanzhen Gao
gaoguanzhen@zjgsu.edu.cn

Specialty section:

This article was submitted to
Nutrition and Microbes,
a section of the journal
Frontiers in Nutrition

Received: 24 November 2021

Accepted: 25 February 2022

Published: 29 March 2022

Citation:

Wang H, Huang J, Ding Y, Zhou J,
Gao G, Han H, Zhou J, Ke L, Rao P,
Chen T and Zhang L (2022)
Nanoparticles Isolated From Porcine
Bone Soup Ameliorated Dextran
Sulfate Sodium-Induced Colitis
and Regulated Gut Microbiota
in Mice. *Front. Nutr.* 9:821404.
doi: 10.3389/fnut.2022.821404

Daily foods contain a great number of self-assembled nanoparticles (NPs) which were incidentally produced during food processing. These food incidental NPs can directly access the human gastrointestinal tract in high frequency and large quantities. Limited reports were focused on whether and how these food incidental NPs affected the gastrointestinal tissues and gut microbiota. In the present study, bone soup and its NPs both significantly ameliorated colitis symptoms in dextran sulfate sodium (DSS)-induced mice and inhibited the release of pro-inflammatory cytokines. They also restored intestinal microbiota dysbiosis by improving the diversity and richness of intestinal microbiota and regulating community composition, such as a remarkable increase in *Muribaculaceae*, *Alistipes*, and *Alloprevotella*, and a decrease in *Helicobacter*. Moreover, the correlation analysis showed that pro-inflammatory cytokines were negatively correlated with *Muribaculaceae*, *Alloprevotella*, and *Alistipes*, but positively correlated with *Helicobacter*. These findings suggest that the food incidental NPs can influence human health through regulating the inflammation of the gastrointestinal tissues and the gut microbiota.

Keywords: bone soup, nanoparticles, ulcerative colitis, gut microbiota, inflammation, dextran sulfate sodium

INTRODUCTION

In the past decades, nanoparticles (NPs) have been increasingly used in food processing and food packaging for improving the quality of food, extending the shelf-life, enhancing the uptake, absorption, and bioavailability of food nutrients, etc (1, 2). These engineered NPs could have direct access to the human gastrointestinal tract (GIT) along with the food (3) and subsequently affect the gut microbiota and the GIT itself, such as GIT inflammation and redox homeostasis (4).

Abbreviations: NPs, nanoparticles; GIT, gastrointestinal tract; DSS, dextran sulfate sodium; DHA, docosahexaenoic acid; EPA, eicosatetraenoic acid; IBD, inflammatory bowel disease; BS, bone soup; BSNPs, bone soup nanoparticles; SASP, sulfasalazine; 16S rRNA, 16S ribosomal RNA; DAI, disease activity index; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; IL-6, interleukin 6; IL-1 β , interleukin 1 beta; TNF- α , tumor necrosis factor alpha; PCoA, principal coordinates analysis; NMDS, non-metric multidimensional scaling; LEfSe, linear discriminant analysis effect size; OTUs, operational taxonomic units; LDA, linear discriminant analysis; NF- κ B, nuclear factor kappa B.

Several studies revealed that these NPs, such as TiO₂ (5), SiO₂ (6), CuO (7), Ag (8), could induce intestinal inflammation and disruption of gut microbiota. On the contrary, some engineered NPs, e.g., Ce oxide (9), ZnO (10), and Ag (11) exerted an anti-inflammatory activity. The underlying mechanisms of the “nanoparticle-GIT” interaction were not fully understood, thus warranting further studies.

Apart from the engineered NPs in the food system, daily foods contained abundant amounts of incidental NPs, too, which were the assemblies of polar or non-polar food components derived by physical and chemical forces during the food processing (heating, cooling, and emulsification, etc.) (12). For instance, the proteoglycan–lipid NPs with an average diameter of 50–67 nm were generated during the heating processing of Freshwater Clams (*Corbicula fluminea* Muller) soup (12). Numerous spherical NPs were found in green tea infusions (13). A large number of self-assembly micro/nanoparticles containing docosahexaenoic acid (DHA) and eicosatetraenoic acid (EPA) have been observed in bigeye tuna (*Thunnus obesus*) head soup (14). Moreover, porcine bone soup, as a traditional nourishing food for preventing immune malfunction and inflammatory bowel disease (IBD) (15, 16), has been demonstrated to contain amounts of nanoscale colloidal particles with an average size of ca. 200 nm, and the formation of these incidental NPs has been investigated (17, 18). These food incidental NPs not only played a crucial role in the stability and texture characterization of food, but also made great contributions to the physiological effects of food due to the unique physicochemical properties at nanoscale. Similar to the engineered NPs, these food incidental NPs may directly encounter cells in human GIT and alter the intestinal microenvironment. To date, however, little has been known on these incidental food NPs and their biological impacts on the gut.

As we previously reported, the spherical NPs mainly consisted of lipids and proteins with an average size of ca. 200 nm and ζ -potential of ca. -15 mV derived from porcine bone soup were isolated and directly interacted with macrophages with preventing cells from peroxyl radical-induced membrane hyperpolarization, mitochondrial malfunction, and phagocytosis suppression (18). Moreover, these NPs derived from the porcine bone soup attenuated oxidative stress-induced intestinal barrier injury in the Caco-2 cell monolayer model (17). In this study, we set off to evaluate the alleviative effects of NPs derived from porcine bone soup on dextran sulfate sodium (DSS) induced colitis in mice, by modulating intestinal inflammation and gut microbiota.

MATERIALS AND METHODS

Preparation of Nanoparticles Derived From Porcine Bone Soup

Fresh porcine bones (Landrace pigs, *Sus scrofa*) were obtained from the fresh local market. The bone soup was prepared, and the NPs were isolated according to our previous reports (17–19). Briefly, 1 kg of fresh porcine bones was cooked with deionized water (1:3, w/v) for 3 h in a boiling water bath after removing the blood residues. Filtered through four layers of cotton gauze

to remove the solid residues and the soup was centrifuged at 400 g for 10 min. The supernatant of bone soup was collected and named as BS. Then BS was applied to a pre-equilibrated size-exclusive chromatographic column. The fractions with strong light scattering intensity were collected and dialyzed against water, and the obtained NPs samples were freeze-dried and designated as BSNPs. Before the animal experiment, BSNPs were dissolved with deionized water to the same derived count rate as that of BS.

Animal Experiments

The 5-week-old specific pathogen-free (SPF) female BALB/c mice were supplied by Zhejiang Center of Laboratory Animals (Hangzhou, China). The mice were housed under the standard conditions ($21 \pm 1^\circ\text{C}$, $55.5 \pm 5\%$ relative humidity) with a controlled light–dark cycle of 12/12 h. The mice were fed on standard chow and distilled water *ad libitum*. After 1 week of adaptive feeding, the mice were randomly divided into five groups with six mice each. Acute colitis in BALB/c mice was established according to the previous study (20). The groups were as follows: (1) Control group, mice were fed with standard chow and distilled water *ad libitum*, daily administered normal saline (50 ml/kg/day) by oral gavage for 7 days. (2) DSS group, provided free access to 5% DSS (36–50 kDa; MP Biomedicals, Irvine, CA, United States) in distilled water and daily administered normal saline (50 ml/kg/day) by oral gavage for 7 days. (3) Sulfasalazine (SASP) group, provided free access to 5% DSS in distilled water and daily administered SASP (250 mg/kg, Shanghai Zhongxi Sanwei Pharmaceutical Co., Ltd., Shanghai, China) by oral gavage for 7 days. (4) The BS group provided free access to 5% DSS in distilled water and administered BS at 50 ml/kg/day by oral gavage for 7 days. (5) BSNPs group provided free access to 5% DSS in distilled water and administered BSNPs at 50 ml/kg/day by oral gavage for 7 days. On the 8th day, mice were sacrificed by cervical dislocation. The fecal samples were collected for 16S ribosomal RNA (16S rRNA) analysis. The colon samples were excised quickly and rinsed with ice-cold 0.9% NaCl. Their length and weight were subsequently measured. At last, colons were divided into three parts for histopathological analysis and real-time polymerase chain reaction (PCR).

All animal experiments were conducted according to the NIH Guidance for the care and use of laboratory animals and approved by the Animal Care and Welfare Committee of Zhejiang Center of Laboratory Animals, China.

Disease Activity Index

The severity of colitis was evaluated daily according to the DAI scoring system as described previously (21), such as body weight loss, stool consistency, and gross rectal bleeding.

Histological Analysis

For histological evaluation, hematoxylin and eosin (H&E) stained colon tissues were performed, and histological changes were evaluated by two independent investigators in a blinded manner according to a scoring system as described previously (22). The sum of each score reflected the colonic histological damage.

TABLE 1 | GenBank accession numbers, sequences of forward and reverse primers, and fragment sizes used for real-time PCR.

Target	GenBank number	Primer sequence	Size, bp
<i>TNF-α</i>	NM_013693.3	F:5' TTGTCTACTCCAGGTTCTCT3' R: 5' GAGGTTGACTTTTCCTGGTATG3'	107
<i>IL-6</i>	NM_031168.2	F:5' CTTCCATCCAGTTGCCTTCT3' R:5' CTCCGACTTGTGAAGTGGTATAG3'	134
<i>IL-1β</i>	NM_008361.4	F:5' CCACCTCAATGGACAGAATATCA3' R:5' CCCAAGGCCACAGGTATTT3'	96
<i>GAPDH</i>	NM_001289726.1	F:5' AACAGCAACTCCCACTCTTC3' R:5' CCTGTTGCTGTAGCCGTATT3'	111

RNA Extract and Real-Time PCR

The mRNA expression of inflammatory cytokines, such as interleukin 6 (IL-6), interleukin 1 beta (IL-1 β), and tumor necrosis factor-alpha (TNF- α), were determined in ileum mucosa by real-time PCR. Total RNA was extracted from the colon tissue using with miRNeasy Mini Kit (QIAGEN, 217004, Hilden, North Rhine-Westphalia, Germany), its concentration and purity were measured using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, United States). cDNA was synthesized using RevertAid First Strand cDNA Synthesis Kit (Thermo Fisher Scientific, K1622, Waltham, MA, United States). The real-time PCR was conducted using Platinum SYBR Green qPCR SuperMix-UDG with ROX (Invitrogen, 11744-500, Carlsbad, CA, United States); the reaction solution contained 10 μ l of 2 \times SYBR Green, 1 μ l of each primer (10 μ M), 1 μ l of reverse transcription product, and 8 μ l of RNase/DNase-free water (total volume 20 μ l). Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as an internal reference gene to normalize target gene transcript levels and the primers sequences of target genes (Sangon Biotech (Shanghai) Co., Shanghai, China) are presented in **Table 1**. The reaction conditions were as follows: 2 min at 95°C, 40 cycles for 15 s at 95°C, 30 s at 60°C, 30 s at 72°C. All the results were normalized to the internal reference gene: GAPDH. The relative expression of mRNA level was calculated with the $2^{-\Delta\Delta C_t}$ method.

16S rRNA Analysis

The gut microbiota composition of mice feces was determined by 16S rRNA gene amplification. Briefly, MagPure Stool DNA KF Kit B (Magen Biotechnology Co., Guangzhou, China) was used to extract genomic DNA from feces. DNA concentration and integrity were measured by a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, United States) and agarose gel electrophoresis, respectively. The 16S rRNA gene V3–V4 region was amplified from the genomic DNA in a 25 μ l reaction using the universal bacterial primers: (343F, 5'-TACGGRAGGCAGCAG-3', and 798R, 5'-AGGGTATCTAATCCT-3'). The PCR products were purified with Agencourt AMPure XP beads (Beckman Coulter Co., Brea, CA, United States) and quantified by using a Qubit dsDNA assay kit (Thermo Fisher Scientific, Waltham, MA, United States). Sequencing was performed on an Illumina NovaSeq 6000 with two paired-end read cycles of 250 bases each (Illumina

Inc., San Diego, CA, United States; OE Biotech Co., Ltd., Shanghai, China).

The Trimmomatic software was used to analyze the raw data. After trimming, operational taxonomic units (OTUs) were generated by using VSEARCH 2.7.1 with a 97% similarity cutoff. The representative read of each OTU was selected by using the Quantitative Insights into Microbial Ecology (QIIME) package. All representative reads were annotated and blasted against the Silva database (Version 132) using the RDP classifier (confidence threshold was 70%). The microbial richness and diversity in fecal content samples were estimated using the alpha diversity that includes Chao1 index, Observed species index, Simpson index, and Shannon index. The UniFrac distance matrix performed by QIIME software was used for the unweighted UniFrac Principal coordinates analysis (PCoA), Non-metric multidimensional scaling (NMDS) analysis, and phylogenetic tree construction. Linear discriminant analysis effect size (LEfSe) based linear discriminant analysis (LDA) and cladogram were generated to assess differentially abundant microbial taxa. The 16S rRNA gene amplicon sequencing and analysis were conducted by OE Biotech Co., Ltd. (Shanghai, China).


Statistical Analysis

Data were expressed as mean \pm standard error of the mean (SEM) and performed with GraphPad Prism 7.0 (GraphPad Software, Inc., San Diego, CA, United States). The significant differences were analyzed using one-way analysis of variance (ANOVA) followed by Tukey's multiple comparisons in multiple groups. Correlations were analyzed by using Spearman's correlation analysis. The value of $p < 0.05$ was considered to be statistically significant.

RESULTS

Bone Soup Nanoparticles Alleviated the Symptoms of Dextran Sulfate Sodium-Induced Colitis in Mice


The influences of the BS or its BSNPs on the symptoms of DSS-induced colitis were investigated. As shown in **Figures 1A–E**, the DSS-treated group showed a significant reduction in bodyweight, increase in DAI score, reduction of colon length, and increment of colon weight/length ratio compared with the healthy control group ($p < 0.05$). The BS or BSNPs treatment, similar to that



of SASP, significantly alleviated the bodyweight loss, the DAI score, and increment in the colon length. A significant reduction of colon weight/length ratio was observed in the BS/BSNPs-treated mice compared with the DSS group ($p < 0.05$). Moreover, no significant difference was observed between BS and BSNPs treatment, implying that the alleviative effects of BS in DSS-induced colitis might be mainly attributed to its NP components.

Bone Soup Nanoparticles Attenuated the Inflammation of Colonic Mucosa

The histological changes of colonic mucosa were consistent with the general symptoms described above, determined with H&E staining. As shown in **Figures 2A,B**, the DSS treatment induced inflammatory cell infiltration and significantly elevated histological score ($p < 0.05$). The treatment of BS or BSNPs



significantly inhibited the cell infiltration and histological score ($p < 0.05$).

In line with the histological changes, the pro-inflammatory cytokines (e.g., TNF- α , IL-6, and IL-1 β) were boosted by DSS but downregulated by BS or BSNPs to an extent similar to that of SASP (Figures 2C–E), judging by quantitative real-time PCR determination of mRNA abundance of cytokines. The BS and BSNPs exhibited the same level of inhibition on colonic inflammation, elucidating the NPs are the principal active components of the bone soup in treating DSS-induced colitis.

Bone Soup Nanoparticles Restored Gut Microbial Diversity

The alpha diversity and beta diversity analysis of the gut microbial community in each group were assessed using a set of statistical analysis indexes. The gut microbial community richness index was calculated with Chao1 and Observed species index, while Shannon index and Simpson index were used to estimate the microbial diversity. As shown in Figures 3A–D, the Chao1 index, Observed species index, Shannon index, and Simpson index of the DSS group were found to be significantly lower than the normal mice, suggesting DSS reduced the microbial richness and diversity, disrupted the microbiota structure. The administration of either BS or BSNPs restored the microbial richness and diversity, in similar effectiveness with SASP. No significant difference was observed between the BS and BSNPs groups.

Besides, the PCoA of the microbial community showed that the microbial evolution of DSS-treated mice was vastly different from the normal mice, while the BS and BSNPs-treated mice were rather similar to the normal mice (Figure 3E). It is in line with the NMDS analysis (Figure 3F). In addition, the unweighted pair-group technique with arithmetic mean (UPGMA) was employed to construct a circular hierarchical clustering tree (Figure 3G), which indicated that the administration of BS, BSNPs, and SASP caused much fewer changes in microbial communities than that of DSS.

Modulation of Gut Microbiota by Bone Soup Nanoparticles

The gut microbiota community structure distribution and relative abundance of taxa were analyzed. At the phylum level, the gut microbiota composition in each group was shown in Figure 4A. Compared to the normal mice, mice in the DSS group exhibited the higher relative abundances of *Campilobacterota* (33.43 vs. 1.26%) and lower relative abundances of *Bacteroidetes* (25.53 vs. 63.95%). The DSS induced an elevated *Firmicutes/Bacteroidetes* (F/B) ratio, too ($p < 0.05$, Figure 4C). The administration of BSNPs or BS significantly reversed these microbiota composition changes at the phylum level and resorted the F/B ratio to a normal level.

Consistently, the BSNPs or BS administration also restored the microflora balance at the genus level (Figure 4B), characterized by significantly increased *Muribaculaceae* (Figure 4D), *Alistipes* (Figure 4H), and *Alloprevotella* (Figure 4G), markedly decreased *Helicobacter* and *Lachnospiraceae_NK4A136_group* (Figures 4E,F) ($p < 0.05$). Furthermore, a heatmap based on the





FIGURE 3 | Analysis of the alpha diversity and beta diversity of the differential microbial community among the control group, DSS group, SASP, BS, and BSNPs-treated group. Violin plots showing Chao1 index (A), Observed species index (B), Simpson index (C), Shannon index (D). (E) UniFrac distance of unweighted PCoA analysis, (F) UniFrac distance of unweighted NMDS analysis, (G) the unweighted pair-group technique with arithmetic mean (UPGMA) analysis the differences between samples. Means without a common letter are significantly different at $p < 0.05$ using one-way ANOVA followed by Tukey's test for *post hoc* analysis.

relative abundance of the top 15 abundant genera of bacteria, a phylogenetic tree, and a heatmap showing relative abundances of OTUs together with a corresponding phylogenetic tree were performed (Figures 4I,J). These results confirmed that BS or



BSNPs treatment could counteract the DSS-induced changes in the gut microbiota.

The LfSe was performed to compare the difference in the abundance of gut microbiota compositions among different groups of mice (**Figures 5A,B**). From the phylum level to the genus level, there were 43 significant taxa in all groups. At the genus levels, *Lachnospiraceae_NK4A136_group* and *Helicobacter* were enriched in the DSS-treated mice, while 11 other taxa, such as *Muribaculaceae*, *Alloprevotella*, *Alistipes*, *Odoribacter*, *Clostridia_vadinBB60_group*, and *Rikenellaceae_RC9_gut_group*, were enriched in normal mice and SASP/BS/BSNPs-treated mice.

Correlations Between the Expressions of Pro-inflammatory Cytokines With Microbiota

The possible correlations between gut microbiota and pro-inflammatory cytokines were examined with Spearman's

correlation analyses (23). As shown in **Figure 6**, six bacteria were negatively correlated with pro-inflammatory cytokines ($p < 0.05$), such as *Muribaculaceae*, *Alloprevotella*, *Alistipes*, *Odoribacter*, *Clostridia_vadinBB60_group*, and *Rikenellaceae_RC9_gut_group*. Notably, these six bacteria were separately enriched in BSNPs, BS, and control group (**Figure 5A**), and *Muribaculaceae*, *Alloprevotella*, and *Alistipes* were significantly lower in the DSS group than other groups (**Figures 4D,G,H**). Furthermore, *Helicobacter* and *Colidextribacter* were significantly positively correlated with the pro-inflammatory cytokines ($p < 0.05$). Intriguingly, *Helicobacter* was enriched in DSS treatment (**Figure 5A**), and the treatment of BS, BSNPs, or SASP can significantly decrease the abundances of *Helicobacter* (**Figure 4E**). These results revealed that the anti-inflammatory effects of BSNPs and BS on DSS-induced colitis might be correlated to the increased abundance in *Muribaculaceae*, *Alloprevotella*, and





FIGURE 5 | Linear discriminant analysis (LDA) and effect size (LEfSe) analysis of bacterial among the different groups. **(A)** LDA indicates the most differential abundant bacterial taxa in specific samples. **(B)** Cladogram illustrating highly abundant taxa across various treatments.



Alistipes, and the decreased abundance in *Helicobacter* in the microbiota of mice.

DISCUSSION

The present study demonstrated that BS and BSNPs (the NPs isolated from bone soups) alleviated the pathological symptoms and inflammatory cytokines of DSS-induced colitis in mice. There was no significant difference between BS and BSNPs treatment, implying that BSNPs may exert protective effects against DSS-induced bowel disease in mice. BSNPs are mainly consisted of lipids and proteins, including collagen (18). It has been reported that the hybrid NPs of lipid-protein were resistant to hydrolysis of enzymes in saliva, stomach, and intestinal fluids and protected its inherent bioactive components from degradation (24, 25). The lipid-protein complex of BSNPs might make the particles resistant to gastrointestinal digestion, as was found in grapefruit-derived NPs (25). In our previous study, BSNPs have been demonstrated impressively high stability under harsh conditions of 4°C for 5 days (18). This indicated that BSNPs might as intact NP form to exert the anti-inflammatory effect in the gut.

Some inorganic NPs [such as Ce oxide (9), ZnO (10), and Ag (11)] could exhibit the therapeutic effects in the experimental colitis model *via* the anti-oxidative mechanism. Other NPs fabricated with natural biopolymers, such as chitosan and silk fibroin, acted as nanoscale drug delivery systems to alleviate ulcerative colitis (4, 26–28). The alleviative effects of BSNPs on intestinal inflammation might be attributed to its antioxidant activity and its constitutive collagen (17–19).

The cellular protective effects of BSNPs have been demonstrated in previous studies (17–19). According to the *in vitro* study, the BSNPs could be engulfed by oral and peritoneal macrophages, possess intracellular antioxidant activities, and suppress the membrane hyperpolarization induced by peroxy radical on macrophage (18, 19). Such suppression could facilitate the anti-inflammatory effects (29). The healthier macrophages would promote the gut homeostasis and recovery of IBD (30). In addition, the disruption of the intestinal barrier is associated with disease of the GIT. Emerging evidence showed inadequate epithelial permeability resulted in IBD, IBS, UC, etc (31, 32). BSNPs could restore the intestinal barrier function of Caco-2 cells (17), which may also contribute to its alleviating effects on colitis.

Ulcerative colitis is accompanied by the dysbiosis of gut microbiota. Our results also revealed that BS or BSNPs could restore gut microbiota dysbiosis induced by the DSS. BS or BSNPs treatment increased the richness or diversity of gut microbial communities, which is badly damaged in IBD patients and DSS-induced colitis mice (33). The BS or BSNPs treatment markedly increased the abundance of *Muribaculaceae*, *Alistipes*, and *Alloprevotella*, while decreasing that of *Helicobacter*. The decrease of *Muribaculaceae* has been identified as the key to the development of colitis (34). *Alistipes* was reported to have protective effects against some diseases, including colitis (35). *Alloprevotella* was demonstrated to be lower in mice with DSS-induced colitis (36). *Helicobacter* was pathogens implicated in the formation of peptic ulcers and IBD (37). Moreover, the BS or BSNPs treatment decreased the F/B ratio. The ratio would be increased in colitis and other inflammatory diseases. A lower F/B ratio is connected to recovery or the health status (38, 39). Notably, our study demonstrated that *Muribaculaceae*, *Alloprevotella*, and *Alistipes*

were negatively correlated with TNF- α , IL-6, and IL-1 β . Some previous studies have shown that *Muribaculaceae*, *Alloprevotella*, and *Alistipes* were negatively correlated with inflammation, cause they promoted SCFA production and improved the intestinal tissue's tolerance to the immune stimulations (34, 36, 40). In contrast, *Helicobacter* was positively correlated with three pro-inflammatory cytokines. Consistently, other reports have also described that *Helicobacter* genus could stimulate the immune response and cause the production of pro-inflammatory cytokines via activating proinflammatory nuclear factor kappa B (NF- κ B) pathway (36, 41). These findings suggested that the protective effects of BSNPs on colitis might be attributed to its regulatory effects of gut microbiota correlated with immune responses and inflammation.

It has been reported that nanosized-food additives can bind to bacteria to form the NP-bacteria complex primarily via van der Waals attraction (42). This complex can induce the degradation of bacterial cell wall components, the alteration of metabolic pathways, oxidation of cellular components, and DNA damage, leading to the influence on the microbiome (43). In addition, naturally occurring NPs from beer could also bind to bacteria to form NP-bacteria complexation and then affect the microbiome (42).

However, it remains unknown whether the BSNPs could have the chance and ability to bind to the microorganisms in the alimentary tract and form a hybrid complex. The BSNPs may have been partially or completely digested before they reach the colon. The constitutive lipid and protein of BSNPs may also contribute to the bioactivities. The underlying mechanism of the interaction between NPs, their digestion products, and microbiota has not been fully understood and warrants further investigation.

CONCLUSION

Altogether, our study demonstrated that the isolated NPs from bone soup showed a preventing effect against DSS-induced colitis. The treatment of BSNPs alleviated the severity of colitis with reduction of the release of inflammatory cytokines, restored gut microbiota dysbiosis by redirecting the bacterial shift to a normal state with increasing the richness or diversity of gut microbial communities and regulating community composition, such as a remarkable increase in *Muribaculaceae*, *Alistipes*, and *Alloprevotella*, and a decrease in *Helicobacter*. It indicated

that the food-derived NPs could influence human health by regulating the inflammation of the gastrointestinal tissues and the gut microbiota.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found below: <https://www.ncbi.nlm.nih.gov/search/all/?term=PRJNA779815>, Sequence Read Archive (SRA) database (PRJNA779815).

ETHICS STATEMENT

The animal study was reviewed and approved by the Animal Care and Welfare Committee of Zhejiang Center of Laboratory Animals, China.

AUTHOR CONTRIBUTIONS

HW contributed to conceptualization, data curation, methodology, validation, formal analysis, and writing the original draft. JH and HH contributed to formal analysis, data curation, and methodology. YD and JRZ contributed to methodology, formal analysis, and investigation. JWZ contributed to formal analysis, supervision, and writing, reviewing, and editing the manuscript. GG contributed to conceptualization, methodology, supervision, investigation, validation, funding acquisition, project administration, and writing, reviewing, and editing the manuscript. LK contributed to formal analysis, project administration, and writing, reviewing, and editing the manuscript. PR contributed to conceptualization and writing, reviewing, and editing the manuscript. TC reviewed and edited the manuscript. LZ contributed to writing, reviewing, and editing the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by the National Key R&D Program of China (2016YFD0400202) and Fundamental Special Funds for the Provincial Universities of Zhejiang (Grant No. JDK21005).

REFERENCES

- Nile SH, Baskar V, Selvaraj D, Nile A, Xiao J, Kai G. *Nanotechnologies in Food Science: Applications, Recent Trends, and Future Perspectives*. Singapore: Springer (2020). doi: 10.1007/s40820-020-0383-9
- Singh T, Shukla S, Kumar P, Wahla V, Bajpai VK. Application of nanotechnology in food science: perception and overview. *Front Microbiol.* (2017) 8:1501. doi: 10.3389/fmicb.2017.01501
- Ghebretatios M, Schaly S, Prakash S. Nanoparticles in the food industry and their impact on human gut microbiome and diseases. *Int J Mol Sci.* (2021) 22:1942. doi: 10.3390/ijms22041942
- Bouwmeester H, van der Zande M, Jepson MA. Effects of food-borne nanomaterials on gastrointestinal tissues and microbiota. *Wiley Interdiscip Rev Nanomed Nanobiotechnol.* (2018) 10:e1481. doi: 10.1002/wnan.1481
- Mu W, Wang Y, Huang C, Fu Y, Li J, Wang H, et al. Effect of long-term intake of dietary titanium dioxide nanoparticles on intestine inflammation in mice. *J Agric Food Chem.* (2019) 67:9382–9. doi: 10.1021/acs.jafc.9b02391
- Ogawa T, Okumura R, Nagano K, Minemura T, Izumi M, Motoooka D, et al. Oral intake of silica nanoparticles exacerbates intestinal inflammation. *Biochem Biophys Res Commun.* (2021) 534:540–6. doi: 10.1016/j.bbrc.2020.11.047

7. Ude VC, Brown DM, Stone V, Johnston HJ. Time dependent impact of copper oxide nanomaterials on the expression of genes associated with oxidative stress, metal binding, inflammation and mucus secretion in single and co-culture intestinal in vitro models. *Toxicol Vitro*. (2021) 74:105161. doi: 10.1016/j.tiv.2021.105161
8. Williams K, Milner J, Boudreau MD, Gokulan K, Cerniglia CE, Khare S. Effects of subchronic exposure of silver nanoparticles on intestinal microbiota and gut-associated immune responses in the ileum of Sprague-Dawley rats. *Nanotoxicology*. (2015) 9:279–89. doi: 10.3109/17435390.2014.921346
9. Asgharzadeh F, Hashemzadeh A, Rahmani F, Yaghoubi A, Nazari SE, Avan A, et al. Cerium oxide nanoparticles acts as a novel therapeutic agent for ulcerative colitis through anti-oxidative mechanism. *Life Sci*. (2021) 278:119500. doi: 10.1016/j.lfs.2021.119500
10. Li J, Chen H, Wang B, Cai C, Yang X, Chai Z, et al. ZnO nanoparticles act as supportive therapy in DSS-induced ulcerative colitis in mice by maintaining gut homeostasis and activating Nrf2 signaling. *Sci Rep*. (2017) 7:43126. doi: 10.1038/srep43126
11. Siczek K, Zatorski H, Chmielowiec-Korzeniowska A, Pulit-Prociak J, Śmiech M, Kordek R, et al. Synthesis and evaluation of anti-inflammatory properties of silver nanoparticle suspensions in experimental colitis in mice. *Chem Biol Drug Des*. (2017) 89:538–47. doi: 10.1111/cbdd.12876
12. Gao G, Wang H, Zhou J, Rao P, Ke L, Lin J, et al. Isolation and characterization of bioactive proteoglycan-lipid nanoparticles from freshwater clam (*Corbicula fluminea* Muller) soup. *J Agric Food Chem*. (2021) 69:1610–8. doi: 10.1021/acs.jafc.0c02402
13. Lin X, Gao X, Chen Z, Zhang Y, Luo W, Li X, et al. Spontaneously assembled nano-aggregates in clear green tea infusions from *Camellia ptilophylla* and *Camellia sinensis*. *J Agric Food Chem*. (2017) 65:3757–66. doi: 10.1021/acs.jafc.7b00068
14. Zhang J, Lin L, Tao N, Zhu Z, Wang X, Wang M. Effect of big eye tuna (*Thunnus obesus*) head soup with different colloidal particle size on TG and TC deposition in FFA-exposed HepG2 cells. *Food Sci Nutr*. (2021) 9:1143–51. doi: 10.1002/fsn3.2092
15. Siebecker A. Traditional bone broth in modern health and disease. *Townsend Lett*. (2005) 259/260:74–81.
16. Morell SF, Daniel KT. *Nourishing Broth: An Old-Fashioned Remedy for the Modern World*. New York, NY: Hachette Book Group Special Markets Department (2014).
17. Gao G, Zhou J, Jin Y, Wang H, Ding Y, Zhou J, et al. Nanoparticles derived from porcine bone soup attenuate oxidative stress-induced intestinal barrier injury in Caco-2 cell monolayer model. *J Funct Foods*. (2021) 83:104573. doi: 10.1016/j.jff.2021.104573
18. Wang H, Gao G, Ke L, Zhou J, Rao P, Jin Y, et al. Isolation of colloidal particles from porcine bone soup and their interaction with murine peritoneal macrophage. *J Funct Foods*. (2019) 54:403–11. doi: 10.1016/j.jff.2019.01.021
19. Ke L, Wang H, Gao G, Rao P, He L, Zhou J. Direct interaction of food derived colloidal micro/nano-particles with oral macrophages. *Npj Sci Food*. (2017) 1:3. doi: 10.1038/s41538-017-0003-3
20. Cançado GGL, Fiuza JA, De Paiva NCN, Lemos LDCD, Ricci ND, Gazzinelli-Guimarães PH, et al. Hookworm products ameliorate dextran sodium sulfate-induced colitis in BALB/c mice. *Inflamm Bowel Dis*. (2011) 17:2275–86. doi: 10.1002/ibd.21629
21. Hidalgo-Cantabrana C, Algieri F, Rodriguez-Nogales A, Vezza T, Martínez-Cambor P, Margolles A, et al. Effect of a rosy exopolysaccharide-producing *Bifidobacterium animalis* subsp. lactis strain orally administered on DSS-induced colitis mice model. *Front Microbiol*. (2016) 7:868. doi: 10.3389/fmicb.2016.00868
22. Tong LC, Wang Y, Wang ZB, Liu WY, Sun S, Li L, et al. Propionate ameliorates dextran sodium sulfate-induced colitis by improving intestinal barrier function and reducing inflammation and oxidative stress. *Front Pharmacol*. (2016) 7:253. doi: 10.3389/fphar.2016.00253
23. Hu R, Wu S, Li B, Tan J, Yan J, Wang Y, et al. Dietary ferulic acid and vanillic acid on inflammation, gut barrier function and growth performance in lipopolysaccharide-challenged piglets. *Anim Nutr*. (2022) 8:144–52. doi: 10.1016/j.aninu.2021.06.009
24. Yang C, Merlin D. Can naturally occurring nanoparticle-based targeted drug delivery effectively treat inflammatory bowel disease? *Expert Opin Drug Deliv*. (2020) 17:1–4. doi: 10.1080/17425247.2020.1698543
25. Wang B, Zhuang X, Deng ZB, Jiang H, Mu J, Wang Q, et al. Targeted drug delivery to intestinal macrophages by bioactive nanovesicles released from grapefruit. *Mol Ther*. (2014) 22:522–34. doi: 10.1038/mt.2013.190
26. Gou S, Huang Y, Sung J, Xiao B, Merlin D. Silk fibroin-based nanotherapeutics: application in the treatment of colonic diseases. *Nanomedicine*. (2019) 14:2373–8. doi: 10.2217/nnm-2019-0058
27. Gou S, Huang Y, Wan Y, Ma Y, Zhou X, Tong X, et al. Multi-bioresponsive silk fibroin-based nanoparticles with on-demand cytoplasmic drug release capacity for CD44-targeted alleviation of ulcerative colitis. *Biomaterials*. (2019) 212:39–54. doi: 10.1016/j.biomaterials.2019.05.012
28. Iglesias N, Galbis E, Díaz-Blanco MJ, Lucas R, Benito E, De-Paz MV. Nanostructured chitosan-based biomaterials for sustained and colon-specific resveratrol release. *Int J Mol Sci*. (2019) 20:398. doi: 10.3390/ijms20020398
29. Li C, Levin M, Kaplan DL. Bioelectric modulation of macrophage polarization. *Sci Rep*. (2016) 6:21044. doi: 10.1038/srep21044
30. Na YR, Stakenborg M, Seok SH, Matteoli G. Macrophages in intestinal inflammation and resolution: a potential therapeutic target in IBD. *Nat Rev Gastroenterol Hepatol*. (2019) 16:531–43. doi: 10.1038/s41575-019-0172-4
31. Antoni L, Nuding S, Wehkamp J, Stange EF. Intestinal barrier in inflammatory bowel disease. *World J Gastroenterol*. (2014) 20:1165–79. doi: 10.3748/wjg.v20.i5.1165
32. Camilleri M, Madsen K, Spiller R, Van Meerveld BG, Verne GN. Intestinal barrier function in health and gastrointestinal disease. *Neurogastroenterol Motil*. (2012) 24:503–12. doi: 10.1111/j.1365-2982.2012.01921.x
33. Manichanh C, Borruel N, Casellas F, Guarner F. The gut microbiota in IBD. *Nat Rev Gastroenterol Hepatol*. (2012) 9:599–608. doi: 10.1038/nrgastro.2012.152
34. Shang L, Liu H, Yu H, Chen M, Yang T, Zeng X, et al. Core altered microorganisms in colitis mouse model: a comprehensive time-point and fecal microbiota transplantation analysis. *Antibiotics*. (2021) 10:643. doi: 10.3390/antibiotics10060643
35. Parker BJ, Wearsch PA, Veloo ACM, Rodriguez-Palacios A. The genus *Alistipes*: gut bacteria with emerging implications to inflammation, cancer, and mental health. *Front Immunol*. (2020) 11:906. doi: 10.3389/fimmu.2020.00906
36. Li AL, Ni WW, Zhang QM, Li Y, Zhang X, Wu HY, et al. Effect of cinnamon essential oil on gut microbiota in the mouse model of dextran sodium sulfate-induced colitis. *Microbiol Immunol*. (2020) 64:23–32. doi: 10.1111/1348-0421.12749
37. Kaur CP, Vadivelu J, Chandramathi S. Impact of *Klebsiella pneumoniae* in lower gastrointestinal tract diseases. *J Dig Dis*. (2018) 19:262–71. doi: 10.1111/1751-2980.12595
38. Grigor'eva IN. Gallstone disease, obesity and the firmicutes/bacteroidetes ratio as a possible biomarker of gut dysbiosis. *J Pers Med*. (2021) 11:13. doi: 10.3390/jpm11010013
39. Tong L, Zhang X, Hao H, Liu Q, Zhou Z, Liang X, et al. *Lactobacillus rhamnosus* gg derived extracellular vesicles modulate gut microbiota and attenuate inflammatory in dss-induced colitis mice. *Nutrients*. (2021) 13:3319. doi: 10.3390/nu13103319
40. Wu M, Li P, An Y, Ren J, Yan D, Cui J, et al. Phloretin ameliorates dextran sulfate sodium-induced ulcerative colitis in mice by regulating the gut microbiota. *Pharmacol Res*. (2019) 150:104489. doi: 10.1016/j.phrs.2019.104489
41. Hu Y, Liu JP, Zhu Y, Lu NH. The importance of toll-like receptors in NF- κ B signaling pathway activation by *Helicobacter pylori* infection and the regulators of this response. *Helicobacter*. (2016) 21:428–40. doi: 10.1111/hel.12292
42. Siemer S, Hahlbrock A, Vallet C, McClements DJ, Balszuweit J, Voskuhl J, et al. Nanosized food additives impact beneficial and pathogenic bacteria in the human gut: a simulated gastrointestinal study. *Npj Sci Food*. (2018) 2:22. doi: 10.1038/s41538-018-0030-8
43. Gangadoo S, Nguyen H, Rajapaksha P, Zreiqat H, Latham K, Cozzolino D, et al. Inorganic nanoparticles as food additives and their influence on the human gut microbiota. *Environ Sci Nano*. (2021) 8:1500–18. doi: 10.1039/d1en00025j

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Wang, Huang, Ding, Zhou, Gao, Han, Zhou, Ke, Rao, Chen and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.