**QUEEN'S
UNIVERSITY
BELFAST**

Testing quasar unification: radiative transfer in clumpy winds

Matthews, J. H., Knigge, C., Long, K. S., Sim, S. A., Higginbottom, N., & Mangham, S. W. (2016). Testing quasar unification: radiative transfer in clumpy winds. *Monthly Notices of the Royal Astronomical Society*, 458(1), 293-305. <https://doi.org/10.1093/mnras/stw323>

Published in:
Monthly Notices of the Royal Astronomical Society

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
This article has been accepted for publication in Monthly Notices of the Royal Astronomical Society.
Copyright 2016 The Authors. Published by Oxford University Press on behalf of the Royal Astronomical Society.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Testing quasar unification: radiative transfer in clumpy winds

J. H. Matthews,¹★ C. Knigge,¹ K. S. Long,² S. A. Sim,³ N. Higginbottom¹
and S. W. Mangham¹

¹*School of Physics and Astronomy, University of Southampton, Highfield, Southampton SO17 1BJ, UK*

²*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, UK*

³*School of Mathematics and Physics, Queen's University Belfast, University Road, Belfast BT7 1NN, UK*

Accepted 2016 February 8. Received 2016 February 8; in original form 2015 December 23

ABSTRACT

Various unification schemes interpret the complex phenomenology of quasars and luminous active galactic nuclei (AGN) in terms of a simple picture involving a central black hole, an accretion disc and an associated outflow. Here, we continue our tests of this paradigm by comparing quasar spectra to synthetic spectra of biconical disc wind models, produced with our state-of-the-art Monte Carlo radiative transfer code. Previously, we have shown that we could produce synthetic spectra resembling those of observed broad absorption line (BAL) quasars, but only if the X-ray luminosity was limited to 10^{43} erg s^{−1}. Here, we introduce a simple treatment of clumping, and find that a filling factor of ~ 0.01 moderates the ionization state sufficiently for BAL features to form in the rest-frame UV at more realistic X-ray luminosities. Our fiducial model shows good agreement with AGN X-ray properties and the wind produces strong line emission in, e.g., Ly α and C IV 1550 Å at low inclinations. At high inclinations, the spectra possess prominent LoBAL features. Despite these successes, we cannot reproduce all emission lines seen in quasar spectra with the correct equivalent-width ratios, and we find an angular dependence of emission line equivalent width despite the similarities in the observed emission line properties of BAL and non-BAL quasars. Overall, our work suggests that biconical winds can reproduce much of the qualitative behaviour expected from a unified model, but we cannot yet provide quantitative matches with quasar properties at all viewing angles. Whether disc winds can successfully unify quasars is therefore still an open question.

Key words: accretion, accretion discs – radiative transfer – methods: numerical – galaxies: active – quasars: absorption lines – quasars: emission lines.

1 INTRODUCTION

The spectra of quasars and luminous active galactic nuclei (AGN) typically exhibit a series of strong emission lines with an underlying blue continuum – the so-called ‘big blue bump’ (BBB). The BBB is often attributed to emission from a geometrically thin, optically thick accretion disc surrounding the central black hole (BH), similar to that described by Shakura & Sunyaev (1973). In addition to the *inflowing* accreting material, *outflows* are ubiquitous in AGN and quasars (Kellermann et al. 1989; Ganguly & Brotherton 2008). These outflows can take the form of highly collimated radio jets (e.g. Hazard, Mackey & Shimmins 1963; Potash & Wardle 1980; Perley, Dreher & Cowan 1984; Marscher 2006) or mass-loaded ‘winds’ emanating from the accretion disc (Weymann et al. 1991; Turner & Miller 2009). Outflows in AGN offer a potential feedback mechanism through which the central source can affect its environ-

ment (King 2003, 2005; Fabian 2012) – feedback that is required in models of galaxy evolution (Springel, Di Matteo & Hernquist 2005) and may explain the ‘*M*– σ ’ relation (Silk & Rees 1998; Häring & Rix 2004).

Perhaps the clearest evidence of outflows in AGN is the blueshifted ($\sim 0.1c$) broad absorption lines (BALs) in the ultraviolet seen in approximately 20 per cent of quasars (Weymann et al. 1991; Knigge et al. 2008; Allen et al. 2011). The simplest explanation for the incidence of BAL quasars (BALQSOs) is in terms of an accretion disc wind. According to this paradigm, a biconical wind rises from the accretion disc and the BALQSO fraction is associated with the covering factor of the outflow. Polarization studies suggest that the wind is roughly equatorial (Cohen et al. 1995; Goodrich & Miller 1995), although there is also evidence for polar BAL outflows in radio-loud sources (Zhou et al. 2006; Ghosh & Punj 2007).

Due to their ubiquitous nature, disc winds offer a natural way to *unify* much of the diverse phenomenology of luminous AGN and quasars (e.g. Murray et al. 1995; Elvis 2000). Depending on viewing angle, an observer may see a BALQSO or normal ‘Type 1’

*E-mail: jm8g08@soton.ac.uk

quasar. Within this geometric unification framework, the broad-line region (BLR) can correspond either to the dense wind base or clumps embedded in the outflow. A biconical wind model can also readily explain the various sub-classifications of BALQSOs: HiBALQSOs, which only exhibit high-ionization line absorption; LoBALQSOs, which also show absorption in lower ionization state species such as Mg II and Al III; and FeLoBALQSOs, which show further absorption in Fe II and III. In unified geometric models, this is generally attributed to ionization stratification of the outflow (e.g. Elvis 2000).

Despite the importance of disc winds in shaping quasar and AGN spectra, much of the underlying outflow physics remains uncertain. Several driving mechanisms have been proposed, including thermal pressure (Weymann et al. 1982; Begelman, de Kool & Sikora 1991), magnetocentrifugal forces (Blandford & Payne 1982; Pelletier & Pudritz 1992) and radiation pressure on spectral lines ('line-driving'; Lucy & Solomon 1970; Shlosman, Vitello & Shaviv 1985; Murray et al. 1995). Of these, line-driving is possibly the most attractive, as strong absorption lines are already seen in BALQSOs and the X-ray spectra of AGN (Reeves, O'Brien & Ward 2003; Pounds & Reeves 2009; Tombesi et al. 2010). The efficiency of line-driving is crucially dependent on the ionization state of the outflow, and in AGN, unlike O stars, the presence of strong X-ray emission can overionize the wind, causing it to 'fail'. Murray et al. (1995) proposed a potential solution: a region of 'hitchhiking gas' that could shield the wind from the central X-ray source. An additional or alternative solution is that the wind is clumped (e.g. Junkkarinen, Burbidge & Smith 1983; Weymann, Turnshek & Christiansen 1985; Hamann et al. 2013) possibly on multiple scalelengths. Local density enhancements could lower the ionization parameter of the plasma without requiring excessively large total column densities and mass-loss rates.

Evidence for dense substructures in AGN winds is widespread. BALQSOs show complex absorption line profiles (Ganguly et al. 2006; Simon & Hamann 2010) and exhibit variability in these profile shapes (Capellupo et al. 2011, 2012; Capellupo, Hamann & Barlow 2014). AGN generally show variability in X-ray absorption components (e.g. Risaliti, Elvis & Nicastro 2002), and many models for the BLR consist of clumps embedded in an outflow (Krolik, McKee & Tarter 1981; Emmering, Blandford & Shlosman 1992; de Kool & Begelman 1995; Cassidy & Raine 1996). Krolik et al. (1981) showed that BLR clouds would be short-lived, suggesting that a confining mechanism is required, such as magnetic confinement (e.g. de Kool & Begelman 1995). Alternatively, the 'line deshadowing instability' (LDI) inherent in line-driven winds (Lucy & Solomon 1970; MacGregor, Hartmann & Raymond 1979; Carlberg 1980; Owocki & Rybicki 1984) may be responsible for smaller scale clumping, as is observed in O-star winds (Fullerton 2011, and references therein). Indeed, simple models of clumping have successfully explained the electron scattering wings of emission lines formed in these line-driven flows (Hillier 1991). Complex substructures are also produced in simulations of line-driven outflows in AGN, although on very different scales to LDIs (Proga, Stone & Kallman 2000; Proga & Kallman 2004; Proga & Kurosawa 2010; Proga et al. 2014). In summary, clumpy winds offer an observationally motivated and theoretically predicted way to lower the ionization state of a plasma, possibly in tandem with a shielding scenario. However, it is equally important to note that there is no consensus view of the size or location of the clumps in the outflow.

We have been engaged in a project to determine whether it is possible to simulate the properties of the spectra of AGN, including

BALQSOs, using simple kinematic prescriptions for biconical disc winds. To address this question, we use a Monte Carlo radiative transfer (MCRT) code that calculates the ionization structure of the wind and simulates the spectra from such a system (Sim et al. 2008, 2010; Higginbottom et al. 2013, 2014, hereafter H13 and H14). The results have been encouraging in the sense that in H13, we showed we could produce simulated spectra that resembled that of BALQSOs, as long as the luminosity of the X-ray source was relatively low, of the order of 10^{43} erg s⁻¹ and the mass-loss rate was relatively high, of the order of the mass accretion rate. However, at higher X-ray luminosities, the wind was so ionized that UV absorption lines were not produced. In addition, and in part due to limitations in our radiative transfer code, the model failed to produce spectra with strong emission lines at any inclination angle.

Here we attempt to address both of these issues, by introducing clumping into our model and a more complete treatment of H and He into our radiative transfer calculations. The remainder of this paper is organized as follows. In Section 2, we describe some of the important photoionization and MCRT aspects of the code. We then outline the model in Section 3, including a description of our clumping implementation and success criteria. Section 4 contains the results from a clumped model, with comparisons to observational data, as well as some discussion. Finally, we summarize our findings in Section 5.

2 IONIZATION AND RADIATIVE TRANSFER

For this study, we use the MCRT code PYTHON¹ we have developed to carry out our radiative transfer and photoionization simulations in non-local thermodynamic equilibrium (non-LTE). The code can be used to model a variety of disc wind systems; applications have included accreting white dwarfs (Long & Knigge 2002, hereafter LK02; Noebauer et al. 2010; Matthews et al. 2015, hereafter M15), young stellar objects (Sim, Drew & Long 2005) and quasars/AGN (H13; H14).

The code operates as follows. This outflow is discretized into $n_x \times n_z$ cells in a 2.5D cylindrical geometry with azimuthal symmetry. From some initial conditions in each cell (typically $T_e \sim 40\,000$ K in AGN, with dilute LTE abundances), the code first calculates the ionization structure of the wind in a series of iterations. Each iteration in an 'ionization cycle' consists of generating photons, actually photon packets, from an accretion disc and central object, and calculating how these photon bundles scatter through the wind (eventually escaping the outflow or hitting the disc). The ionization and temperature structure is updated based on the properties of the radiation field in each cell, and the process is repeated. The radiative transfer and thermal balance are carried out according to the principle of radiative equilibrium, which assumes both statistical equilibrium and that radiative heating balances radiative cooling. Once the ionization structure has converged, it is held fixed, and synthetic spectra are generated at specific inclination angles in a series of 'spectral cycles'. LK02 provide a more detailed description of the original code; various improvements have been made since then and are described by Sim et al. (2005), H13 and M15. We focus here on the specific changes made for this study intended to improve the ionization calculation of H and He and to allow for clumping in the wind.

¹ Named *c. 1995*, predating the inexorable rise of a certain programming language.

2.1 Line transfer

Our approach to line transfer is based upon the macro-atom implementation developed by Lucy (2002, 2003), in which the energy flows through the system are described in terms of indivisible energy quanta of radiant or kinetic energy ('*r*-packets' and '*k*-packets', respectively; see also Section 3.1), which are reprocessed by 'macro-atoms' with associated transition probabilities. We use the Sobolev approximation (e.g. Sobolev 1957, 1960; Rybicki & Hummer 1978) to compute the location and optical depth of line interactions. In our case, for reasons of computational efficiency, we adopt the hybrid macro-atom scheme described by M15. In this scheme, the energy packets interact with either two-level 'simple atoms' or full macro-atoms. This allows one to treat non-LTE line transfer in radiative equilibrium without approximation for elements that are identified as full macro-atoms, while maintaining the fast 'two-level' treatment of resonance lines when elements are identified as simple atoms (see M15). In this study, only H and He are treated as macro-atoms, because the process is computationally intensive, and we expect recombination to be particularly important in determining their level populations and resultant line emission. We are also especially interested in the contribution to AGN spectra of Lyman α . H13 treated all atoms in a two-level approximation.

2.2 Ionization and excitation treatment

Macro-atoms have their ion and level populations derived from MC rate estimators as described by Lucy (2002, 2003). Previously (LK02; H13; M15), we used a modified Saha approach to calculate the ionization fractions of simple atoms. As part of this effort, we have now improved PYTHON to explicitly solve the rate equations between ions in non-LTE. This dispenses with a number of small assumptions made in the modified Saha approach, is more numerically stable, and prepares for a more accurate treatment of other physical processes in future.

In order to calculate the photoionization rate, we model the spectral energy distribution (SED) in a grid cell using the technique described by H13. In this scheme, the mean intensity, J_ν , in a series of n bands is modelled as either a power law or exponential in frequency ν , with the fit parameters deduced from band-limited radiation field estimators. This allows the calculation of a photoionization rate estimator. Ion abundances are then calculated by solving the rate equations between ions. We include collisional ionization and photoionization balanced with radiative, dielectronic and collisional (three-body) recombination. As in M15, we use a dilute Boltzmann approximation to calculate the populations of levels for simple atoms. This dilute approximation is not required for macro-atom levels.

2.3 Physical processes

We include free-free, bound-free and bound-bound heating and cooling processes in the model. For radiative transfer purposes, we treat electron scattering in the Thomson limit, but take full account of Compton heating and cooling when calculating the thermal balance of the plasma (see H13). Adiabatic cooling in a cell is calculated using the estimator

$$C_A = f_V k_B T_e V (\nabla \cdot \mathbf{v}) \left(n_e + \sum_{i=1}^{\text{ions}} N_i \right), \quad (1)$$

where V is the cell volume, T_e is the electron temperature, n_e is the electron density, N_i is the number density of ion i and $\nabla \cdot \mathbf{v}$ is the

divergence of the velocity field at the centre of the cell. The sum is over all ions included in the simulation. For consistency, k -packets then have a destruction probability of

$$P_{k,A} = \frac{C_A}{C_{\text{tot}}}, \quad (2)$$

where C_{tot} is the sum over all cooling rates in the cell.

2.4 Atomic data

We use the same atomic data described by LK02 as updated by H13 and M15. As in M15, macro-atom level information is from TOPBASE (Cunto et al. 1993). Photoionization cross-sections are from TOPBASE (Cunto et al. 1993; Verner et al. 1996). Dielectronic and radiative recombination rate coefficients for simple atoms are taken from the CHIANTI data base version 7.0 (Dere et al. 1997; Landi et al. 2012). We use ground-state recombination rates from Badnell (2006) where available, and otherwise default to calculating recombination rates from the Milne relation. Free-free Gaunt factors are from Sutherland (1998).

3 A CLUMPY BICONICAL DISC WIND MODEL FOR QUASARS

Our kinematic prescription for a biconical disc wind model follows Shlosman & Vitello (1993, hereafter SV93), and is described further by LK02, H13 and M15. The purpose of this purely kinematic wind model is to provide a simple tool for exploring different outflow geometries. A schematic is shown in Fig. 1, with key aspects marked. The general biconical geometry is similar to that invoked by Murray et al. (1995) and Elvis (2000) to explain the phenomenology of quasars and BALQSOs. However, we do not include the initial vertical component present in the Elvis (2000) model, which was invoked as a qualitative explanation for narrow absorption lines.

3.1 Photon sources

We include two sources of *r*-packets in our model: an accretion disc and a central X-ray source. The accretion disc is assumed to be geometrically thin, but optically thick. Accordingly, we treat the disc as an ensemble of blackbodies with a Shakura & Sunyaev (1973)

Figure 1. A cartoon showing the geometry and some key parameters of our biconical wind model.

effective temperature profile. The emergent SED is then determined by the specified accretion rate ($\dot{m}$) and central BH mass (M_{BH}). All photon sources in our model are opaque, meaning that r -packets that strike them are destroyed. The inner radius of the disc extends to the innermost stable circular orbit (ISCO) of the BH. We assume a Schwarzschild BH with an ISCO at $6r_{\text{G}}$, where $r_{\text{G}} = GM_{\text{BH}}/c^2$ is the gravitational radius. For a $10^9 M_{\odot}$ BH, this is equal to 8.8×10^{14} cm or $\sim 10^{-4}$ pc.

The X-ray source is treated as an isotropic sphere at the ISCO, which emits r -packets according to a power law in flux with index α_{X} , of the form

$$F_{\text{X}}(\nu) = K_{\text{X}} \nu^{\alpha_{\text{X}}}. \quad (3)$$

The normalization, K_{X} , of this power law is such that it produces the specified 2–10 keV luminosity, L_{X} . Photons, or r -packets, produced by the accretion disc and central X-ray source are reprocessed by the wind. This reprocessing is dealt with by enforcing strict radiative equilibrium (*modulo* adiabatic cooling; see Section 2.3) via an indivisible energy packet constraint (see Lucy 2002; M15).

3.2 Kinematics and geometry

In the SV93 model, a biconical disc wind rises from the accretion disc between launch radii r_{min} and r_{max} . The opening angles of the wind are set to θ_{min} and θ_{max} . The poloidal velocity along each individual streamline at a poloidal distance l is then given by (e.g. Rybicki & Hummer 1978)

$$v_l = v_0 + [v_{\infty}(r_0) - v_0] \frac{(l/R_v)^{\alpha}}{(l/R_v)^{\alpha} + 1}, \quad (4)$$

where v_0 is the velocity at the base of the streamline, α is an exponent governing how quickly the wind accelerates and R_v is the ‘acceleration length’, defined as the distance at which the outflow reaches half of its terminal velocity, v_{∞} . The terminal velocity is set to a fixed multiple of the escape velocity, v_{esc} , at the base of the streamline (radius r_0). The rotational velocity, v_{ϕ} , is initially Keplerian ($v_{\phi} = [GM/r_0]^{1/2}$), and the wind conserves specific angular momentum, such that

$$v_{\phi} r = v_{\phi} r_0. \quad (5)$$

The velocity law is crucial in determining the output spectra, as it affects not only the projected velocities along the line of sight, but also the density and ionization state of the outflow. A wind that accelerates more slowly will have a denser wind base with correspondingly different ionization and emission characteristics.

3.3 A simple approximation for clumping

In our previous modelling efforts, we assumed a smooth outflow, in which the density at a given point was determined only by the kinematic parameters and mass-loss rate. However, as already discussed, AGN winds exhibit significant substructure – the outflow is expected to be *clumpy*, rather than smooth, and probably on a variety of scales. A clumpy outflow offers a possible solution to the so-called ‘overionization problem’ in quasar and AGN outflows (e.g. Junkkarinen et al. 1983; Weymann et al. 1985; Hamann et al. 2013). This is the main motivation for incorporating clumping into our model.

Deciding on how to implement clumping into our existing wind models was not straightforward. First, and most importantly, the physical scalelengths and density contrasts in AGN outflows are not well constrained from observations or theory. As a result, while one

could envision in principle, clouds with a variety of sizes and density contrasts varying perhaps as a function of radius, there would have been very little guidance on how to set nominal values of the various parameters of such a model. Secondly, there are significant computational difficulties associated with adequately resolving and realistically modelling a series of small-scale, high-density regions with an MCRT – or for that matter, a hydrodynamical – code. Given the lack of knowledge about the actual type of clumping, we have adopted a simple approximation used successfully in stellar wind modelling, known as *microclumping* (e.g. Hamann & Koesterke 1998; Hillier & Miller 1999; Hamann, Oskinova & Feldmeier 2008).

The underlying assumption of microclumping is that clump sizes are much smaller than the typical photon mean free path, and thus the clumps are both geometrically and optically thin. This approach allows one to treat clumps only in terms of their volume filling factor, f_v , instead of having to specify separately their size and density distributions. In our model, f_v is independent of position (see Section 4.4). The inter-clump medium is modelled as a vacuum, although the outflow is still non-porous and axisymmetric. This approach therefore assumes that the inter-clump medium is unimportant in determining the output spectrum, which we expect to be true only when density contrasts are large and the inter-clump medium is both very ionized and of low emissivity and opacity. The density of the clumps is multiplied by the ‘density enhancement’ $D = 1/f_v$. Opacities, κ , and emissivities, ϵ , can then be expressed as

$$\kappa = f_v \kappa_C(D); \quad \epsilon = f_v \epsilon_C(D). \quad (6)$$

Here the subscript C denotes that the quantity is calculated using the enhanced density in the clump. The resultant effect is that, *for fixed temperature*, processes that are linear in density, such as electron scattering, are unchanged, as f_v and D will cancel out. However, any quantity that scales with the square of density, such as collisional excitation or recombination, will increase by a factor of D . In our models, the temperature is not fixed, and is instead set by balancing heating and cooling in a given cell. In the presence of an X-ray source, this thermal balance is generally dominated by bound–free heating and line cooling. The main effect of including clumping in our modelling is that it moderates the ionization state due to the increased density. This allows an increase in the ionizing luminosity, amplifying the amount of bound–free heating and also increasing the competing line cooling term (thermal line emission).

The shortest length-scale in a Sobolev MCRT treatment such as ours is normally the Sobolev length, given by

$$l_s = \frac{v_{\text{th}}}{|dv/ds|}. \quad (7)$$

This is typically $\sim 10^{13}$ cm near the disc plane, increasing outwards. We use the mean density to calculate the Sobolev optical depth, which assumes that l_s is greater than the typical clump size. Thus, for the microclumping assumption to be formally correct, clumps should be no larger than $\sim 10^{12}$ cm. This size scale is not unreasonable for quasar outflows, as de Kool & Begelman (1995) suggest that BAL flows may have low filling factors with clump sizes of $\sim 10^{11}$ cm.

Our clumping treatment is necessarily simple; it does not adequately represent the complex substructures and stratifications in ionization state we expect in AGN outflows. Nevertheless, this parametrization allows simple estimates of the effect clumping has on the ionization state and emergent line emission.

Table 1. The grid points used in the parameter search. The sensitivity to some of these parameters is discussed further in Section 4.5.

Parameter	Grid point values			
$r_{\min}$	$60r_g$	$180r_g$	$300r_g$	
$\theta_{\min}$	55°	70°		
R_v	10^{18} cm	10^{19} cm		
α	0.5	0.6	0.75	1.5
f_V	0.01	0.1		

3.4 The simulation grid

Using this prescription, we conducted a limited parameter search over a five-dimensional parameter space involving the variables $r_{\min}$, $\theta_{\min}$, f_V , α and R_v . The grid points are shown in Table 1. The aim here was to first fix M_{BH} and $\dot{m}$ to their H13 values, and increase L_X to 10^{45} erg s $^{-1}$ (a more realistic value for a quasar of $10^9 M_\odot$ and an Eddington fraction of 0.2; see Section 4.3).

We then evaluated these models based on how closely their synthetic spectra reproduced the following properties of quasars and BALQSOs:

- (i) UV absorption lines with BI > 0 at ~ 20 per cent of viewing angles (e.g. Knigge et al. 2008);
- (ii) line emission emerging at low inclinations, with equivalent width (EW) ~ 40 Å in C IV 1550 Å (e.g. Shen et al. 2011);
- (iii) H recombination lines with EW ~ 50 Å in Ly α (e.g. Shen et al. 2011);
- (iv) Mg II and Al III (LoBAL) absorption features with BI > 0 at a subset of BAL viewing angles;
- (v) verisimilitude with quasar composite spectra.

Here BI is the ‘balnicity index’ (Weymann et al. 1991), given by

$$\text{BI} = \int_{3000 \text{ km s}^{-1}}^{25000 \text{ km s}^{-1}} C \left(1 - \frac{f(v)}{0.9} \right) dv. \quad (8)$$

The constant $C = 0$ everywhere, unless the normalized flux has satisfied $f(v) < 0.9$ continuously for at least 2000 km s $^{-1}$, whereby C is set to 1.

In the next section, we present one of the most promising models, which we refer to as the fiducial model, and discuss the various successes and failures with respect to the above criteria. This allows us to gain insight into fundamental geometrical and physical constraints and assess the potential for unification. We then discuss the sensitivity to key parameters in Section 4.5. The full grid, including output synthetic spectra and plots, can be found at jhmattews.github.io/quasar-wind-grid/.

4 RESULTS AND DISCUSSION FROM A FIDUCIAL MODEL

Here we describe the results from a fiducial model, and discuss these results in the context of the criteria presented in Section 3.4. The parameters of this model are shown in Table 2. Parameters differing from the benchmark model of H13 are highlighted with an asterisk. In this section, we examine the physical conditions of the flow, and present the synthetic spectra, before comparing the X-ray properties of this particular model to samples of quasars and luminous AGN.

4.1 Physical conditions and ionization state

Fig. 2 shows the physical properties of the wind. The wind rises slowly from the disc at first, with densities within clumps of $n_H \sim$

Table 2. Wind geometry parameters used in the fiducial model, as defined in the text and Fig. 1. Parameters differing from the benchmark model of H13 are highlighted with an asterisk.

Fiducial model parameters	Value
M_{BH}	$1 \times 10^9 M_\odot$
$\dot{m}_{\text{acc}}$	$5 M_\odot \text{ yr}^{-1} \simeq 0.2 \dot{M}_{\text{Edd}}$
α_X	-0.9
L_X	$10^{45} \text{ erg s}^{-1}$ *
$r_{\text{disc}}(\min) = r_X$	$6r_g = 8.8 \times 10^{14} \text{ cm}$
$r_{\text{disc}}(\max)$	$3400r_g = 5 \times 10^{17} \text{ cm}$
$\dot{m}_{\text{wind}}$	$5 M_\odot \text{ yr}^{-1}$
$r_{\min}$	$300r_g = 4.4 \times 10^{16} \text{ cm}$
$r_{\max}$	$600r_g = 8.8 \times 10^{16} \text{ cm}$
$\theta_{\min}$	70°
$\theta_{\max}$	82°
$v_\infty(r_0)$	$v_{\text{esc}}(r_0)$
R_v	10^{19} cm *
α	0.5 *
f_V	0.01 *
n_x	100
n_z	200

10^{11} cm^{-3} close to the disc plane, where n_H is the local number density of H. The flow then accelerates over a scalelength of $R_v = 10^{19} \text{ cm}$ up to a terminal velocity equal to the escape velocity at the streamline base ($\sim 10\,000 \text{ km s}^{-1}$). This gradual acceleration results in a wind that exhibits a stratified ionization structure, with low-ionization material in the base of the wind giving way to highly ionized plasma further out. This is illustrated in Fig. 2 by the panels showing the ion fraction $F = n_j/n_{\text{tot}}$ of some important ions. With a clumped wind, we are able to produce the range of ionization states observed in quasars and BALQSOs, while adopting a realistic 2–10 keV X-ray luminosity of $L_X = 10^{45} \text{ erg s}^{-1}$. Without clumping, this wind would be overionized to the extent that opacities in e.g. C IV would be entirely negligible (see H13).

One common way to quantify the ionization state of a plasma is through the ionization parameter, U_H , given by

$$U_H = \frac{4\pi}{n_{\text{HC}}} \int_{13.6 \text{ eV}/h}^{\infty} \frac{J_\nu d\nu}{h\nu}, \quad (9)$$

where ν denotes photon frequency. Shown in Fig. 2, the ionization parameter is a useful measure of the global ionization state, as it represents the ratio of the number density of H ionizing photons to the local H density. It is, however, a poor representation of the ionization state of species such as C IV as it encodes no information about the shape of the SED. In our case, the X-ray photons are dominant in the photoionization of the UV resonance line ions. This explains why a factor of 100 increase in X-ray luminosity requires a clumping factor of 0.01, even though the value of U_H decreases by only a factor of ~ 10 compared to H13.

The total line luminosity also increases dramatically compared to the unclumped model described by H13. This is because the denser outflow can absorb the increased X-ray luminosity without becoming overionized, leading to a hot plasma which produces strong collisionally excited line emission. This line emission typically emerges on the edge of the wind nearest to the central source. The location of the line emitting regions is dependent on the ionization state, as well as the incident X-rays. The radii of these emitting regions are important, and can be compared to observations. The line luminosities, L , shown in the figure correspond to the

Figure 2. Contour plots showing the logarithm of some important physical properties of the outflow. The spatial scales are logarithmic, and the x and z scales are not the same. Symbols are defined in the text. The solid black line marks a sphere at $1000 r_G$. The dotted lines show the 72° and 78° sightlines to the centre of the system, and illustrate that different sightlines intersect material of different ionization states. The line luminosities, L , represent the luminosity of photons escaping the Sobolev region for each line. These photons do not necessarily escape to infinity.

Figure 3. Synthetic spectra at four viewing angles for the fiducial model. At 20° and 60° , we show a comparison to an SDSS quasar composite from Reichard et al. (2003). At 73° and 76° , we show a comparison to an *HST* Space Telescope Imaging Spectrograph (STIS) spectrum of the high-balnicity BALQSO PG0946+301 (Arav et al. 2001), and an SDSS spectrum of the LoBAL quasar SDSS J162901.63+453406.0, respectively. The dotted line shows a disc only continuum to show the effect of the outflow on the continuum level. All the spectra are scaled to the model flux at $2000 \text{ \AA}$, except for the *HST* STIS spectrum of PG0946+301, which is scaled to $1350 \text{ \AA}$ due to the incomplete wavelength coverage.

luminosity in erg s^{-1} of photons escaping the Sobolev region for each line. As shown in Fig. 2, the $\text{C IV } 1550 \text{ \AA}$ line in the fiducial model is typically formed between 100 and $1000 r_G$ ($\sim 10^{17}$ – 10^{18} cm). This is in rough agreement with the reverberation mapping results of Kaspi et al. (2007) for the $2.6 \times 10^9 M_\odot$ quasar S5 0836+71, and also compares favourably with microlensing measurements of the size of the $\text{C IV } 1550 \text{ \AA}$ emission line region in the BALQSO H1413+117 (O’Dowd et al. 2015).

4.2 Synthetic spectra: comparison to observations

Fig. 3 shows the synthetic spectrum in the UV from the fiducial model. To assess the ability of the synthetic spectra to match real quasar spectra, we also show Sloan Digital Sky Survey (SDSS) quasar composites from Reichard et al. (2003), normalized to the flux at $2000 \text{ \AA}$ for low inclinations. Unfortunately, the wide variety of line profile shapes and internal trough structure in BALQSOs tend to ‘wash out’ BAL troughs in composite spectra to the extent that BALQSO composites do not resemble typical BALQSOs. Because of this, we instead compare to a *Hubble Space Telescope* (*HST*) STIS spectrum of the high-balnicity BALQSO PG0946+301 (Arav

Figure 4. A cartoon describing the broad classes of sightline in the fiducial model, illustrating how geometric effects lead to the different emergent spectra. The colour gradient is approximate, but indicates the stratified ionization structure, from highly ionized (yellow) to low-ionization (purple) material.

et al. 2001), and an SDSS spectrum of the LoBAL quasar SDSS J162901.63+453406.0, for the angles of 73° and 76° , respectively. We show a cartoon illustrating how geometric effects determine the output spectra in Fig. 4.

4.2.1 Broad absorption lines ('BALQSO-like' angles)

The UV spectrum is characterized by strong BAL profiles at high inclinations ($>70^\circ$). This highlights the first success of our model: clumping allows the correct ionization state to be maintained in the presence of strong X-rays, resulting in large resonance line opacities. At the highest inclinations, the cooler, low-ionization material at the base of the wind starts to intersect the line of sight. This produces multiple absorption lines in species such as Mg II, Al III and Fe II. The potential links to LoBALQSOs and FeLoBALQSOs are discussed in Section 2.4.

The high-ionization BAL profiles are often saturated, and the location in velocity space of the strongest absorption in the profile varies with inclination. At the lowest inclination BAL sightlines, the strongest absorption occurs at the red edge, whereas at higher inclinations (and for the strongest BALs) the trough has a sharp edge at the terminal velocity. This offers one potential explanation for the wide range of BALQSO absorption line shapes (see e.g. Trump et al. 2006; Knigge et al. 2008; Filiz Ak et al. 2014).

The absorption profiles seen in BALQSOs are often non-black, but saturated, with flat bases to the absorption troughs (Arav et al. 1999a,b). This is usually explained either as partial covering of the continuum source or by scattered contributions to the BAL troughs, necessarily from an opacity source not co-spatial with the BAL forming region. The scattered light explanation is supported by spectropolarimetry results (Lamy & Hutsemékers 2000). Our spectra do not show non-black, saturated profiles. We find black, saturated troughs at angles $i > 73^\circ$, and the BALs are non-saturated at lower inclinations. The reasons for this are inherent in the construction of our model. First, the microclumping assumption does not allow for porosity in the wind, meaning that it does not naturally produce a partial covering absorber. To allow this, an alternative approach such as *macroclumping* would be required (e.g. Hamann et al. 2008; Šurlan et al. 2012). Secondly, our wind does not have a significant scattering contribution along sightlines which do not pass through the BAL region, meaning that any scattered component to the BAL troughs is absorbed by line opacity. This suggests that either the scattering cross-section of the wind must be increased (with higher mass-loss rates or covering factors) or that an additional source of electron opacity is required, potentially in a polar direction above the disc. We note that the scattering contribution from plasma in polar regions is significant in some 'outflow-from-inflow' simulations (Kurosawa & Proga 2009; Sim et al. 2012).

4.2.2 Broad emission lines ('quasar-like' angles)

Unlike H13, we now find significant collisionally excited line emission emerges at low inclinations in the synthetic spectra, particular in the C IV and N V lines. We also find a strong Ly α line and weak He II 1640 Å line as a result of our improved treatment of recombination using macro-atoms. In the context of unification, this is a promising result, and shows that a biconical wind can produce significant emission at 'quasar-like' angles. To demonstrate this further, we show line luminosities and monochromatic continuum luminosities from the synthetic spectra in Table 3. These are compared to mean values from a subsample of the SDSS DR7 quasar catalogue (Shen et al. 2011) with BH mass and Eddington fraction estimates similar to the fiducial model values (see the caption). The spectra do not contain the strong C III] 1909 Å line seen in the quasar composite spectra, but this is due to a limitation of our current treatment of C; semi-forbidden (intercombination) lines are not included in our modelling.

Table 3. Some derived spectral properties of the fiducial model, at 20° , compared to observations. The observed values are taken from the Shen et al. (2011) SDSS DR7 quasar catalogue, and correspond to mean values with standard deviations in log space from a subsample with $8.5 > \log(M_{\text{BH}}) < 9.5$ and $1.5 < \log(L_{\text{bol}}/L_{\text{Edd}}) < 0$, where the BH mass is a C IV virial estimate. Units are logarithms of values in erg s^{-1} .

Property	Synthetic, 20°	Observed (S11)
$\log L[\text{C IV}]$	44.60	44.42 ± 0.32
$\log L[\text{Mg II}]$	43.92	43.54 ± 0.28
$\log(\nu L_\nu)_{1350}$	46.42	46.01 ± 0.30
$\log(\nu L_\nu)_{3000}$	46.18	45.79 ± 0.30

In Fig. 5, we show an F_ν spectrum with broader waveband coverage that includes the optical, showing that our synthetic spectra also exhibit H α and H β emission. In this panel, we include a low inclination and also a very high inclination spectrum, which looks underneath the wind cone. This model shows strong line emission with very similar widths and line ratios to the quasar composites, and the Balmer lines are double peaked, due to velocity projection effects. Such double-peaked lines are seen in so-called 'disc emitter' systems (e.g. Eracleous & Halpern 1994) but not the majority of AGN. The line EWs increase at high inclination due to a weakened continuum from wind attenuation, disc foreshortening and limb darkening. This effect also leads to a redder continuum slope, as seen in quasars, which is due to Balmer continuum and Balmer and Fe II line emission.

The 89° viewing angle cannot represent a typical quasar within a unified model, as extreme inclinations should be extremely under-represented in quasar samples. This is in part due to the Compton-thick absorber or 'torus' expected at high inclinations (e.g. Antonucci & Miller 1985; Martínez-Sansigre et al. 2007). However, this spectrum does show that a wind model can naturally reproduce quasar emission lines if the emissivity of the wind is increased *with respect to the disc continuum*. In addition, it neatly demonstrates how a stratified outflow can naturally reproduce the range of ionization states seen in quasars.

Despite a number of successes, there are some properties of the synthetic spectra that are at odds with the observations. First, the ratios of the EW of the Ly α and Mg II 2800 Å lines to the EW of C IV 1550 Å are much lower than in the composite spectra. Similar problems have also been seen in simpler photoionization models for the BLR (Netzer 1990). It may be that a larger region of very dense ($n_e \sim 10^{10} \text{cm}^{-3}$) material is needed, which could correspond to a disc atmosphere or 'transition region' (see e.g. Murray et al. 1995; Knigge & Drew 1997). While modest changes to geometry may permit this, the initial grid search did not find a parameter space in which the Ly α or Mg II EWs were significantly higher (see Section 4.5). Secondly, we find that EWs increase with inclination (see Figs 3 and 5; also Fig. 7), to the extent that, even though significantly denser models can match the line EWs fairly well at low inclinations, they will then possess overly strong red wings to the BAL P-Cygni profiles at high inclinations. The fact that the EW increase in our model is directly related to limb darkening and foreshortening of the continuum. This appears to contradict observations, which show remarkably uniform emission line properties in quasars and BALQSOs (Weymann et al. 1991; DiPompeo et al. 2012). The angular distribution of the disc continuum and line emission is clearly crucially important in determining the emergent broad-line EWs, as suggested by, e.g., the analysis of Risaliti, Salvati & Marconi (2011). We shall explore this question further in a future study.

Figure 5. Synthetic spectra at two viewing angles, this time in frequency space and including the optical band, compared to the non-BAL SDSS quasar composite. The spectra are normalized to the flux at 2000 Å, then an offset of 2 is applied per spectrum for clarity – the dotted lines show the zero-point of F_v/F_{2000} in each case.

Figure 6. X-ray (2 keV) luminosity of our clumped model (red squares) and the H13 model (blue squares), plotted against monochromatic luminosity at 2500 Å. The points are labelled according to inclination; angles $>70^\circ$ correspond to BALs in our scheme (see Fig. 4). Also plotted are measurements from the COMBI-7 AGN and the BQS samples (Steffen et al. 2006) and the Saez et al. (2012) sample of BALQSOs. The dotted line shows the best-fitting relation for non-BALQSOs from Steffen et al. (2006).

4.3 X-ray properties

The main motivation for adding clumping to the model was to avoid overionization of the wind in the presence of strong X-rays. Having verified that strong BALs appear in the synthetic spectra, it is also important to assess whether the X-ray properties of this fiducial model agree well with quasar and BALQSO samples for the relevant inclinations.

Fig. 6 shows the emergent monochromatic luminosity (L_v) at 2 keV and plotted against L_v at 2500 Å for a number of different viewing angles in our model. The monochromatic luminosities are calculated from the synthetic spectra and thus include the effects of wind reprocessing and attenuation. In addition to model outputs, we also show the BALQSO sample of Saez et al. (2012) and luminous AGN and quasar samples from Steffen et al. (2006). The best-fitting relation from Steffen et al. (2006) is also shown. For low inclination,

‘quasar-like’ viewing angles, we now find excellent agreement with AGN samples. The slight gradient from 20° to 60° in our models is caused by a combination of disc foreshortening and limb darkening (resulting in a lower L_{2500} for higher inclinations), and the fact that the disc is opaque, and thus the X-ray source subtends a smaller solid angle at high inclinations (resulting in a lower $L_{2\text{keV}}$ for higher inclinations).

The high-inclination, ‘BALQSO-like’ viewing angles show moderate agreement with the data, and are X-ray weak due to bound-free absorption and electron scattering in the wind. Typically, BALQSOs show strong X-ray absorption with columns of $N_H \sim 10^{23} \text{ cm}^{-2}$ (Green & Mathur 1996; Mathur et al. 2000; Green et al. 2001; Grupe, Mathur & Elvis 2003). This is often cited as evidence that the BAL outflow is shielded from the X-ray source, especially as sources with strong X-ray absorption tend to exhibit deep BAL

troughs and high outflow velocities (Brandt, Laor & Wills 2000; Laor & Brandt 2002; Gallagher et al. 2006). Our results imply that the clumpy BAL outflow itself can be responsible for the strong X-ray absorption, and supports Hamann et al.’s (2013) suggestion that geometric effects explain the weaker X-ray absorption in mini-BALs compared to BALQSOs.

4.4 LoBALs and ionization stratification

At high inclinations, the synthetic spectra exhibit blueshifted BALs in Al III and Mg II – the absorption lines seen in LoBALQSOs, and we even see absorption in Fe II at the highest inclinations. Line profiles in velocity space for C IV, Al III and Mg II are shown in Fig. 7 for a range of BALQSO viewing angles. We find that ionization stratification of the wind causes lower ionization material to have a smaller covering factor, as demonstrated by Figs 2 and 7. This confirms the behaviour expected from a unification model such as Elvis (2000). LoBALs are only present at viewing angles close to edge-on ($i > 75^\circ$), as predicted by polarization results (Brotherton et al. 1997). As observed in a BALQSO sample by Filiz Ak et al. (2014), we find that BAL troughs are wider and deeper when low-ionization absorption features are present, and high-ionization lines have higher blue-edge velocities than the low-ionization species.

There is also a correlation between the strength of LoBAL features and the amount of continuum attenuation at that sightline, particularly bluewards of the Lyman edge as the low-ionization base intersects the line of sight. A model such as this therefore predicts that LoBALQSOs and FeLoBALQSOs have stronger Lyman edge absorption and are more Compton-thick than HiBALQSOs and Type 1 quasars. An edge-on scenario also offers a potential explanation for the rarity of LoBAL and FeLoBAL quasars, due to a foreshortened and attenuated continuum, although BAL fraction

inferences are fraught with complex selection effects (Goodrich 1997; Krolik & Voit 1998).

4.5 Parameter sensitivity

Having selected an individual fiducial model from the simulation grid, it is important to briefly explore how specialized this model is, and how small parameter changes can affect the synthetic spectra. Fig. 8 shows the EW for C IV 1550 Å and Mg II 2800 Å at a low inclination, and BI for C IV 1550 Å at a high inclination for the simulation grid. We find that almost all the models with $f_V = 0.1$ are overionized, and fail to produce strong C IV BALs or emission lines. However, the models with $f_V = 0.01$ generally produce C IV BALs and emission lines. The fiducial model is representative of this family of models and the spectra are generally similar, although it is clear from the figure that we have selected a model near the upper end of the EW distributions.

We find that it is difficult to significantly increase line emission while keeping the luminosity and mass-loss rate of the system fixed. We show an additional point in Fig. 7 corresponding to a model with an order of magnitude higher X-ray luminosity and double the mass-loss rate. As expected, this results in far higher line EWs, but fails to produce BALs because the collisionally excited emission swamps the BAL profile. In addition, this model would lie well above the expected $L_{2\text{keV}}-L_{2500}$ relation in Fig. 5. Such a high X-ray luminosity could therefore not be the cause of the strong line emission seen in all Type 1 quasars.

The parameter search presented here is by no means exhaustive, and we may be limited by the specific parametrization of the outflow kinematics we have used. Nevertheless, we suggest that the angular distribution of both the line and continuum emission is perhaps the crucial aspect to understand. With this in mind, obtaining reliable

Figure 7. C IV, Mg II, Al III and Fe II line profiles for viewing angles from 72° – 79° . The profiles are plotted relative to the local continuum with an offset applied for clarity. Lower ionization profiles appear at a subset of high inclinations, compared to the ubiquitous C IV profile.

Figure 8. The EW of the C IV 1550 Å line at 20° plotted against (a) the BI of C IV 1550 Å at 75°, (b) the EW of the Mg II 2800 Å line at 20° and (c) the EW of Ly α at 20°. The circles correspond to the simulation grid for two different values of f_V , and the fiducial model is marked with an orange star. We also show a higher X-ray luminosity model and a higher mass-loss rate with a red star.

orientation indicators appears to be a crucial observational task if we are to further our understanding of BAL outflows and their connection, or lack thereof, to the BLR.

5 SUMMARY AND CONCLUSIONS

We have carried out MCRT simulations using a simple prescription for a biconical disc wind, with the aim of expanding on the work of H13. To do this, we introduced two main improvements: First, we included a simple treatment of clumping, and secondly, we improved the modelling of recombination lines by treating H and He as ‘macro-atoms’. Having selected a fiducial model from an initial simulation grid, we assessed the viability of such a model for geometric unification of quasars, and found the following main points.

(i) Clumping the wind with a volume filling factor of 0.01 moderates the ionization state sufficiently to allow for the formation of strong UV BALs while agreeing well with the X-ray properties of luminous AGN and quasars.

(ii) A clumpy outflow model naturally reproduces the range of ionization states expected in quasars, due to its stratified density and temperature structure. LoBAL line profiles are seen at a subset of viewing angles, and Fe II absorption is seen at particularly high inclinations.

(iii) The synthetic spectra show a Ly α line and weak He II 1640 Å line as a result of our improved treatment of recombination using macro-atoms. We also see Balmer emission lines and a Balmer recombination continuum in the optical spectrum, but this is only really significant at high inclination where the continuum is suppressed.

(iv) The higher X-ray luminosity causes a significant increase in the strength of the collisionally excited emission lines produced by the model. However, the EW ratios of the emission lines do not match observations, suggesting that a greater volume of dense ($n_e \sim 10^{10} \text{ cm}^{-3}$) material may be required.

(v) The line EWs in the synthetic spectra increase with inclination. BAL and non-BAL quasar composites have comparable EWs, so our model fails to reproduce this behaviour. If the BLR emits fairly isotropically, then for a foreshortened, limb-darkened accretion disc it is not possible to achieve line ratios at low inclinations that are comparable to those at high inclinations. We suggest that understanding the angular distribution of line and continuum emission is a crucial question for theoretical models.

Our work confirms a number of expected outcomes from a geometric unification model, and suggests that a simple biconical geometry such as this can come close to explaining much of the phenomenology of quasars. However, our conclusions pose some challenges to a picture in which BALQSOs are explained by an *equatorial* wind rising from a classical thin disc, and suggest that the angular distribution of emission is important to understand if this geometry is to be refuted or confirmed. We suggest that obtaining reliable observational orientation indicators and exploring a wider parameter space of outflow geometries in simulations are obvious avenues for future work.

ACKNOWLEDGEMENTS

The work of JHM, SWM, NSH and CK is supported by the Science and Technology Facilities Council (STFC), via two studentships and a consolidated grant, respectively. CK also acknowledges a Leverhulme fellowship. We would like to thank the anonymous referee for a helpful and constructive report. We would also like to thank Omer Blaes, Ivan Hubeny and Shane Davis for their assistance with AGNSPEC. We are grateful to Mike Brotherton, Mike DiPompeo, Sebastien Hoenig and Frederic Marin for helpful correspondence regarding polarization measurements and orientation indicators. We would also like to thank Daniel Proga, Daniel Capellupo, Sam Connolly and Dirk Grupe for useful discussions. Simulations were conducted using PYTHON version 80, and made use of the IRIDIS High Performance Computing Facility at the University of Southampton. Figures were produced using the matplotlib plotting library (Hunter 2007). This work made use of the Sloan Digital Sky Survey. Funding for the Sloan Digital Sky Survey has been provided by the Alfred P. Sloan Foundation, the US Department of Energy Office of Science and the Participating Institutions.

REFERENCES

- Allen J. T., Hewett P. C., Maddox N., Richards G. T., Belokurov V., 2011, *MNRAS*, 410, 860
- Antonucci R. R. J., Miller J. S., 1985, *ApJ*, 297, 621
- Arav N., Korista K. T., de Kool M., Junkkarinen V. T., Begelman M. C., 1999a, *ApJ*, 516, 27
- Arav N., Becker R. H., Laurent-Muehleisen S. A., Gregg M. D., White R. L., Brotherton M. S., de Kool M., 1999b, *ApJ*, 524, 566
- Arav N. et al., 2001, *ApJ*, 561, 118
- Badnell N. R., 2006, *ApJS*, 167, 334

- Begelman M., de Kool M., Sikora M., 1991, *ApJ*, 382, 416
- Blandford R. D., Payne D. G., 1982, *MNRAS*, 199, 883
- Brandt W. N., Laor A., Wills B. J., 2000, *ApJ*, 528, 637
- Brotherton M. S., Tran H. D., van Breugel W., Dey A., Antonucci R., 1997, *ApJ*, 487, L113
- Capellupo D. M., Hamann F., Shields J. C., Rodríguez Hidalgo P., Barlow T. A., 2011, *MNRAS*, 413, 908
- Capellupo D. M., Hamann F., Shields J. C., Rodríguez Hidalgo P., Barlow T. A., 2012, *MNRAS*, 422, 3249
- Capellupo D. M., Hamann F., Barlow T. A., 2014, *MNRAS*, 444, 1893
- Carlberg R. G., 1980, *ApJ*, 241, 1131
- Cassidy I., Raine D. J., 1996, *A&A*, 310, 49
- Cohen M. H., Ogle P. M., Tran H. D., Vermeulen R. C., Miller J. S., Goodrich R. W., Martel A. R., 1995, *ApJ*, 448, L77
- Cunto V., Mendoza C., Ochsenbein F., Zeppen C. J., 1993, *A&A*, 275, L5
- de Kool M., Begelman M. C., 1995, *ApJ*, 455, 448
- Dere K. P., Landi E., Mason H. E., Monsignori Fossi B. C., Young P. R., 1997, *A&AS*, 125, 149
- DiPompeo M. A., Brotherton M. S., Cales S. L., Runnoe J. C., 2012, *MNRAS*, 427, 1135
- Elvis M., 2000, *ApJ*, 545, 63
- Emmering R. T., Blandford R. D., Shlosman I., 1992, *ApJ*, 385, 460
- Eracleous M., Halpern J. P., 1994, *ApJS*, 90, 1
- Fabian A. C., 2012, *ARA&A*, 50, 455
- Filiz Ak N. et al., 2014, *ApJ*, 791, 88
- Fullerton A. W., 2011, in Neiner C., Wade G., Meynet G., Peters G., eds, *Proc. IAU Symp. 272, Active OB Stars: Structure, Evolution, Mass Loss, and Critical Limits*. Cambridge Univ. Press, Cambridge, p. 136
- Gallagher S. C., Brandt W. N., Chartas G., Priddey R., Garmire G. P., Sambruna R. M., 2006, *ApJ*, 644, 709
- Ganguly R., Brotherton M. S., 2008, *ApJ*, 672, 102
- Ganguly R., Sembach K. R., Tripp T. M., Savage B. D., Wakker B. P., 2006, *ApJ*, 645, 868
- Ghosh K. K., Punsly B., 2007, *ApJ*, 661, L139
- Goodrich R. W., 1997, *ApJ*, 474, 606
- Goodrich R. W., Miller J. S., 1995, *ApJ*, 448, L73
- Green P. J., Mathur S., 1996, *ApJ*, 462, 637
- Green P. J., Aldcroft T. L., Mathur S., Wilkes B. J., Elvis M., 2001, *ApJ*, 558, 109
- Grupe D., Mathur S., Elvis M., 2003, *AJ*, 126, 1159
- Hamann F., Chartas G., McGraw S., Rodríguez Hidalgo P., Shields J., Capellupo D., Charlton J., Eracleous M., 2013, *MNRAS*, 435, 133
- Hamann W.-R., Koesterke L., 1998, *A&A*, 335, 1003
- Hamann W.-R., Oskinova L. M., Feldmeier A., 2008, in Hamann W.-R., Feldmeier A., Oskinova L. M., eds, in *Proc. Int. Workshop, Clumping in Hot-Star Winds*. p. 75
- Häring N., Rix H.-W., 2004, *ApJ*, 604, L89
- Hazard C., Mackey M. B., Shimmings A. J., 1963, *Nature*, 197, 1037
- Higginbottom N., Knigge C., Long K. S., Sim S. A., Matthews J. H., 2013, *MNRAS*, 436, 1390 (H13)
- Higginbottom N., Proga D., Knigge C., Long K. S., Matthews J. H., Sim S. A., 2014, *ApJ*, 789, 19 (H14)
- Hillier D. J., 1991, *A&A*, 247, 455
- Hillier D. J., Miller D. L., 1999, *ApJ*, 519, 354
- Hunter J. D., 2007, *Comput. Sci. Eng.*, 9, 90
- Junkkarinen V. T., Burbidge E. M., Smith H. E., 1983, *ApJ*, 265, 51
- Kaspi S., Brandt W. N., Maoz D., Netzer H., Schneider D. P., Shemmer O., 2007, *ApJ*, 659, 997
- Kellermann K. I., Sramek R., Schmidt M., Shaffer D. B., Green R., 1989, *AJ*, 98, 1195
- King A., 2003, *ApJ*, 596, L27
- King A., 2005, *ApJ*, 635, L121
- Knigge C., Drew J. E., 1997, 486, 445
- Knigge C., Scaringi S., Goad M. R., Cottis C. E., 2008, *MNRAS*, 386, 1426
- Krolik J. H., Voit G. M., 1998, *ApJ*, 497, L5
- Krolik J. H., McKee C. F., Tarter C. B., 1981, *ApJ*, 249, 422
- Kurosawa R., Proga D., 2009, *ApJ*, 693, 1929
- Lamy H., Hutsemékers D., 2000, *A&A*, 356, L9
- Landi E., Del Zanna G., Young P. R., Dere K. P., Mason H. E., 2012, *ApJ*, 744, 99
- Laor A., Brandt W. N., 2002, *ApJ*, 569, 641
- Long K. S., Knigge C., 2002, *ApJ*, 579, 725 (LK02)
- Lucy L. B., 2002, *A&A*, 384, 725
- Lucy L. B., 2003, *A&A*, 403, 261
- Lucy L. B., Solomon P. M., 1970, *ApJ*, 159, 879
- MacGregor K. B., Hartmann L., Raymond J. C., 1979, *ApJ*, 231, 514
- Marscher A. P., 2006, in Hughes P. A., Bregman J. N., eds, *AIP Conf. Ser. Vol. 856, Relativistic Jets: The Common Physics of AGN, Microquasars, and Gamma-Ray Bursts*. Am. Inst. Phys., New York, p. 1
- Martínez-Sansigre A. et al., 2007, *MNRAS*, 379, L6
- Mathur S. et al., 2000, *ApJ*, 533, L79
- Matthews J. H., Knigge C., Long K. S., Sim S. A., Higginbottom N., 2015, *MNRAS*, 450, 3331 (M15)
- Murray N., Chiang J., Grossman S. A., Voit G. M., 1995, *ApJ*, 451, 498
- Netzer H., 1990, in Blandford R. D., Netzer H., Woltjer L., Courvoisier T. J.-L., Mayor M., eds, *Saas-Fee Advanced Course of the Swiss Society for Astrophysics and Astronomy: Active Galactic Nuclei*. p. 57
- Noebauer U. M., Long K. S., Sim S. A., Knigge C., 2010, *ApJ*, 719, 1932
- O'Dowd M. J., Bate N. F., Webster R. L., Labrie K., Rogers J., 2015, *ApJ*, 813, 62
- Owocki S. P., Rybicki G. B., 1984, *ApJ*, 284, 337
- Pelletier G., Pudritz R. E., 1992, *ApJ*, 394, 117
- Perley R. A., Dreher J. W., Cowan J. J., 1984, *ApJ*, 285, L35
- Potash R. I., Wardle J. F. C., 1980, *ApJ*, 239, 42
- Pounds K. A., Reeves J. N., 2009, *MNRAS*, 397, 249
- Proga D., Kallman T. R., 2004, *ApJ*, 616, 688
- Proga D., Kurosawa R., 2010, in Maraschi L., Ghisellini G., Della Ceca R., Tavecchio F., eds, *ASP Conf. Ser. Vol. 427, Accretion and Ejection in AGN: A Global View*. Astron. Soc. Pac., San Francisco, p. 41
- Proga D., Stone J. M., Kallman T. R., 2000, *ApJ*, 543, 686
- Proga D., Jiang Y.-F., Davis S. W., Stone J. M., Smith D., 2014, *ApJ*, 780, 51
- Reeves J. N., O'Brien P. T., Ward M. J., 2003, *ApJ*, 593, L65
- Reichard T. A. et al., 2003, *AJ*, 126, 2594
- Risaliti G., Elvis M., Nicastro F., 2002, *ApJ*, 571, 234
- Risaliti G., Salvati M., Marconi A., 2011, *MNRAS*, 411, 2223
- Rybicki G. B., Hummer D. G., 1978, *ApJ*, 219, 654
- Saez C., Brandt W. N., Gallagher S. C., Bauer F. E., Garmire G. P., 2012, *ApJ*, 759, 42
- Shakura N. I., Sunyaev R. A., 1973, *A&A*, 24, 337
- Shen Y. et al., 2011, *ApJS*, 194, 45
- Shlosman I., Vitello P., 1993, *ApJ*, 409, 372 (SV93)
- Shlosman I., Vitello P. A., Shaviv G., 1985, *ApJ*, 294, 96
- Silk J., Rees M. J., 1998, *A&A*, 331, L1
- Sim S. A., Drew J. E., Long K. S., 2005, *MNRAS*, 363, 615
- Sim S. A., Long K. S., Miller L., Turner T. J., 2008, *MNRAS*, 388, 611
- Sim S. A., Miller L., Long K. S., Turner T. J., Reeves J. N., 2010, *MNRAS*, 404, 1369
- Sim S. A., Proga D., Kurosawa R., Long K. S., Miller L., Turner T. J., 2012, *MNRAS*, 426, 2859
- Simon L. E., Hamann F., 2010, *MNRAS*, 409, 269
- Sobolev V. V., 1957, *SvA*, 1, 678
- Sobolev V. V., 1960, *Moving Envelopes of Stars*. Harvard Univ. Press Cambridge, MA
- Springel V., Di Matteo T., Hernquist L., 2005, *ApJ*, 620, L79
- Steffen A. T., Strateva I., Brandt W. N., Alexander D. M., Koekemoer A. M., Lehmer B. D., Schneider D. P., Vignali C., 2006, *AJ*, 131, 2826
- Šurlan B., Hamann W.-R., Kubát J., Oskinova L. M., Feldmeier A., 2012, *A&A*, 541, A37
- Sutherland R. S., 1998, *MNRAS*, 300, 321
- Tombesi F., Cappi M., Reeves J. N., Palumbo G. G. C., Yaqoob T., Baito V., Dadina M., 2010, *A&A*, 521, A57
- Trump J. R. et al., 2006, *ApJS*, 165, 1
- Turner T. J., Miller L., 2009, *A&AR*, 17, 47

- Verner D. A., Ferland G. J., Korista K. T., Yakovlev D. G., 1996, *ApJ*, 465, 487
- Weymann R. J., Scott J. S., Schiano A. V. R., Christiansen W. A., 1982, *ApJ*, 262, 497
- Weymann R. J., Turnshek D. A., Christiansen W. A., 1985, in Miller J. S., ed., *Astrophysics of Active Galaxies and Quasi-Stellar Objects*. University Science Books, Mill Valley, CA, p. 333

- Weymann R. J., Morris S. L., Foltz C. B., Hewett P. C., 1991, *ApJ*, 373, 23
- Zhou H., Wang T., Wang H., Wang J., Yuan W., Lu Y., 2006, *ApJ*, 639, 716

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.