

Monogenic conditions and central nervous system anomalies: a prospective study, systematic review and meta-analysis

Blayney, G., Laffan, E., Jacob, P., Baptiste, C., Gabriel, H., Sparks, T., Yaron, Y., Norton, M., Diderich, K., Wang, Y., Chong, K., Chitayat, D., Saini, N., Aggarwal, S., Pauta, M., Borrell, A., Gilmore, K., Chandler, N., Allen, S., ... Mone, F. (2023). Monogenic conditions and central nervous system anomalies: a prospective study, systematic review and meta-analysis. Prenatal Diagnosis. Advance online publication. https://doi.org/10.1002/pd.6466

Published in:

Prenatal Diagnosis

Document Version:

Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:

Link to publication record in Queen's University Belfast Research Portal

Publisher rights Copyright 2023 the authors.

This is an open access Creative Commons Attribution-NonCommercial License (https://creativecommons.org/licenses/by-nc/4.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. - Share your feedback with us: http://go.qub.ac.uk/oa-feedback

ORIGINAL ARTICLE

PRENATAL **WILEY**

Monogenic conditions and central nervous system anomalies: A prospective study, systematic review and meta-analysis

Gillian V. Blayney¹ | Eoghan Laffan² | Preethi A. Jacob³ | Caitlin D. Baptiste⁴ | Heinz Gabriel⁵ | Teresa N. Sparks⁶ | Yuval Yaron⁷ | Mary E. Norton⁶ | Karin Diderich⁸ | Yiming Wang⁹ | Karen Chong^{9,10} | David Chitayat^{9,10} | Neelam Saini¹¹ | Shagun Aggarwal¹¹ | Montse Pauta¹² | Antoni Borrell¹² | Kelly Gilmore¹³ | Natalie J. Chandler¹⁴ | Stephanie Allen¹⁵ | Neeta Vora¹³ | Abdul Noor^{16,17} | Caitriona Monaghan¹ | Mark D. Kilby^{18,19} | Ronald J. Wapner⁴ | Lyn S. Chitty^{14,20} | Fionnuala Mone²¹ |

¹Fetal Medicine Department, Royal Jubilee Maternity Service, Belfast Health and Social Care Trust, Belfast, UK

²Department of Radiology, Children' Health Ireland at Crumlin, Dublin, Ireland

³Northampton General Hospital, Northampton, UK

⁴Columbia University, New York, New York, USA

⁵Praxis für Humangenetik Tübingen, Tübingen, Germany

⁶Department of Obstetrics, Gynaecology & Reproductive Sciences, University of California San Francisco, San Francisco, California, USA

⁷Prenatal Genetic Diagnosis Unit, Genetic Institute, Tel Aviv Sourasky Medical Center, Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

⁸Department of Clinical Genetics, Erasmus Medical Centre, Rotterdam, the Netherlands

⁹Division of Clinical and Metabolic Genetics, Department of Paediatrics, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada

¹⁰The Prenatal Diagnosis and Medical Genetics Program, Department of Obstetrics & Gynecology, Mount Sinai Hospital, University of Toronto, Toronto, Ontario, Canada

¹¹Department of Medical Genetics, Nizam's Institute of Medical Sciences, Hyderabad, India

¹²Insitut d'Investigacions Biomèdiques August Pi I Sunyer (IDIBAPS), BCNatal, Barcelona, Spain

¹³Department of Obstetrics and Gynaecology, Division of Maternal-Fetal Medicine, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

¹⁴North Thames Genomic Laboratory Hub, NHS Foundation Trust, London, UK

¹⁵West Midlands Regional Genetics Laboratory, South and Central Genomic Laboratory Hub, Birmingham, UK

¹⁶Division of Diagnostic Medical Genetics, Department of Pathology and Laboratory Medicine, Mount Sinai Hospital, Toronto, Ontario, Canada

¹⁷Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario, Canada

¹⁸Institute of Metabolism and Systems Research, College of Medical & Dental Sciences, University of Birmingham, Birmingham, UK

¹⁹Fetal Medicine Centre, Birmingham Women's and Children's NHS Foundation Trust, Birmingham, UK

²⁰Genetics and Genomic Medicine, UCL GOS Institute of Child Health, London, UK

²¹Centre for Public Health, Queen's University Belfast, Belfast, UK

LC-18 Central nervous system anomalies and the incremental yield with prenatal exome sequencing: a systematic review and meta-analysis oral presentation 20th June 2023 ISPD Edinburgh.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

 $\ensuremath{\mathbb{C}}$ 2023 The Authors. Prenatal Diagnosis published by John Wiley & Sons Ltd.

Correspondence

Fionnuala Mone, Centre for Public Health, Queen's University, Institute of Clinical Sciences, Royal Victoria Hospital, Belfast, Northern Ireland, UK Email: f.mone@qub.ac.uk

Abstract

Objectives: Determine the incremental diagnostic yield of prenatal exome sequencing (pES) over chromosome microarray (CMA) or G-banding karyotype in fetuses with central nervous system (CNS) abnormalities.

Methods: Data were collected via electronic searches from January 2010 to April 2022 in MEDLINE, Cochrane, Web of Science and EMBASE. The NHS England prenatal exome cohort was also included. Incremental yield was calculated as a pooled value using a random-effects model.

Results: Thirty studies were included (n = 1583 cases). The incremental yield with pES for any CNS anomaly was 32% [95%CI 27%-36%; $l^2 = 72\%$]. Subgroup analysis revealed apparent incremental yields in; (a) isolated CNS anomalies; 27% [95%CI 19%-34%; $l^2 = 74\%$]; (b) single CNS anomaly; 16% [95% CI 10%-23%; $l^2 = 41\%$]; (c) more than one CNS anomaly; 31% [95% CI 21%-40%; $l^2 = 56\%$]; and (d) the anatomical subtype with the most optimal yield was Type 1 malformation of cortical development, related to abnormal cell proliferation or apoptosis, incorporating microcephalies, megalencephalies and dysplasia; 40% (22%-57%; $l^2 = 68\%$). The commonest syndromes in isolated cases were Lissencephaly 3 and X-linked hydrocephalus.

Conclusions: Prenatal exome sequencing provides a high incremental diagnostic yield in fetuses with CNS abnormalities with optimal yields in cases with multiple CNS anomalies, particularly those affecting the midline, posterior fossa and cortex.

key points

What is already known about this topic?

- Prenatal next-generation sequencing increases the incremental diagnostic yield in fetuses with sonographic structural abnormalities and a normal G-banding karyotype and/or chromosome microarray.
- Published diagnostic yields specific to central nervous system abnormalities are variable, highlighting the need for a systematic review.

What does this study add?

- This is the first systematic review and meta-analysis of the literature available to date in this area with sub-classification by a pediatric neuroradiologist
- A subgroup analysis provides the incremental diagnostic yield for specific anatomical CNS anomalies

1 | INTRODUCTION

Congenital structural anomalies affect 2.2% of births, many of which have an underlying genetic etiology.¹ Anomalies affecting the fetal central nervous system (CNS) contribute substantially to this figure, occurring in 0.26%–0.31% of all births and 3%–6% of stillbirths.^{2,3} CNS anomalies pose a specific challenge related to the need for deep phenotyping using additional imaging modalities such as fetal magnetic resonance imaging (MRI), assessment of the phenotypic evolution as pregnancy progresses and ambiguity related to counseling regarding long-term outcomes.⁴ Obtaining a unifying genetic

diagnosis can prove invaluable to women and practitioners in guiding pregnancy management, treatment, delivery plans and postnatal management, enabling prognostication and providing information on the index and subsequent pregnancies.⁵

Conventional genetic testing, namely G-banding karyotype and chromosome microarray analysis (CMA), are limited to identifying aneuploidy, structural aberrations and copy number variation (CNV), yielding a unifying diagnosis in one-fifth of fetal CNS anomalies.^{6–8} Identification of pathogenic single gene variants using next-generation sequencing (NGS) technologies, namely prenatal exome sequencing (pES) has been demonstrated to increase this yield.^{9,10}

The NHS England pES pathway recommends the provision of trio pES in instances of major CNS anomaly, excluding neural tube defects; however, what should be included within this major category has not yet been specified.¹¹ Hence, the objectives of this prospective study, systematic review and meta-analysis are to determine the incremental yield of pES over and above karyotype and/or CMA for; (i) isolated or multisystem, (ii) single and multiple CNS anomalies , and (iii) CNS anomalies based on anatomical subtype as classified by a pediatric neuroradiologist.

2 | METHODS

2.1 | Protocol and registration

We developed a review protocol aligned to the recommended methods for systematic review and PRISMA guidance, which was prospectively registered with the PROSPERO international systematic review database (PROSPERO No. CRD42022328851 registration 04.05.22).¹²⁻¹⁴

2.2 | Eligibility criteria

Inclusion criteria detailed any retrospective or prospective studies which: (i) included 10 or more pregnancies with a prenatal diagnosis of any CNS anomaly based upon ultrasound with or without fetal MRI; (ii) which underwent NGS (whole, clinical, targeted ES or whole genome sequencing (WGS)); (iii) in which CMA and/or karyotype was non-diagnostic; (iv) in which NGS was initiated based upon prenatal phenotype only; and (v) results of genetic testing were known. Where adequate information was not obtainable from the manuscript the corresponding author was contacted and data were requested. In addition to the search, prospectively collected data from the NHS England pES pathway, commencing October 2021 to February 2023, where pES was performed in instances of a 'major' CNS phenotype (excluding neural tube defects) were included. The methodology of this pathway has been described previously and data used here was collected as part of a registered audit.11

If NGS was initiated postnatally, studies were included only if testing was based solely on prenatal phenotype. Conference abstracts, case series and case reports were also subjected to eligibility criteria assessment. In studies that were not specific to CNS anomalies but detailed such cases, the data was extracted either from the paper or following author correspondence and data sharing. Where available, extended datasets were included. Variants were deemed diagnostic if they were causative of the phenotype and classified as class IV (likely pathogenic) or V (pathogenic) according to the American College of Medical Genetics and Genomics (AMCG) or the Association for Clinical Genomic Science (ACGS).¹⁵ Class III variants of uncertain significance (VUS) and incidental findings (IFs) were also recorded.

PRENATAL DIAGNOSIS-WILEY___

2.3 | Information sources and search strategy

Databases including MEDLINE, Web of Science, Cochrane Library and EMBASE were electronically searched for relevant citations from 1st January 2010 (inception of pES) to first May 2022. The search strategy consisted of relevant Medical Subject Headings (MeSH) terms, keywords and word variants for 'prenatal', 'exome sequencing', and 'abnormality' were used with alternative terms encompassing 'fetus'. fetal', 'prenatal diagnosis', 'antenatal', 'whole exome sequencing', 'exome', 'whole genome sequencing', 'genome human', 'sequence analysis. DNA', 'anomaly' and 'defect'. All study abstracts, were reviewed and full manuscripts were subsequently retrieved for further analysis if they met inclusion criteria. Manuscripts were excluded if they were duplicates; did not meet the inclusion criteria; or if there was inadequate phenotypic information of positive and/or negative cases. In the latter scenario, the corresponding authors were contacted and the study included if further information was available.

2.4 | Data extraction and assessment of risk of bias

Data on study characteristics and outcomes were independently extracted from each study by two reviewers (G.V.B, and P.A.J.) and any conflicts were resolved by a senior reviewer (F.M.). Categories included prenatal phenotype based on ultrasound and when available, fetal MRI; gestation at testing; the source of DNA; sequencing approach; variants reported including gene, clinical syndrome; inheritance pattern; turnaround time; pregnancy outcome; and type of initial non-diagnostic genetic testing (karyotype or CMA). The categorisation of neurological abnormalities and/or disruptions was reviewed and verified by a pediatric neuroradiologist (E.L.). Study characteristics and outcome data were logged under a generated case number and categorized as isolated and non-isolated and under categories including (1) developmental for example, neural tube defects, (2) posterior fossa anomalies for example, Dandy-Walker variants and Chiari II malformations, (3) ventricular, (4) midline for example, holoprosencephaly, agenesis of corpus callosum, (5) malformations of cortical development; A. Abnormal cell proliferation or apoptosis for example, microcephaly, megalencephaly, dysplasia; B. Abnormal cell migration for example, heterotopia, lissencephaly (heterotopia/ cobblestone), schizencephaly, C. Abnormal post-migrational development for example, polymicrogyria or D. Miscellaneousporencephaly, tumors, intracranial haemorrhage.^{17,18}

Quality assessment of the included studies was performed using modified Standards for Reporting of Diagnostic Accuracy (STARD) criteria.¹⁶

3

MILEY-DIAGNOSI

2.5 | Data synthesis

The primary outcome of interest was the incremental yield of pES over CMA/karyotype expressed as a risk difference. This was estimated by pooling risk differences from each included study using a random effects model, using a previously published method with adjustment for 'zero' values from negative karyotype/CMA testing.¹⁹⁻²¹ Results were displayed as forest plots with corresponding 95% confidence intervals (CIs) and pooled for all studies in a meta-analysis using a random effects model. Both the overall yield and yield for isolated CNS anomalies were calculated. A subgroup analysis for the key neurological categories with greater than or equal to n = 5 cases previously listed were used to investigate the effect on incremental yield of pre-test case selection for higher likelihood of monogenic disease. Between-study heterogeneity was assessed graphically within the forest plot and statistically using 'Higgins' I². Publication bias was assessed graphically using funnel plots. Statistical analysis was performed using RevMan version 5.3.4 (Review Manager®, The Cochrane Collaboration, Copenhagen, Denmark) statistical software.

3 | RESULTS

3.1 | Study selection and characteristics

The study selection process is demonstrated in the PRISMA flow diagram (Figure 1). Thirty studies fulfilled the eligibility criteria and were suitable for meta-analysis (1583 cases).^{4,7,9,10,22-47} This included data from the NHS England prenatal exome sequencing pathway.³⁹ For studies that met the inclusion criteria but provided inadequate phenotypic information, corresponding authors were contacted to request further data (n = 57) of which 15 (26.3%) responded. Eleven studies provided extended data sets.^{4,9,10,27,29,37,40,43,46-48} Supplementary Table 1 highlights the characteristics of the included studies and Figure 2 shows the overall quality assessment.

3.2 | Synthesis of results

Twenty-three studies were included in the sub-analysis (1264 cases). All cases underwent G-banding karyotype or CMA prior to pES with



FIGURE 1 PRISMA flow diagram (*Authors contacted for further information **Includes unpublished audit of NHS England cases).



FIGURE 2 Quality assessment of 29 studies included in systematic review, using modified Standards for Reporting of Diagnostic Accuracy criteria (ACMG, American College of Medical Genetics and Genomics; ES, exome sequencing).

52% having both. The median maternal age at testing was 31 years (range 19-49). Ethnicity was known in 21% (n = 265) of cases, of which 64% (n = 170) were Caucasian. The source of fetal DNA was specified in 41% (n = 516) of cases, of which the majority, 53% (n = 274), was from amniocentesis. In the cases that documented turnaround time for prenatal sequencing (n = 755) the median time was 34 days (range 6-1001 days). Pregnancy outcome was known in 63% (n = 792) of cases, of which 65% (n = 513) ended in the termination of pregnancy. Multi-system abnormalities (CNS plus at least one other system) occurred in 50% (n = 627/1264) of cases. Most frequent extra-CNS anomalies included those affecting the extremities (30%), heart (29%) or face (28%). A single CNS abnormality occurred in 57% (n = 717) of cases, with the remainder (n = 547) classified as multiple/complex CNS abnormalities. Only four (27.5%) studies and the NHS England pES series clearly documented whether fetal MRI was used as an additional imaging modality, hence sub-analysis of this group was not possible.

Systematic review of pathogenic variants 3.3

In total 454 cases were identified as having a causative pathogenic or likely pathogenic (P/LP) variant with pES with an incremental yield of 32% (95% Cl 27–36; $I^2 = 72\%$) for any CNS anomaly; 35% (95% Cl 27– 44; $I^2 = 77\%$) for cases of CNS anomaly as part of a multisystem anomaly; 27% (95% Cl 19-34; $l^2 = 74\%$) for isolated CNS anomaly

(Table 1 and Supplementary Figures 1-5). Incremental yields for a single isolated CNS anomaly of 16% (95% Cl 10-23; $l^2 = 41\%$) and a multiple isolated CNS anomaly of 31% (95% Cl 21-40; $l^2 = 56\%$) are also recorded. Incremental yields from the sub-analysis for specific phenotypes are demonstrated in Table 2 with most optimal yields for anomalies of the posterior fossa; 36% [95% Cl 28-43, $I^2 = 50\%$], the midline; 35% [95% Cl 27-44, $l^2 = 77\%$] and the cortex; 35% [95% Cl 26-44, $I^2 = 32\%$], with the greatest yield in those with Type 1 malformations of cortical development, related to abnormal cell proliferation or apoptosis incorporating microcephaly, megalencephaly and dysplasia; 40% (22%–57%; $I^2 = 68\%$) (Supplementary Figures 6-10).

PRENATAL

A list of clinical syndromes caused by Class IV or V causative variants included within the final meta-analysis is outlined in Tables 1 and 2. Where documented, the most common genetic syndromes in isolated CNS anomalies were Lissencephaly 3 (TUBA1A), Coffin-Siris syndrome (ARID1A/B) and congenital X-linked hydrocephalus (L1CAM). In cases in which the inheritance pattern was clearly documented (n = 159); (i) 102 (64.2%) were autosomal dominant, (ii) 34 (21.4%) were autosomal recessive, and (iii) 23 (14.4%) were Xlinked. Women with causative class IV and V variants identified on pES were more likely to terminate their pregnancy (70%; n = 181/259 of known outcomes) than those in which a causative variant was not identified (61.6%; n = 332/539 of known outcomes) p = 0.02. The pooled incremental yield for VUS was 4% (95% CI, 2-6; $l^2 = 57\%$) with the number of incidental findings reported too small to derive a pooled value.

TABLE 1 The incremental yield of prenatal exome sequencing over chromosome microarray and/or G-banding karyotype in prenatally identified CNS anomalies.

CNS anomaly	Incremental yield (%) [95% CI]	Most common affected genes	Corresponding syndromes	VUS (%) [95% CI]
All CNS anomalies	32% [27%-36%] I ² = 72%	TUBA1A 4% (n = 15/375) ^a	Lissencephaly 3	4% [2%-6%] I ² = 57%
		ARID1A/B 3.2% (n = 12/375)	Coffin-Siris	
		TUBB 2.7% (n = 10/375)	Cortical dysplasia, complex, with other brain malformations	
		CEP290 2.7% (n = 10/375)	Joubert or Meckel	
Cases of CNS anomaly with multisystem anomaly	35% [27%-44%] I ² = 77%	CC2D2A or CEP290 or TCTN2 or TMEM 10% (n = 22/222)	Joubert or Meckel	5% [2%-8%] $l^2 = 43\%$
		BRAF or PTPN11 5% (n = 11/222)	Noonan	
		ARID1A/B 3.2% (n = 7/222)	Coffin-Siris	
Cases of isolated CNS anomaly	27% [19%-34%] I ² = 74%	TUBA1A 10.1% (N = 14/139)	Lissencephaly 3	3% [1%-4%] l ² = 0%
		ARID1A/B 3.6% (N = 5/139)	Coffin Siris	
		L1CAM 4.3% (N = 6/139)	Hydrocephalus, congenital, X-linked	

Abbreviations: CNS, central nervous system; VUS, variant of uncertain significance. ^aDenominator is where a pathogenic variant genotype was recorded.

TABLE 2 The incremental yield of prenatal exome sequencing over chromosome microarray and/or G-banding karyotype in prenatally identified CNS anomalies according to sub-analysis for specific phenotypes.

CNS anomaly	Incremental yield (%) [95% Cl]	Most common affected genes	Corresponding syndromes
Developmental	19% [7%-31%] <i>I</i> ² = 38%	CC2D2A 21.1% (n = 4/19) ^a CEP290 21.1% (n = 4/19)	Joubert or Meckel
Ventricular	32% [25%-40%] <i>I</i> ² = 68%	TUBA1A 4.4% (n = 7/158)	Lissencephaly 3
		ARID1A/B 3.8% (n = 6/158)	Coffin-Siris
		POMT1/2 3.8% (n = 6/158)	Muscular dystrophy-dystroglycanopathy (congenital with brain and eye anomalies)
Mild VM	20% [4%-36%] $l^2 = 60\%$	N/S	
Moderate VM	22% [5%-39%] I ² = 49%	N/S	
Severe VM	20% [13%-27%] $l^2 = 0\%$	L1CAM 20% (n = 2/10)	Hydrocephalus, congenital, X-linked
Posterior fossa	36% [28%-43%] I ² = 50%	CEP290 or TMEM67/138 11.9% (n = 16/135)	Joubert or Meckel
		TUBA1A 8.1% (n = 11/135)	Lissencephaly 3
		CHD7 4.4% (n = 6/135)	CHARGE
Midline	35% [27%-44%] I ² = 77%	TUBA1A 6.8% (n = 11/161)	Lissencephaly 3
		TUBB 5.6% (n = 9/161)	Cortical dysplasia, complex, with other brain malformations
		ARID1A/B 4.3% (n = 7/161)	Coffin-Siris
Absent corpus callosum	36% [28%-44%] I ² = 46%	TUBA1A 7.5% (n = 8/107)	Lissencephaly 3
		ARID1A/B 6.5% (n = 7/107)	Coffin-Siris
		TUBB 4.7% (n = 5/107)	Cortical dysplasia, complex, with other brain malformations

TABLE 2 (Continued)



CNS anomaly	Incremental yield (%) [95% Cl]	Most common affected genes	Corresponding syndromes
Holoprosencephaly	34% [15%-53%] <i>I</i> ² = 28%	SHH 38.5% (n = 5/13)	Holoprosencephaly 3
Malformation of cortical development	35% [26%-44%] I ² = 55%	TUBA1A 9.8% (n = 9/92)	Lissencephaly 3
		TUBB 6.5% (n = 6/92)	Cortical dysplasia, complex, with other brain malformations
Type 1 cortical	40% [22%-57%] I ² = 68%	COL4A1 7.9% (n = 3/38)	Microangiopathy and leukoencephalopathy, pontine autosomal dominant
(Abnormal cell proliferation and apoptosis)			
Type 2 cortical	31% [15%-49%] <i>I</i> ² = 63%	TUBB 15.4% (N = 4/26)	Cortical dysplasia, complex, with other brain malformations
(Abnormal cell migration)		TUBA1A 11.5% (N = 3/26)	Lissencephaly 3
Type 3 cortical	32% [21%-44%] I ² = 4%	TUBB 13% (N = 3/23)	Cortical dysplasia, complex, with other brain malformations
(Abnormal post-migrational development)		TUBA1A 8.7% (N = 2/23)	Lissencephaly 3

Abbreviations: CNS, central nervous system; N/S, not specified; VM, ventriculomegaly; VUS, variant of uncertain significance. ^aDenominator is where a pathogenic variant genotype was recorded.

4 | DISCUSSION

We report a systemic review and meta-analysis assessing the incremental yield of pES over standard genomic testing strategies for fetuses with isolated CNS anomalies with an overall incremental diagnostic yield of 27%. Unlike previous studies, we have categorised cases as isolated single CNS anomaly, multiple CNS anomalies and CNS with multisystem anomalies, thereby allowing us to review specific CNS anomaly sub-categories. For isolated CNS anomalies, the commonest syndromes and causative genes were Lissencephaly 3 (*TUBA1A*), Coffin-Siris syndrome (*ARID1A/B*), both having an autosomal dominant inheritance pattern, and congenital X-linked hydrocephalus (*L1CAM*).

Previous studies report variable diagnostic rates of pES in fetal CNS abnormalities ranging from 13% to 71%.^{9,25,31,35,44,48,49} A recent systematic review by Mellis *et al.* reported an incremental yield of 17% (95% Cl, 12%–22%).²¹ Variation in yield may be due to differences in sample size or depth of fetal phenotyping. Notably, our research details a large dataset with high diagnostic rates. This may be due to the selected nature of cases, strict review inclusion criteria and the majority of cases adopting a trio exome approach.

The incremental yield increases when there is more than one CNS anomaly. However, we demonstrate an apparent incremental yield even in less severe single CNS anomalies such as isolated mild ventriculomegaly. We report a 20% incremental yield with isolated severe ventriculomegaly, which is lower than a recent systematic review by Mustafa *et al.*, who reported a yield of 35% although our incremental yield with isolated ACC (36%) was similar to that of a further study (30%) by the same group.^{50,51} Of note, these reviews included fewer case numbers and studies with >3 cases compared to

our limitation of including >10 cases in an attempt to minimise selection bias. Furthermore, the definition of CNS anomalies ideally requires fetal MRI or advanced neurosonography as many CNS anomalies are not readily detectable by ultrasound alone. Thus, cases of mild ventriculomegaly and ACC, for example, may have underlying cortical abnormalities only detectable by MRI, thus inflating the "isolated" CNS anomaly category.

This systematic review reports Joubert syndrome, Meckel syndrome, Noonan syndrome, Lissencephaly and Coffin-Siris syndrome as the commonest syndromes identified when a causative pathogenic variant was recorded with any CNS anomaly. The most common genes included CEP290, TUBA1A, L1CAM and ARID1A. ARID1A, L1CAM and CEP290 were all reported by Mustafa et al., in cases of bilateral severe ventriculomegaly, both isolated and with extracranial anomalies or other brain malformations.⁵⁰ TUBA1A and L1CAM were also the genes with the overall highest frequency in cases of ACC.⁵¹ Of note, many of the syndromes identified, which would typically present prenatally with a multisystem phenotype, were reported as an isolated CNS anomaly in 42% (n = 5) of Coffin-Siris Syndrome; 33% (n = 9) of Joubert or Meckel Syndrome and 15% (n = 2) of Noonan's Syndrome. This demonstrates the incomplete prenatal phenotyping, or poor reporting, offered from prenatal imaging and highlights the need for a low threshold to perform pES in cases of apparently isolated CNS anomaly.

It is important to highlight that the natural history of many neurological condition abnormalities is such that changes may not be detected until later in gestation. In this review, 200 cases (16%) were identified after 24 weeks of gestation and 106 (8%) after 30 weeks. This indicates the evolving nature of a CNS phenotype as pregnancy progresses and the need for deep phenotyping. Within the limitations of reporting, advanced neurosonography was not specifically reported and fetal MRI was documented only as having been performed in under a third of cases, which is likely a significant under-representation.^{52,53} Interestingly, Type 1 cortical anomalies demonstrated the greatest yield, highlighting the importance of investigating a primary or secondary microcephaly (defined prenatally as a head circumference (HC) more than 3 standard deviations (SDs) below the mean for gestational age) which may only manifest in the third trimester and the challenges regarding megalencephaly (HC more than 2SDs above the mean for gestational age) which rarely present prenatally but have a strong association with genes within the P13L-AKT-mTOR pathway and often lead to developmental delay, intellectual disability and early onset seizures.⁵⁴ It is important to consider that the cause may also be benign/familial in nature hence it is always useful to measure the parental occipital frontal circumference.⁵⁵ Furthermore, whilst prenatal diagnosis may not guide management in the index pregnancy in cases of serious CNS abnormality, such as Lissencephaly or small cerebellum, it can be extremely useful for guiding future pregnancies.

Key strengths of this study include the global-scale contribution depicted through the included studies which have contributed to this large-scale review of over 1500 cases of fetal CNS anomaly and the subsequent classification of the CNS phenotype by a neuroradiologist. Limitations include the fact that the phenotype was limited to what was provided by the authors and the phenotypic information detailed within the included studies, including the lack of fetal MRI results. Additionally, heterogeneity was high although we attempted to minimise this by applying a random effects model and sub-analysis limited to studies with $n \geq 5$ cases.

5 | CONCLUSION

The findings of this review reveal a high incremental yield for fetal CNS anomalies with pES over and above standard genomic testing strategies, most notably where there are multiple CNS anomalies, particularly those affecting the midline, posterior fossa and cortex. Prenatal exome sequencing in CNS anomalies can assist with prenatal genetic counseling, providing parents with more information on prognosis and inheritance and assisting clinicians in developing targeted management plans. Although one should always strive to obtain a deep phenotype, pES can facilitate in establishing a diagnosis where this is not feasible or where the CNS phenotype appears mild.

ACKNOWLEDGMENTS

We thank the Rapid fetal exomes sequencing, Bioinformatics team and the Translational research teams at the North Thames and South and Central Genomic Laboratory Hubs. University of North Carolina at Chapel Hill High Throughput Sequencing Facility and the University of North Carolina at Chapel Hill Biospecimen Processing Facility. LSC is partially funded by the NIHR Biomedical Research Center at Great Ormond Street Hospital. All research at Great Ormond Street Hospital

10970223, 0, Downloaded from https://obgyn.onlinelibrary.wiley.com/doi/10.1002/pd.6466 by Queen'S University Belfast, Wiley Online Library on [08/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.co anc ns) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

NHS Foundation Trust and UCL Great Ormond Street Institute of Child Health is made possible by the NIHR Great Ormond Street Hospital Biomedical Research Center. The National Institute of Child Health and Human Development R01HD105868. K23HD088742; PI: Vora; National Center for Advancing Translational Sciences R21TR002779; PI: Vora. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the UK Department of Health.

CONFLICT OF INTEREST STATEMENT

LSC has grant funding from the UK NIHR to evaluate the implementation of fetal sequencing in the English National Health Service. NV received supplies in kind from Illumina for an NIH funded grant on genome sequencing. MDK is a Senior Principal Clinical Scientist in the Medical Genomics Research Group, Illumina, Cambridge, UK.

DATA AVAILABILITY STATEMENT

The anonymized dataset is available from the corresponding author on reasonable request.

ORCID

Caitlin D. Baptiste https://orcid.org/0000-0002-6585-8265 Teresa N. Sparks https://orcid.org/0000-0002-8593-2186 Yuval Yaron https://orcid.org/0000-0001-8622-7668 Karin Diderich https://orcid.org/0000-0003-2637-9998 Yiming Wang https://orcid.org/0000-0001-8031-6184 Karen Chong https://orcid.org/0000-0003-2580-7532 Shagun Aggarwal https://orcid.org/0000-0001-8738-9888 Montse Pauta https://orcid.org/0000-0002-5740-4415 Natalie J. Chandler https://orcid.org/0000-0003-1396-0740 Stephanie Allen https://orcid.org/0000-0003-1511-613X Neeta Vora https://orcid.org/0000-0002-2504-9455 Abdul Noor https://orcid.org/0000-0002-4892-5876 Mark D. Kilby https://orcid.org/0000-0001-9987-4223 Lyn S. Chitty https://orcid.org/0000-0002-4857-7138 Fionnuala Mone https://orcid.org/0000-0002-0718-7547

REFERENCES

- NHS Digital 2022. NCARDRS Congenital Anomaly Official Statistics Report; 2020. Available at URL: NCARDRS Congenital Anomaly Official Statistics Report, 2020 – NDRS (digital.nhs.uk). Accessed 08/12/22.
- Onkar D, Onkar P, Mitra K. Evaluation of fetal central nervous system anomalies by ultrasound and its anatomical co-relation. J Clin Diagn Res. 2014;8(6):Ac05-7. https://doi.org/10.7860/jcdr/2014/ 8052.4437
- 3. European Surveillance of Congenital Anomalies (EUROCAT). Cases and prevalence (per 10,000 births) for all full member registries from 2013-2019, prevalence tables. Available at URL: Prevalence charts and tables | EU RD Platform (europa.eu)
- Mone F, Subieh HA, Doyle S, et al. Evolving fetal phenotypes and clinical impact of progressive prenatal exome sequencing pathways: cohort study. Ultrasound Obstet Gynaecol. 2022;59(6):723-730. https://doi.org/10.1002/uog.24842
- 5. Dempsey E, Haworth A, Ive L, et al. A report on the impact of rapid prenatal exome sequencing on the clinical management of 52 ongoing

pregnancies: a retrospective review. BJOG. 2021;128(6):1012-1019. https://doi.org/10.1111/1471-0528.16546

- de Wit MC, Srebniak MI, Govaerts LC, Van Opstal D, Galjaard RJH, Go ATJI. Additional value of prenatal genomic array testing in fetuses with isolated structural ultrasound abnormalities and a normal karyotype: a systematic review of the literature. Ultrasound Obstet Gynaecol. 2014;43(2):139-146. https://doi.org/10.1002/uog.12575
- Fu F, Li R, Li Y, et al. Whole exome sequencing as a diagnostic adjunct to clinical testing in fetuses with structural abnormalities. Ultrasound Obstet Gynaecol. 2018;51(4):493-502. https://doi.org/10. 1002/uog.18915
- Song T, Xu Y, Li Y, et al. Detection of submicroscopic chromosomal aberrations by chromosome microarray analysis for the prenatal diagnosis of central nervous system abnormalities. J Clin Lab Anal. 2020;34(10):e23434. https://doi.org/10.1002/jcla.23434
- Yaron Y, Ofen Glassnet V, Mory A, et al. Exome sequencing as firsttier test for fetuses with severe central nervous system structural anomalies. *Ultrasound Obstet Gynaecol*. 2022;60(1):59-67. https://doi. org/10.1002/uog.24885
- Baptiste C, Mellis R, Aggarwal V, et al. Fetal central nervous system anomalies: when should we offer exome sequencing? *Prenat Diagn*. 2022;42(6):736-743. https://doi.org/10.1002/pd.6145
- R21 criteria NHS England. Rapid Exome Sequencing Service for Fetal Anomalies Testing; 2021:376. Accessed from 377. https://labs. gosh.nhs.uk/media/1396340/rapid_prenatal_exome_sequencing_ r21_f378aq_v1
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLOS Med.* 2009;6(7):e10000100. https://doi.org/10.1371/journal. pmed.1000100
- Stroup DF, Berline JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Metaanalysis of Observational Studies in Epidemiology (MOOSE) group. JAMA. 2000;283(15):2008-2012. https://doi.org/10.1001/jama.283. 15.2008
- 14. PROSPERO Systematic Review Registry. www.crd.york.ac.uk/ PROSPERO/
- Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants@ a joint consensus recommendation of the American College of medical genetics and genomics and the association for molecular pathology. *Genet Med.* 2015;17(5): 405-424. https://doi.org/10.1038/gim.2015.30
- Bossuyt PM, Reitsma JB, Burns DE, et al. Standards for Reporting of Diagnostic Accuracy. Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD initiative. Standards for Reporting of Diagnostic Accuracy. *Clin Chem.* 2003;49(1):1-6. https:// doi.org/10.1373/49.1.1
- Cater SW, Boyd BK, Ghate SV. Abnormalities of the fetal central nervous system: prenatal US diagnosis with postnatal correlation. *Radiographics*. 2020;40(5):1458-1472. https://doi.org/10.1148/rg. 2020200034
- Desikan RS, Barkovich AJ. Malformations of cortical development. Ann Neurol. 2016;80(6):797-810. https://doi.org/10.1002/ana.24793
- 19. Mone F, Eberhardt RY, Morris RK, et al. Congenital heart disease and the Diagnostic yield with Exome sequencing (CODE) study: prospective cohort study and systematic review. *Ultrasound Obstet Gynaecol.* 2020;57(1):43-51. https://doi.org/10.1002/uog.22072
- Mone F, Eberhardt RY, Hurles ME, et al. Fetal hydrops and the incremental yield of Next-generation sequencing over standard prenatal Diagnostic testing (FIND) study: prospective cohort study and meta-analysis. Ultrasound Obstet Gynaecol. 2021;58(4):509-518. https://doi.org/10.1002/uog.23652
- 21. Mellis R, Oprych K, Scotchman E, Hill M, Chitty LS. Diagnostic yield of exome sequencing for prenatal diagnosis of fetal structural

anomalies: a systematic review and meta-analysis. *Prenat Diagn*. 2022;42(6):662-685. https://doi.org/10.1002/pd.6115

- Boissel S, Fallet-Bianco C, Chitayat D, et al. Genomic study of severe fetal anomalies and discovery of GREB1L mutations in renal agenesis. *Genet Med.* 2018;20(7):745-753. https://doi.org/10.1038/gim. 2017.173
- Chen M, Chen J, Wang C, et al. Clinical application of medical exome sequencing for prenatal diagnosis of fetal structural anomalies. *EJ Obstet Gynaecol Reprod Bio*. 2020;251:119-124. https://doi.org/10. 1016/j.ejogrb.2020.04.033
- Daum H, Meiner V, Elpeleg O, Harel T. Fetal exome sequencing yield and limitations in a tertiary referral centre. Ultrasound Obstet Gynaecol. 2019;53(1):80-86. https://doi.org/10.1002/uog.19168
- De Koning M, Hoffer M, Nibbeling E, et al. Prenatal exome sequencing: a useful tool for the fetal neurologist. *Clin Genet*. 2022; 101(1):65-77. https://doi.org/10.1111/cge.14070
- Deden C, Neveling K, Zafeiropopoulou D, et al. Rapid whole exome sequencing in pregnancies to identify the underlying genetic cause in fetuses with congenital anomalies detected by ultrasound imaging. *Prenat Diagn.* 2020;40(8):972-983. https://doi.org/10.1002/pd.5717
- Diderich K, Romijn K, Joosten M, et al. The potential diagnostic yield of whole exome sequencing in pregnancies complicated by fetal ultrasound anomalies. Acta Obstet Gynecol Scand. 2021;100(6): 1106-1115. https://doi.org/10.1111/aogs.14053
- Dufke A, Hoopman M, Waldmüller S, et al. A single center experience of prenatal parent-fetus trio exome sequencing for pregnancies with congenital anomalies. *Prenat Diagn.* 2022;42(7):901-910. https://doi.org/10.1002/pd.6170
- Gabriel H, Korinth D, Ritthaler M, et al. Trio exome sequencing is highly relevant in prenatal diagnostics. *Prenat Diagn*. 2022;42(7): 845-851. https://doi.org/10.1002/pd.6081
- Greenbaum L, Pode-Shakked B, Eisenberg-Barzilai S, et al. Evaluation of diagnostic yield in fetal whole-exome sequencing: a report on 45 consecutive families. *Front Genet*. 2019;10:425. https://doi.org/ 10.3389/fgene.2019.00425
- Heide S, Spentchian M, Valence S, et al. Prenatal exome sequencing in 65 fetuses with abnormality of the corpus callosum: contribution to further diagnostic delineation. *Genet Med.* 2020;22(11):1887-1891. https://doi.org/10.1038/s41436-020-0872-8
- 32. Laio Y, Yang Y, Wen H, et al. Abnormal Sylvian fissure at 20-30 weeks as indicator of malformations of cortical development: role of prenatal whole-genome sequencing. *Ultrasound Obstet gynaecol.* 2022;59:550-562.
- Lei L, Zhou L, Xiong J.-J, et al. Whole-exome sequencing increases the diagnostic rate for prenatal fetal structural anomalies. J Med Genet. 2021;64(9):104288. https://doi.org/10.1016/j.ejmg.2021. 104288
- Lei T.-Y, Qin S, Fu F, et al. Prenatal exome sequencing in fetuses with callosal anomalies. *Prenat Diagn*. 2022;42(6):744-752. https://doi. org/10.1002/pd.6107
- Li L, Fu F, Li R, et al. Genetic tests aid in counselling of fetuses with cerebellar vermis defects. *Prenat Diagn*. 2020;40(10):1228-1238. https://doi.org/10.1002/pd.5732
- Normand E, Braxton A, Nassef S, et al. Clinical exome sequencing for fetuses with ultrasound abnormalities and a suspected Mendelian disorder. *Genome Med.* 2018;10(1):74. https://doi.org/10.1186/ s13073-018-0582-x
- Pauta M, Campos B, Segura-Puimedon M, et al. Next-generation sequencing gene panels and "solo" clinical exome sequencing applied in structurally abnormal fetuses. *Fetal Diagn Ther.* 2021; 48(10):746-756. https://doi.org/10.1159/000519701
- Qi Q, Jiang Y, Zhou X, et al. Simultaneous detection of CNVs and SNVs improves the diagnostic yield of fetuses with ultrasound anomalies and normal karyotypes. *Genes.* 2020;11(12):1397. https:// doi.org/10.3390/genes11121397

PRENATAL DIAGNOSIS-WILEY____

9

10

- R21 rapid prenatal exome sequencing. Accessed from: https://www. genomicseducation.hee.nhs.uk/genotes/knowledge-hub/r21-rapidprenatal-exome-sequencing/ 2023
- Saini N, Venkatapuram VS, Vineeth VS, et al. Fetal phenotypes of Mendelian disorders: a descriptive study from India. *Prenat Diagn*. 2022;42(7):911-926. https://doi.org/10.1002/pd.6172
- Shamseldin H, Kurdi W, Almusafri F, et al. Molecular autopsy in maternal-fetal medicine. *Genet Med.* 2018;20(4):420-427. https:// doi.org/10.1038/gim.2017.111
- Smogavec M, Bujalkova M, Lehney R, et al. Singleton exome sequencing of 90 fetuses with ultrasound anomalies revealing novel disease-causing variants and genotype-phenotype correlations. *Eur J Hum Genet*. 2022;30(4):428-438. https://doi.org/10.1038/s41431-021-01012-7
- Sparks T, Adami L, Holliman P, et al. Exome sequencing for prenatal diagnosis in nonimmune hydrops fetalis. N J Med. 2020;383(18): 1746-1756. https://doi.org/10.1056/nejmoa2023643
- Tan H, Xie Y, Chen F, et al. Novel and recurrent variants identified in fetuses with central nervous system abnormalities by trios-medical exome sequencing. *Clin Chim Acta*. 2020;510:599-604. https://doi. org/10.1016/j.cca.2020.08.018
- 45. Tolusso L, Hazelton P, Wong B, Swarr D. Beyond diagnostic yield: prenatal exome sequencing results in maternal, neonatal, and familial clinical management changes. *Genet Med.* 2021;23(5):909-917. https://doi.org/10.1038/s41436-020-01067-9
- Vora N, Gilmore K, Brandy A, et al. An approach to integrating exome sequencing for fetal structural anomalies into clinical practice. Genet Med. 2020;22(5):954-961. https://doi.org/10.1038/ s41436-020-0750-4
- Wang Y, Greenfeld E, Watkins N, et al. Diagnostic yield of genome sequencing for prenatal diagnosis of fetal structural anomalies. *Prenat Diagn*. 2022;42(7):1-9. https://doi.org/10.1002/pd.6108
- Reches A, Hiersch L, Simchoni S, et al. Wole-exome sequencing in fetuses with central nervous system abnormalities. *J Perinatol.* 2018; 38(10):1301-1308. https://doi.org/10.1038/s41372-018-0199-3
- 49. Weitensteiner V, Zhang R, Bungenberg J, et al. Exome sequencing in syndromic brain malformations identifies novel mutations in ACTB, and SLC(A6, and suggests BAZ1A as a new candidate gene. *Birth Defects Res.* 2018;110(7):587-597. https://doi.org/10.1002/bdr2.1200

- Mustafa H, Sambatur E, Barbera J, et al. Diagnostic yield with exome sequencing in pre-natal severe bilateral ventriculomegaly: a systematic review and meta-analysis. *Am J Obstet Gynaecol MFM*. 2023; 5(9):101048. https://doi.org/10.1016/j.ajogmf.2023.101048
- Mustafa H, Sambatar E, Mohammad-Hossein H, et al. Prenatal agenesis of corpus callosum and diagnostic yield with exome sequencing, systematic review and meta-analysis. *AJOG*. 2023;S320:483. [conference abstract].
- Paladini D, Malinger G, Birnbaum R, et al. ISUOG Practice Guidelines (updated): sonographic examination of the fetal central nervous system. Part 2: performance of targeted neurosonography. *Ultrasound Obstet Gynecol.* 2021;57(4):661-671. https://doi.org/10.1002/ uog.23616
- Mone F, Homfray T, Kagan KO, Kilby MD. Enhancement of fetal phenotyping in investigation of fetus using next-generation sequencing. Ultrasound Obstet Gynecol. 2023;62:459-461. Epub ahead of print. PMID: 37401773. https://doi.org/10.1002/uog.26301
- Leibovitz Z, Lerman-Sagie T, Haddad L. Fetal brain development: regulating processes and related malformations. *Life (Basel)*. 2022; 12(6):809. https://doi.org/10.3390/life12060809
- 55. Hanzlik E, Gigante J. Microcephaly. Child (Basel). 2017;4(6):47. PMCID: PMC5483622. https://doi.org/10.3390/children4060047

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Blayney GV, Laffan E, Jacob PA, et al. Monogenic conditions and central nervous system anomalies: A prospective study, systematic review and meta-analysis. *Prenat Diagn.* 2023;1-10. https://doi.org/10.1002/pd.6466