



**QUEEN'S
UNIVERSITY
BELFAST**

The effect of a group-based mindfulness and acceptance training on psychological flexibility and adherence to antiretroviral therapy among adolescents in Uganda: an open-label randomized trial

Musanje, K., Kanya, M. R., Kasujja, R., Vanderplasschen, W., Sinclair, D. L., Baluku, M. M., Odokonyero, R. F., Namisi, C. P., Mukisa, J., White, R. G., & Camlin, C. S. (2024). The effect of a group-based mindfulness and acceptance training on psychological flexibility and adherence to antiretroviral therapy among adolescents in Uganda: an open-label randomized trial. *Journal of the International Association of Providers of AIDS Care*, 23. <https://doi.org/10.1177/23259582241236260>

Published in:

Journal of the International Association of Providers of AIDS Care

Document Version:

Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2024 the authors.

This is an open access Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits use, distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

The Effect of a Group-Based Mindfulness and Acceptance Training on Psychological Flexibility and Adherence to Antiretroviral Therapy Among Adolescents in Uganda: An Open-Label Randomized Trial

Journal of the International
Association of Providers of AIDS Care
Volume 23: 1-11
© The Author(s) 2024
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/23259582241236260
journals.sagepub.com/home/jia



Khamisi Musanje, MSc^{1,2} , Moses R. Kamyu, PhD³, Rosco Kasujja, PhD⁴,
Wouter Vanderplasschen, PhD⁵, Deborah L. Sinclair, PhD⁵,
Martin M. Baluku, PhD², Raymond F. Odokonyero, MSc⁶,
Charles P. Namisi, MSc⁷, John Mukisa, MSc⁷, Ross G. White, PhD⁸,
and Carol S. Camlin, PhD⁹

Abstract

Adherence to antiretroviral therapy (ART) is lower in adolescents with HIV (AWH) than in any other age group, partly due to self-regulatory challenges during development. Mindfulness and acceptance training have been shown to support psychological flexibility, a self-regulatory skill that potentially improves adolescent adherence to medication. We assessed the effect of weekly group-based mindfulness and acceptance training sessions on ART adherence among older adolescents (15–19 years) in Kampala, Uganda. One hundred and twenty-two AWH (median age 17, range 15–19 years, 57% female) receiving care at a public health facility in Kampala were randomized 1:1 to receive 4 weekly 90-min group sessions facilitated by experienced trainers or standard-of-care ART services. The training involved (Session 1) clarifying values, (Session 2) skillfully relating to thoughts, (Session 3) allowing and becoming aware of experiences non-judgmentally, and (Session 4) exploring life through trial and error. At baseline, postintervention, and 3-month follow-up, psychological flexibility was measured using the Avoidance and Fusion Questionnaire for Youth (AFQ-Y8), and self-reported ART adherence was assessed using the Morisky Medication Adherence Scale (MMAS-8). At baseline, the intervention and standard-of-care arms had similar psychological flexibility (AFQ-Y8 score: 15.45 ± 0.82 ; 15.74 ± 0.84) and ART adherence (MMAS-8 score: 5.32 ± 0.24 ; 5.13 ± 0.23). Retention through the study was moderate (71%). Completion of mindfulness and acceptance training was associated with a significant reduction in psychological inflexibility at the 3-month follow-up (AFQ-Y8 score: 12.63 ± 1.06 ; 14.05 ± 1.07 , $P = .006$). However, no significant differences were observed in self-reported adherence to ART at the 3-month follow-up (MMAS-8 score: 5.43 ± 0.23 ; 4.90 ± 0.33 , $P = .522$). Group-based mindfulness and acceptance training improved psychological flexibility in this population of adolescents on ART in Uganda but did not significantly improve ART adherence.

¹ Clinical Epidemiology Unit, Makerere University, Kampala, Uganda

² Department of Educational, Social and Organizational Psychology, Makerere University, Kampala, Uganda

³ Department of Medicine, Makerere University, Kampala, Uganda

⁴ Department of Mental Health and Community Psychology, Makerere University, Kampala, Uganda

⁵ Department of Special Needs Education, Ghent University, Belgium

⁶ Department of Psychiatry, Makerere University, Kampala, Uganda

⁷ Department of Epidemiology and Biostatistics, Makerere University, Kampala, Uganda

⁸ School of Psychology, Queens University, Belfast, Northern Ireland

⁹ Department of Obstetrics, Gynecology & Reproductive Sciences, University of California, San Francisco, CA, USA

Corresponding Author:

Khamisi Musanje: Makerere, University School of Psychology, P.O Box 7062, Kampala, Uganda.

Email: khamisi.musanje@mak.ac.ug



Future research should explore integrated approaches that combine behavioral management training with other empowerment aspects to improve ART adherence among AWH.

Keywords

HIV, adherence, mindfulness, mental health, health outcomes, psychological flexibility

Date received: 27 September 2023; revised: 5 February 2024; accepted: 12 February 2024.

Introduction

Antiretroviral therapy (ART) has improved HIV treatment and changed the face of a once deadly infectious disease to a manageable chronic condition.^{1,2} Antiretroviral therapy can effectively suppress the HIV viral load, improve the immune function, and control the risk of opportunistic infections, thus, enabling a stable and healthy life for people living with HIV/AIDS (PLWHA).³ To maximize the clinical benefits of ART, adherence to the treatment plan is important.³ Successful ART programs are highly dependent on optimal adherence and sustained care engagement⁴; further, slowed progression to AIDS, sustained viral load suppression (VLS), reductions to ART resistance, and lengthened survival rates are all achieved through maintenance of a >95% daily oral dosing of ART.⁵

While the benefits of ART in the treatment of HIV are well known, adherence to treatment plans remains a challenge among PLWHA,^{1,6,7} particularly among adolescents with HIV (AWH),^{8–10} who are approximated to be 1.7 million in number globally.³ Adherence to ART is even poorer among older adolescents (15–19 years).¹¹ Older AWH interrupt treatment more often than adults, are more difficult to retain in care compared to young adolescents (10–14 years),¹² and are at a higher risk of HIV-related mortality.³ In Uganda, an estimated 170,000 young people are living with HIV,¹³ of which about 90% have acquired it vertically through their parents and 10% through sexual exposure,¹⁴ yet they are also the only group where HIV-related mortality is on the rise, amidst a decline in other groups such as children and adults with HIV.^{15–17} Of the individual, interpersonal, community, and health systems challenges, the increase in HIV-related mortality is largely attributed to lapsed adherence to ART.^{16,18,19} Young people are less adherent to medical regimens compared to adults⁸ and those with vertically acquired HIV report poorer adherence than those with behaviorally acquired HIV.¹⁴ Studies using VLS data, a key indicator of medication success among PLWHA, show that in 2017, only 39.6% of young people in Uganda achieved VLS, compared to 74.2% of adults.²⁰ Several other studies have also reported lapsed adherence to ART among AWH in Uganda.^{15,17,21,22}

Adherence to ART is affected by several variables ranging from socio-economic factors such as poverty and lack of food, psychosocial and individual factors such as discrimination, stigma, anxiety, and mode of transmission to treatment-related factors such as late initiation, long-term use, pill burden, and

drug fatigue.^{18,23–25} In addition to the structural factors that might be beyond individual control, lapsed adherence to ART among AWH is further attributed to the developmental changes at the onset of puberty which affects health-seeking behaviors.^{2,26}

Adolescence as a life stage influences perceptions, beliefs, attitudes, and evaluations of the severity of disease²⁷ and, is marked with heightened vulnerability to risk, reckless behaviors, and insecurities.^{28,29} Concurrently, adolescence coincides with changes in the social environment, such as spending more time with peers, as well as an increase in autonomy.³⁰ For AWH, adolescence also means taking personal responsibility for one's health.^{29,31} When such unprecedented life changes combine with the stigma that arises out of living with HIV, AWH experience cognitive dissonance characterized by self-hate, blame, entanglement with thoughts (cognitive fusion), and obsessing about the suppression of negative thought experiences (experiential avoidance).³² Since the mind is the doorway through which individuals experience life events,³³ such mental experiences affect psychosocial pathways to care among AWH.^{31,34–36}

Individual attributes such as self-control and emotion regulation help build resilience which is needed when dealing with challenges that accompany living with a chronic condition.^{33,37,38} However, during adolescence, such executive attributes are still underdeveloped, which presents self-management challenges that affect health-seeking behaviors among AWH.³⁹ Thus, interventions that promote self-regulation can benefit AWH.^{19,40} Furthermore, such approaches should also be able to promote resilience and self-acceptance which are functional responses that influence mental health and HIV treatment outcomes.¹⁹

Mindfulness and acceptance-based interventions (MABIs) have the potential to promote self-regulation, resilience, and self-acceptance among young people^{41,42} through a mechanism called psychological flexibility, the ability to live consciously aware of life challenges but deliberately making choices that are personally meaningful.^{43–45} MABIs, through targeting psychological flexibility, address cognitive fusion (a state where individual behaviors are dominated by particular thoughts) and experiential avoidance (obsessing about suppressing negative thoughts),^{41,46} processes that can impact adherence to medication among young people.^{32,47} Furthermore, there is growing evidence that directly connects psychological flexibility to adherence to lifelong treatments.^{48,49} Psychologically flexible people deal with unpleasant experiences associated with daily treatment openly and elaborately, making choices that serve

personal values rather than suppressing such experiences.^{49–52} To improve adherence to ART among AWH, approaches that promote behavioral management skills such as MABIs need to be evaluated and possibly included in the standard-of-care (SOC) offered to AWH.

One particular model based on mindfulness and acceptance philosophy that is specifically designed for young people is the Discoverer-Noticer-Advisor-values (DNA-v) intervention. The DNA-v is derived from Acceptance and Commitment Therapy (ACT), which is a cognitive behavioral approach that helps people relate better with painful and unhelpful thoughts and experiences over which they may have little or no control.^{32,53} The DNA-v uses age-appropriate metaphors, language, and techniques to support the development of 3 functional classes of behavior coined in the acronym DNA: the *Discoverer* involves learning to explore the world through trial and error, the *Noticer* involves developing better awareness and self-awareness, and the *Advisor* involves becoming familiar with and developing a relationship with one's inner voice.⁴¹ All 3 behaviors promote psychological flexibility and work in service of values.⁴¹ The DNA-v can be offered as individual therapy, one-on-one sessions between the provider and the client focused on individual needs, or as group therapy involving 2 or more clients focusing on common concerns.⁴¹ In HIV treatment and care, group programs are encouraged because they promote experience sharing, reduce stigma, and improve self-esteem.⁵⁴

Interventions based on mindfulness and acceptance philosophies have growing evidence supporting the mental health of people living with chronic conditions^{55–57} including adolescents.^{58–60} However, literature linking such psychotherapies to adherence to medication among people with HIV is lacking. Thus this study aims to evaluate the effect of the DNA-v model on improving adherence to ART among AWH through an open-label group treatment trial. We hypothesize that the intervention group will report higher psychological flexibility and improved adherence to ART at the end of the trial.

Methods

Study Design and Setting

We conducted an open-label group treatment trial with individuals randomly assigned to either the DNA-v intervention or to the current SOC procedures and collected data on self-reported ART adherence and psychological flexibility at 3 time points: baseline, 4 weeks postintervention, and at 3-month follow-up. Participants were recruited from Kisenyi Health Center IV (KHC), an urban public health facility in the Kisenyi zone, Rubaga division of the Kampala district. The KHC is administratively managed by the Kampala Capital City Authority and serves a catchment population of about 1 million people within the central business district. The KHC offers free HIV care services and serves approximately 590 AWH.

Participants

The participants were older adolescents receiving care at the KHC. Inclusion criteria were: (a) being 15–19 years old, (b) an HIV-positive diagnosis, (c) attending care at the participating study clinic, and (d) the ability to speak and understand Luganda or English (the most commonly spoken languages in Kampala City). AWH with cognitive impairments or disabilities, those planning to move out of the catchment area during the study period, or those participating in an HIV care study were excluded. Recruitment of participants began on 1 November 2022 and ended on 3 December 2022.

Procedures

Once administrative clearance was offered by the city authority, we were able to access anonymous medical records to screen participants for eligibility. A list of eligible AWH was generated and were contacted telephonically by clinic staff supporting the study. Adolescents who expressed interest in the study were scheduled for individual appointments to inform and obtain consent/assent which was followed by a group briefing and later baseline assessment. Participants were then randomly assigned to the intervention or control group using allocation sequences prepared by an external bio-statistician. Allocation concealment was done using sealed serialized brown opaque envelopes that were kept by the statistician.

Randomization. Participants were randomly allocated to the intervention or control group using computer-generated random numbers. The block randomization method was used for allocating AWH to the intervention or control groups and, to achieve a 1:1 randomization ratio, we used a block size of 6 with equal individuals assigned to the 2 arms per block. Participants allocated to the intervention arm received the DNA-v training and the SOC, while participants who were allocated to the control group continued with the SOC only. To minimize contamination, the research team worked closely with healthcare providers to schedule sessions on non-clinic days to reduce the possibility of unplanned meet-ups among participants, furthermore, the team also encouraged participants not to discuss the training with non-group members until it was completed. This was an open-label randomized study because healthcare providers (counselors at the adolescent clinic) participated in recruitment, administration of the intervention, and the assessment of outcomes. The intervention flow process is shown in Figure 1.

Study Intervention

The DNA-v. The DNA-v intervention was delivered to AWH as life skills training responding to the developmental needs of young people.⁴¹ We split participants in the intervention arm into 4 sub-groups ranging between 11 and 12 participants. After creating sub-groups, we designed a session schedule to guide the implementation of the intervention. The first author, who is trained in using the DNA-v model, and




Figure 1. Flow diagram of the intervention process.

the second author, who is trained in the therapeutic use of mindfulness and acceptance, delivered 4 90-minute sessions to each sub-group across 4 weeks, meeting once a week at the clinic premises. We conducted measures at baseline, postintervention (at 4 weeks), and follow-up (after 3 months). Participants were reimbursed \$10 (37,000 Uganda shillings) every week to compensate for their time and transport costs. The session content is detailed in Table 1.

Standard-of-Care. The SOC involved a combination of clinic-based strategies routinely used to support AWH. The services are offered during adolescent clinic days or as weekend programs by trained healthcare providers or young adolescent peer leaders. Services include psychosocial assessment (using the HEADSS tool for gathering information on home, education, eating, employment, activity, drugs, sex, sexuality, and suicidal ideation), and support (either individual or group-based, led by a healthcare worker to discuss disclosure of status, stigma, discrimination, treatment, loss and

bereavement, and transitioning for AWH turning 18).⁶² Psychosocial support is combined with health education such as information about ART (available care packages, benefits, side effects, and managing lifelong treatment), information on growth and development such as sexual reproductive health (antenatal care, safe deliveries, and postnatal care), life skills education, counseling on substance use, nutrition services, pregnancy testing, and contraceptive methods.⁶² The SOC also involved participation in teen drama clubs which often run during school holidays and intensive adherence counseling administered to AWH with ≥ 1000 copies/mL after ≥ 6 months of ART.

Measures and Outcomes

We measured adherence to ART using the Morisky Medication Adherence Scale (MMAS-8).⁶³ The MMAS-8 is comprised of 8 items designed to facilitate the recognition of

Table 1. Content and Activities of the DNA-v Training with AWH.⁶¹

Training session	Description of content
Session 1	Overview of the DNA-v program and its main goal (life skills training that supports meaning-driven living, forming better relationships, and becoming who you want to be regardless of the situation). Self-introductions and bonding, agreeing to the group behaviors of respect and confidentiality, introducing the DNA-v model as a play. <i>Values:</i> Identifying what brings meaning to you as a person (participants engaged in a values card sort exercise). Living a values-led life (small steps that matter). Navigating the challenges of living or struggling to fit in.
Session 1	<i>The Noticer:</i> Normalizing the Noticer (games), naming emotions (using cards depicting emotions), practicing awareness, naming and describing emotions, and creating new rules for noticing.
Session 3	<i>The Advisor:</i> Normalizing the inner voice (we all have an “advisor” that keeps talking to us; games to become familiar with the Advisor), appreciating the helpfulness of the inner voice, detaching from the inner voice (games to “unhook”), choosing when to listen to the inner voice and when not to depending on the situation. Creating new rules for relating with the inner voice (Advisor).
Session 4	<i>The Discoverer:</i> Trying, tracking, and building strength to serve values. Small steps build big action, reflecting and learning from what has worked and what has not. Putting D-N-A together while moving toward values: acting out the DNA with humor as we end the training.

Abbreviations: DNA-v, Discoverer-Noticer-Advisor-values; AWH, adolescents with HIV.

Table 2. Background Characteristics of Participants at Enrollment Into the Trial.

Variable	Control (n = 61)	Intervention (n = 61)	P value
Age, mean (\pm SD)	16.95 (\pm 1.59)	17.03 (\pm 1.28)	.85
Sex, n (%)			
Female	35 (57.38)	37 (60.66)	.71
Male	26 (42.62)	24 (39.34)	
School status, n(%)			
In school	33 (54.10)	33 (54.10)	1.00
Out of school	28 (45.90)	28 (45.90)	

barriers to and behaviors associated with (intentional and unintentional) adherence to chronic medication. Response options for the MMAS-8 are “yes” or “no” for items 1–7, while item 8 has a 5-point Likert response scale ranging from 0 to 4. In scoring the MMAS-8, each “no” is rated as 1 and each “yes” is rated as 0 except for item 5 which is reverse coded. Item 8 is divided by 4 when calculating a summed score. Total scores range from 0 to 8, representing low adherence ≤ 6 , medium adherence = 6 to < 8 , and high adherence = 8. The MMAS-8 has demonstrated acceptable reliability when used with PLWHA ($\alpha = .66$).⁶⁴

We measured psychological flexibility using the Avoidance and Fusion Questionnaire for Youth (AFQ-Y8).⁶⁵ The AFQ-Y8 is an 8-item instrument where participants rate their level of agreement with statements targeting cognitive fusion and experiential avoidance such as “my thoughts and feelings mess up my life.” The AFQ-Y8 is scored on a 5-point Likert scale underpinned by a single factor. Scores range from 0 to 24, with higher scores indicating higher levels of psychological inflexibility and lower scores indicating psychological flexibility. The AFQ-Y8 has excellent internal consistency ($\alpha = .90$), and test–retest validity ($r = .80$)⁶⁶ and is recommended for measuring psychological flexibility among adolescents.⁶⁶

The study questionnaire was administered in either English or Luganda and the translation involved 2 bilingual native speakers of both languages. One translated the original English tool into Luganda and the other back-translated the Luganda version into English. The back-translated and original English tools were compared and inconsistencies were addressed. Both versions of the tool were then pilot tested with adolescents in another public health facility which was not part of the study.

Statistical Analysis

The sample size was based on a Swedish and Australian study on the effectiveness of ACT for adolescent mental health where a mean difference of 4.09 was considered ($SD = 5.13$).⁶⁷ Confidence interval and power of 80% was used leading to a sample size of 52 participants in each group. To compensate for a 10% loss to follow-up, the sample size was increased to 116 (58 in each group). Baseline characteristics were compared using the Student’s *t*-test for continuous variables while the categorical variables were compared using the Fisher’s exact and chi-square tests. We compared the intervention and control groups at 3 time points: baseline, postintervention, and 3-month follow-up. The means and standard errors of the 2 outcomes of psychological flexibility and adherence were summarized for each time point in both the control and intervention groups. We performed repeated measures analysis of variance (ANOVA) after testing for the homogeneity of variances assumption (sphericity) using Mauchly’s test. We then performed the repeated measures test with a *P* value of .05 considered as statistical significance. We conducted post hoc pairwise comparisons of means for psychological flexibility using the Bonferonni adjustment and adjusted the *P* value. Participants who withdrew before the 3-month follow-up assessment were excluded from the final analysis. Analyses

were conducted using SPSS version 25.0 and followed an intention-to-treat analysis approach.

Ethical Approval and Informed Consent

Ethics approval was granted by the Makerere University School of Medicine Research and Ethics Committee (Mak-SOMREC-2021-176), and the Uganda National Council for Science and Technology (HS1656ES). We also obtained administrative clearance from the Kampala Capital City Authority, the body that manages all public health centers in the city. AWH who were 18 years and above provided written informed consent while those below 18 years of age provided assent, with written consent provided by their caregivers. Caregivers were not treated as participants and did not provide study data. The consent form was in both Luganda and English languages and participants were given sufficient time to read or be read the form in their language of preference and to ask any questions. The consent/assent process was conducted in a private room at the clinic with a healthcare provider present to witness that participation was voluntary.

Study methods and results followed the 2010 CONSORT checklist for reporting parallel group randomized trials.⁶⁸ The study protocol was registered on ClinicalTrials.gov (NCT05010317) and a full study protocol is available from the corresponding author.

Results

Description of the Study Population

We randomly allocated 61 participants to each study arm and conducted evaluation at 3 time points (baseline, postintervention, and 3-month follow-up). The mean age of the participants, sex, and school status (whether in school or not) did not differ significantly across groups at baseline as shown in Table 2. Furthermore, the level of psychological flexibility and self-reported adherence to ART also did not differ significantly between the 2 groups at baseline, while at 3-month

follow-up, significant differences were recorded in psychological flexibility but not in self-reported adherence to ART as shown in Table 3.

Outcome Measures

From the repeated measures analysis, we found that there was a significant difference in mean scores for psychological flexibility across the 3 time points ($F=5.240$, $P=.006$) as shown in Table 4. For adherence, there was no statistically significant difference between mean scores across the 3 time points irrespective of the initial randomization ($F=0.652$, $P=.522$) as shown in Table 4. We further performed a post hoc analysis for psychological flexibility as shown in Table 5.

For psychological flexibility as a mechanism of change underlying MABIs, in the intervention arm, we found that Mauchly's test for sphericity was fulfilled with $P=.761$. We performed post hoc pairwise comparisons of the means using the Bonferroni adjustment, which showed that there was a significant difference between the baseline psychological flexibility mean scores and postintervention scores (15.45, 13.52, $P=0.020$) and also baseline psychological flexibility scores and follow-up scores (15.45, 12.63, $P=.022$) as shown in Table 5. There were no significant differences noted between the comparison of the mean scores between the postintervention scores and 3-month follow-up scores (13.52, 12.63, $P=1.00$) as shown in Table 4. In the control arm, we found that Mauchly's test for sphericity was fulfilled with $P=.368$. Repeated measures ANOVA determined that mean psychological flexibility did not differ significantly across the 3 time points ($F=1.991$, $P=.140$) as shown in Table 4. No post hoc comparisons of the means were performed for this group.

For adherence to ART, overall, there were no significant differences between the intervention and control arms at all 3 time points. In the intervention arm, we found that Mauchly's test for sphericity was fulfilled with a $P=1.00$. Repeated measures ANOVA determined that mean adherence did not differ significantly across the 3 time points ($F=0.652$, $P=.522$) as shown in Table 4. No post hoc pairwise comparisons of the means were performed using the Bonferroni adjustment. Similarly, in the control arm, Mauchly's test for sphericity was fulfilled with a $P=.813$. Repeated measures ANOVA determined that mean adherence did not differ significantly across the 3 time

Table 3. Group Comparisons at 3 Time Points Using Repeated Measures ANOVA.

	Intervention (N = 44) mean (\pm standard error)	Control (N = 43) mean (\pm standard error)
Psychological inflexibility		
Baseline	15.45 (\pm 0.82)	15.74 (\pm 0.84)
Posttest	13.52 (\pm 0.99)	13.52 (\pm 1.10)
Follow-up (3 months)	12.63 (\pm 1.06)	14.05 (\pm 1.07)
Adherence to ART		
Baseline	5.32 (\pm 0.24)	5.13 (\pm 0.23)
Posttest	5.64 (\pm 0.26)	5.14 (\pm 0.32)
Follow-up (3 months)	5.43 (\pm 0.23)	4.90 (\pm 0.33)

Abbreviations: ANOVA, analysis of variance; ART, antiretroviral therapy.

Table 4. Repeated Measures Analysis of Variance (ANOVA) With Randomization as a Between-Subjects Effect.

Characteristic	df	F value	P value
Model 1 for flexibility			
Psychological flexibility	2	5.240	.006
Psychological flexibility \times randomization	2	1.992	.140
Intercept/error (flexibility)	168		
Model 2 for adherence			
Adherence	2	.652	.522
Adherence \times randomization	2	.112	.894
Intercept/error (adherence)	170		

Table 5. Post hoc Pairwise Comparisons of the Psychological Flexibility Means Using the Bonferroni Adjustment.

(I) Flexibility	(J) Flexibility	Mean difference		Sig. ^b	95% confidence interval for difference ^b	
		(I - J)	Std. error		Lower bound	Upper bound
1	2	2.117 ^a	.760	.020	.261	3.974
	3	1.948 ^a	.708	.022	.219	3.677
2	1	-2.117 ^a	.760	.020	-3.974	-.261
	3	-.169	.712	1.000	-1.908	1.569
3	1	-1.948 ^a	.708	.022	-3.677	-.219
	2	.169	.712	1.000	-1.569	1.908

Based on estimated marginal means.

^aThe mean difference is significant at the .05 level.

^bAdjustment for multiple comparisons: Bonferroni.

points ($F = 0.112$, $P = .894$), and no post hoc pairwise comparisons of the means were performed.

Discussion

In this study, we evaluated the effect of the DNA-v on psychological flexibility and adherence to ART among AWH. Results from the study contribute to the evidence that is needed to support the integration of mindfulness and acceptance-based strategies into care services offered to AWH in Uganda. We compared the effect of the DNA-v to the SOC on improving psychological flexibility and self-reported adherence to ART among older AWH (15–19 years) receiving care at an urban clinic in Kampala. We hypothesized that participants receiving the DNA-v intervention would report higher levels of psychological flexibility and improved adherence to ART at postintervention and 3-month follow-up compared to the SOC group. Results from the study show that the intervention group reported higher levels of psychological flexibility as hypothesized, however, both groups did not differ significantly in self-reported adherence. That is, results suggest that the DNA-v as an MABI has the potential to improve psychological flexibility among AWH but does not necessarily result in improved adherence to ART. To our knowledge, this is the first study to test the effect of the DNA-v on improving psychological flexibility and adherence to ART among AWH in a low-resource context through a controlled trial. Although similar studies have used MABIs with adults living with HIV and have tested for mental health outcomes,⁶⁹ studies conducted with AWH are rare and highlight our critical contribution to the literature.

Higher levels of psychological flexibility reported in the intervention group but not in the control group at the follow-up stage indicate that psychological flexibility as a self-regulation skill can be trained and improved among AWH. This observation is important for HIV treatment and care of AWH who experience developmental changes that require enhanced levels of self-management for coping. Psychological flexibility denotes becoming open to life experiences non-judgmentally, dealing with difficult situations with awareness, and deliberately making choices that bring meaning to an individual.^{32,44} Since living with HIV presents numerous challenges,

developing skills that help one approach challenges with openness, is therapeutic. Our findings are consistent with literature that identifies psychological flexibility as a skill that can improve with time through deliberate training or psychotherapy.^{43–45} Moreover, the results concur with studies that found improved levels of psychological flexibility among adolescents who were exposed to DNA-v training.^{42,67} By improving psychological flexibility among AWH, our results confirm that MABIs are transdiagnostic and can address a range of emotional and behavioral challenges across conditions.⁷⁰

The study did not find a significant difference between groups on self-reported adherence to ART. Results suggest that while the DNA-v training could have helped AWH develop self-regulatory skills by improving psychological flexibility, such a transformation did not impact adherence to ART. This could mean that adherence to ART among AWH is influenced by a range of factors that may extend beyond behavioral regulation. The study findings diverge from the literature that suggests that through psychological flexibility and values clarification, MABIs help patients make active treatment choices including adhering to treatments.⁷¹ Results further deviate from research findings indicating that interventions that promote self-efficacy and self-acceptance which are core components of psychological flexibility result in improved adherence among people with HIV.⁷² Furthermore, results do not align with studies that found ACT to yield improved adherence to medication among patients with chronic conditions.⁷³ Nevertheless, it is worth noting that while MABIs may empower patients to make values-consistent choices such as adhering to a treatment regimen, such individual-level choices may be compromised by high-level social and economic factors that impact the entire cascade of care. In low-resource settings such as Uganda, factors such as abject poverty limit access to food, while high levels of stigma and discrimination are suffered at home, in the community, and at school due to misconceptions about HIV. The far distances to health centers are an access barrier, while unreliable family support due to the loss of biological parents, and school arrangements which limit privacy¹⁵ may affect health-seeking choices AWH make regardless of their levels of resilience. This may partially explain why the vast majority of studies reporting a positive

association between mindfulness and acceptance skills and medication adherence have been conducted in resource-rich contexts.

The study findings, however, are consistent with the literature which highlights that behavioral change approaches that target adherence often register limited success.⁷⁴ The findings are also in line with the literature that suggested that PLWHA interventions that promote psychological flexibility only improve overall well-being but do not explain intentional or unintentional non-adherence to ART.⁶⁹ Furthermore, our results confirm earlier findings that adherence to ART is affected by a range of factors that may go beyond individual control and have the power to compromise individual choices or meaning.⁷⁵

Although there were no significant differences between groups on self-reported adherence to ART, within-group mean differences showed that participants in the intervention arm had improved adherence scores at posttest and 3-month follow-up adjusting for baseline scores compared to the SOC group. As adherence is a salient challenge among AWH, this minor improvement is still promising and can inform follow-up studies. Furthermore, since the study is based on self-reported measures which can be subject to bias, more objective measures such as pill counts or electronic monitoring systems could provide alternative findings. This observation is supported by a systematic review that found large–medium effect sizes among studies that measured adherence by pill count or electronic event monitoring systems.⁷⁴ Nevertheless, subjective assessments are still valid measures of adherence and have widely been used across HIV studies.^{64,76}

Strength and Limitations

The study had notable strengths. First, randomizing participants helped to balance the study groups and thus controlled for possible confounders that could have affected the interpretation of the results. The use of this rigorous design lowered the risk of bias, which is critical when generating evidence that can influence the uptake of health interventions. Secondly, adding a follow-up period to the study was another strength. The 3-month follow-up helped to gauge whether the impact of the intervention could be sustained.

Notwithstanding these strengths, the study had some limitations. Foremost, the results may have been affected by the high rate of attrition, given that more vulnerable participants may have dropped out. However, since groups were almost equivalent at baseline and attrition did not differ significantly between conditions at 3-month follow-up, its effect is likely to be minimal. Nevertheless, smaller numbers in randomized conditions could still provide useful insights. Furthermore, while our 3-month follow-up is still considered sufficient to identify consistency in the behavioral patterns, multiple follow-up assessments (eg, at 6, 9, and 12 months) would have given a clearer picture of the sustainability of the intervention effect. Taken together, the study provides a starting point for future

studies with larger samples and longer follow-ups to generate additional evidence.

Conclusion

Adherence to ART remains a significant challenge among AWH and may best be addressed by a comprehensive strategy that incorporates various components of care. The results from this study demonstrate that while an MABI for young people did not improve self-reported adherence to ART among AWH, it led to improvements in levels of psychological flexibility. Although adherence was the primary outcome measure, improvements in psychological flexibility are promising for addressing care concerns that may improve adherence in the long run. Given that psychological flexibility represents a set of self-regulation components (self-acceptance, self-efficacy, and resilience), strengthening it as a skill among AWH can result in better clinical outcomes. Since the literature connecting MABIs to adherence to ART among adolescents is still in its infancy, future studies are needed with larger samples, and longer follow-up periods evaluating psychological flexibility as a mechanism of change underlying MABIs with objective measures of adherence.

Acknowledgments

The authors wish to thank Louise Hayes, Joseph Ciarrochi, and Ann Bailey for permitting us to use the DNA-v model. We also thank the Institution for Adherence for licensing the use of the adherence scale, the management and staff of Kisenyi Health Center IV in Kampala for allowing the study to be conducted there, the adolescents who participated in the study, the Makerere University Behavioral Social Science project (Mak-BSSR), and the research team: Kelly Gonzaga Kyagaba, Joseph Balikuddembe, and Hope Kirabo.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Research reported in this publication was supported by the Fogarty International Center (FIC), the National Institute of Alcohol Abuse and Alcoholism (NIAAA), and the National Institute of Mental Health (NIMH), of the National Institutes of Health (NIH) under the Award Number D43 TW011304. The study was also supported by the Makerere University Research and Innovations Fund (Mak-RIF). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health (<https://www.nih.gov/>) and the Makerere University Research and Innovations Fund (Mak-RIF).

ORCID iD

Khamisi Musanje  <https://orcid.org/0000-0003-2825-9105>

References

1. WHO. *Global update on HIV treatment 2013: results, impact and opportunities*. WHO; 2013.
2. Okonji EF, Mukumbang FC, Orth Z, Vickerman-Delpont SA, Van Wyk B. Psychosocial support interventions for improved adherence and retention in ART care for young people living with HIV (10–24 years): a scoping review. *BMC Public Health*. 2020;20(1):1841.
3. WHO. *Global HIV program. Treatment and care in children and adolescents 2023*, Accessed September 20, 2023. <https://www.who.int/teams/global-hiv-hepatitis-and-stis-programmes/hiv/treatment/treatment-and-care-in-children-and-adolescents>
4. Ranjit YS, Krishnan A, Ghosh D, Cravero C, Zhou X, Altice FL. Mhealth intervention to improve treatment outcomes among people with HIV who use cocaine: protocol for a pilot randomized controlled trial. *JMIR Res Protoc*. 2022;11(3):e28332.
5. Paterson DL, Potoski B, Capitano B. Measurement of adherence to antiretroviral medications. *J Acquir Immune Defic Syndr*. 2002;31(Suppl 3):S103–S106.
6. Willie TC, Overstreet NM, Sullivan TP, Sikkema KJ, Hansen NB. Barriers to HIV medication adherence: examining distinct anxiety and depression symptoms among women living with HIV who experienced childhood sexual abuse. *Behav Med*. 2016;42(2):120–127.
7. van Wyk BE, Davids LC. Challenges to HIV treatment adherence amongst adolescents in a low socio-economic setting in Cape Town. *South Afr J HIV Med*. 2019;20(1):1002.
8. Soomro N, Fitzgerald G, Seeley J, Schatz E, Nachega JB, Negin J. Comparison of antiretroviral therapy adherence among HIV-infected older adults with younger adults in Africa: systematic review and meta-analysis. *AIDS Behav*. 2019;23(2):445–458.
9. Sekatawa E. *Adolescents and the HIV/AIDS epidemic in Uganda*. Pathfinder International; 2000.
10. UNICEF. HIV and AIDS in adolescents. Monitoring the situation of children and women 2021, Accessed September 20, 2023. <https://data.unicef.org/topic/hiv-aids/>
11. Gitahi-Kamau N, Wahome S, Bukusi EA, Ngure K. Determinants of antiretroviral therapy adherence among older adolescents living with HIV in Kenya during the transition to adult care; an observational study. *J AIDS HIV Res (Online)*. 2020;12(2):24–33.
12. Van Wyk B, Kriel E, Mukumbang F. Retention in care for adolescents who were newly initiated on antiretroviral therapy in the Cape Metropole in South Africa. *South African Journal of HIV Medicine*. 2020;21(1):1–8.
13. UNAIDS. *Global and regional data. Building on two decades of progress against AIDS*. 2021.
14. Mugabi B, Nanyingi M, Kabanda R, et al. Prevalence and factors associated with sexual and reproductive health literacy among youth living with HIV in Uganda: a cross-sectional study. *BMC Public Health*. 2023;23(1):1497.
15. Nabukeera-Barungi N, Elyanu P, Asire B, et al. Adherence to antiretroviral therapy and retention in care for adolescents living with HIV from 10 districts in Uganda. *BMC Infect Dis*. 2015;15:520.
16. Ministry of Health Uganda. Release of preliminary results of the 2020 Uganda population-based HIV impact assessment. Kampala 2022.
17. Natukunda J, Kirabira P, Ong KIC, Shibanuma A, Jimba M. Virologic failure in HIV-positive adolescents with perfect adherence in Uganda: a cross-sectional study. *Trop Med Health*. 2019;47(1):8.
18. Adejumo OA, Malee KM, Ryscavage P, Hunter SJ, Taiwo BO. Contemporary issues on the epidemiology and antiretroviral adherence of HIV-infected adolescents in sub-Saharan Africa: a narrative review. *J Int AIDS Soc*. 2015;18(1):20049–20049.
19. Mental Health and Antiretroviral Treatment Adherence among Adolescents Living with HIV. Evidence on risk pathways and protective factors [press release]. 2021.
20. UPHIA. Uganda population-based HIV impact assessment 2016–2017. Summary sheet: preliminary findings. 2017.
21. Kizito S, Namuwonge F, Brathwaite R, et al. Monitoring adherence to antiretroviral therapy among adolescents in southern Uganda: comparing Wisepill to self-report in predicting viral suppression in a cluster-randomized trial. *J Int AIDS Soc*. 2022;25(9):e25990.
22. Bermudez LG, Jennings L, Ssewamala FM, Nabunya P, Mellins C, McKay M. Equity in adherence to antiretroviral therapy among economically vulnerable adolescents living with HIV in Uganda. *AIDS Care*. 2016;28(Suppl 2):83–91.
23. Inzaule SC, Hamers RL, Kityo C, Rinke de Wit TF, Roura M. Long-term antiretroviral treatment adherence in HIV-infected adolescents and adults in Uganda: a qualitative study. *PLoS One*. 2016;11(11):e0167492.
24. Hudelson C, Cluver L. Factors associated with adherence to antiretroviral therapy among adolescents living with HIV/AIDS in low- and middle-income countries: a systematic review. *AIDS Care*. 2015;27(7):805–816.
25. MacCarthy S, Saya U, Samba C, Birungi J, Okobo S, Linnemayr S. “How am I going to live?”: Exploring barriers to ART adherence among adolescents and young adults living with HIV in Uganda. *BMC Public Health*. 2018;18(1):1158.
26. Naswa S, Marfatia YS. Adolescent HIV/AIDS: issues and challenges. *Indian J Sex Transm Dis AIDS*. 2010;31(1):1–10.
27. Murphy DA, Sarr M, Durako SJ, Moscicki AB, Wilson CM, Muenz LR. Barriers to HAART adherence among human immunodeficiency virus-infected adolescents. *Arch Pediatr Adolesc Med*. 2003;157(3):249–255.
28. Steinberg L. A social neuroscience perspective on adolescent risk-taking. *Dev Rev*. 2008;28(1):78–106.
29. Rana Y, Haberer J, Huang H, et al. Short message service (SMS)-based intervention to improve treatment adherence among HIV-positive youth in Uganda: focus group findings. *PLoS One*. 2015;10(4):e0125187.
30. Gutgesell ME, Payne N. Issues of adolescent psychological development in the 21st century. *Pediatr Rev*. 2004;25(3):79–85.
31. Vreeman RC, McCoy BM, Lee S. Mental health challenges among adolescents living with HIV. *J Int AIDS Soc*. 2017;20(Suppl 3):100–109.
32. Hayes SC. Acceptance and commitment therapy, relational frame theory, and the third wave of behavioral and cognitive therapies. *Behav Ther*. 2003;35(4):639–665.

33. Equality UIoH. *Psychosocial pathways and health outcomes: informing action on health inequalities*. London 2017.
34. Katz IT, Ryu AE, Onuegbu AG, et al. Impact of HIV-related stigma on treatment adherence: systematic review and meta-synthesis. *J Int AIDS Soc*. 2013;16(3 Suppl 2):18640.
35. Michliga GJ, Westergaard RP, Lama Y, et al. Avoidance, meaning and grief: psychosocial factors influencing engagement in HIV care. *AIDS Care*. 2018;30(4):511–517.
36. Kerrigan D, Grieb SM, Ellen J, Sibinga E. Exploring the dynamics of ART adherence in the context of a mindfulness instruction intervention among youth living with HIV in Baltimore, Maryland. *AIDS Care*. 2018;30(11):1400–1405.
37. Nguyen N, Lovero KL, Falcao J, et al. Mental health and ART adherence among adolescents living with HIV in Mozambique. *AIDS Care*. 2023;35(2):182–190.
38. Nabunya P, Ssewamala FM, Bahar OS, et al. Suubi4Stigma study protocol: a pilot cluster randomized controlled trial to address HIV-associated stigma among adolescents living with HIV in Uganda. *Pilot Feasibility Stud*. 2022;8(1):95.
39. Bell R. *Psychosocial pathways and health outcomes: informing action on health inequalities*. PHE Publications; 2017.
40. UNAIDS. *Integration of mental health and HIV interventions 2022*.
41. Hayes LL, Ciarrochi JV. Growing up flexibly: DNA-v as a contextual behavioural science approach to working with young people. In: Michael E, Levin MPT, Krafft J, eds. *Innovations in acceptance and commitment therapy: clinical advancements and applications in ACT*. Vol. 2020. New Harbinger Publications; 2020:296.
42. Musanje K, Camlin CS, Kanya MR, et al. Culturally adapting a mindfulness and acceptance-based intervention to support the mental health of adolescents on antiretroviral therapy in Uganda. *PLOS Global Public Health*. 2023;3(3):e0001605.
43. Falkner S, O'Dell S, Golden J. Targeting Psychological Flexibility, sleep hygiene, and physical activity in high school students using DNA-V model. Paper presented at: Annual conference for the Association of Contextual Behavioral Science; July 24–29, 2018; Montréal, Québec, Canada.
44. Fletcher L, Hayes SC. Relational frame theory, acceptance and commitment therapy, and a functional analytic definition of mindfulness. *J Ration Emot Cogn Behav Ther*. 2005;23(4):315–336.
45. Hafeman DM, Ostroff AN, Feldman J, et al. Mindfulness-based intervention to decrease mood lability in at-risk youth: preliminary evidence for changes in resting state functional connectivity. *J Affect Disord*. 2020;276:23–29.
46. Hayes L. Acceptance and commitment therapy for the treatment of adolescent depression: a pilot study in a psychiatric outpatient setting. *Mindfulness*. 2011;2:86–94.
47. Kim SH, Gerver SM, Fidler S, Ward H. Adherence to antiretroviral therapy in adolescents living with HIV: systematic review and meta-analysis. *AIDS*. 2014;28(13):1945–1956.
48. Moitra E, Herbert JD, Forman EM. Acceptance-based behavior therapy to promote HIV medication adherence. *AIDS Care*. 2011;23(12):1660–1667.
49. Graham CD, Gouick J, Krahe C, Gillanders D. A systematic review of the use of acceptance and commitment therapy (ACT) in chronic disease and long-term conditions. *Clin Psychol Rev*. 2016;46:46–58.
50. Tatta J, Willgens AM, Palombaro KM. Mindfulness and acceptance-based interventions in physical therapist practice: the time is now. *Phys Ther*. 2022;102(3):1–10.
51. Pielech M, Vowles KE, Wicksell R. Acceptance and commitment therapy for pediatric chronic pain: theory and application. *Children*. 2017;4(2):10.
52. Mahrer NE, Sommer VS, Berman DI, Ngo N, Gold JI. Virtual acceptance and commitment therapy (vACT) for youth with chronic illnesses: a case report. *Med Res Arch*. 2023;11(6):1–12.
53. Coyne LW, McHugh L, Martinez ER. Acceptance and commitment therapy (ACT): advances and applications with children, adolescents, and families. *Child Adolesc Psychiatr Clin N Am*. 2011;20(2):379–399.
54. Bateganya MH, Amanyeiwe U, Roxo U, Dong M. Impact of support groups for people living with HIV on clinical outcomes: a systematic review of the literature. *J Acquir Immune Defic Syndr*. 2015;68(Suppl 3(3)):S368–S374.
55. Zhao C, Lai L, Zhang L, et al. The effects of acceptance and commitment therapy on the psychological and physical outcomes among cancer patients: a meta-analysis with trial sequential analysis. *J Psychosom Res*. 2021;140:110304.
56. Depping MK, Uhlenbusch N, Löwe B. Acceptance and commitment therapy meets peer-support. Development of a supportive self-care intervention for patients with rare diseases: a multistage development process. *BMJ open*. 2021;11(7):e042856.
57. Hadlandsmyth K, White KS, Nesin AE, Greco LA. Proposing an acceptance and commitment therapy intervention to promote improved diabetes management in adolescents: a treatment conceptualization. *Int J Behav Consult Therapy*. 2013;7(4):12–15.
58. Woidneck MR, Morrison KL, Twohig MP. Acceptance and commitment therapy for the treatment of posttraumatic stress among adolescents. *Behav Modif*. 2014;38(4):451–476.
59. Na E, Lee K, Jeon BH, et al. Acceptance and commitment therapy for destructive experiential avoidance (ACT-DEA): a feasibility study. *Int J Environ Res Public Health*. 2022;19(24):16434.
60. Viskovich S, Pakenham KI. Pilot evaluation of a web-based acceptance and commitment therapy program to promote mental health skills in university students. *J Clin Psychol*. 2018;74(12):2047–2069.
61. Musanje K, Kanya MR, Kasujja R, et al. Acceptability of an adapted mindfulness and acceptance-based intervention to support adolescents with HIV: a qualitative study with Ugandan health care providers. *J Contextual Behav Sci*. 2023;29:160–168.
62. Ministry of Health. *Consolidated guidelines for the prevention and treatment of HIV/AIDS in Uganda*. In: Kampala 2020.
63. Morisky DE, Ang A, Krousel-Wood M, Ward HJ. Predictive validity of a medication adherence measure in an outpatient setting. *J Clin Hyperten*. 2008;10(5):348–354.
64. Aguiar A, Piñeiro C, Serrão R, Duarte R. The 8-item Morisky medication adherence scale: validation of a Portuguese version in HIV patients. *Eur J Public Health*. 2020;30(5):v246–v247.

65. Greco LA, Lambert W, Baer RA. Psychological inflexibility in childhood and adolescence: development and evaluation of the avoidance and fusion questionnaire for youth. *Psychol Assess*. 2008;20(2):93–102.
66. Jenny C, Sandra W, JoAnne D, Gustaf LA. Preliminary validation of the Swedish short version of the avoidance and fusion questionnaire for youth (AFQ-Y8) for children and adolescents with cancer. *J Context Behav Sci*. 2018;10:103–107.
67. Livheim F, Hayes L, Ghaderi A, et al. The effectiveness of acceptance and commitment therapy for adolescent mental health: Swedish and Australian pilot outcomes. *J Child Fam Stud*. 2015;24:1016–1030.
68. Schulz KF, Altman DG, Moher D. [CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials (Chinese version)]. *Zhong xi yi jie he xue bao. J Chin Integr Med*. 2010;8(7):604–612.
69. Harrison A, Scott W, Timmins L, Graham CD, Harrison AM. Investigating the potentially important role of psychological flexibility in adherence to antiretroviral therapy in people living with HIV. *AIDS Care*. 2021;33(3):337–346.
70. Hayes SC, Hofmann SG, Ciarrochi J. A process-based approach to psychological diagnosis and treatment: the conceptual and treatment utility of an extended evolutionary meta model. *Clin Psychol Rev*. 2020;82:101908.
71. Graham CD, McCracken LM, Harrison A, Walburn J, Weinman J. Outlining an acceptance and commitment therapy approach to treatment non-adherence. *Br J Health Psychol*. 2022;27(1):1–12.
72. Johnson MO, Neilands TB, Dilworth SE, Morin SF, Remien RH, Chesney MA. The role of self-efficacy in HIV treatment adherence: validation of the HIV treatment adherence self-efficacy scale (HIV-ASES). *J Behav Med*. 2007;30(5):359–370.
73. As'hab PP, Keliat BA, Wardani IY. The effects of acceptance and commitment therapy on psychosocial impact and adherence of MDR-TB patients. *J Public Health Res*. 2021;11(2):2737.
74. Conn VS, Ruppap TM, Enriquez M, Cooper P. Medication adherence interventions that target subjects with adherence problems: systematic review and meta-analysis. *Res Social Admin Pharm*. 2016;12(2):218–246.
75. Lantche MW, Fokam J, Cheudjui AJN, et al. Factors associated with non-adherence to antiretroviral therapy among HIV-infected adolescents aged 15-19 years: a snapshot from the mother and child center in Yaounde, Cameroon. *Pan Afr Med J*. 2021;39(1):154.
76. Biney IJK, Kyei KA, Ganu VJ, et al. Antiretroviral therapy adherence and viral suppression among HIV-infected adolescents and young adults at a tertiary hospital in Ghana. *Afr J AIDS Res*. 2021;20(4):270–276.