

Abiotic and biotic correlates of the occurrence, extent and cover of invasive aquatic Elodea nuttallii

Crane, K., Kregting, L., Coughlan, N. E., Cuthbert, R. N., Ricciardi, A., MacIsaac, H. J., Dick, J. T. A., & Reid, N. (2022). Abiotic and biotic correlates of the occurrence, extent and cover of invasive aquatic Elodea nuttallii. *Freshwater Biology*, *67*(9), 1559-1570. https://doi.org/10.1111/fwb.13960

Published in:

Freshwater Biology

Document Version: Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:

Link to publication record in Queen's University Belfast Research Portal

Publisher rights

Copyright 2022 the authors.

This is an open access article published under a Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: http://go.qub.ac.uk/oa-feedback

ORIGINAL ARTICLE

Freshwater Biology WILEY

Abiotic and biotic correlates of the occurrence, extent and

cover of invasive aquatic Elodea nuttallii

Revised: 22 May 2022

Kate Crane^{1,2} | Louise Kregting^{2,3} | Neil E. Coughlan^{1,2,4} | Ross N. Cuthbert^{1,2,5} Anthony Ricciardi⁶ | Hugh J. MacIsaac⁷ | Jaimie T.A. Dick^{1,2,8} | Neil Reid^{1,8}

¹School of Biological Sciences, Queen's University Belfast, Belfast, UK

²Queen's University Marine Laboratory, Portaferry, UK

³School of Natural and Built Environment, Queen's University Belfast, Belfast, UK

⁴School of Biological, Earth and Environmental Science, University College Cork, Cork, Ireland

⁵GEOMAR Helmholtz-Zentrum für Ozeanforschung Kiel, Kiel, Germany

⁶Redpath Museum, McGill University, Montreal, QC, Canada

⁷Great Lakes Institute for Environmental Research, University of Windsor, Windsor, ON, Canada

⁸Institute for Global Food Security (IGFS), Queen's University Belfast, Belfast, UK

Correspondence

Louise Kregting, Queen's University Marine Laboratory, 12-13 The Strand, Portaferry BT22 1PF, UK. Email: I.kregting@qub.ac.uk

Funding information

KC was supported through contributions from Queen's University Belfast, the University of Windsor, McGill University and Waterways Ireland. NEC and JTAD were supported by the Irish Environmental Protection Agency (EPA) research grant 2015-NC-MS-4. RNC acknowledges support from the Alexander von Humboldt Foundation and Department for the Economy (DfE), Northern Ireland. AR and HJM acknowledge support from Natural Sciences and Engineering Research Council of Canada (NSERC).

Abstract

- 1. Biological invasions, especially invasive alien aquatic plants, are a major and growing ecological and socioeconomic problem worldwide. Freshwater systems are particularly vulnerable to invasion, where impacts of invasive alien species can damage ecological structure and function. Identifying abiotic and biotic factors that mediate successful invasions is a management priority. Our aim was to determine the environmental correlates of Elodea nuttallii; a globally significant invasive aquatic species.
- 2. Elodea nuttallii presence/absence (occurrence), extent (patch area) and percentage cover (density) was visually assessed from a boat throughout Lough Erne (approximately 144 km²), County Fermanagh, Northern Ireland during the active summer growth season (July-September). In addition, substrate type and zebra mussel Dreissena polymorpha occurrence was recorded. Fourteen water chemistry variables were collected monthly from 12 recording stations throughout the lake during the 9 years before the survey to spatially interpolate values and establish temporal trajectories in their change. Shoreline land use was derived from CORINE land cover maps. Environmental associations between E. nuttallii, substrate, D. polymorpha, water chemistry and land use were assessed.
- 3. Elodea nuttallii occurrence was positively associated with water conductivity, alkalinity, suspended solids, phosphorus (both total and soluble) and chlorophyll-a concentrations, but negatively associated with pH and total oxidised nitrogen. E. nuttallii patch extent and proportional cover were positively associated, to varying degrees, with the presence of *D. polymorpha*, biological oxygen demand, water clarity and soft substrate, but negatively associated with urban development and ammonium.
- 4. Elodea nuttallii displayed high levels of phenotypic plasticity in response to environmental variation, allowing it to adapt to a wide range of conditions and potentially gain competitive advantage over native or other invasive macrophytes.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited. © 2022 The Authors. Freshwater Biology published by John Wiley & Sons Ltd.

5. It is evident that multiple abiotic and biotic factors, including facilitation by cooccurring invasive dreissenid mussels, interact to influence the distribution and abundance of *E. nuttallii*. Thus, it is necessary to consider a more comprehensive environmental context when planning *Elodea* management strategies.

KEYWORDS

competition, Dreissena polymorpha, freshwater ecosystems, macrophytes, nutrients

1 | INTRODUCTION

Biological invasions continue to alter and, in some cases, detrimentally impact ecosystem function worldwide, resulting in a plethora of environmental, economic and social problems (Haubrock et al., 2021; Simberloff et al., 2013). Rates of biological invasion continue to increase, with intensifying trade and transport networks driving introductions from disparate biogeographical regions (Bailey et al., 2020; Seebens et al., 2017, 2021). Management strategies for invasive alien species are reliant on evidence-based, data-driven science (Dick et al., 2017), yet have hitherto been largely insufficient in reducing ecological and economic damages (Cuthbert, Pattison, et al., 2021a). Specifically, understanding the abiotic and biotic correlates of biological invasions, whose status can be quantified using numerous metrics such as occurrence, extent or proportional cover, requires a holistic approach in considering potential drivers of invasion throughout the total environment. Therefore, the consideration of multiple variables in concert is increasingly required to better understand and predict invasions worldwide.

Aquatic ecosystems are regarded as particularly vulnerable to biological invasions and their impacts (Ricciardi & MacIsaac, 2011). Invasive aquatic macrophytes often are recorded to be ecologically and economically damaging, for example, through increased flood risk, devaluation of property, the disruption of navigation, water abstraction, irrigation and recreational activities (Hussner et al., 2017; Oreska & Aldridge, 2011). Studies typically have focused on a small number of variables to explain invasion success, including the relationships between macrophytes and spatiotemporal patterns in water quality and the surrounding land use (Lougheed et al., 2001; Sass et al., 2010). A more comprehensive understanding of invasion drivers, including the emergent effects of other species impacts and their abiotic interactions, is needed to enhance predictive capacities for future invasions.

Elodea nuttallii (Planch.) St John is a perennial, submerged aquatic macrophyte which is native to North America and invasive in Europe, Asia and Australasia (Barrat-Segretain et al., 2002; Jones et al., 2000; Zehnsdorf et al., 2015). Introduced to Great Britain and Ireland in 1966, *E. nuttallii* typically inhabits lakes, ponds and slow-moving rivers (Barrat-Segretain et al., 2002; Champion et al., 2010), often displacing *Elodea canadensis*, a congeneric invasive macro-phyte (Simpson, 1990). Notably, *E. nuttallii* can grow in highly eutrophic, turbid waters (Cook & Urmi-König, 1985; Ozimek et al., 1993; Thiébaut & Muller, 1999), and can rapidly dominate invaded ecosystems through the formation of dense monospecific stands that outcompete native macrophytes and significantly alter freshwater

communities (Angelstein & Schubert, 2008; Champion et al., 2010; Thouvenot & Thiébaut 2018; Zehnsdorf et al., 2015). Furthermore, it appears that *E. nuttallii* can be transported long distances overland by anthropogenic vectors (Coughlan et al., 2018; Kelly et al., 2014).

Interestingly, E. nuttallii often co-occurs with invasive dreissenid bivalves (Crane et al., 2020), such as the zebra mussel Dreissena polymorpha (Pallas 1771), itself a prolific Ponto-Caspian species that frequently dominates invaded ecosystems, causing myriad ecological and socioeconomic impacts (Sousa et al., 2014). Dreissena polymorpha impacts include displacement of native mussels, increased water clarity, altered nutrient cycling, and shifts in filamentous algal blooms and macrophyte assemblage composition (Ricciardi, 2003; Ricciardi et al., 1998; Rosell et al., 1999; Ward & Ricciardi, 2007). Given increasing accumulations of invaders in aquatic ecosystems (Ricciardi, 2015), understanding how multiple invasive alien species interact and respond to variability in environmental conditions is essential to understanding the spatiotemporal dynamics of invasions, which inform the scale of any management actions required. Empirical data indicate that the presence of D. polymorpha can promote facilitative interactions in favour of E. nuttallii (Crane et al., 2020). However, the extent of potential facilitations between invasive alien species in field-based conditions has not been rigorously tested, nor compared with additional environmental variables that might mediate invasion dynamics.

The aim of this study was to identify the environmental correlates of a major *E. nuttallii* invasion in Lough Erne; a large (c. 144 km²) freshwater lake in Northern Ireland, capturing the invasion status using three metrics: occurrence (presence/absence), extent (area in hectares) and proportional cover (percentage cover). The objective was to assess environmental correlates of *E. nuttallii* including spatial and temporal trends in water quality variables, as well as surrounding land cover and land use. In addition, we included the co-occurrence of *E. nuttallii* with *D. polymorpha* within our analyses with the objective of assessing the potential role played by interspecific interactions in the distribution of *E. nuttallii* alongside the environmental variation.

2 | METHODS

2.1 | Study site

We assessed environmental correlates of the *E. nuttallii* invasion of Lough Erne, County Fermanagh, Northern Ireland, UK. *Elodea nuttallii* was first recorded in Lough Erne in 2006, and by 2010 it had spread extensively, causing significant ecological, navigational and recreational problems, while competitively displacing E. canadensis that has invaded previously (Kelly et al., 2015). Lough Erne is a naturally eutrophic lake consisting of two major waterbodies connected by a 10-km stretch of the River Erne (O'Higgins, 2018). Lower Lough Erne (109 km²) is the third largest lake in the UK, with a mean depth of 11.9 m. Upper Lough Erne (34.5 km²) has a mean depth of 2.3 m (Lawrie et al., 1992) and comprises an array of wide river channels, open lake and numerous islands, creating a highly complex sinuous shoreline. The water chemistry of the two lakes reflects the underlying geology of limestone and sandstone which gives rise to both carbonate-rich and softly acidic waters. The catchment has a sparse human population and is largely agricultural, with mature, seminatural riparian woodland. Designated as a Ramsar Site (DOE, 1997), Special Protection Area (DOE, 1997) and including ten Areas of Special Scientific Interest (DAERA, 2018), Upper Lough Erne is further classified as a Special Area of Conservation (JNCC, 2015), and the Lough is an internationally important area of conservation concern

2.2 | Surveys

Upper and Lower Lough Erne were circumnavigated by boat and a visual distribution survey was carried out for mature E. nuttallii at the peak of the growing season, July-September 2015. Immature plants at depth can be surveyed using rake samples, but this was not undertaken here as a consequence of the near ubiquity of obvious and mature patches of *Elodea* within the study system. High water clarity allowed for the visual detection of stands/swards of plants. A bathyscope and waterproof GoPro Hero 5 were used to identify the extent of each patch of macrophytes. The boundary of each patch was recorded using a handheld Trimble Yuma Rugged GPS unit (model YMA-FGS6AS-00) and mapped using ARCMAP 10.5 (ESRI, California, USA). Multiple GPS points were used to ensure that the boundaries of patches were recorded accurately. The proportional cover of E. nuttallii relative to other vegetation within the patch was visually estimated by eye using an ordinal categorical scale: "0" = 0% E. nuttallii, 100% native macrophytes; "1" = 0%-25% E. nuttallii; "2" = 26%-50% E. nuttallii; "3" = 51%-75% E. nuttallii; "4" = 76%-100% E. nuttallii.

2.3 | Environmental variables

During the surveys, substrate type was visually estimated as being either hard (fine and coarse gravel/shingle, and larger stones and/or rock) or soft (silt, sand and mud). *Dreissena polymorpha* presence or absence (also visually estimated) was recorded for each vegetation patch. Other invasive bivalves are not known to reside within the study site, and there is a paucity of native bivalves. Presence was recorded on a binary scale, as visual survey from a boat precluded robust determination of abundance. For logistical and practical Freshwater Biology –WILEY

purposes, substrate samples as well as water quality measures were not measured empirically at each surveyed macrophyte patch. Instead, 14 water quality variables were obtained from the Northern Ireland Environment Agency (NIEA) Water Management Unit, who have carried out routine, standardised, lake water monitoring and surveillance since 2006 in compliance with the Water Framework Directive, National Water Quality Monitoring Programme (Table S1). Water samples were collected monthly by NIEA staff from 12 recording stations throughout the lake system (Figure S1). Recording stations were c. 5 km apart, much further than the distance between surveyed macrophyte patches. Thus, the 14 water chemistry variables were Kriged using the Spatial Analyst toolbox in ArcGIS to interpolate values for regions between recording stations, and to extrapolate values out to the lake's edge to create a heat map of spatial variation in each variables (Figure S1). Kriging used an ordinary spherical semivariogram model with a search radius of three points; as recording stations were distributed linearly from the lake's source to mouth, this emulated a moving window averaging water chemistry values across a combination of three stations (one upstream, one in the middle, one downstream), providing ecologically plausible smoothed heat map surfaces for each variable (Figure S1). In addition to spatial interpolation, temporal trends in water chemistry variables also were captured. Data for each variable were plotted separately against time for each recording station, and a linear regression fitted to obtain the regression coefficient, i.e. the slope of the line (β -value) representing the direction of any trend in that variable at each station. Raw data were standardised to have a $\bar{x} = 0$ and $\sigma = 1$ so that regression coefficients were directly comparable. For each variable, spatiotemporal variation was again captured by Kriging the spatially explicit regression coefficients as above to create a heat map of change in each variable from 2006 to 2015 (Figure S1). Imputed values for both the mean and regression coefficient for each variable were extracted from the heat map surfaces for the centroid of each macrophyte patch of E. nuttallii mapped during the survey. It should be noted that variation in water chemistry values between adjacent macrophyte patches was, necessarily, minimised by the nature of the interpolation methods which is further commented on in the Discussion.

Land cover and land use were derived from the Land Cover Map (2007), produced by the Centre for Ecology and Hydrology from analysis of satellite or aerial imagery (www.ceh.ac.uk/services/land-cover-map-2007). Each patch of *E. nuttallii* was buffered to a distance of 500m, and the extent (area in hectares) of terrestrial land cover or land use within each was extracted representing variation in adjacent habitats. In total, 31 environmental variables were measured or estimated for each patch of *E. nuttallii* (Table 1).

2.4 | Data analyses

Principal components analysis (PCA) with varimax rotation was used to reduce an initially large set of putative explanatory variables, exhibiting a high degree of multicollinearity, to a much-reduced set. ILEY- Freshwater Biology

Statistical and ecological convention is that all principal component axes with an eigenvalue >1.0 are retained for further analyses (following, e.g., Kaiser, 1960, Franklin et al., 1995 and Björklund, 2019).

Generalised linear mixed models (GLMMs) were used to examine three dependent variables: *E. nuttallii* (i) occurrence was fitted as a binomial distribution with a logit link function (accounting for its binary [0/1] distribution); (ii) extent, i.e. area in hectares fitted as a Gamma distribution with a logit link function (accounting for its positive right-skew, i.e., a high frequency of small patches and a low frequency of large patches); and (iii) proportional cover, i.e. estimated percentage cover within each patch fitted using a Gaussian distribution and identity link function (accounting for its normal bellcurve distribution). Model structure was identical for all three models: Lough_ID (i.e., Upper or Lower) was fitted as a two-level random factor to account for the pseudoreplication of multiple observations per lake whilst *D. polymorpha* presence or absence (0/1) and substrate type (hard or soft) were fitted as two-level fixed factors. Six principal component axes representing the water quality variables

TABLE 1 Principal components analysis (PCA) of environmental variables describing water quality, trends in water quality and land cover/ use.

		Principal component axes					
Variable groupings		PC1	PC2	PC3	PC4	PC5	PC6
	Eigenvalue	10.277	4.490	4.152	3.066	1.718	1.640
	% variance	33.150	14.484	13.395	9.892	5.541	5.291
	Cumulative % variance	33.150	47.635	61.082	70.920	76.461	81.752
Water quality, Mean values	1. pH	-0.926	0.227	-0.196	0.096	-0.132	0.091
(2006–15)	2. ALK	0.804	0.318	0.121	0.304	-0.069	-0.034
	3. COND	0.935	0.093	-0.144	0.208	0.084	0.144
	4. COL	-0.226	-0.254	0.789	0.352	-0.036	-0.293
	5. SS	0.954	0.098	-0.005	0.197	0.087	0.049
	6. P(SOL)	0.878	-0.267	0.221	0.141	0.099	-0.054
	7. P(TOT)	0.755	-0.530	-0.040	0.166	0.123	0.072
	8. NH ₄	0.299	0.178	0.862	0.223	0.060	-0.187
	9. NO ₂	0.299	-0.051	-0.032	0.394	-0.504	-0.357
	10. TOxN	-0.926	0.267	-0.064	-0.137	-0.070	0.106
	11. CHL A	0.939	-0.231	-0.211	-0.009	0.096	0.059
	12. BOD	-0.195	-0.269	-0.753	-0.319	-0.046	0.263
	13. pH	-0.024	-0.886	-0.183	-0.119	0.036	-0.083
Temporal trends (2006–15)	14. ALK	0.263	-0.651	0.166	0.642	0.126	0.098
	15. COND	0.479	0.015	0.103	0.755	0.085	0.111
	16. COL	-0.203	0.877	0.007	-0.329	-0.007	-0.032
	17. SS	-0.188	0.849	0.258	0.262	-0.043	0.078
	18. P(SOL)	-0.890	0.403	0.157	-0.008	-0.092	-0.033
	19. P(TOT)	0.013	0.448	0.486	0.038	0.359	0.110
	20. NH ₄	-0.528	0.117	-0.601	-0.529	-0.106	0.038
	21. NO ₂	-0.211	-0.307	-0.343	-0.773	0.151	0.095
	22. TOxN	0.915	-0.121	0.212	0.250	0.042	-0.119
	23. CHL A	0.413	-0.585	0.018	-0.207	0.014	0.246
	24. BOD	0.836	0.222	0.395	0.221	0.130	-0.022
Land cover/use	25. Other farmland	0.264	-0.173	-0.258	-0.156	0.650	-0.207
	26. Improved grassland	0.071	0.043	0.107	0.005	0.282	-0.869
	27. Conifer plantations	-0.126	-0.010	-0.129	-0.212	-0.587	0.108
	28. Deciduous woodland	0.327	-0.007	-0.410	0.276	-0.135	0.445
	29. Other woody habitats	-0.479	0.185	-0.198	-0.108	-0.171	0.288
	30. Bog, fen & moor	0.117	-0.015	0.067	-0.031	0.520	-0.090
	31. Urban/suburban areas	0.038	0.171	0.812	-0.143	-0.010	0.183

Bold values indicate the presence of substantial principle component axis loadings (r > 0.5)

and land cover or land use were fitted as covariates. The residuals of each model were examined using Kolmogorov–Smirnov tests to ensure normality and adherence to the assumptions of each test and its selected distribution. Models were used to assess the environmental correlates *E. nuttallii* and were not used for prediction, i.e. there was no creation of probability heat surfaces or extrapolation of values to unsurveyed sites. For this reason, the information theoretic approach of all subset regression with model selection using the Akaike Information Criterion (AIC) was not used. Instead, fitting the global model was favoured for the purposes of comparability, i.e. fitting a standard model structure for each independent variable allowing the effects of each variable to be explicitly reported, even at p > 0.05.

Each principal component axis was associated with a large number of collinear variables and not all may have been related to E. nuttallii. However, a variety of individual environmental variables associated with substantial principal component axis loadings (r > 0.5; Table 1) were found to be significantly associated (p < 0.05) with each of the three dependent variables across different model axes using GLMM (i.e., PC1 for binary E. nuttallii occurrence and PC3 for E. nuttallii area and proportional cover; Table 2). Most environmental variables were nonparametric. Thus, Mann-Whitney U-tests were used to determine significant differences in PC1 variables between sites where E. nuttallii was present compared to those where it was absent. Spearman's rank correlations were used to determine significant relationships between PC3 variables and E. nuttallii extent (patch size) and Kruskal-Wallis tests were used to determine significant relationships between PC3 variables and E. nuttallii proportional cover category. Probability values from multiple tests were Bonferroni-corrected for multiple comparisons. Statistical analyses were performed using SPSS v25 (IBM) and all significant relationships were plotted using SIGMAPLOT v12 (Systat Software Inc.).

3 | RESULTS

3.1 | Distribution

Elodea nuttallii was found to be widespread throughout the Lough Erne (Figure 1) but occurred most frequently and with greater proportional cover in Upper reaches of the system (Figure 2a). Elodea nuttallii proportional cover was positively related to patch extent (i.e., size of area in hectares; Kruskal-Wallis $\chi^2_{df=4} = 33.306$, p < 0.001; Figure 2b), thereby indicating that larger patches of *E. nuttallii* tended to have a higher dominance by the invader within the community. Patch extent varied from 0.04 to 50.6 ha. Proportional cover category "0" representing 0% *E. nuttallii* and 100% native macrophyte had a mean patch size of ~2 ha. Proportional cover category "1" representing $\leq 25\%$ *E. nuttallii* cover had a mean patch extent increased with proportional cover, with category '4' (i.e., 75%–100% *E. nuttallii* cover) having a mean patch size of ~12 ha.

Freshwater Biology –WILEY

A total of 31 environmental variables were reduced to six principal component axes which captured 81.752% of variation in the dataset (Table 1). Of these, two contributed significantly to subsequent GLMMs: PC1 accounted for 33.150% and PC3 for 13.395% of environmental variation (Table 1). *Elodea nuttallii* occurrence was positively associated with PC1 (Table 2a), while the extent of each patch was negatively associated with PC3 and positively associated with *D. polymorpha* presence (Table 2b). Patch proportional cover was negatively associated with PC3 and varied with substrate type (Table 2c). Not all variables that contributed to PC1 and PC3 had loadings >0.5 (Table 1) with most, but not all, significantly associated with *E. nuttallii* in univariate tests (Table 3).

3.2 | Occurrence

Sites invaded by *E. nuttallii* had significantly higher values for suspended solids, phosphorus (P, both total and soluble), chlorophyll*a*, conductivity and alkalinity (and thus had lower values for pH) than uninvaded sites (Figure 3a). Patches of *E. nuttallii* were associated with lower total oxidised nitrogen (N) than native macrophyte patches. Values for soluble P and total oxidised N exhibited declines throughout the lake system from 2006 to 2015 (i.e., had negative temporal trends). However, declines of soluble P were greater (i.e., had a steeper slope) and declines of total oxidised N were lower (i.e., had a shallower slope) where *E. nuttallii* had invaded compared to where it was absent. Likewise, presence of *E. nuttallii* accelerated increases in biological oxygen demand (BOD) (Figure 3a).

3.3 | Extent

Elodea nuttallii patches were larger (~5–6 ha) in extent when *D. polymorpha* was present than when it was absent (1–3 ha: Figure 3bi). Of those variables associated with PC3, five exhibited significant relationships with *E. nuttallii* patch extent (Table 3). Landscapes with a high proportion of urban or suburban development had the smallest patches of *E. nuttallii*, whereas the largest patches (>20 ha in size) were found in parts of the lake system that had no such development. No other land cover or land use variables were associated with *E. nuttallii*. Large patches had higher BOD than smaller ones and had lower values for Hazen scale water colour (i.e., the water within large patches was clearer). Ammonium (NH₄) declined throughout the study system from 2006 to 2015, but values were lower and its decline faster within large *E. nuttallii* patches.

3.4 | Cover

High proportional cover of *E. nuttallii* was likely to occur on soft rather than hard substrates, whereas native macrophyte patches (proportional cover category "0") were slightly more likely to occur on the latter (Figure 3c). As for extent, the proportional cover of WILEY- Freshwater Biology

Model; Explanatory variables	$\beta \pm SE$	F	n.df.	d.df.	р		
(a) <i>E. nuttallii</i> presence (F _{9,195} = 1.017, <i>p</i> = 0.428, AUC = 0.750)							
Depth	-0.075 ±0.239	0.098	1	195	0.755		
D. polymorpha presence	-0.154 ± 0.400	0.149	1	195	0.700		
Substrate type	0.262 ±0.207	1.593	1	195	0.208		
PC1	0.628 ±0.293	4.606	1	195	0.033		
PC2	0.220 ± 0.284	0.602	1	195	0.439		
PC3	-0.329 ±0.267	1.519	1	195	0.219		
PC4	0.044 ± 0.196	0.051	1	195	0.821		
PC5	0.175 ± 0.212	0.682	1	195	0.410		
PC6	0.072 ±0.238	0.092	1	195	0.762		
(b) <i>E. nuttallii</i> area (<i>F</i> _{9,195} = 4.43	12, $p < 0.001$, $r^2 = 0.03$	87)					
Depth	0.002 ± 0.105	0.042	1	195	0.837		
D. polymorpha presence	0.178 ± 0.090	3.912	1	195	0.049		
Substrate type	-0.155 ±0.092	2.848	1	195	0.093		
PC1	0.106 ±0.089	1.408	1	195	0.237		
PC2	-0.080 ±0.088	0.825	1	195	0.365		
PC3	-0.450 ±0.102	19.619	1	195	<0.001		
PC4	0.166 ±0.089	3.465	1	195	0.064		
PC5	0.143 ±0.089	2.609	1	195	0.108		
PC6	-0.007 ±0.091	0.006	1	195	0.940		
(c) E. nuttallii proportional cover ($F_{9,195} = 4.528, p < 0.001, r^2 = 0.206$)							
Depth	0.057 ±0.093	0.378	1	195	0.539		
D. polymorpha presence	0.061 ± 0.079	0.592	1	195	0.443		
Substrate type	-0.298 ±0.081	13.680	1	195	<0.001		
PC1	0.104 ±0.157	0.439	1	195	0.508		
PC2	0.054 ±0.094	0.324	1	195	0.570		
PC3	-0.407 ±0.089	20.830	1	195	< 0.001		
PC4	-0.069 ±0.086	0.641	1	195	0.424		
PC5	-0.020 ±0.079	0.064	1	195	0.801		
PC6	0.043 ± 0.080	0.289	1	195	0.592		

TABLE 2 Generalised linear mixed model (GLMM) results for *Elodea nuttallii* (a) occurrence, (b) extent (i.e., area in ha) and (c) proportional cover (i.e., % cover where lough (Upper and Lower) was fitted as a random factor

Note: Regression slope (β) values are directly comparable. Categorical variables (*D. polymorpha* and substrate type) were converted to 0/1 and standardised; thus β -values represent the slope between the mean values of the dependent variable between each category.

E. nuttallii was greatest in areas with lower levels of urban development, and higher covers were associated with lower water colour (i.e., lower Hazen values) and reduced NH_4 . Greater coverage of *E. nuttallii* intensified BOD, but also was associated with a reduction in the rate of decline of NH_4 in the study systems at highest cover (proportional cover categories "3" and "4"; Figure 3c).

4 | DISCUSSION

The present study identified significant abiotic and biotic environmental determinants of the distribution and impacts of the globally widespread, highly invasive, aquatic macrophyte *E. nuttallii*. The presence of *E. nuttallii* was correlated with water chemistry variables and substrate type, and patch size was associated with invasive *D. polymorpha* presence. Using a time series of spatial data from our study system, we also found evidence that *E. nuttallii* has altered key chemical characteristics of the system through time, specifically through accelerating rates of soluble P decline, increasing biological demands for dissolved oxygen, reducing total oxidised N losses, and slight reductions in NH_4 .

The differences in significant principal components indicate that *E. nuttallii* occurrence, extent and proportional cover were each associated with different abiotic and biotic variables, with complex combinations thus influencing invasion status. Proportional cover of patches was significantly related to the areal extent of *E. nuttallii*, such that the larger the area colonised the greater the proportional coverage – as would be expected from expansion from a central





FIGURE 2 (a) Frequency of Elodea nuttallii patches in relation to the proportional cover category for Lower and Upper Lough Erne and (b) relationship between proportional cover and mean patch extent (i.e., area in ha)

inoculum. The positive association between the extent of E. nuttallii swards and the presence of invasive D. polymorpha empirically corroborated laboratory findings that indicated a mutualism between these species (but not with the benign co-occurring alien macrophyte E. canadensis; Crane et al., 2020). Although it might be thought that mussel beds provide a complex, compact, biogenic reef-like substrate that offers an anchor for attachment, given that patch proportional cover was associated with soft (not hard) substrates, it seems that the driver of the association between patch proportional cover and D. polymorpha may be independent of substrate type and influenced more by environmental correlates, such as water quality variables. Therefore, in an absence of suitable substrates, D. polymorpha could use E. nuttallii as attachment surfaces, indeed, many of the larger patches of E. nuttallii were found in silty, sheltered bays where uprooting forces inhibiting its colonisation would be minimal, whereas native macrophytes were more commonly associated with hard substrates. However, the lack of significant relationship between E. nuttallii occurrence/proportional cover and D. polymorpha may indicate that this mutualism only effectively promotes extent

at later stages of invasion (i.e., after establishment), where it exacerbates rates of spread - and thus aerial extent.

In clear waters (i.e., low turbidity), E. nuttallii can maximise growth and extend its range into deeper water (Barko & Smart, 1981; Dar et al., 2014; Maberly & Madsen, 2002; Szabó et al., 2018). The filtering activity of *D. polymorpha* increases this water clarity (Chambers & Kalff, 1985; Higgins et al., 2008), thereby allowing deeper light penetration and potentially increasing the extent over which macrophytes can grow. Furthermore, experimental data indicate that E. nuttallii growth is strongly enhanced by the presence of D. polymorpha (Crane et al., 2020). Accordingly, when considering the high abundance of D. polymorpha in Upper Lough Erne (9.5 billion individuals; 2,238 tonnes; Maguire & Gibson, 2005), where E. nuttallii colonisation has been most successful, this mutualism potentially has a major capacity to mediate invasion dynamics at large scales. Reciprocally, there appears to be a mutualistic relationship whereby, in the field, mussels utilise E. nuttallii as a settling substrate (Bodamer & Ostrofsky, 2010; Crane et al., 2020; Horvath & Lamberti, 1997; Maclsaac, 1996). As such, mussel recruitment, coupled with macrophyte growth facilitation, may represent a positive feedback cycle that drives invasion success between the two groups.

The presence of D. polymorpha significantly influenced the extent of E. nuttallii, yet other environmental correlates were potentially determinants of macrophyte invasion dynamics. Conductivity and alkalinity were significantly higher at sites invaded by E. nuttallii than in uninvaded sites. Elodea nuttallii was, however, present where pH was lowest, preferring near-neutral conditions. This is consistent with the results of Jones et al. (2000), who found that E. nuttallii photosynthesis is reduced at >pH 7, in part due to low availability of CO₂ under alkaline conditions. Sites invaded by E. nuttallii also had significantly higher values for suspended solids. A possible explanation for this relationship might be that E. nuttallii was most commonly found on soft (i.e., silty) substrates, where suspended particles can be dislodged within dense mats of macrophytes.

1565

Dependent variable; test statistic	PCA	Independent variable	Statistic	р
(a) E. nuttallii occurrence	PC1	CHL A	-3.226	0.001
Mann-Whitney Z		TOxN trend	-3.032	0.002
		pН	-2.802	0.005
		SS	-2.734	0.006
		COND	-2.695	0.007
		TOxN	-2.562	0.010
		P(SOL)	-2.416	0.016
		P(SOL) trend	-2.228	0.026
		P(TOT)	-2.047	0.043
		ALK	-2.020	0.043
		BOD trend	-1.987	0.047
		NH ₄ trend	-0.238	0.812
		Other woody habitats	-0.180	0.857
(b) E. nuttallii extent	PC3	COL	-0.257	<0.001
Spearman's r		NH ₄ trend	0.179	0.010
		BOD	0.160	0.022
		NH ₄	-0.159	0.023
		Urban land cover	-0.156	0.026
(c) E. nuttallii proportional cover	PC3	COL	22.821	<0.001
Kruskal–Wallis χ^2		NH ₄ trend	17.528	0.002
		BOD	15.390	0.004
		NH ₄	13.911	0.008
		Urban land cover	5.877	0.209

CRANE ET AL.

TABLE 3 Univariate tests of the relationship between *Elodea nuttallii* (a) occurrence, (b) extent and (c) proportional cover with the individual explanatory variables making up the principal component axes with which they were related in Table 2

variable.	vote: The	values in the s	tatistic column re	effect the test st	atistic named und	er the dependent
	/ariable.					

However, a more detailed assessment of substrate types in relation to grain size and composition should be considered, rather than the straightforward visual estimation of substrate used in the present study. Nevertheless, sites invaded by *E. nuttallii* had significantly higher values for total and soluble phosphorus. Elodea species reguire moderate-to-high nutrient levels within an ecosystem, and as such prefer eutrophic waters (Baldy et al., 2015; Melzer, 1999). The P content in the sediment can be taken up via E. nuttallii roots and released back into the water column via leakage from the plant or decomposition (Barko & Smart, 1980; Carignan & Kalff, 1980; van Donk et al., 1993), which would in turn support higher values of P within E. nuttallii patches and increase growth rates. This increased growth may afford E. nuttallii a competitive advantage in canopy formation, thus inhibiting competition from co-occurring species such as E. canadensis (Kelly et al., 2015). If plant biomass is high, there also can be significant effects on P and N cycling (Frankouich et al., 2006; James et al., 1999; Ozimek et al., 1993). Indeed, values for soluble P declined throughout the lake system over the study period, and particularly in the presence of *E. nuttallii*. Nonetheless, the importance of pollution regulation measures should not be discounted for the general decline in P from agricultural runoff, and E. nuttallii appeared

to perform better in rural, agricultural areas as compared to those with greater urbanisation.

Ammonium in the system also declined over the study period, and was lower, and its decline faster, within large E. nuttallii patches. Equally, total oxidised N was lower in E. nuttallii patches, but declined more slowly, when E. nuttallii was present. Elodea nuttallii takes up NH₄-N to generate plant biomass (Bishop & Eighmy, 1989; Dendéne et al., 1993; Rolland & Trémolières, 1995), and biomass increases as nutrient concentration rises (Rolland & Trémolières, 1995). Therefore, the lower values and greater temporal declines in NH₄ could be attributed to uptake by E. nuttallii during expansion and colonisation. Furthermore, a laboratory study has shown that where E. nuttallii is present with high D. polymorpha densities there is a greater depletion of NH₄ than in the absence of D. polymorpha (Crane et al., 2020), even though D. polymorpha can excrete significant concentrations of NH₄ (Gardner et al., 1995). This suggests that NH₄ uptake by E. nuttallii increases at higher D. polymorpha densities, and thus their co-occurrence in the present study may have synergistically altered such nutrient levels. Nevertheless, these data were generated from spatially kriged (so called "rubber-sheeted") values between recording stations throughout the lake system.

WILEY- Freshwater Biology



FIGURE 3 Relationships between Elodea nuttallii measured as (a) occurrence (x-axis a = absence, P = presence), (b) extent (i.e., area in ha) and (c) proportional cover (categorical 0<1<2<3<4) against water quality variables identified as significant independent correlates of each dependent variable in Table 3. For definitions and units see Table S1 and Figure S1. Values are means ± 1 SE

The average distance between recording stations was 5 km whilst the average distance between *E. nuttallii* patches was only 0.35 km. Spatially kriged values from heat maps extracted at a higher spatial resolution than that of the input data, derived from interpolated and extrapolated averages across space, are likely to produce small absolute differences as a result of how they were generated. This should not be interpreted as rendering the relationships established here as less meaningful, but rather support the found relationships probably being indicative of greater biological and ecological significance if only real-time values had been available for each patch of E. nuttallii synchronously with the survey, which was not just impractical but impossible. It also should be noted that this study surveyed mature plants only and that immature plants may have occurred at the study sites at some depth (thus, being missed by surveys). If sufficiently numerous, these immature plants also may have impacted water chemistry variables.

Sites invaded by E. nuttallii additionally had significantly higher values for chlorophyll-a, indicative of higher productivity,

most notably of epiphytic algae associated with E. nuttallii (Kelly et al., 2015). Like the increased amounts of suspended solids within dense E. nuttallii patches, the reduced surface water turbidity where E. nuttallii patches reach the water surface may provide suitable conditions for rapid, localised algal growth. In turn, sites invaded by E. nuttallii and D. polymorpha had significantly higher values for BOD than uninvaded sites, and the larger the patch of E. nuttallii, the greater the oxygen demand. The BOD generally increased throughout the system over the duration of the study period. The vast volumes of weed occurring in the lake will, cyclically, die and decay, a process which utilises oxygen (Asaeda et al., 2001; Jewell, 1971; Pereira et al., 1994). Concerningly, the lowering of dissolved oxygen can impact other aquatic life and potentially affect the species composition, having significant knock-on effects on the biological functioning of shallow, freshwater systems (Buscemi, 1958; Dickey et al., 2021).

From a management perspective, although the prevention of further invader spread generally is considered to be the most efficient

1567

WILEY- Freshwater Biology

management approach (Coughlan et al., 2020), an improved understanding of how invasive macrophytes interact with the abiotic and biotic components of ecosystems can aid decision makers in prioritising the allocation of resources for ongoing control of populations of invasive macrophytes. A broad range of management options have been trialled for E. nuttallii populations, including biological control with herbivorous fishes, mechanical control via cutting, shading and water drawdown, as well as chemical control via herbicide applications (Zehnsdorf et al., 2015). Effective methods for the prevention of introduction and further spread of invasive macrophytes, such as thermal shock (e.g., steam; Crane et al., 2019), can mitigate ongoing ecological and socioeconomic impacts that arise from longterm post-invasion management (Cuthbert, Pattison, et al. 2021a; Cuthbert, Diagne, et al. 2021b; Zehnsdorf et al., 2015). In addition, an understanding of environmental determinants of invasion success can help to pinpoint resources for preventative measures for invasive macrophytes, such as via biosecurity (Coughlan et al., 2020). Likewise, careful management of existing populations of invasive macrophytes could lead to improved water quality, such as through the removal of excess P. Accordingly, future manipulative studies should be carried out to complement the observational fieldwork to assess abiotic and biotic correlates with the occurrence, extent and cover for E. nuttallii, as well as other invasive macrophytes to enable strategic management practices. Such approaches could include controlled environmental studies to determine the direct and interacting effects of environmental variables found to influence invasion observations here and in other regions. In particular, an improved understanding of interactive effects between bivalves and macrophytes under a range different abiotic conditions is needed (Crane et al., 2020). To achieve this, basic laboratory mesocosm experiments can be scaled to the level of infestation observed at an invaded site (Coughlan et al., 2021), which will support cost-benefit assessments for management interventions. In addition, this paper presents data from surveys from a single system collected during summer months. To confirm that our findings are not specific to Lough Erne, studies incorporating a range of locations and seasonal assessments also should be conducted.

5 | CONCLUSIONS

We show that invasion of freshwater lakes by *E. nuttallii* is associated with a range of water quality variables, their change over time, the influence of surrounding land cover and use, as well as mutualism with other invasive species. It is evident that many abiotic and biotic factors influence the distribution and abundance of *E. nuttallii*. Accordingly, it is necessary to consider the total environment when planning management strategies. Increased nutrients have a considerable influence on the growth of *E. nuttallii*, therefore, reductions in these would be beneficial. *Elodea nuttallii* was positively associated with more rural locations with lower urbanisation, perhaps suggesting that it can consolidate its growth in locations less disturbed by river activities including boating or by utilising

the excess nutrient runoff from nearby farmland. Within this system, the removal of excess *E. nuttallii* biomass, with its accumulated P, could be a helpful alternative technique to other eutrophication control strategies. Improved management of facilitative invaders such as *D. polymorpha* also could be beneficial. Nevertheless, further assessments are required to elucidate the mechanisms that underpin invasion dynamics and nutrient cycling amongst these freshwater invaders.

AUTHOR CONTRIBUTIONS

Conceptualisation: KC, AR, HJM, JTAD, NR. Developing methods: KC, LK, NEC, RNC, JTAD, NR. Conducting the research: KC, NR. Data analysis: KC, NR. Preparation of figures and tables: KC, LK, NEC, RNC, NR. Data interpretation: KC, LK, NEC, RNC, NR. Writing: KC, LK, NEC, RNC, AR, HJM, JTAD, NR. Manuscript revision: KC, LK, NEC, RNC, NR. All authors give final approval for publication of this study. All authors agree to act as guarantors for the accuracy and integrity of this study.

ACKNOWLEDGMENTS

We particularly thankful to Michael Brown, Luke Taggart, Deirdre Spillane, Simon Exley, Emma Healey, Gillian Riddell, James Dickey, Elle O'Dea and Maurice Collins for their helpful contributions. We also thank the Northern Ireland Environment Agency (NIEA), Department of Agriculture, Australian Government, Environment and Rural Affairs (DAERA), Northern Ireland for their support and the contribution of long-term water monitoring dataset from their Water Management Unit (WMU).

DATA AVAILABILITY STATEMENT

The data are available from the authors upon reasonable request.

ORCID

Louise Kregting b https://orcid.org/0000-0001-7217-0146 Ross N. Cuthbert b https://orcid.org/0000-0003-2770-254X

REFERENCES

- Angelstein, S., & Schubert, H. (2008). Elodea nuttallii: Uptake, translocation and release of phosphorus. Aquatic Biology, 3, 209–216.
- Asaeda, T., Trung, V. K., Manatunge, J., & Bon, T. V. (2001). Modelling macrophyte-nutrient-phytoplankton interactions in shallow eutrophic lakes and the evaluation of environmental impacts. *Ecological Engineering*, 16, 341–357.
- Bailey, S. A., Brown, L., Campbell, M. L., Canning-Clode, J., Carlton, J. T., Castro, N., Chainho, P., Chan, F. T., Creed, J. C., Curd, A., Darling, J., Fofonoff, P., Galil, B. S., Hewitt, C. L., Inglis, G. J., Keith, I., Mandrak, N. E., Marchini, A., McKenzie, C. H., ... Zhan, A. (2020). Trends in the detection of aquatic non-indigenous species across global marine, estuarine and freshwater ecosystems: A 50-year perspective. *Diversity and Distributions*, 26, 1780–1797.
- Baldy, V., Thiébaut, G., Fernandez, C., Sagova-Mareckova, M., Korboulewsky, N., Monnier, Y., Perez, Tl., & Tremolieres, M. (2015). Experimental assessment of the water quality influence on the phosphorus uptake of an invasive aquatic plant: Biological responses throughout its phenological stage. *PLoS One*, 10, e0118844.

Freshwater Biology –WILEY

- Barko, J. W., & Smart, R. M. (1980). Mobilization of sediment phosphorus by submersed freshwater macrophytes. *Freshwater Biology*, 10, 229–238.
- Barko, J. W., & Smart, R. M. (1981). Comparative influences of light and temperature on the growth and metabolism of selected submersed freshwater macrophytes. *Ecological Monographs*, 51, 219–236.
- Barrat-Segretain, M. H., Elger, A., Sagnes, P., & Puijalon, S. (2002). Comparison of three life-history traits of invasive *Elodea canadensis* Michx. and *Elodea nuttallii* (planch.) H. St. John. *Aquatic Botany*, 74, 299-313.
- Bishop, P. L., & Eighmy, T. T. (1989). Aquatic wastewater treatment using Elodea nuttallii. Water Pollution Control Federation, 61, 641–648.
- Björklund, M. (2019). Be careful with your principal components. *Evolution*, 73(10), 2151–2158.
- Bodamer, B. L., & Ostrofsky, M. L. (2010). The use of aquatic plants by populations of the zebra mussel (*Dreissena polymorpha*) (Bivalvia: Dreissenidae) in a small glacial lake. *Nautilus*, 124, 100–106.
- Buscemi, P. A. (1958). Littoral oxygen depletion produced by a cover of *Elodea canadensis*. Oikos, 9, 239–245.
- Carignan, R., & Kalff, J. (1980). Phosphorus sources for aquatic weeds: Water or sediments? *Science*, 207, 987–989.
- Chambers, P. A., & Kalff, J. (1985). Depth distribution and biomass of submerged aquatic macrophyte communities in relation to Secchi depth. Canadian Journal of Fisheries and Aquatic Sciences, 42, 701–709.
- Champion, P. D., Clayton, J. S., & Hofstra, D. E. (2010). Nipping aquatic plant invasion in the bud: Weed risk assessment and the trade. *Hydrobiologia*, 656, 167–172.
- Cook, C. D. K., & Urmi-König, K. (1985). A revision of the genus *elodea* (Hyrdocharitacaea). Aquatic Botany, 21, 111–156.
- Coughlan, N. E., Cunningham, E. M., Cuthbert, R. N., Joyce, P. W. S., Anastácio, P., Banha, F., Bonel, N., Bradbeer, S. J., Briski, E., Butitta, V. L., Čadková, Z., Dick, J. T. A., Douda, K., Eagling, L. E., Ferreira-Rodríguez, N., Hünicken, L. A., Johansson, M. L., Kregting, L., Labecka, A. M., ... Sylvester, F. (2021). Biometric conversion factors as a unifying platform for comparative assessment of invasive freshwater bivalves. Journal of Applied Ecology, 58, 1945–1956. https://doi.org/10.1111/1365-2664.13941
- Coughlan, N. E., Cuthbert, R. N., & Dick, J. T. A. (2020). Aquatic biosecurity remains a damp squib. *Biodiversity and Conservation*, 29, 3091–3093.
- Coughlan, N. E., Cuthbert, R. N., Kelly, T. C., & Jansen, M. A. K. (2018). Parched plants: Survival and viability of invasive aquatic macrophytes following exposure to various desiccation regimes. *Aquatic Botany*, 150, 9–15.
- Crane, K., Coughlan, N. E., Cuthbert, R. N., Dick, J. T. A., Kregting, L., Ricciardi, A., MacIsaac, H. J., & Reid, N. (2020). Friends of mine: An invasive freshwater mussel facilitates growth of invasive macrophytes and mediates their competitive interactions. *Freshwater Biology*, 65, 1063–1072.
- Crane, K., Cuthbert, R. N., Dick, J. T. A., Kregting, L., Maclsaac, H. J., & Coughlan, N. E. (2019). Full steam ahead: Direct steam exposure to inhibit spread of invasive aquatic macrophytes. *Biological Invasions*, 21, 1311–1321. https://doi.org/10.1007/s10530-018-1901-2
- Cuthbert, R. N., Pattison, Z., Taylor, N. G., Verbrugge, L., Diagne, C., Ahmed, D. A., Leroy, B., Angulo, E., Briski, E., Capinha, C., Catford, J. A., Dalu, T., Essl, F., Gozlan, R. E., Haubrock, P. J., Kourantidou, M., Kramer, A. M., Renault, D., Wasserman, R. J., & Courchamp, F. (2021a). Global economic costs of aquatic invasive alien species. *Science of the Total Environment*, 775, 145238.
- Cuthbert, R.N., Diagne, C., Hudgins, E.J., Turbelin, A., Ahmed, D.A., Albert, C., Bodey, T.W., Briski, E., Essl, F., Haubrock, P.J., Gozlan, R.E., Kirichenko, N., Kourantidou, M., Kramer, A.M., Courchamp, F. (2021b). Biological invasion costs reveal insufficient proactive management worldwide. SSRN, pre-print. https://doi.org/10.2139/ ssrn.3993359

- DAERA (2018). Areas of special scientific interest. Department of Agriculture, environment and rural affairs website. Accessed 06/02/2018. www.daera-ni.gov.uk/topics/land-and-landscapes/ areas-special-scientific-interest.
- Dar, N. A., Pandit, A. K., & Ganai, B. A. (2014). Factors affecting the distribution patterns of aquatic macrophytes. *Limnological Review*, 14, 75–81.
- Dendéne, M. A., Rolland, T., Trémolières, M., & Carbiener, R. (1993). Effect of ammonium ions on the net photosynthesis of three species of *elodea*. Aquatic Botany, 46, 301–315.
- Dick, J. T. A., Laverty, C., Lennon, J. J., Barrios-O'Neill, D., Mensink, P. J., et al. (2017). Invader relative impact potential: A new metric to understand and predict the ecological impacts of existing, emerging and future invasive alien species. *Journal of Applied Ecology*, 54, 1259–1267.
- Dickey, J. W. E., Coughlan, N. E., Dick, J. T. A., Médoc, V., McCard, M., Leavitt, P. R., Lacroix, G., Fiorini, S., Millot, A., & Cuthbert, R. N. (2021). Breathing space: Deoxygenation of aquatic environments can drive differential ecological impacts across biological invasion stages. *Biological Invasions*, 23, 2831–2847. https://doi. org/10.1007/s10530-021-02542-3
- DOE (1997) Upper Lough Erne Ramsar site. Report published by the Department of Environment, Environment and Heritage Service. Downloaded from https://www.daera-ni.gov.uk/ sites/default/files/publications/daera/ULE%20Citation%20and %20Map.pdf
- Franklin, S. B., Gibson, D. J., Robertson, P. A., Pohlmann, J. T., & Fralish, J. S. (1995). Parallel analysis: A method for determining significant principal components. *Journal of Vegetation Science*, 6(1), 99–106.
- Frankouich, T. A., Gainer, E. E., Zieman, J. C., & Wachnick, A. H. (2006). Spatial and temporal distribution of epiphytic diatoms growing on *thalassic testudinum* banks ex Konigh. Relationship to waters quality. *Hydrobiology*, 560, 259–271.
- Gardner, W. S., Cavaletto, J. F., Johengen, T. H., Johnson, J. R., Heath, R. T., & Cotner, J. B. Jr (1995). Effects of the zebra mussel, Dreissena polymorpha, on community nitrogen dynamics in Saginaw Bay, Lake Huron. Journal of Great Lakes Research, 21, 529–544. https:// doi.org/10.1016/S0380-1330(95)71064-4
- Haubrock, P. J., Turbelin, A. J., Cuthbert, R. N., Novoa, A., Taylor, N. G., Angulo, E., Ballesteros-Mejia, L., Bodey, T. W., Capinha, C., Diagne, C., Essl, F., Golivets, M., Kirichenko, N., Kourantidou, M., Leroy, B., Renault, D., Verbrugge, L., & Courchamp, F. (2021). Economic costs of invasive alien species across Europe. *NeoBiota*, *67*, 153–190.
- Higgins, T. M., Grennan, J. M., & McCarthy, T. K. (2008). Effects of recent zebra mussel invasion on water chemistry and phytoplankton production in a small Irish lake. *Aquatic Invasions*, *3*, 14–20.
- Horvath, T. G., & Lamberti, G. A. (1997). Drifting macrophytes as a mechanism for zebra mussel (*Dreissena polymorpha*) invasion of lakeoutlet streams. *American Midland Naturalist*, 138, 29–36.
- Hussner, A., Stiers, I., Verhofstad, M. J. J. M., Bakker, E. S., Grutters, B. M. C., Haury, J., van Valkenburg, J. L. C. H., Brundu, G., Newman, J., Clayton, J. S., Anderson, L. W. J., & Hofstra, D. (2017). Management and control methods of invasive alien freshwater aquatic plants: A review. Aquatic Botany, 136, 112–137.
- James, C. S., Eaton, J. W., & Hardwick, K. (1999). Competition between three submerged macrophytes, *Elodea canadensis* Michx, *Elodea nuttallii* (planch.) St John and *Lagarosiphon major* (Ridl.) Moss. *Hydrobiologia*, 415, 35-40.
- Jewell, W. J. (1971). Aquatic weed decay: Dissolved oxygen utilization and nitrogen and phosphorus regeneration. Water Pollution Control Federation, 43, 1457–1467.
- JNCC (2015). Upper Lough Erne special areas of conservation. Standard data form by the Department of Environment, Environment and Heritage Service. Downloaded from http://jncc.defra.gov.uk/Prote ctedSites/SACselection/n2kforms/UK0016614.pdf

ILEY- Freshwater Biology

- Jones, J. I., Eaton, J. W., & Keith Hardwick, K. (2000). The effect of changing environmental variables in the surrounding water on the physiology of *Elodea nuttallii*. *Aquatic Botany*, *66*, 115–129.
- Kaiser, H. F. (1960). The application of electronic computers to factor analysis. Educational and Psychological Measurement, 20, 141–151.
- Kelly, R., Harrod, C., Maggs, C. A., & Reid, N. (2015). Effects of *Elodea* nuttallii on temperate freshwater plants, microalgae and invertebrates: Small differences between invaded and uninvaded areas. *Biological Invasions*, 17, 2123–2138.
- Kelly, R., Lundy, M. G., Mineur, F., Harrod, C., Maggs, C. A., Humphries, N. E., Sims, D. W., & Reid, N. (2014). Historical data reveal powerlaw dispersal patterns of invasive aquatic species. *Ecography*, *37*, 581–590.
- Lawrie, E.W., Wolfe-Murphy, S.A., Gibson, C.E. (1992). Large lakes A botanical survey of the eight largest lakes in Northern Ireland. Section 7. A report by the Northern Ireland Lakes Survey. Countryside and Wildlife, Department of Environment (NI) Report.
- Lougheed, V. L., Crosbie, B., & Chow-Fraser, P. (2001). Primary determinants of macrophyte community structure in 62 marshes across the Great Lakes basin: Latitude, land use, and water quality effects. *Canadian Journal of Fisheries and Aquatic Sciences*, 58, 1603–1612.
- Maberly, S. C., & Madsen, T. V. (2002). Freshwater angiosperm carbon concentrating mechanisms: Processes and patterns. *Functional Plant Biology*, 29, 393–405.
- MacIsaac, H. J. (1996). Potential abiotic and biotic impacts of zebra mussels on the inland waters of North America. *American Zoologist*, *36*, 287–299.
- Maguire, C., & Gibson, C. (2005). Ecological change in Lough Erne: Influence of catchment changes and species invasions. *Freshwater Forum*, 24, 38–58.
- Melzer, A. (1999). Aquatic macrophytes as tools for lake management. Hydrobiologia, 395(396), 181–190.
- O'Higgins, T. (2018). Management and impact of invasive alien species (IAS) in Lough Erne in Ireland. Aquacross D9.2 case study 4 report.
- Oreska, M. P. J., & Aldridge, D. C. (2011). Estimating the financial costs of freshwater invasive species in Great Britain: A standardized approach to invasive species costing. *Biological Invasions*, 13, 305–319.
- Ozimek, T., Van Donk, E., & Gulati, R. D. (1993). Growth and nutrient uptake by two species of *elodea* in experimental conditions and their role in nutrient accumulation in a macrophyte dominated lake. *Hydrobiologia*, 251, 13–18.
- Pereira, A., Tassin, B., & Jörgensen, S. E. (1994). A model for decomposition of the down vegetation in an Amazonian reservoir. *Ecological Modelling*, 75(76), 447–458.
- Ricciardi, A. (2003). Predicting the impacts of an introduced species from its invasion history: An empirical approach applied to zebra mussel invasions. *Freshwater Biology*, 48, 972–981.
- Ricciardi, A. (2015). Ecology of invasive alien invertebrates. In J. Thorp & D. C. Rogers (Eds.), *Thorp and Covich's freshwater invertebrates* (pp. 83–91). Elsevier.
- Ricciardi, A., Neves, R. J., & Rasmussen, J. B. (1998). Impending extinctions of north American freshwater mussels (Unionoida) following the zebra mussel (Dreissena polymorpha) invasion. Journal of Animal Ecology, 67, 613–619.
- Ricciardi, A., & MacIsaac, H. J. (2011). Impacts of biological invasions on freshwater ecosystems. In D. M. Richardson (Ed.), Fifty years of invasion ecology: The legacy of Charles Elton (pp. 211–224). Wiley-Blackwell.
- Rolland, T., & Trémolières, M. (1995). The role of ammonium nitrogen in the distribution of two species of *elodea*. *Acta Botanica Gallica*, 142, 733–739.
- Rosell, R. S., Maguire, C. M., & McCarthy, T. K. (1999). First reported settlement of zebra mussels *Dreissena polymorpha* in the Erne system, Co. Fermanagh, Northern Ireland. *Biology and Environment*, 98, 191–193.

- Sass, L. L., Bozek, M. A., Hauxwell, J. A., Wagner, K., & Knight, S. (2010). Response of aquatic macrophytes to human land use perturbations in the watersheds of Wisconsin lakes, U.S.A. Aquatic Botany, 93, 1–8.
- Seebens, H., Blackburn, T. M., Dyer, E. E., Genovesi, P., Hulme, P. E., Jeschke, J. M., Pagad, S., Pyšek, P., Winter, M., Arianoutsou, M., Bacher, S., Blasius, B., Brundu, G., Capinha, C., Celesti-Grapow, L., Dawson, W., Dullinger, S., Fuentes, N., Jäger, H., ... Essl, F. (2017). No saturation in the accumulation of alien species worldwide. *Nature Communications*, 8, 1–9.
- Seebens, H., Bacher, S., Blackburn, T. M., Capinha, C., Dawson, W., Dullinger, S., Genovesi, P., Hulme, P. E., van Kleunen, M., Kühn, I., & Jeschke, J. M. (2021). Projecting the continental accumulation of alien species through to 2050. *Global Change Biology*, 27(5), 970–982.
- Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D. A., Aronson, J., Courchamp, F., Galil, B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E., & Vilà, M. (2013). Impacts of biological invasions: what's what and the way forward. *Trends in Ecology and Evolution*, 28, 58-66.
- Simpson, D. A. (1990). Displacement of Elodea canadensis Michx by Elodea nuttallii (planch.) H. St John in the British Isles. Watsonia, 18, 173-177.
- Sousa, R., Novais, A., & Strayer, D. L. (2014). Invasive bivalves in fresh waters: Impacts from individuals to ecosystems and possible control strategies. *Hydrobiologia*, 735, 231-251.
- Szabó, S., Peeters, E. T. H. M., Várbíró, G., Borics, G., & Lukács, B. A. (2018). Phenotypic plasticity as a clue for invasion success of the submerged aquatic plant *Elodea nuttallii*. *Plant Biology*, 21, 54–63.
- Thiébaut, G., & Muller, S. (1999). A macrophyte communities sequence as an indicator of eutrophication and acidification levels in weakly mineralised streams in North-Eastern France. *Hydrobiologia*, 410, 17–24.
- Thouvenot, L., & Thiébaut, G. (2018). Regeneration and colonization abilities of the invasive species Elodea canadensis and Elodea nuttallii under a salt gradient: implications for freshwater invasibility. *Hydrobiologia*, 817, 193–203. https://doi.org/10.1007/s1075 0-018-3576-1
- van Donk, E., Gulati, R. D., ledema, A., & Meulemans, J. T. (1993). Macrophyte-related shifts in the nitrogen and phosphorus contents of the different trophic levels in a biomanipulated shallow lake. *Hydrobiologia*, 251, 19-26.
- Ward, J. M., & Ricciardi, A. (2007). Impacts of *Dreissena* invasions on benthic macroinvertebrate communities: A meta-analysis. *Diversity and Distributions*, 13, 155–165.
- Zehnsdorf, A., Hussner, A., Eismann, F., Rönicke, H., & Melzer, A. (2015). Management options of invasive *Elodea nuttallii* and *Elodea canadensis*. *Limnologica*, 51, 110–111.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Crane, K., Kregting, L., Coughlan, N. E., Cuthbert, R. N., Ricciardi, A., MacIsaac, H. J., Dick, J. T., & Reid, N. (2022). Abiotic and biotic correlates of the occurrence, extent and cover of invasive aquatic *Elodea nuttallii*. *Freshwater Biology*, *67*, 1559–1570. https://doi.org/10.1111/fwb.13960