



**QUEEN'S
UNIVERSITY
BELFAST**

Serum Fatty Acid Binding Protein 4 (FABP4) Predicts Pre-eclampsia in Women with Type 1 Diabetes

Wotherspoon, A. C., Young, I. S., McCance, D. R., Patterson, C. C., Maresh, M. J. A., Pearson, D. W. M., Walker, J. D., & Holmes, V. A. (2016). Serum Fatty Acid Binding Protein 4 (FABP4) Predicts Pre-eclampsia in Women with Type 1 Diabetes. *Diabetes Care*, 39(10), 1827-1829. <https://doi.org/10.2337/dc16-0803>

Published in:
Diabetes Care

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2016 by the American Diabetes Association.

This is an author-created, uncopyedited electronic version of an article accepted for publication in *Diabetes Care*. The American Diabetes Association (ADA), publisher of *Diabetes Care*, is not responsible for any errors or omissions in this version of the manuscript or any version derived from it by third parties. The definitive publisher-authenticated version will be available in a future issue of *Diabetes Care* in print and online at <http://care.diabetesjournals.org/content/39/10/1827>

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Serum Fatty Acid Binding Protein 4 (FABP4) Predicts Pre-eclampsia in Women with Type 1 Diabetes

Running Title: FABP4 Predicts Pre-eclampsia

Authors: Amy C Wotherspoon¹, Ian S Young¹, David R McCance², Chris C Patterson¹, Michael J.A. Maresh (MD),³ Donald W.M. Pearson (MD),⁴ James D. Walker (MD),⁵ Valerie A Holmes¹, for the Diabetes and Pre-eclampsia Intervention Trial (DAPIT) Study Group

¹Centre for Public Health, Queen's University Belfast, U.K.

²Regional Centre for Endocrinology and Diabetes, Royal Victoria Hospital, Belfast, U.K.

³Department of Obstetrics, St Mary's Hospital, Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK;

⁴Department of Diabetes, Aberdeen Royal Infirmary, Aberdeen, UK;

⁵Department of Diabetes, St John's Hospital at Howden, West Lothian, UK

Corresponding Author: Valerie A Holmes

Address: Centre for Public Health, Queen's University Belfast, Institute of Clinical Sciences Block B, Royal Victoria Hospital, Grosvenor Road, Belfast, BT12 6BA.

Word Count: 1,346 including abstract subheadings (Limit 1,350)

Number of Tables: 1

Number of Figures: 0

Abstract

Objective

To examine the association between Fatty Acid Binding Protein 4 (FABP4) and pre-eclampsia risk in women with type 1 diabetes.

Research Design and Methods

Serum FABP4 was measured in 710 women from the DAPIT study in early pregnancy and in the second trimester (median 14 and 26 weeks gestation, respectively).

Results

FABP4 was significantly elevated in early pregnancy (geometric mean [interquartile range]: 15.8ng/ml [11.6-21.4] vs. 12.7ng/ml [9.6-17]; $P<0.001$) and the second trimester (18.8ng/ml [13.6-25.8] vs. 14.6ng/ml [10.8-19.7]; $P<0.001$) in women who later developed pre-eclampsia. Elevated second trimester FABP4 was independently associated with pre-eclampsia (OR 2.87 (95%CI 1.24, 6.68), $P=0.03$). Addition of FABP4 to established risk factors significantly improved Net Reclassification Improvement at both time-points and Integrated Discrimination Improvement in the second trimester.

Conclusions

Increased second trimester FABP4 independently predicted pre-eclampsia and significantly improved reclassification and discrimination. FABP4 shows potential as a novel biomarker for pre-eclampsia prediction in women with type 1 diabetes.

Word Count: 153 including subheadings (limit 150)

Introduction

Pre-eclampsia, defined as new onset hypertension and proteinuria occurring after 20 weeks gestation, is associated with significant perinatal morbidity and mortality(1, 2).

Women with type 1 diabetes are 2 to 4 more times more likely develop pre-eclampsia than the background population(3, 4).

Fatty Acid Binding Protein 4 (FABP4), or adipocyte fatty acid binding protein, is an intracellular lipid chaperone involved in co-ordination of lipid transportation(5).

FABP4 is expressed mainly in adipocytes and can be released into the circulation(6).

In the non-pregnant state FABP4 is associated with known pre-eclampsia risk factors: obesity, hypertension and diabetes(6, 7).

Several studies have reported elevated FABP4 in women with pre-eclampsia(8, 9) or who later developed pre-eclampsia(10), compared to those who did not, although all studies excluded women with diabetes.

Our objective was to examine the role of FABP4 as a potential biomarker for pre-eclampsia alone, and in tandem with established clinical risk factors, in women with type 1 diabetes.

Research design and Methods

The population comprised 710 of the 762 women from the Diabetes and Pre-eclampsia Intervention Trial (DAPIT), a randomized multicenter double-blind placebo-controlled trial investigating vitamin C and E supplementation and risk of pre-eclampsia in women with type 1 diabetes(11). Pre-eclampsia, the primary endpoint, was defined using International guidelines(1, 12). Women provided written

informed consent. West-Midlands Multicenter Research Ethics Committee approved the study (MREC 02/7/016).

Blood samples were collected at randomization (8-22 weeks; median 14 weeks) and in the second trimester (median 26 weeks). Serum samples were batch-analyzed, blind to pre-eclampsia status, using a commercially available ELISA (Biovendor, Modrice, Czech Republic). Inter-assay and intra-assay CVs were 8.3% and 3.8%, respectively.

Intervention and control groups were analyzed together as supplementation did not reduce pre-eclampsia risk (risk ratio: 0.81; 95% CI: 0.59, 1.12, $P=0.20$)(11); FABP4 levels were not different between groups, and the effect of supplementation on pre-eclampsia risk did not depend on FABP4.

Statistical analysis

FABP4 concentrations were positively skewed and therefore logarithmically transformed. Values are reported as geometric mean and interquartile range (IQR). Chi-squared (χ^2) and independent samples t -tests were used for group comparisons. Logistic regression analysis investigated the association between FABP4 and pre-eclampsia before and after adjusting for the following risk factors: age, BMI, gestational age, blood pressure, parity, duration of diabetes, smoking status, history of pre-eclampsia, HbA_{1c}, renal status, and DAPIT treatment group. The area under the receiver operating characteristic curve (AUROC) was used to assess the predictive value of FABP4 for pre-eclampsia(13). Integrated Discrimination Improvement (IDI) and Net Reclassification Improvement (NRI) indices were calculated to determine the clinical utility of the addition of FABP4 to established risk factors and the ability of FABP4 to improve pre-eclampsia prediction (for calculation

AUROC, NRI and IDI, see Supplementary Data)(14). Statistical analysis was performed using SPSS version 20 (IBM Corp., Armonk, NY), Stata release 14 (StataCorp, College Station, TX) and the Hmisc package in R version 3.1.3 (R Core Team, Vienna, Austria).

Results

Serum from one or both time-points was available for 710 women, of whom 120 (17%) developed pre-eclampsia (for maternal and clinical characteristics see Supplementary Table 1).

FABP4 levels at randomization (median 14 weeks) and 26 weeks were significantly higher in women who later developed pre-eclampsia compared with those who did not (Table 1). Of the 319 women with blood samples at 13 weeks or less, FABP4 levels were elevated in those who later developed pre-eclampsia compared to those who did not ($P=0.02$).

For each doubling of serum FABP4, adjusted risk of pre-eclampsia increased by 40% (OR 1.4 (95%CI 1.0, 2.0), $P=0.031$) and 60% (1.6 (1.1, 2.3), $P=0.017$) at 14 and 26 weeks, respectively. The independent ability of FABP4 to predict preeclampsia, in relation to quarters of FABP4 in combination with established risk factors, is shown in Supplementary Table 2. In the second trimester, the highest quarter of FABP4 was a significant independent predictor of pre-eclampsia compared with the lowest (OR 2.87; 95%CI 1.24, 6.68), $P=0.03$).

AUROC for FABP4 alone were 0.622 and 0.646 at 14 and 26 weeks, respectively (Table 1). When FABP4 was added to the model containing established risk factors,

AUROC were 0.801 and 0.825 at 14 and 26 weeks, respectively (both non-significant from established risk factors alone).

The NRI statistic showed that the addition of FABP4 to established risk factors significantly increased correct reclassification of cases and non-cases at both 14 and 26 weeks. The IDI statistic showed that FABP4 significantly increased discrimination between cases and non-cases in the second trimester (Table 1).

Conclusions

We believe this is the first study to date to investigate FABP4 levels in relation to development of pre-eclampsia and to investigate its clinical utility in pregnant women with type 1 diabetes. Maternal FABP4 levels were significantly elevated in women who later developed pre-eclampsia compared with those who did not both at 14 and 26 weeks gestation. Furthermore, the addition of FABP4 to established risk factors improved the reclassification index at both time-points and discrimination index in the second trimester.

Only one previous study has measured FABP4 prior to the onset of pre-eclampsia. In line with our findings, Scifres and colleagues(10) reported significantly elevated FABP4 before 13 weeks and at 26 weeks gestation in non-diabetic women who later developed pre-eclampsia.

FABP4 may play a role in pre-eclampsia development. Shangguan and colleagues(9) reported increased third trimester FABP4 levels in women with pre-eclampsia compared with healthy pregnant and non-pregnant women, suggesting no effect of pregnancy on FABP4, thus implicating FABP4 in pre-eclampsia development.

This study explores the both predictive properties and clinical utility of FABP4 in relation to pre-eclampsia in women with type 1 diabetes. FABP4 remained significantly associated with pre-eclampsia after controlling for established risk factors. FABP4, when added to established risk factors, did not significantly increase the AUROC. However, the NRI and IDI indices, which are proposed as superior methods to determine clinical utility of a biomarker(14), showed that FABP4, in addition to clinical risk factors, improved the prediction of pre-eclampsia. This was significant for NRI at 14 weeks and for both NRI and IDI at 26 weeks.

The strengths of this study include the study size, with DAPIT being the largest prospective dataset reported of thoroughly characterized women with type 1 diabetes, together with the strict definition of pre-eclampsia, as each potential case was reviewed independently by three clinicians(11, 15). In addition, women with essential hypertension or any existing renal disease were included, making our study more representative of the typical population. The study has the limitation that trial participants may not be entirely representative of all pregnant women with type 1 diabetes, as they were self-selecting. Randomization samples also represented a range of gestational ages from 8 weeks to 22 weeks.

In summary, we believe this is the first study to investigate FABP4 in relation to the development of pre-eclampsia in pregnant women with type 1 diabetes. FABP4 levels in both the first and second trimester were significantly associated with the development of pre-eclampsia. Elevated second trimester FABP4 independently predicted pre-eclampsia, significantly improving reclassification and discrimination. While further studies are now needed to replicate these findings, our data suggest that FABP4 shows potential as a novel biomarker for pre-eclampsia prediction in women with type 1 diabetes.

Acknowledgements

This study was funded by grants 067028/Z/02/Z and 083145/Z/07/Z from The Wellcome Trust (registered charity number 210183) and from the Department of Employment and Learning, Northern Ireland. A.C.W. carried out laboratory analysis, researched and wrote the manuscript. I.S.Y., D.R.M., C.C.P., M.J.A.M., D.W.M.P, J.D.W and V.A.H reviewed and edited the manuscript. C.C.P. also supervised the statistical analysis. No potential conflicts of interest relevant to this article were reported. V.A.H. is the guarantor of this work and as such, had full access to all the data and the study and takes responsibility for the integrity of the data and accuracy of the data analysis. Parts of this study have been accepted for presentation at the 76th Scientific Sessions of the American Diabetes Association, New Orleans, 10-14th June 2016.

The authors thank Cyril McMaster from the Centre for Public Health, Queen's University Belfast, United Kingdom, for his assistance with laboratory analysis, the patients who took part in the DAPIT study, the DAPIT research midwives who collected the data and the collaborators at each center.

References

1. Brown MA, Lindheimer MD, de Swiet M, Assche AV, Moutquin J. The classification and diagnosis of the hypertensive disorders of pregnancy: statement from the International Society for the Study of Hypertension in Pregnancy (ISSHP). *Hypertension in Pregnancy*. 2001;20(1):ix-xiv.
2. Roberts JM, Gammill HS. Preeclampsia: recent insights. *Hypertension*. 2005;46(6):1243-9.
3. Persson M, Norman M, Hanson U. Obstetric and Perinatal Outcomes in Type 1 Diabetic Pregnancies A large, population-based study. *Diabetes Care*. 2009;32(11):2005-9.
4. Jensen DM, Damm P, Moelsted-Pedersen L, Ovesen P, Westergaard JG, Moeller M, et al. Outcomes in Type 1 Diabetic Pregnancies A nationwide, population-based study. *Diabetes Care*. 2004;27(12):2819-23.
5. Furuhashi M, Ishimura S, Ota H, Miura T. Lipid chaperones and metabolic inflammation. *Int J Inflam*. 2011;2011:642612.
6. Xu A, Wang Y, Xu JY, Stejskal D, Tam S, Zhang J, et al. Adipocyte fatty acid-binding protein is a plasma biomarker closely associated with obesity and metabolic syndrome. *Clin Chem*. 2006;52(3):405-13.
7. Yamada M, Mochizuki K, Honma K, Miyauchi R, Kasezawa N, Tohyama K, et al. Serum Fatty Acid Binding Protein 4 Concentrations Are Positively and Independently Associated with Blood Pressure and Abdominal Fat among Parameters in Health Check-Ups in Ordinary Middle-Aged Japanese Males. *J Nutr Sci Vitaminol (Tokyo)*. 2015;61(4):291-8.
8. Fasshauer M, Seeger J, Waldeyer T, Schrey S, Ebert T, Kratzsch J, et al. Serum levels of the adipokine adipocyte fatty acid-binding protein are increased in preeclampsia. *Am J Hypertens*. 2008;21(5):582-6.
9. Shangguan X, Liu F, Wang H, He J, Dong M. Alterations in serum adipocyte fatty acid binding protein and retinol binding protein-4 in normal pregnancy and preeclampsia. *Clinica Chimica Acta*. 2009;407(1):58-61.
10. Scifres CM, Catov JM, Simhan H. Maternal serum fatty acid binding protein 4 (FABP4) and the development of preeclampsia. *J Clin Endocrinol Metab*. 2011;97(3):E349-56.
11. McCance DR, Holmes VA, Maresh MJ, Patterson CC, Walker JD, Pearson DW, Young IS. Diabetes and Pre-eclampsia Intervention Trial (DAPIT) Study Group. Vitamins C and E for prevention of pre-eclampsia in women with type 1 diabetes (DAPIT): a randomised placebo-controlled trial. *Lancet*. 2010;376(9737):259-66.
12. National High Blood Pressure Education Program Working Group on High Blood Pressure in Pregnancy. Report of the National High Blood Pressure Education

Program Working Group on High Blood Pressure in Pregnancy. *Obstet Gynecol.* 2000 7;183(1):s1-s22.

13. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics.* 1988:837-45.

14. Pencina MJ, D'Agostino RB, Vasan RS. Evaluating the added predictive ability of a new marker: from area under the ROC curve to reclassification and beyond. *Stat Med.* 2008;27(2):157-72.

15. Holmes VA, Young IS, Patterson CC, Maresh MJ, Pearson DW, Walker JD, et al. The Role of Angiogenic and Antiangiogenic Factors in the Second Trimester in the Prediction of Preeclampsia in Pregnant Women With Type 1 Diabetes. *Diabetes Care.* 2013;36(11):3671-7.

Table 1:

Serum FABP4 concentrations at randomization and in the second trimester, with area under the receiver operating characteristic curve (AUROC), Net Reclassification Improvement (NRI) and Integrated Discrimination Improvement (IDI) analyses

	Maternal serum FABP4 concentration* (ng/ml)		Area under receiver operating characteristic curve (AUROC)				NRI (<i>P</i>)	IDI (<i>P</i>)
	Pre-eclampsia	No pre-eclampsia	FABP4 alone	Established risk factors without FABP4	Established risk factors with FABP4	Incremental area (<i>P</i>) [†]		
Randomization	15.8 (11.6-21.4)	12.7 (9.6-17.0)	0.622	0.793	0.801	0.008 (0.17)	0.306 (0.003)	0.009 (0.11)
Second trimester	18.8 (13.6-25.8)	14.6 (10.8-19.7)	0.648	0.819	0.825	0.006 (0.34)	0.251 (0.02)	0.012 (0.048)

*Reported as geometric mean and interquartile range

[†] Comparison of AUROCs: Established risk factors without FABP4 vs Established risk factors with FABP4