



**QUEEN'S
UNIVERSITY
BELFAST**

Estimating quantum chromatic numbers

Paulsen, V. I., Severini, S., Stahlke, D., Todorov, I. G., & Winter, A. (2016). Estimating quantum chromatic numbers. *Journal of Functional Analysis*, 270(6), 2180-2222. <https://doi.org/10.1016/j.jfa.2016.01.010>

Published in:
Journal of Functional Analysis

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2016 Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/> which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

ESTIMATING QUANTUM CHROMATIC NUMBERS

VERN I. PAULSEN, SIMONE SEVERINI, DANIEL STAHLKE, IVAN G. TODOROV,
AND ANDREAS WINTER

ABSTRACT. We develop further the new versions of quantum chromatic numbers of graphs introduced by the first and fourth authors. We prove that the problem of computation of the commuting quantum chromatic number of a graph is solvable by an SDP algorithm and describe an hierarchy of variants of the commuting quantum chromatic number which converge to it. We introduce the tracial rank of a graph, a parameter that gives a lower bound for the commuting quantum chromatic number and parallels the projective rank, and prove that it is multiplicative. We describe the tracial rank, the projective rank and the fractional chromatic numbers in a unified manner that clarifies their connection with the commuting quantum chromatic number, the quantum chromatic number and the classical chromatic number, respectively. Finally, we present a new SDP algorithm that yields a parameter larger than the Lovász number and is yet a lower bound for the tracial rank of the graph. We determine the precise value of the tracial rank of an odd cycle.

1. INTRODUCTION

We assume that the reader is familiar with some concepts from graph theory and refer the reader to the text [13] for any terminology that we do not explain.

In [12, 1, 3, 27] the concept of the quantum chromatic number $\chi_q(G)$ of a graph G was developed and inequalities for estimating this parameter, as well as methods for its computation, were presented. In [24] several new variants of the quantum chromatic number, denoted $\chi_{qc}(G)$, $\chi_{qa}(G)$ and $\chi_{qs}(G)$, were introduced, as well as $\chi_{\text{vect}}(G)$. The motivation behind them came from conjectures of Tsirelson and Connes and the fact that the set of correlations of quantum experiments may possibly depend on which set of quantum mechanical axioms one chooses to employ. Given a graph G , the

Date: 10 September 2015.

2010 Mathematics Subject Classification. Primary 46L07, 05C15; Secondary 47L25, 81R15.

Key words and phrases. operator system, tensor product, chromatic number.

This work supported in part by NSF (USA), EPSRC (United Kingdom), the Royal Society, the European Commission (STREP “RAQUEL”), the ERC (Advanced Grant “IRQUAT”), the Spanish MINECO (project FIS2008-01236, with the support of FEDER funds), and the Isaac Newton Institute for Mathematical Sciences during the semester *Mathematical Challenges in Quantum Information*, Aug-Dec 2014.

aforementioned chromatic numbers satisfy the inequalities

$$\chi_{\text{vect}}(G) \leq \chi_{\text{qc}}(G) \leq \chi_{\text{qa}}(G) \leq \chi_{\text{qs}}(G) \leq \chi_{\text{q}}(G) \leq \chi(G),$$

where $\chi(G)$ denotes the classical chromatic number of the graph G .

The motivation of [24] for defining and studying these new chromatic numbers comes from the fact that if Tsirelson’s conjecture is true, then $\chi_{\text{qc}}(G) = \chi_{\text{q}}(G)$ for every graph G , while if Connes’ Embedding Problem has an affirmative answer, then $\chi_{\text{qc}}(G) = \chi_{\text{qa}}(G)$ for every graph G . Thus, computing these invariants gives a means to test the corresponding conjectures.

In [5] it was shown that

$$\lceil \vartheta^+(\overline{G}) \rceil = \chi_{\text{vect}}(G),$$

where $\lceil r \rceil$ denotes the least integer greater than or equal to r and ϑ^+ is Szegedy’s [30] variant of Lovász’s [20] ϑ -function. Furthermore, this identity was used to give the first example of a graph for which $\chi_{\text{vect}}(G) \neq \chi_{\text{q}}(G)$. Also, since ϑ^+ is defined by an SDP, the aforementioned result shows that $\chi_{\text{vect}}(G)$ is computable by an SDP.

In this paper we show that for each size of graph, $\chi_{\text{qc}}(G)$ is also computable by an SDP. Our proof builds on ideas borrowed from the “NPA hierarchy” exhibited in [21]. It uses a compactness argument to show that for the purposes of computing this integer the hierarchy terminates, but does not yield the stage at which it does so. Thus, while we can say that it is computable by one of the SDP’s in the hierarchy, we cannot explicitly determine the size of this SDP. It is known that $\chi(G)$ is computable by an SDP, but it is still not known if the same is true for $\chi_{\text{qa}}(G)$, $\chi_{\text{qs}}(G)$ and $\chi_{\text{q}}(G)$. If the Tsirelson and Connes conjectures hold true then these three quantities must also be computable by SDP’s as they then are all equal to $\chi_{\text{qc}}(G)$.

D. Roberson and L. Mančinska [26] introduced a Hilbert space variant of the fractional chromatic number $\chi_f(G)$ of a graph G , called the projective rank, and denoted $\xi_f(G)$. They proved that $\xi_f(G)$ is a lower bound for $\chi_{\text{q}}(G)$; this estimate was critical for identifying a graph G with $\chi_{\text{vect}}(G) \neq \chi_{\text{q}}(G)$. However, it is still not known if $\xi_f(G)$ is a lower bound for the variants of the quantum chromatic number studied in [24].

In the present paper, we introduce a new variant of the projective rank, which we call *tracial rank* and show that it is a lower bound for $\chi_{\text{qc}}(G)$. We also give a new interpretation of the projective rank in terms of traces on finite dimensional C*-algebras. These parameters and their properties allow us to give the first example of a graph for which $\chi_{\text{vect}}(G) \neq \chi_{\text{qc}}(G)$.

Finally, we present a new SDP that yields a parameter larger than Szegedy’s bound (i.e., it is an *SDP relaxation* of the latter), and which is still a lower bound on $\chi_{\text{qc}}(G)$. En route, we show that our tracial rank is multiplicative.

To put our work into a broader context, recently a number of graph parameters, including clique, chromatic, and independence numbers, have

been generalized to “quantum” versions by relating the respective number to attaining maximum probability 1 in a so-called non-local game. The present work shows that some of the techniques to bound and separate these numbers, originally developed for the exact, finite-dimensional and tensor product case, can be extended to the approximate and relativistic setting, and that they can be bounded by SDP’s without going directly to the NPA hierarchy.

2. QUANTUM CORRELATIONS AND VARIANTS OF THE QUANTUM CHROMATIC NUMBER

In this section we summarize the properties of some of the variants of the quantum chromatic number introduced in [24] from the viewpoint of correlations and derive a few additional properties of the corresponding models that will be essential later. We introduce a new notation that we hope serves to clarify and unify many of the ideas from [24].

Let $G = (V, E)$ be a graph; here, $V = V(G)$ is the set of vertices, while $E = E(G)$ is the set of edges, of G . If $(v, w) \in E$, we write $v \sim w$. In [1] and [3], the authors considered a “graph colouring game”, where two players, Alice and Bob, try to convince a referee that they have a colouring of G ; the referee inputs a pair (v, w) of vertices of G , and each of the players produces an output, according to a previously agreed “quantum strategy”, that is, a probability distribution derived from an entangled state and collections of positive operator-valued measures, POVM’s. To formalise this, recall that a POVM is a collection $(E_i)_{i=1}^k$ of positive operators acting on a Hilbert space H with $\sum_{i=1}^k E_i = I$ (as usual, here we denote by I the identity operator). If, in addition, each E_i is a projection, then the collection is called a projection-valued measure, or PVM. When $H = \mathbb{C}^p$ is finite dimensional, we identify the operators on H with elements of the algebra M_p of all $p \times p$ complex matrices. Given POVM’s $(E_{v,i})_{i=1}^c \subseteq M_p$ and $(F_{w,j})_{j=1}^c \subseteq M_q$, where $v, w \in V$, and a unit vector $\xi \in \mathbb{C}^p \otimes \mathbb{C}^q$, one associates with each pair (v, w) of vertices of G the probability distribution

$$(1) \quad p(i, j|v, w) = \langle (E_{v,i} \otimes F_{w,j})\xi, \xi \rangle, \quad 1 \leq i, j \leq c,$$

where, for an input (v, w) from the referee, $\langle (E_{v,i} \otimes F_{w,j})\xi, \xi \rangle$ is the probability for Alice producing an output i and Bob – an output j .

The set of all $nc \times nc$ matrices that are obtained by allowing p and q to vary through all the natural numbers, is called the set of *quantum correlations* and is generally denoted $Q(n, c)$; when n and c are clear from the context, we simply write Q . For reasons that will be clear shortly, we shall adopt the notation: $C_q(n, c) = Q(n, c)$.

Alice and Bob can thus convince the referee that they have a c -colouring if the following conditions are satisfied:

$$(2) \quad \forall v, \forall i \neq j, \langle (E_{v,i} \otimes F_{v,j})\xi, \xi \rangle = 0, \\ \forall (v, w) \in E, \forall i, \langle (E_{v,i} \otimes F_{w,i})\xi, \xi \rangle = 0.$$

If this happens, we say that the graph G admits a *quantum c -colouring*; the smallest positive integer c for which G admits a quantum c -colouring was called in [3] the *quantum chromatic number* of G and denoted by $\chi_q(G)$.

One can interpret the quantum chromatic number in terms of a linear functional on the set of quantum correlations.

Definition 2.1. Let $G = (V, E)$ be a graph, let $|V| = n$ and fix $c \in \mathbb{N}$. The graph functional, $L_{G,c} : M_{nc} \rightarrow \mathbb{C}$ is defined by

$$L_{G,c}((a_{v,i,w,j})) = \sum_{i \neq j, v} a_{v,i,v,j} + \sum_{i, v \sim w} a_{v,i,w,i}.$$

Proposition 2.2. Let G be a graph on n vertices. Then $\chi_q(G) \leq c$ if and only if there exists $A \in C_q(n, c)$ such that $L_{G,c}(A) = 0$.

Proof. Since the entries of every correlation in $C_q(n, c)$ are non-negative, it follows that $L_{G,c}(A) = 0$ if and only if $a_{v,i,w,j} = 0$ whenever $v = w$ and $i \neq j$ and whenever (v, w) is an edge and $i = j$. Thus, there exists $A \in C_q(n, c)$ precisely when conditions (2) are satisfied. \square

Several variants of the quantum chromatic number were introduced in [24]. We focus on two denoted by $\chi_{qc}(G)$ and $\chi_{qa}(G)$, and called the *commuting quantum chromatic number*, and the *approximate quantum chromatic number*, respectively. Both of them can be obtained as in Proposition 2.2 by varying the sets of correlations that can be considered in place of $C_q(n, c)$.

Let $C_{qa}(n, c) := C_q(n, c)^-$ be the closure of $C_q(n, c)$; we note that it is not known if $C_q(n, c)$ is closed. In [24] it was shown that $C_{qa}(n, c)$ can be identified with the state space of a certain minimal tensor product and that consequently, $\chi_{qa}(G) = \chi_{qmin}(G)$, where this latter quantity was originally given a different definition.

We let $C_{qc}(n, c)$ denote the set of correlations obtained using relativistic quantum field theory. To be precise, instead of assuming that the POVM's $(E_{v,i})$ and $(F_{w,j})$ act on two finite dimensional Hilbert spaces and forming the tensor product of those spaces, we assume instead that they act on a common, possibly infinite dimensional, Hilbert space and that the E 's and F 's mutually commute. Thus, $C_{qc}(n, c)$ is the set of all $nc \times nc$ matrices of the form

$$p(i, j|v, w) = (\langle E_{v,i} F_{w,j} \xi, \xi \rangle)_{v,i,w,j},$$

where $E_{v,i}, F_{w,j}$ are positive operators on a Hilbert space \mathcal{H} satisfying

$$(3) \quad \sum_{i=1}^c E_{v,i} = \sum_{j=1}^c F_{w,j} = I, \forall v, w,$$

$$E_{v,i} F_{w,j} = F_{w,j} E_{v,i}, \forall v, w, i, j,$$

and $\xi \in \mathcal{H}$ is a unit vector.

As in the finite dimensional case, by enlarging the Hilbert space, one may assume without loss of generality, that all of these operators are orthogonal projections (see [24, Theorem 2.9] and Theorem 5.3 below).

Finally, $\text{Loc}(n, c)$ denotes the set of all *classical correlations*, also called *local correlations*, that is, the set of matrices $(p(i, j|v, w))_{v,i,w,j}$ which are in the closed convex hull of the product distributions $p(i, j|v, w) = p^1(i|v)p^2(j|w)$ where $p^1(i|v) \geq 0$ and $\sum_{i=1}^c p^1(i|v) = 1$ is a set of c -outcome probability distributions indexed by $v \in V$ with $|V| = n$ and, similarly, $p^2(j|w)$ is a set of c -outcome probability distributions indexed by $w \in V$. Since both of these sets of probability distributions define compact sets in \mathbb{R}^{nc} by Caratheodory's theorem, given any element of $\text{Loc}(n, c)$, there exist at most $n^2 c^2 + 1$ probability distributions $(p_k^1(i|v))_{i=1}^c$, $v \in V$ (resp. $(p_k^2(j|w))_{i=1}^c$, $w \in V$), and non-negative scalars λ_k such that

$$(4) \quad p(i, j|v, w) = \sum_k \lambda_k p_k^1(i|v) p_k^2(j|w), \quad i, j = 1, \dots, c, \quad v, w \in V.$$

For consistency of notation, we set $C_{\text{loc}}(n, c) := \text{Loc}(n, c)$.

There is another useful characterisation of the set $C_{\text{loc}}(n, c)$, discussed in [24, p. 5]. Let \mathcal{D} denote the tensor product of n copies of the abelian C^* -algebra ℓ_c^∞ , which is $*$ -isomorphic to the space of all (continuous) functions on nc points and set $e_{v,i} = 1 \otimes \dots \otimes 1 \otimes e_i \otimes 1 \dots \otimes 1$, where e_i denotes the i -th canonical basis vector for ℓ_c^∞ and it occurs in the v -th term of the tensor product, $1 \leq v \leq n$. Then $(p(i, j|v, w)) \in C_{\text{loc}}(n, c)$ if and only if there is a state $s : \mathcal{D} \otimes \mathcal{D} \rightarrow \mathbb{C}$ such that $p(i, j|v, w) = s(e_{v,i} \otimes e_{w,j})$. To see this fact, note that in the above formula each $p_k^1(i|v) = s_k^1(e_{v,i})$ for a state $s_k^1 : \mathcal{D} \rightarrow \mathbb{C}$, while $p_k^2(j|w) = s_k^2(e_{w,j})$. Thus, (4), for a typical element of $C_{\text{loc}}(n, c)$, becomes $p(i, j|v, w) = \sum_k \lambda_k s_k^1 \otimes s_k^2(e_{v,i} \otimes e_{w,j})$, which is a convex combination of product states. The fact that convex combinations of product states yields all states on $\mathcal{D} \otimes \mathcal{D}$ follows from another application of Caratheodory's theorem.

It is easy to see that

$$C_{\text{loc}}(n, c) \subseteq C_{\text{q}}(n, c) \subseteq C_{\text{qa}}(n, c) \subseteq C_{\text{qc}}(n, c).$$

Note that the correlations belonging to $C_{\text{loc}}(n, c)$ can be realised as in (1), but with the POVM's $(E_{v,i})_{i=1}^c$ and $(F_{w,j})_{j=1}^c$ consisting of mutually commuting operators.

These various sets of correlations allowed [24] to generalise and unify the definitions of the quantum chromatic number of a graph G (on n vertices) by

setting $\chi_x(G)$ equal to the least integer c such that there exists $p(i, j|v, w) \in C_x(n, c)$ satisfying:

$$(5) \quad \forall v, \forall i \neq j, p(i, j|v, v) = 0, \\ \forall (v, w) \in E, \forall i, p(i, i|v, w) = 0,$$

where $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$. In [3] it is shown that $\chi_{\text{loc}}(G)$ is equal to the usual chromatic number of a graph, $\chi(G)$.

Since $p(i, j|v, w) \geq 0$ for elements of each of these correlation sets, the proof of the following is identical to the proof of the last proposition:

Proposition 2.3. *Let G be a graph on n vertices and let $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$. Then $\chi_x(G) \leq c$ if and only if there exists $A \in C_x(n, c)$ such that $L_{G,c}(A) = 0$.*

From the above containments and proposition, we immediately see [24] that

$$\chi_{\text{qc}}(G) \leq \chi_{\text{qa}}(G) \leq \chi_{\text{q}}(G) \leq \chi(G).$$

Remark 2.4. By Proposition 2.3, and the fact that $\text{Loc}(n, c)$ is compact and convex, each graph G with $\chi_{\text{q}}(G) = c < \chi(G)$ yields a graph functional $L_{G,c}$ that is strictly positive on the set $C_{\text{loc}}(n, c) = \text{Loc}(n, c)$ of classical correlations and vanishes on the set $C_{\text{q}}(n, c) = Q(n, c)$. Thus, each such graph gives a functional $L_{G,c}$ that gives a Bell-type inequality separating local from quantum.

Remark 2.5. By Proposition 2.3(1), determining if the minimum of $L_{G,c}$ over the polytope $\text{Loc}(n, c)$ is 0, gives a LP to determine if $\chi(G) \leq c$.

Remark 2.6. The statement $C_{\text{q}}(n, c) = C_{\text{qc}}(n, c)$ for all n and c is often referred to as the (*strong bivariate*) *Tsirelson conjecture*. Thus, if the Tsirelson conjecture holds true, then necessarily, $\chi_c(G) = \chi(G)$ for every graph G . In addition, work of N. Ozawa [22] shows that the Connes Embedding Conjecture is equivalent to the statement that $C_{\text{qa}}(n, c) = C_{\text{qc}}(n, c)$ for all n and c . Thus, if the Connes Embedding Conjecture holds true, then necessarily $\chi_{\text{qc}}(G) = \chi_{\text{qa}}(G)$ for every graph G . We shall refer to the equality $C_{\text{qa}}(n, c) = C_{\text{q}}(n, c)$, $\forall n, c \in \mathbb{N}$, as the *closure conjecture*. Thus, if the closure conjecture is true, then $\chi_{\text{q}}(G) = \chi_{\text{qa}}(G)$ for every graph G .

Hence, determining if these chromatic numbers are always equal or can be separated would shed some light on these two conjectures. This was the original motivation for introducing these new parameters in [24].

3. A HIERARCHY OF CHROMATIC NUMBERS

In this section we revisit the ideas of the NPA hierarchy [21] and use them to construct a descending sequence of state spaces $C^N(n, c)$ that converges in an appropriate sense to $C_{\text{qc}}(n, c)$. These will allow us to construct a sequence of “chromatic” numbers $\chi_{\text{qc}}^N(G)$ each of which is computable by an

SDP and which converges to $\chi_{\text{qc}}(G)$. To accomplish this, we need to review a certain C*-algebra intrinsic in the definition of $\chi_{\text{qc}}(G)$.

Let $n, c \in \mathbb{N}$, $F(n, c) = \mathbb{Z}_c * \cdots * \mathbb{Z}_c$ (n copies), where $\mathbb{Z}_c = \{0, 1, \dots, c-1\}$ is the cyclic group with c elements, and $C^*(F(n, c))$ is the (full) C*-algebra of $F(n, c)$. The C*-algebra of \mathbb{Z}_c is canonically *-isomorphic, via Fourier transform, to the (abelian) C*-algebra $\ell_c^\infty = \{(\lambda_i)_{i=1}^c : \lambda_i \in \mathbb{C}, i = 1, \dots, c\}$. Thus, $C^*(F(n, c))$ is canonically *-isomorphic to the free product C*-algebra, amalgamated over the unit, $\ell_c^\infty *_1 \cdots *_1 \ell_c^\infty$ (n copies).

We denote by V the set $\{1, 2, \dots, n\}$. Let $e_{v,i}$ denote the element of $C^*(F(n, c))$ that is in the v -th copy of ℓ_c^∞ and is the vector that is 1 in the i -th component and 0 elsewhere. Alternatively, if we regard $F(n, c)$ as generated by unitaries, u_v with $u_v^c = 1$, then $e_{v,j}$ corresponds to the spectral projection of u_v onto the eigenspace of ω^j where $\omega = e^{2\pi i/c}$. In particular, $e_{v,j}$ belongs to the group algebra of $F(n, c)$ and because $u_v = \sum_{i=1}^c \omega^i e_{v,i}$ the collection $\{e_{v,i} : 1 \leq i \leq c, v \in V\}$ is another set of generators of the group algebra.

We let $\mathcal{S}(n, c) = \ell_c^\infty \oplus_1 \cdots \oplus_1 \ell_c^\infty$ (n copies) be the corresponding operator system coproduct (see, e.g., [17]). By [8] or since each generator u_v is in the span of $e_{v,i}, 1 \leq i \leq c$, $\mathcal{S}(n, c)$ can be identified with the span of the generators of the group $F(n, c)$ inside the C*-algebra $C^*(F(n, c))$. Then

$$\mathcal{S}(n, c) = \text{span}\{e_{v,i} : v \in V, 1 \leq i \leq c\}.$$

We note the relations

$$(6) \quad e_{v,j}^2 = e_{v,j} = e_{v,j}^*, \quad e_{v,i}e_{v,j} = 0, i \neq j, \quad \sum_{i=1}^c e_{v,i} = 1, \quad v \in V, 1 \leq j \leq c.$$

Because the left regular representation of $F(n, c)$ is faithful on the group algebra, the C*-algebra $C^*(F(n, c))$ can thus be viewed as the universal C*-algebra generated by the set $\mathcal{E} = \{e_{v,i} : v \in V, 1 \leq i \leq c\}$ satisfying (6).

A word in \mathcal{E} is an element of the form

$$(7) \quad \alpha = e_{v_1, i_1} e_{v_2, i_2} \cdots e_{v_k, i_k}$$

where $v_j \in V$ and $1 \leq i_j \leq c, j = 1, \dots, k$. The length $|\alpha|$ of a word α is the smallest k for which w can be written in the form (7). A polynomial of degree k is an element p of $C^*(F(n, c))$ of the form $p = \sum_{j=1}^m \lambda_j \alpha_j$, where α_j are words, $\lambda_j \in \mathbb{C}$, and $\text{deg}(p) \stackrel{\text{def}}{=} \max_{j=1, \dots, m} |\alpha_j| = k$.

For a given $N \in \mathbb{N}$, let

$$\mathcal{P}_N = \text{span}\{p : \text{a polynomial with } \text{deg}(p) \leq N\},$$

and

$$\mathcal{P} = \text{span}\{\mathcal{P}_N : N \in \mathbb{N}\}.$$

We note that \mathcal{P}_N is an operator subsystem of $C^*(F(n, c))$ and that $\mathcal{P}_N \subseteq \mathcal{P}_{N+1}, N \in \mathbb{N}$. We also note that \mathcal{P} is a (dense) *-subalgebra of $C^*(F(n, c))$ and hence possesses a canonical induced operator system structure.

Note that

$$C^*(F(n, c)) \otimes_{\max} C^*(F(n, c)) = C^*(F(n, c) \times F(n, c)),$$

up to a (canonical) *-isomorphism. Let

$$\mathcal{A} = \mathcal{P} \otimes \mathcal{P} \subseteq C^*(F(n, c) \times F(n, c)).$$

We have that $C^*(F(n, c) \times F(n, c))$ is the universal C*-algebra generated by two families of elements, $\mathcal{E} = \{e_{v,i} : v \in V, 1 \leq i \leq c\}$ and $\mathcal{F} = \{f_{w,j} : w \in V, 1 \leq j \leq c\}$, each of which satisfies relations (6), as well as the commutativity relations

$$(8) \quad e_{v,i} f_{w,j} = f_{w,j} e_{v,i}, \quad v, w \in V, 1 \leq i, j \leq c.$$

We define polynomials on $\mathcal{E} \cup \mathcal{F}$ in a similar fashion – note that, due to (8), each such polynomial is a sum of products of the form $\alpha\beta$, where α (resp. β) is a word on \mathcal{E} (resp. \mathcal{F}).

Let

$$\Gamma_N = \{\gamma : \text{a word in } \mathcal{E} \cup \mathcal{F}, |\gamma| \leq N\},$$

$\mathcal{A}_N = \text{span} \Gamma_N$ and

$$S_N = S_{N,n,c} = \{s : \mathcal{A} \rightarrow \mathbb{C} : s(1) = 1, s(p^*p) \geq 0 \text{ for all } p \in \mathcal{A}_N\}.$$

The functionals s above, as well as all functionals appearing hereafter, are assumed to be linear. Set $\Gamma = \bigcup_{N=1}^{\infty} \Gamma_N$. Note that

$$\mathcal{A} = \text{span} \Gamma = \text{span}\{\mathcal{A}_N : N \in \mathbb{N}\}.$$

Lemma 3.1. *Let $s : \mathcal{A} \rightarrow \mathbb{C}$ be a linear functional. Then $s \in S_N$ if and only if $(s(\beta^*\alpha))_{\alpha, \beta \in \Gamma_N}$ is a positive matrix.*

Proof. By definition, $s \in S_N$ if and only if $s(p^*p) \geq 0$ for all $p \in \mathcal{A}_N$, if and only if

$$s \left(\left(\sum_{k=1}^m \overline{\lambda_k} \alpha_k^* \right) \left(\sum_{k=1}^m \lambda_k \alpha_k \right) \right) \geq 0,$$

for all $\alpha_k \in \Gamma_N$, $\lambda_k \in \mathbb{C}$, $k = 1, \dots, m$, if and only if

$$\sum_{k=1}^m \overline{\lambda_k} \lambda_l \alpha_k^* \alpha_l \geq 0,$$

for all $\alpha_k \in \Gamma_N$, $\lambda_k \in \mathbb{C}$, $k = 1, \dots, m$, if and only if the matrix $(s(\beta^*\alpha))_{\alpha, \beta \in \Gamma_N}$ is positive. \square

Every functional on \mathcal{A} can, after restriction, be considered as a functional on \mathcal{A}_N . Letting

$$S = S_{n,c} = \{s : \mathcal{A} \rightarrow \mathbb{C} : s(1) = 1, s(p^*p) \geq 0 \text{ for all } p \in \mathcal{A}\},$$

we clearly have

$$S = \{s : \mathcal{A} \rightarrow \mathbb{C} : s|_{\mathcal{A}_N} \in S_N, \text{ for all } N \in \mathbb{N}\}.$$

Lemma 3.2. *$s \in S$ if and only if there exists a state \tilde{s} of the C*-algebra $\mathcal{C} = C^*(F(n, c) \times F(n, c))$ with $\tilde{s}|_{\mathcal{A}} = s$.*

Proof. The statement follows from a standard construction of GNS type; the detailed arguments are omitted. \square

Let \mathcal{R} be the ideal of all polynomials on the set of non-commuting variables $\mathcal{E} \cup \mathcal{F}$ generated by the elements

$$e_{v,j}^2 - e_{v,j}, \quad e_{v,j}^* - e_{v,j}, \quad \sum_{k=1}^c e_{v,k} - 1, \quad e_{v,i}f_{w,j} - f_{w,j}e_{v,i},$$

where $v, w \in V, 1 \leq i, j \leq c$. Set $\mathcal{R}_N = \mathcal{R} \cap \mathcal{A}_N$. For example, \mathcal{R} contains the elements $\beta_1^* \alpha_1 - \beta_2^* \alpha_2$ if $\beta_1^* \alpha_1 = \beta_2^* \alpha_2$ in Γ . We let

$$\mathcal{M} = \left\{ (c_{\alpha,\beta})_{\alpha,\beta \in \Gamma} : \sum_{k=1}^m \lambda_k \beta_k^* \alpha_k \in \mathcal{R} \implies \sum_{k=1}^m \lambda_k c_{\alpha_k, \beta_k} = 0 \right\}.$$

We also let \mathcal{M}_N be the set of all compressions of matrices in \mathcal{M} to $\Gamma_N \times \Gamma_N$. In particular, if $(c_{\alpha,\beta})_{\alpha,\beta \in \Gamma_N}$ then $c_{\alpha_1, \beta_1} = c_{\alpha_2, \beta_2}$ whenever $\beta_1^* \alpha_1 = \beta_2^* \alpha_2$.

Lemma 3.3. *There is a bijective correspondence between \mathcal{M} and the set of all linear functionals on \mathcal{A} , sending an element $(c_{\alpha,\beta})_{\alpha,\beta \in \Gamma}$ of \mathcal{M} to the functional $f : \mathcal{A} \rightarrow \mathbb{C}$ given by $f(\beta^* \alpha) = c_{\alpha,\beta}$, $\alpha, \beta \in \Gamma$.*

Proof. The only thing that needs to be checked is that, given $(c_{\alpha,\beta})_{\alpha,\beta \in \Gamma} \in \mathcal{M}$, the mapping defined on the generators of \mathcal{A} by $f(\beta^* \alpha) = c_{\alpha,\beta}$, $\alpha, \beta \in \Gamma$, and extended by linearity, is well-defined. This follows from the definition of \mathcal{M} . \square

For $N \in \mathbb{N}$ we now let

$$C^N(n, c) = \{(s(e_{v,i}f_{w,j}))_{v,i,w,j} : s \in S_N\}.$$

Note that

$$C_{\text{qc}}(n, c) = \{(s(e_{v,i} \otimes f_{w,j}))_{v,i,w,j} : s \in S(\mathcal{S}(n, c) \otimes_c \mathcal{S}(n, c))\},$$

the set of all relativistic quantum correlations.

Theorem 3.4. (i) *A matrix $A = (a_{v,i,w,j})_{v,i,w,j}$ belongs to $C_{\text{qc}}(n, c)$ if and only if it can be completed to a positive matrix $B = (b_{\alpha,\beta})_{\alpha,\beta \in \Gamma} \in \mathcal{M}$ (meaning that every finite submatrix of B is positive) with $b_{1,1} = 1$.*

(ii) *A matrix $A = (a_{v,i,w,j})_{v,i,w,j}$ belongs to $C^N(n, c)$ if and only if it can be completed to a positive matrix $B = (b_{\alpha,\beta})_{\alpha,\beta \in \Gamma_N} \in \mathcal{M}_N$ with $b_{1,1} = 1$.*

Proof. (i) Let $s \in S(\mathcal{S}(n, c) \otimes_c \mathcal{S}(n, c))$ be a state such that $A = (s(e_{v,i} \otimes f_{w,j}))_{v,i,w,j}$. Let H be a Hilbert space, $\pi : C^*(F(n, c)) \otimes_{\max} C^*(F(n, c)) \rightarrow \mathcal{B}(H)$ be a *-representation and $\xi \in H$ be a unit vector such that $s(T) = \langle \pi(T)\xi, \xi \rangle$, $T \in C^*(F(n, c)) \otimes_{\max} C^*(F(n, c))$. Thus,

$$s(\beta^* \alpha) = \langle \pi(\alpha)\xi, \pi(\beta)\xi \rangle,$$

and every finite submatrix of the matrix $(s(\beta^* \alpha))_{\alpha,\beta \in \Gamma}$ is positive.

Conversely, suppose that $B = (b_{\alpha,\beta})_{\alpha,\beta \in \Gamma} \in \mathcal{M}$ is a completion of A that has the property that all of its finite submatrices are positive. It follows from

Lemma 3.1 that the linear functional $s : \mathcal{A} \rightarrow \mathbb{C}$ given by $s(\alpha) = b_{\alpha,1}$ (and well-defined by Lemma 3.3) is positive. Note that $s(1) = 1$. By Lemma 3.2, s is the restriction to \mathcal{A} of a state of $C^*(F(n,c) \times F(n,c))$. Thus, $A \in C_{\text{qc}}(n,c)$.

(ii) Suppose that $A = (a_{v,i,w,j})_{v,i,w,j}$ belongs to $C^N(n,c)$. Then there exists $s \in S_N$ such that $a_{v,i,w,j} = s(e_{v,i}f_{w,j})$, $v,w \in V$, $i,j = 1, \dots, c$. By Lemma 3.1, $(s(\beta^*\alpha))_{\alpha,\beta \in \Gamma_N}$ is a positive completion of A that lies in \mathcal{M}_N .

Conversely, suppose that B is a positive completion of A that lies in \mathcal{M}_N . By Lemma 3.3, B is the compression to $\Gamma_N \times \Gamma_N$ of a matrix of the form $(s(\beta^*\alpha))_{\alpha,\beta \in \Gamma}$ and, by Lemma 3.1, $s \in S_N$. It follows that $A \in A_N$. \square

The following corollary follows directly from Lemma 3.3; we omit the detailed proof.

Corollary 3.5. *Let $G = (V, E)$ be a graph. We have that $\chi_{\text{qc}}(G) \leq c$ if and only if there exists $s \in S$ such that*

$$(9) \quad \begin{aligned} \forall v, \forall i \neq j, s(e_{v,i}f_{v,j}) &= 0, \\ \forall (v, w) \in E, \forall i, s(e_{v,i}f_{w,i}) &= 0. \end{aligned}$$

Lemma 3.6. *Let $s \in S_N$. Then $|s(\gamma)| \leq 1$ for all $\gamma \in \Gamma$ with $|\gamma| \leq 2N$.*

Proof. We first show that $0 \leq s(\gamma^*\gamma) \leq 1$ for all words γ with $|\gamma| \leq N$. To this end, we use induction on $|\gamma|$. We have that $s(e_{v,i}) = s(e_{v,i}^*e_{v,i}) \geq 0$ while $\sum_{i=1}^n s(e_{v,i}) = 1$; it follows that $0 \leq s(e_{v,i}) \leq 1$ for all v, i . By symmetry, $0 \leq s(f_{w,j}) \leq 1$ for all w, j , that is, the claim holds for all γ with $|\gamma| = 1$.

Suppose $|\gamma| \leq k$, for some $k \leq N - 1$. If $v \in V$ then

$$0 \leq s((e_{v,i}\gamma)^*(e_{v,i}\gamma)) = s(\gamma^*e_{v,i}\gamma) \quad \text{and} \quad \sum_{k=1}^c s(\gamma^*e_{v,k}\gamma) = s(\gamma^*\gamma) \leq 1.$$

It follows by induction that $0 \leq s(\gamma^*\gamma) \leq 1$ whenever $|\gamma| \leq N$.

Now suppose $|\gamma| \leq 2N$ and write $\gamma = \beta^*\alpha$ for some words α and β of length at most N . By Lemma 3.1, the matrix $\begin{pmatrix} s(\alpha^*\alpha) & s(\beta^*\alpha) \\ s(\alpha^*\beta) & s(\beta^*\beta) \end{pmatrix}$ is positive. Thus,

$$|s(\gamma)| = |s(\beta^*\alpha)| \leq \max\{s(\alpha^*\alpha), s(\beta^*\beta)\} \leq 1. \quad \square$$

Lemma 3.7. *Let $s_N \in S_N$, $N \in \mathbb{N}$. There exists a subsequence $(s_{N'})_{N'}$ of $(s_N)_N$ which converges pointwise to an element of S .*

Proof. Since $s_N \in S_1$ for all N , Lemma 3.6 implies that $|s_N(\gamma)| \leq 1$ whenever $\gamma \in \Gamma_2$. Thus, there exists a subsequence

$$s_{N_1}^{(1)}, s_{N_2}^{(1)}, s_{N_3}^{(1)}, \dots$$

such that $(s_{N_k}^{(1)}(\gamma))_{k \in \mathbb{N}}$ converges whenever $\gamma \in \Gamma_2$. Deleting the first terms of this subsequence, if necessary, we may assume that $s_{N_k}^{(1)} \in S_2$ for all k .

Using Lemma 3.6, it now follows that there exists a subsequence $(s_{N_k}^{(2)})_{k \in \mathbb{N}}$ of $(s_{N_k}^{(1)})_{k \in \mathbb{N}}$ such that $(s_{N_k}^{(2)}(\gamma))_{k \in \mathbb{N}}$ converges whenever $\gamma \in \Gamma_4$. Deleting the first terms of this subsequence, if necessary, we may assume that $s_{N_k}^{(2)} \in S_3$ for all k .

Continuing inductively, for each m we obtain a sequence $(s_{N_k}^{(m)})_{k \in \mathbb{N}}$, which is a subsequence of $(s_{N_k}^{(m-1)})_{k \in \mathbb{N}}$, and which has the property that $(s_{N_k}^{(m)}(\gamma))_{k \in \mathbb{N}}$ converges whenever $\gamma \in \Gamma_{2m}$.

We claim that $(s_{N_k}^{(k)}(\gamma))_{k \in \mathbb{N}}$ converges whenever $\gamma \in \Gamma$. To see this, let $\gamma \in \Gamma$ and assume that $|\gamma| \leq 2m$ for some m . Then $(s_{N_k}^{(k)}(\gamma))_{k \geq m}$ is a subsequence of $(s_{N_k}^{(m)}(\gamma))_{k \in \mathbb{N}}$ and hence converges.

Set $s(p) = \lim_{k \rightarrow \infty} s_{N_k}^{(k)}(\gamma)$, $p \in \mathcal{A}$. Since s is a pointwise limit of linear maps, it is a linear map itself. Clearly, $s(1) = 1$. We claim that $s \in S$. If $p \in \mathcal{A}$ is a polynomial, then $p \in \mathcal{A}_N$ for some N and hence $s_{N_k}^{(k)}(p^*p) \geq 0$ whenever $N_k \geq N$. It follows that $s(p^*p) \geq 0$. \square

Theorem 3.8. *We have that $\bigcap_{N=2}^{\infty} C^N(n, c) = C_{\text{qc}}(n, c)$.*

Proof. Suppose that $s \in S(\mathcal{S}(n, c) \otimes_c \mathcal{S}(n, c))$. By [24, Lemma 2.8],

$$(10) \quad \mathcal{S}(n, c) \otimes_c \mathcal{S}(n, c) \subseteq_{\text{coi}} C^*(F(n, c)) \otimes_{\text{max}} C^*(F(n, c)),$$

and hence s extends to a state \tilde{s} on $C^*(F(n, c)) \otimes_{\text{max}} C^*(F(n, c))$. Letting $s_N = \tilde{s}|_{\mathcal{A}_N}$, we have that $s_N \in S_N$; clearly, $s(e_{v,i} \otimes f_{w,j}) = s_N(e_{v,i} f_{w,j})$, $N \geq 2$, so $C_{\text{qc}}(n, c) \subseteq \bigcap_{N=2}^{\infty} C^N(n, c)$.

Conversely, suppose that $A \in \bigcap_{N=2}^{\infty} C^N(n, c)$. For each $N \geq 2$, let $s_N \in S_N$ be such that $A = (s_N(e_{v,i} f_{w,j}))_{v,i,w,j}$. By Lemma 3.7, there exists a subsequence of (s_N) which converges pointwise to an element $s \in S$. Since $s_N(e_{v,i} f_{w,j}) = s_M(e_{v,i} f_{w,j})$ for all $N, M \in \mathbb{N}$, we have that $s(e_{v,i} f_{w,j}) = s_N(e_{v,i} f_{w,j})$, $N \in \mathbb{N}$. By Lemma 3.2, s is the restriction of a state \tilde{s} on $C^*(F(n, c) \times F(n, c)) = C^*(F(n, c)) \otimes_{\text{max}} C^*(F(n, c))$. By (10), we have that $A \in C_{\text{qc}}(n, c)$. \square

Definition 3.9. *Let G be a graph on a set V of n vertices with set E of edges. A quantum N, c -colouring of G is a state $s \in S_{N,n,c}$ such that*

$$(11) \quad \begin{aligned} \forall v, \forall i \neq j, s(e_{v,i} f_{v,j}) &= 0, \\ \forall (v, w) \in E, \forall i, s(e_{v,i} f_{w,i}) &= 0. \end{aligned}$$

The N th quantum chromatic number $\chi_{\text{qc}}^N(G)$ of G is the smallest positive integer c for which there exists a quantum N, c -colouring of G .

According to Theorem 3.4, quantum N, c -colourings of G correspond bijectively to matrices $A = (a_{v,i,w,j})_{v,i,w,j} \in \mathcal{C}^N(n, c)$ such that

$$a_{v,i,v,j} = 0, \forall v, \forall i \neq j \quad \text{and} \quad a_{v,i,w,i} = 0, \forall (v, w) \in E, \forall i.$$

Theorem 3.10. *Let G be a graph. We have $\chi_{\text{qc}}^N(G) \leq \chi_{\text{qc}}^{N+1}(G) \leq \chi_{\text{qc}}(G)$, for every $N \in \mathbb{N}$. Moreover, $\lim_{N \rightarrow \infty} \chi_{\text{qc}}^N(G) = \chi_{\text{qc}}(G)$.*

Thus, given $n \in \mathbb{N}$, there exists $N \in \mathbb{N}$ such that $\chi_{\text{qc}}(G) = \chi_{\text{qc}}^N(G)$ for all graphs on at most n vertices.

Proof. Let $c = \chi_{\text{qc}}^{N+1}(G)$ and $s \in S_{N+1,n,c}$ be a quantum N, c -colouring of G . Then $s \in S_{N,n,c}$. Since $\chi_{\text{qc}}^N(G)$ is the minimum of all c for which there exists a quantum N, c -colouring for G , we have that $\chi_{\text{qc}}^N(G) \leq \chi_{\text{qc}}^{N+1}(G)$.

A similar argument, using the fact that $S \subseteq S_{N,n,c}$, shows the second inequality. It follows that the sequence $(\chi_{\text{qc}}^N(G))_{N=1}^{\infty}$ stabilises, that is, there exist $c, N_0 \in \mathbb{N}$ such that $\chi_{\text{qc}}^N(G) = c$ for all $N \geq N_0$. For each $N \geq N_0$, let $s_N \in S_N$ be a quantum N, c -colouring of G . By Lemma 3.7 and Corollary 3.5, $\chi_{\text{qc}}(G) \leq c$.

Finally, since the sequence stabilises for each graph and there are only finitely many graphs on at most n vertices, there exists $N \in \mathbb{N}$ such that $\chi_{\text{qc}}^N(G) = \chi_{\text{qc}}(G)$, for all graphs on at most n vertices. \square

Remark 3.11. Let N_n be the least integer such that $\chi_{\text{qc}}^{N_n}(G) = \chi_{\text{qc}}(G)$ for all graphs on at most n vertices. Because we obtain the integer N_n by a compactness argument, we do not have effective bounds on N_n . In particular, we do not know if $\sup_n N_n$ is bounded or have any information on its growth rate.

4. AN SDP FOR THE COMMUTING QUANTUM CHROMATIC NUMBER

In this section we prove that for each graph $G = (V, E)$ determining if $\chi_{\text{qc}}(G) \leq c$ is decidable by a semidefinite programming problem.

We fix a graph $G = (V, E)$, let $n = |V|$ and let $N := N_n$ be the natural number given by Remark 3.11. Let $c \in \mathbb{N}$. We set $C^N = C^N(n, c) = \{(s(e_{v,i} f_{w,j}) : s \in S_N) \subseteq M_{nc}$.

Recall that a set is called a *spectrahedron* if it can be realised as the intersection of the set of positive semidefinite matrices of some size with an affine subset.

Recall that the graph functional $L_{G,c} : M_{nc} \rightarrow \mathbb{C}$ is defined by

$$L_{G,c}((a_{v,i,w,j})) = \sum_{i \neq j, v} a_{v,i,v,j} + \sum_{i, v \sim w} a_{v,i,w,i}.$$

When the value of c is understood, we will often write L_G for $L_{G,c}$.

Proposition 4.1. *For any $N \in \mathbb{N}$, the problem of minimising $L_{G,c}$ over the set C^N is an SDP.*

Proof. Recall that \mathcal{M}_N is a space of $|\Gamma_N| \times |\Gamma_N|$ matrices defined by some linear constraints. Thus, the set \mathcal{B}_N of positive semidefinite matrices $B \in \mathcal{M}_N$ with $b_{1,1} = 1$ is a spectrahedron. By Theorem 3.4 (ii), every element of C^N is the restriction of such a matrix B in \mathcal{M}_N to some of its components. Thus, if we extend $L_{G,c}$ to a linear functional F on \mathcal{B}_N by setting it equal

to 0 on all the components not included in C^N then we see that minimizing F over \mathcal{B}_N is the same as minimizing $L_{G,c}$ over C^N . \square

Theorem 4.2. *Let $G = (V, E)$ be a graph on n vertices and let $N = \max\{N_n, 2\}$, where N_n is defined in Remark 3.11. Then $\chi_{\text{qc}}(G) \leq c$ if and only if $\inf\{L_{G,c}(A) : A \in C^N(n, c)\} = 0$. Hence, the problem of determining if $\chi_{\text{qc}}(G) \leq c$ is solvable by this SDP.*

Proof. For any $s \in S_N$, any $v, w \in V$, any $1 \leq i, j \leq c$, since $N \geq 2$, we have that

$$0 \leq s((e_{v,ifw,j})^*(e_{v,ifw,j})) = s(e_{v,ifw,j}).$$

Thus, all the elements of C^N are non-negative and, consequently, $L_{G,c}(A) \geq 0$ on C^N .

Since C^N is a compact set the infimum is 0 if and only if it is attained at some matrix $A \in C^N$, but in that case we have that A is the image of a state that defines a quantum N, c -colouring and so $\chi_{\text{qc}}(G) = \chi_{\text{qc}}^N(G) \leq c$. \square

Note that since C^N is a compact set, the above infimum is actually a minimum.

Remark 4.3. It is known that computing $\chi_{\text{q}}(G)$ is an NP-hard problem [14], but it is not known if computing $\chi_{\text{qc}}(G)$ or $\chi_{\text{qa}}(G)$ is NP-hard. A proof that did not rely on Tsirelson's or Connes' conjecture that these are also NP-hard would be interesting. A proof that either of these is of complexity P would be a dramatic result. It would show that either the corresponding conjecture is false or that $P=NP$.

We can strengthen the above result a bit as follows.

Theorem 4.4. *For each $n \in \mathbb{N}$, there is a constant $\epsilon_n > 0$ such that if $G = (V, E)$ is a graph on n vertices and $N = \max\{2, N_n\}$, then $\chi_{\text{qc}}(G) \leq c$ if and only if $\inf\{L_{G,c}(A) : A \in C^N\} < \epsilon_n$.*

Proof. The graphs on n vertices split into two subsets: those for which $\chi_{\text{qc}}(G) \leq c$, and those for which $\chi_{\text{qc}}(G) > c$. For each graph in the latter set, we have that $\inf\{L_{G,c}(A) : A \in C^N\} = b_G > 0$. Since there are only finitely many such graphs, we may set $\epsilon_n = \min\{b_G\}$ over this set of graphs. \square

The above result is not of much computational use without estimates for ϵ_n , but it might be of theoretical use. Since we know that we only need to get the SDP within ϵ_n this may give us a crude operation/complexity count.

5. SYNCHRONOUS STATES AND THE TRACIAL RANK

D. Roberson and L. Mančinska [26] introduced the projective rank $\xi_f(G)$ of a graph G and showed that $\xi_f(G) \leq \chi_{\text{q}}(G)$. This lower bound has been crucial for computing the quantum chromatic numbers of some graphs. Unfortunately, the proof of this estimate uses in a critical way the fact that the

involved representations are finite dimensional and, consequently, we do not know if it is also a lower bound for $\chi_{\text{qc}}(G)$. To ameliorate this situation, we will develop an analogous quantity that is better suited to work with infinite dimensional representations.

Our first result applies to a larger family of games than the graph colouring game, namely games where Alice and Bob have the same set of inputs V and require that when Alice and Bob receive the same input, then they must produce the same output. Given a correlation $(p(i, j|v, w))_{v, i, w, j}$, set

$$p(i = j|v, w) := \sum_{i=1}^c p(i, i|v, w).$$

Then a perfect strategy for this game means $p(i = j|v = w) = 1$, that is, $p(i = j|v, v) = 1$ for all $v \in V$. Our result can be summarised as saying that such correlations always arise from tracial states. Recall that a state s on a C^* -algebra \mathcal{A} is called tracial provided that $s(xy) = s(yx)$ for all $x, y \in \mathcal{A}$. We show that, in fact, the projective rank of a graph can be described by using tracial states on finite dimensional C^* -algebras.

Definition 5.1. For $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$, we call a correlation $(p(i, j|v, w))$ from $C_x(n, c)$ synchronous if it satisfies the condition $p(i = j|v = w) = 1$, and let $C_x^s(n, c) \subseteq C_x(n, c)$ denote the subset of all synchronous correlations.

Note that $C_q^s(n, c) = C_{\text{qc}}^s(n, c) \cap C_q(n, c)$.

Definition 5.2. A realisation of an element $(p(i, j|v, w))_{v, i, w, j}$ of $C_{\text{qc}}(n, c)$ is a tuple $\left(((E_{v,i})_{i=1}^c)_{v \in V}, ((F_{w,j})_{j=1}^c)_{w \in V}, \mathcal{H}, \eta \right)$, where V is an index set of cardinality n , \mathcal{H} is a Hilbert space, $\eta \in \mathcal{H}$ is a unit vector, and $E_{v,i}, F_{w,j} \in \mathcal{B}(\mathcal{H})$ are projections satisfying

- (i) $p(i, j|v, w) = \langle E_{v,i} F_{w,j} \eta, \eta \rangle$, $v, w \in V$, $i, j = 1, \dots, c$;
- (ii) $\sum_{i=1}^c E_{v,i} = \sum_{j=1}^c F_{w,j} = I$, $v, w \in V$;
- (iii) $E_{v,i} F_{w,j} = F_{w,j} E_{v,i}$, $v, w \in V$, $i, j = 1, \dots, c$.

When n and c are understood, to avoid excessive notation, we will often denote a realisation by simply $((E_{v,i}), (F_{w,j}), \mathcal{H}, \eta)$.

Theorem 5.3. A correlation $((p(i, j|v, w))_{v, i, w, j})$ belongs to $C_q(n, c)$ if and only if it has a realisation $((E_{v,i}), (F_{w,j}), \mathcal{H}, \eta)$ for which \mathcal{H} is finite dimensional.

Proof. Suppose that $((p(i, j|v, w))_{v, i, w, j})$ is a correlation which possesses a realisation $((E_{v,i}), (F_{w,j}), \mathcal{H}, \eta)$ for which \mathcal{H} is finite dimensional. We now essentially recall the argument from the unpublished preprint [28, Theorem 1] which allows us to pass to spacial tensoring. Let \mathcal{C} (resp. \mathcal{D}) be the C^* -algebra generated by $\{E_{v,i} : v \in V, i = 1, \dots, c\}$ (resp. $\{F_{w,j} : w \in V, j = 1, \dots, c\}$). Since \mathcal{C} is finite dimensional, we may assume, without loss of generality, that $H = \bigoplus_{s=1}^t \mathbb{C}^{k_s} \otimes \mathbb{C}^{l_s}$ and $\mathcal{C} = \bigoplus_{s=1}^t M_{k_s} \otimes 1_{l_s}$. Since \mathcal{D} and \mathcal{C} commute, we have that \mathcal{D} is contained in the C^* -algebra

$\bigoplus_{s=1}^t 1_{k_s} \otimes M_{l_s}$. Thus, $E_{v,i} = \bigoplus_{s=1}^t E_{v,i}^s \otimes I_{l_s}$ and $F_{w,j} = \bigoplus_{s=1}^t I_{k_s} \otimes F_{w,j}^s$ for some projections $E_{v,i}^s \in M_{k_s}$, $F_{w,j}^s \in M_{l_s}$. Now let $k = \max\{k_1, \dots, k_t\}$ and $l = \max\{l_1, \dots, l_t\}$. Consider the Hilbert space H as a subspace, in the natural way, of $\bigoplus_{s=1}^t \mathbb{C}^k \otimes \mathbb{C}^l$, and identify the latter space with $\mathbb{C}^k \otimes \mathbb{C}^l$. Under these identifications, the projections $E_{v,i}$ (resp. $F_{w,j}$) have the form $E_{v,i} = E'_{v,i} \otimes I_l$ (resp. $F_{w,j} = I_k \otimes F'_{w,j}$), for some projections $E'_{v,i}$ (resp. $F'_{w,j}$) on \mathbb{C}^k (resp. \mathbb{C}^l). It follows that $((p(i,j|v,w))_{v,i,w,j} \in C_q(n,c)$.

Conversely, assume that $(p(i,j|v,w)) \in C_q(n,c)$. Then there exist finite dimensional Hilbert spaces \mathcal{H}_A and \mathcal{H}_B , POVM's $((P_{v,i})_{i=1}^c)_{v \in V}$ on \mathcal{H}_A , POVM's $((R_{w,j})_{j=1}^c)_{w \in V}$ on \mathcal{H}_B and a unit vector $\eta \in \mathcal{H}_A \otimes \mathcal{H}_B$ such that $p(i,j|v,w) = \langle P_{v,i} \otimes R_{w,j} \eta, \eta \rangle$.

For convenience, set $V = \{1, \dots, n\}$. Let $\tilde{\mathcal{H}}_A = \mathcal{H}_A \otimes \mathbb{C}^c$, regarded as the direct sum of c copies of \mathcal{H}_A . Note that $\tilde{\mathcal{H}}_A$ is still finite dimensional and define an inclusion $W : \mathcal{H}_A \rightarrow \tilde{\mathcal{H}}_A$ via $h \rightarrow (P_{1,1}^{1/2}h, \dots, P_{1,c}^{1/2}h)$. The fact that $(P_{1,i})_{i=1}^c$ is a POVM implies that this inclusion is an isometry.

Define operators on $\tilde{\mathcal{H}}_A$ by setting: $\tilde{P}_{1,i} = I_{\mathcal{H}_A} \otimes E_{i,i}$ (where $E_{i,i}$ here denotes the corresponding diagonal matrix unit on \mathbb{C}^c) and for $v \neq 1$, let $\tilde{P}_{v,i}$ be the operator matrix, with (k,l) -entry,

$$\tilde{P}_{v,i} = (P_{1,k}^{1/2} P_{v,i} P_{1,l}^{1/2}), i \neq 1,$$

and

$$\tilde{P}_{v,1} = (P_{1,k}^{1/2} P_{v,1} P_{1,l}^{1/2}) + (I_{\tilde{\mathcal{H}}} - WW^*).$$

Note that this standard dilation trick turns the POVM $(P_{1,i})_{i=1}^c$ into a PVM $(\tilde{P}_{1,i})_{i=1}^c$, and turns each POVM $(P_{v,i})_{i=1}^c$, $v \neq 1$, into a new POVM $(\tilde{P}_{v,i})_{i=1}^c$ on the larger space. Moreover, $W^* \tilde{P}_{v,i} W = P_{v,i}$.

Also, note that for $i \neq 1$,

$$\tilde{P}_{v,i}^2 = \sum_{t=1}^c (P_{1,k}^{1/2} P_{v,i} P_{1,t}^{1/2} P_{1,t}^{1/2} P_{v,i} P_{1,l}^{1/2}) = (P_{1,k}^{1/2} P_{v,i}^2 P_{1,l}^{1/2}).$$

Similarly,

$$\tilde{P}_{v,1}^2 = (P_{1,k}^{1/2} P_{v,1}^2 P_{1,l}^{1/2}) + (I_{\tilde{\mathcal{H}}} - WW^*).$$

Thus, if $P_{v,i}$ is a projection, then its dilation $\tilde{P}_{v,i}$ is also a projection.

Hence, if any $(P_{v,i})_{i=1}^c$ is already a PVM, that property is preserved by the dilation. It follows that if we repeat this standard dilation trick n times, once for each v , then we will obtain a family of PVM's $(\hat{P}_{v,i})_{i=1}^c$, $1 \leq v \leq n$, on the finite dimensional Hilbert space $\hat{\mathcal{H}}_A = \mathcal{H}_A \otimes \mathbb{C}^{2^n c}$ and an isometric embedding $W_A : \mathcal{H}_A \rightarrow \hat{\mathcal{H}}_A$ such that $W_A^* \hat{P}_{v,i} W_A = P_{v,i}$.

Repeating the same process for the POVM's $(R_{w,j})$ on \mathcal{H}_B , we obtain a family of PVM's $(\hat{R}_{w,j})$ on a finite dimensional space $\hat{\mathcal{H}}_B$ and an isometry $W_B : \mathcal{H}_B \rightarrow \hat{\mathcal{H}}_B$ such that $R_{w,j} = W_B^* \hat{R}_{w,j} W_B$.

Finally, $\langle (\hat{P}_{v,i} \otimes I)(I \otimes \hat{R}_{w,j})(W_A \otimes W_B)\eta, (W_A \otimes W_B)\eta \rangle = p(i, j|v, w)$ so that

$$\left((\hat{P}_{v,i} \otimes I), (I \otimes \hat{R}_{w,j}), \hat{\mathcal{H}}_A \otimes \hat{\mathcal{H}}_B, (W_A \otimes W_B)\eta \right)$$

is a realisation of $(p(i, j|v, w))$ by commuting PVM's on a finite dimensional Hilbert space. It now follows that $((p(i, j|v, w))_{v,i,w,j})$ is in $C_q(n, c)$. \square

Remark 5.4. Similarly it can be shown that, $((p(i, j|v, w)) \in C_{\text{loc}}(n, c)$ if and only if the realisation can be chosen such that all the operators commute. We do not know of an analogous characterisation of correlations in $C_{\text{qa}}(n, c)$.

Theorem 5.5. *Let $(p(i, j|v, w)) \in C_{\text{qc}}^s(n, c)$ be a synchronous correlation with realisation $\{(E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta\}$. Then*

- (i) $E_{v,i}\eta = F_{v,i}\eta, \quad v \in V, i = 1, \dots, c;$
- (ii) $p(i, j|v, w) = \langle E_{v,i}E_{w,j}\eta, \eta \rangle = \langle F_{w,j}F_{v,i}\eta, \eta \rangle = p(j, i|w, v)$
- (iii) *The functional $s : X \rightarrow \langle X\eta, \eta \rangle$ is a tracial state on the C^* -algebra generated by the set $\{E_{v,i} : v \in V, i = 1, \dots, c\}$ (resp. $\{F_{w,j} : w \in V, j = 1, \dots, c\}$).*

Conversely, given a family of projections $\{e_{v,i} : v \in V, 1 \leq i \leq c\}$ in a unital C^* -algebra \mathcal{A} such that $\sum_{i=1}^c e_{v,i} = I, v \in V$, and a tracial state s on \mathcal{A} , then $(p(i, j|v, w)) = (s(e_{v,i}e_{w,j}))_{v,i,w,j}$ is in $C_{\text{qc}}^s(n, c)$. That is, there exists a Hilbert space \mathcal{H} , a unit vector $\eta \in \mathcal{H}$ and mutually commuting POVM's $(E_{v,i})_{i=1}^c$ and $(F_{w,j})_{j=1}^c$ on \mathcal{H} which are a realisation of $(s(e_{v,i}e_{w,j}))_{v,i,w,j}$ additionally satisfying

$$(12) \quad s(e_{v,i}e_{w,j}) = \langle E_{v,i}E_{w,j}\eta, \eta \rangle = \langle F_{w,j}F_{v,i}\eta, \eta \rangle = \langle E_{v,i}F_{w,j}\eta, \eta \rangle.$$

Proof. Applying the Cauchy-Schwarz inequality, for every $v \in V$, we have the following chain of identities and inequalities.

$$\begin{aligned} 1 &= \sum_{i,j=1}^c p(i, j|v, v) = \sum_{i=1}^c p(i, i|v, v) = \sum_{i=1}^c \langle E_{v,i}F_{v,i}\eta, \eta \rangle \\ &= \sum_{i=1}^c \langle F_{v,i}\eta, E_{v,i}\eta \rangle \leq \sum_{i=1}^c \|F_{v,i}\eta\| \|E_{v,i}\eta\| \\ &\leq \left(\sum_{i=1}^c \|F_{v,i}\eta\|^2 \right)^{1/2} \left(\sum_{i=1}^c \|E_{v,i}\eta\|^2 \right)^{1/2} \\ &= \left(\sum_{i=1}^c \langle F_{v,i}\eta, \eta \rangle \right)^{1/2} \left(\sum_{i=1}^c \langle E_{v,i}\eta, \eta \rangle \right)^{1/2} = 1. \end{aligned}$$

Thus, we must have equality throughout. In particular, the equality between the 2nd and 3rd lines implies that the vectors $(\|F_{v,1}\eta\|, \dots, \|F_{v,c}\eta\|)$ and $(\|E_{v,1}\eta\|, \dots, \|E_{v,c}\eta\|)$ are equal. Thus, $\|F_{v,i}\eta\| = \|E_{v,i}\eta\|, v \in V, 1 \leq i \leq c$. On the other hand, the equality on the second line implies that $F_{v,i}\eta = \alpha_i E_{v,i}\eta$, for some $|\alpha_i| = 1, i = 1, \dots, c$. If $E_{v,i}\eta \neq 0$ then

$$\alpha_i E_{v,i}\eta = F_{v,i}^2 \eta = F_{v,i}(\alpha_i E_{v,i}\eta) = \alpha_i E_{v,i}F_{v,i}\eta = \alpha_i E_{v,i}(\alpha_i E_{v,i}\eta) = \alpha_i^2 E_{v,i}\eta,$$

which forces $\alpha_i = 1$. Thus,

$$(13) \quad E_{v,i}\eta = F_{v,i}\eta, \quad v \in V, i = 1, \dots, c.$$

and so (i) holds.

To prove (ii), note that, by condition (i) of Definition 5.2, we have

$$p(i, j|v, w) = \langle E_{v,i}F_{w,j}\eta, \eta \rangle = \langle E_{v,i}E_{w,j}\eta, \eta \rangle.$$

Using condition (iii) of Definition 5.2, we have

$$\langle E_{v,i}F_{w,j}\eta, \eta \rangle = \langle F_{w,j}E_{v,i}\eta, \eta \rangle = \langle F_{w,j}F_{v,i}\eta, \eta \rangle.$$

Finally,

$$p(j, i|w, v) = \langle E_{w,j}E_{v,i}\eta, \eta \rangle = \langle \eta, E_{w,j}E_{v,i}\eta \rangle = \langle E_{v,i}E_{w,j}\eta, \eta \rangle = p(i, j|v, w).$$

Combining (i) with commutativity we have that

$$E_{v_1, i_1}E_{v_2, i_2}\eta = E_{v_1, i_1}F_{v_2, i_2}\eta = F_{v_2, i_2}E_{v_1, i_1}\eta = F_{v_2, i_2}F_{v_1, i_1}\eta.$$

Proceeding inductively, we have the following word reversal:

$$(14) \quad E_{v_1, i_1}E_{v_2, i_2} \cdots E_{v_k, i_k}\eta = F_{v_k, i_k} \cdots F_{v_2, i_2}F_{v_1, i_1}\eta.$$

To prove (iii), let W be an operator that is a product of elements of the set $\{E_{v,i} : v \in V, i = 1, \dots, c\}$; then

$$\begin{aligned} s(E_{v,i}W) &= \langle E_{v,i}W\eta, \eta \rangle = \langle W\eta, E_{v,i}\eta \rangle = \langle W\eta, F_{v,i}\eta \rangle \\ &= \langle F_{v,i}W\eta, \eta \rangle = \langle WF_{v,i}\eta, \eta \rangle = \langle WE_{v,i}\eta, \eta \rangle = s(WE_{v,i}). \end{aligned}$$

Thus, we have

$$\begin{aligned} s((E_{v_1, i_1}E_{v_2, i_2})W) &= s(E_{v_1, i_1}(E_{v_2, i_2}W)) = s(E_{v_2, i_2}(WE_{v_1, i_1})) \\ &= s(WE_{v_1, i_1}E_{v_2, i_2}). \end{aligned}$$

The general case follows by induction and the fact that the linear combinations of the words on the set $\{E_{v,i} : v \in V, i = 1, \dots, c\}$ are dense in the C^* -algebra generated by this set.

The proof that s is a tracial state on the C^* -algebra generated by the set $\{F_{w,j} : w \in V, j = 1, \dots, c\}$ is identical.

Finally, assume that we have a unital C^* -algebra \mathcal{A} , a tracial state s , and projections $e_{v,i}$ as above. It is clear that $s(e_{v,i}e_{v,j}) = 0$ whenever $v \in V$ and $i \neq j$; thus, $(p(i, j|v, w))_{v, i, w, j}$ is synchronous. Without loss of generality, we can assume that \mathcal{A} is generated by the set $\{e_{v,i} : v \in V, i = 1, \dots, c\}$. The GNS construction associated with (\mathcal{A}, s) , produces a Hilbert space \mathcal{H} , a unital $*$ -homomorphism $\pi : \mathcal{A} \rightarrow B(\mathcal{H})$ and a unit vector $\eta \in \mathcal{H}$ such that $s(X) = \langle \pi(X)\eta, \eta \rangle$, $X \in \mathcal{A}$. Set $E_{v,i} = \pi(e_{v,i})$, $v \in V$, $i = 1, \dots, c$. Since these operators are the images of projections that sum to 1, they form a PVM. By construction, \mathcal{H} is the cyclic subspace corresponding to η . Thus, every vector in \mathcal{H} can be approximated by a sum of the form

$$\sum_{r=1}^k W_r \eta,$$

where W_1, \dots, W_r are words on $\{E_{v,i} : v \in V, i = 1, \dots, c\}$.

Fix $v \in V$ and $j \in \{1, \dots, c\}$. Using the facts that s is a tracial state and that $E_{v,j}^* = E_{v,j}$, we have that

$$\begin{aligned} \left\| \sum_{r=1}^k W_r E_{v,j} \eta \right\|^2 &\leq \left\| \sum_{r=1}^k W_r E_{v,j} \eta \right\|^2 + \left\| \sum_{r=1}^k W_r (I - E_{v,j}) \eta \right\|^2 \\ &= \sum_{r,l=1}^k \langle E_{v,j}^* W_l^* W_r E_{v,j} \eta, \eta \rangle + \sum_{r,l=1}^k \langle (I - E_{v,j})^* W_l^* W_r (I - E_{v,j}) \eta, \eta \rangle \\ &= \sum_{r,l=1}^k \langle E_{v,j} W_l^* W_r \eta, \eta \rangle + \sum_{r,l=1}^k \langle (I - E_{v,j}) W_l^* W_r \eta, \eta \rangle \\ &= \sum_{r,l=1}^k \langle W_l^* W_r \eta, \eta \rangle = \left\| \sum_{r=1}^k W_r \eta \right\|^2. \end{aligned}$$

Thus, the operator $F_{v,j}$ on \mathcal{H} given by

$$F_{v,j} \left(\sum_{r=1}^k W_r \eta \right) = \sum_{r=1}^k W_r E_{v,j} \eta$$

is a well-defined contraction.

Using that the $E_{v,j}$'s form a PVM, it follows that $F_{v,j}^2 = F_{v,j} = F_{v,j}^*$ and $\sum_{j=1}^c F_{v,j} = I$, i.e., the $F_{v,j}$'s also form a PVM. Clearly, $F_{v,j} \eta = E_{v,j} \eta$. Also,

$$F_{v,j} E_{w,i} (W \eta) = E_{w,i} (W E_{v,j} \eta) = E_{w,i} (F_{v,j} W \eta)$$

whenever W is a word on $\{E_{v,i} : v \in V, i = 1, \dots, c\}$, which shows that $F_{v,j} E_{w,i} = E_{w,i} F_{v,j}$.

The fact that $E_{v,i} F_{w,j} = F_{w,j} E_{v,i}$ easily implies the relations (12). \square

Corollary 5.6. *A correlation $(p(i, j|v, w))_{v,i,w,j}$ belongs to $C_{\text{qc}}^s(n, c)$ (resp. $C_{\text{q}}^s(n, c)$, resp. $C_{\text{loc}}^s(n, c)$) if and only if there exists a C^* -algebra (resp. finite dimensional C^* -algebra \mathcal{A} , resp. abelian C^* -algebra) \mathcal{A} , a tracial state $s : \mathcal{A} \rightarrow \mathbb{C}$ and a generating family $\{e_{v,i} : v \in V, i = 1, \dots, c\}$ of projections satisfying $\sum_{i=1}^c e_{v,i} = 1$, $v \in V$, such that*

$$p(i, j|v, w) = s(e_{v,i} e_{w,j}), \quad v, w \in V, i, j = 1, \dots, c.$$

Proof. The statement concerning $C_{\text{qc}}^s(n, c)$ is immediate from Theorem 5.5. For the second equivalence, notice that if a synchronous correlation belongs to $C_{\text{q}}^s(n, c)$ then it admits a realisation $\{(E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta\}$ for which \mathcal{H} is finite dimensional. Thus, the C^* -algebra generated by $\{E_{v,i} : v \in V, i = 1, \dots, c\}$ is finite dimensional. Conversely, if \mathcal{A} is a finite dimensional C^* -algebra and $s : \mathcal{A} \rightarrow \mathbb{C}$ is any state, then the GNS construction yields a finite dimensional Hilbert space. Thus, the operators $E_{v,i}$ and $F_{w,j}$ from Theorem 5.5 act on a finite dimensional Hilbert space, and the claim now follows from Theorem 5.3.

Finally, if a synchronous correlation belongs to $C_{\text{loc}}(n, c)$, then it has a realisation such that the C^* -algebra generated by $\{E_{v,i} : v \in V, i = 1 \dots c\}$ is abelian. Conversely, if \mathcal{A} is abelian and s is any state, then the GNS construction yields an abelian family of projections $\{E_{v,i}\}$ and one can set $F_{v,i} = E_{v,i}$. \square

Let $\{(E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta\}$ be a realisation of a synchronous correlation $(p(i, j|v, w))_{v,i,w,j}$. Let \mathcal{H}_0 be the smallest closed subspace of \mathcal{H} containing η and invariant under the operators $F_{w,j}$, $w \in V$, $j = 1, \dots, c$. Since $F_{w,j}$ is selfadjoint, it is reduced by \mathcal{H}_0 . Thus, $F_{w,j}$ has a diagonal matrix form with respect to the decomposition $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_0^\perp$. Moreover, since $F_{w,j}$ is a projection, the operator $F_{w,j}^0 = F_{w,j}|_{\mathcal{H}_0}$ is a projection and $\sum_{j=1}^c F_{w,j}^0 = I_{\mathcal{H}_0}$, i.e., $(F_{w,j}^0)_{j=1}^c$ is a PVM on \mathcal{H}_0 for each $w \in V$.

By equation (14), \mathcal{H}_0 reduces the operators $E_{v,i}$. Hence, setting $E_{v,i}^0 = E_{v,i}|_{\mathcal{H}_0}$, we have that $(E_{v,i}^0)_{i=1}^c$ is a PVM on \mathcal{H}_0 ; moreover,

$$E_{v,i}^0 F_{w,j}^0 = F_{w,j}^0 E_{v,i}^0, \quad v, w \in V, i, j = 1, \dots, c.$$

Thus, all the properties of (3) are satisfied for the new family of operators, but in addition η is cyclic for the C^* -algebra generated by $\{F_{w,j} : w \in V, j = 1, \dots, c\}$.

Definition 5.7. *Given a synchronous correlation $(p(i, j|v, w)) \in C_{\text{qc}}^s(n, c)$, we call a realisation $\{(E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta\}$ minimal if η is a cyclic vector for the C^* -algebra generated by the family $\{F_{w,j} : w \in V, j = 1, \dots, c\}$.*

Given a graph $G = (V, E)$, a collection $\{(E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta\}$ will be called a c -realisation of G provided that it is a realisation of a synchronous correlation $(p(i, j|v, w))_{v,i,w,j}$ that belongs to the kernel of the functional $L_{G,c}$. We will refer to the collection as a minimal c -realisation of G provided it is also a minimal realisation.

The discussion preceding Definition 5.7 shows how to obtain a minimal c -realisation from any c -realisation.

Proposition 5.8. *Let $(p(i, j|v, w))_{v,i,w,j} \in C_{\text{qc}}^s(n, c)$ be a synchronous correlation with a minimal realisation $((E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta)$. Then the following are equivalent:*

- (i) $\langle E_{v,i} F_{w,j} \eta, \eta \rangle = 0$,
- (ii) $E_{v,i} E_{w,j} = 0$,
- (iii) $F_{v,i} F_{w,j} = 0$.

If this family is a minimal c -realisation of a graph G on n vertices, then for every edge (v, w) of G we have that

$$E_{v,i} E_{w,i} = F_{v,i} F_{w,i} = 0, \quad i = 1, \dots, c.$$

Proof. We prove the equivalence of (i) and (ii). The equivalence of (i) and (iii) is identical. If (ii) holds then, by (13), $\langle E_{v,i} F_{w,j} \eta, \eta \rangle = \langle E_{v,i} E_{w,j} \eta, \eta \rangle = 0$.

Conversely, if (i) holds, then

$$\begin{aligned} \|E_{v,i}E_{w,j}\eta\|^2 &= \langle E_{v,i}E_{w,j}\eta, E_{v,i}E_{w,j}\eta \rangle = \langle E_{v,i}F_{w,j}\eta, E_{v,i}F_{w,j}\eta \rangle \\ &= \langle E_{v,i}F_{w,j}\eta, \eta \rangle = 0, \end{aligned}$$

since the operators $E_{v,i}$ and $F_{w,j}$ are commuting projections.

Next, for any vector ξ of the form $\xi = F_{w_1,j_1} \cdots F_{w_k,j_k}\eta$, we have that

$$E_{v,i}E_{w,j}\xi = F_{w_1,j_1} \cdots F_{w_k,j_k}E_{v,i}E_{w,j}\eta = 0.$$

Part (ii) now follows by minimality.

Finally, the statement involving graphs follows from the equivalence of (i) and (ii) and the fact that if (v, w) is an edge, then $\langle E_{v,i}F_{w,i}\eta, \eta \rangle = 0$. \square

Note that when $p(i, j|v, w) = \langle E_{v,i}F_{w,j}\eta, \eta \rangle$, then

$$\sum_j p(i, j|v, w) = \langle E_{v,i}\eta, \eta \rangle := p_A(i|v)$$

is independent of w and represents the marginal probability that Alice produces outcome i given input v . Similarly, $\sum_i p(i, j|v, w) = \langle F_{w,j}\eta, \eta \rangle := p_B(j|w)$ represents the marginal probability of Bob producing outcome j given input w .

Proposition 5.9. *Let $G = (V, E)$ be a graph on n vertices that admits a c -realisation. Then there exists a minimal c -realisation $((E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta)$ of G such that the marginal probabilities satisfy*

$$(15) \quad \langle E_{v,i}\eta, \eta \rangle = \langle F_{w,j}\eta, \eta \rangle = \frac{1}{c}$$

for every $v, w \in V$ and every $i, j = 1, \dots, c$.

Moreover, if G admits a c -realisation for which the corresponding synchronous correlation is in $C_x(n, c)$, for $x \in \{\text{loc}, \text{q}, \text{qa}\}$, then a minimal c -realisation $((E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta)$ with marginal probabilities equal to $\frac{1}{c}$ can be chosen so that the corresponding synchronous correlation is in $C_x(n, c)$.

Proof. Let $\{(E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta\}$ be a c -realisation of G . Let $\tilde{\mathcal{H}}$ be the direct sum of c copies of \mathcal{H} and set

$$\tilde{E}_{v,i} = E_{v,1+i} \oplus \cdots \oplus E_{v,c+i} \text{ and } \tilde{F}_{w,j} = F_{w,1+j} \oplus \cdots \oplus F_{w,c+j},$$

where the addition in the set of indices is performed modulo c . Set $\tilde{\eta} = \frac{1}{\sqrt{c}}(\eta \oplus \cdots \oplus \eta)$.

It is easy to check that $\{(\tilde{E}_{v,i})_{i=1}^c, (\tilde{F}_{w,j})_{j=1}^c, \tilde{\mathcal{H}}, \tilde{\eta}\}$ is a c -realisation of G . Moreover, for all $v \in V$ and all $i = 1, \dots, c$, we have

$$\langle \tilde{E}_{v,i}\tilde{\eta}, \tilde{\eta} \rangle = \frac{1}{c} \sum_{k=1}^c \langle E_{v,k}\eta, \eta \rangle = \frac{1}{c};$$

similarly,

$$\langle \tilde{F}_{w,j} \tilde{\eta}, \tilde{\eta} \rangle = \frac{1}{c}$$

for all $w \in V$, $j = 1, \dots, c$. The proof is complete after passing to a minimal c -realisation, as described before Definition 5.7.

Suppose that the original synchronous correlation belongs to $C_q(n, c)$, then it has a c -realisation of G whose Hilbert space is finite dimensional. Then the procedure described in the previous two paragraphs yields a finite dimensional Hilbert space, which shows that the graph G admits a c -realisation that satisfies (15) and whose synchronous correlation belongs to $C_q(n, c)$.

If the original correlation belongs to $C_{\text{loc}}(n, c)$, then all PVM's realising the given correlations can be chosen to commute with each other and the described procedure yields a commuting family of operators, and hence the claim follows.

Finally, suppose that the original synchronous correlation of a c -realisation of G belongs to $C_{\text{qa}}(n, c)$. Let $(p_k)_{k \in \mathbb{N}}$ be a sequence of correlations that belong to $C_q(n, c)$ such that $\lim_k p_k(i, j|v, w) = p(i, j|v, w)$ for all i, j, v, w . By the above construction, the correlations, defined by

$$\tilde{p}_k(i, j|v, w) = \frac{1}{c} \sum_{l=1}^c p_k(i+l, j+l|v, w),$$

belong to $C_q(n, c)$ and have constant marginals. Moreover, $\lim_k \tilde{p}_k(i, j|v, w) = \frac{1}{c} \sum_{l=1}^c p(i, j|v, w) := \tilde{p}(i, j|v, w)$. Thus, $\tilde{p} \in C_{\text{qa}}(n, c)$ has constant marginals. Finally, the fact that $L_{G,c}(p) = 0$ implies that $L_{G,c}(\tilde{p}) = 0$. \square

D. Roberson and L. Mančinska [26] define the *projective rank* $\xi_f(G)$ of a graph G to be the infimum of the numbers $\frac{d}{r}$ such that there exists a Hilbert space of (finite) dimension d and projections E_v , $v \in V$, all of rank r , such that $E_v E_w = 0$ whenever (v, w) is an edge of G ; such a collection is called a *d/r -projective representation* of G . Recall that the functional $\text{tr}(X) = \frac{1}{d} \text{Tr}(X)$, where Tr is the usual trace on M_d , is the unique tracial state on M_d , and that if E_v is a projection of rank r then $\text{tr}(E_v) = \frac{r}{d}$. Thus, $\xi_f(G)^{-1}$ is the supremum of quantities of the form $s(E_v)$, over a set of tracial states s of matrix algebras. This viewpoint motivates the following definition.

Definition 5.10. *Let $G = (V, E)$ be a graph. We define the tracial rank $\xi_{\text{tr}}(G)$ of G to be the reciprocal of the supremum of the set of real numbers u for which there exists a unital C^* -algebra \mathcal{A} , a tracial state s on \mathcal{A} and projections $e_v \in \mathcal{A}$, $v \in V$, such that $e_v e_w = 0$ whenever $(v, w) \in E$ and $s(e_v) = u$ for every $v \in V$.*

Proposition 5.11. *Let G be a graph. Then $\xi_f(G)$ is equal to the reciprocal of the supremum of the set of real numbers u for which there exists a finite dimensional C^* -algebra \mathcal{A} , a tracial state s on \mathcal{A} and projections $e_v \in \mathcal{A}$,*

$v \in V$, such that $e_v e_w = 0$ whenever $(v, w) \in E$ and $s(e_v) = u$ for every $v \in V$.

Proof. Let \mathcal{U} be the set of all positive real numbers u for which there exists a finite dimensional C*-algebra \mathcal{A} , a tracial state s on \mathcal{A} and projections $e_v \in \mathcal{A}$, $v \in V$, such that $e_v e_w = 0$ whenever $(v, w) \in E$ and $s(e_v) = u$ for every $v \in V$. Set $U = \sup \mathcal{U}$. By the paragraph preceding Definition 5.10, we see that each r/d appearing in the definition of $\xi_f(G)$ is in \mathcal{U} , and hence $\xi_f(G)^{-1} \leq U$.

Let $u \in \mathcal{U}$ and \mathcal{A} be a finite dimensional C*-algebra as in the previous paragraph. Then \mathcal{A} is *-isomorphic to a direct sum of matrix algebras, say, $\mathcal{A} \cong \sum_{l=1}^L \oplus M_{d_l}$, and every tracial state on \mathcal{A} has the form

$$s\left(\bigoplus_{l=1}^L X_l\right) = \sum_{l=1}^L p_l \operatorname{tr}(X_l),$$

for some $p_l \geq 0$ with $\sum_{l=1}^L p_l = 1$. Set $q_l = \frac{p_l}{d_l}$, $l = 1, \dots, L$.

Each projection e_v is of the form $e_v = \bigoplus_{l=1}^L e_v^l$, where e_v^l is a projection in M_{d_l} , and

$$u = \sum_{l=1}^L p_l \frac{\operatorname{rank}(e_v^l)}{d_l} = \sum_{l=1}^L q_l \operatorname{rank}(e_v^l).$$

Moreover, $\sum_{l=1}^L q_l d_l = 1$.

Let

$$u' = \max\{t : \text{there exist } q_l \geq 0, l = 1, \dots, L, \text{ such that } \sum_{l=1}^L q_l d_l = 1$$

$$\text{and } \sum_{l=1}^L q_l \operatorname{rank}(e_v^l) = t \text{ for all } v \in V\}.$$

By the previous paragraph, $u \leq u'$.

Since the coefficients of the constraint equations are all integers, the maximum u' will be attained at an L -tuple (q_1, \dots, q_L) whose entries are rational. Writing $q_l = m_l/d$ for some integers d and m_l , $l = 1, \dots, L$, and setting

$$e'_v = \bigoplus_{l=1}^L e_v^l \otimes I_{m_l}$$

we obtain a set of projection matrices of size

$$\sum_{l=1}^L m_l d_l = d$$

satisfying the required relations and such that

$$\operatorname{rank}(e'_v) = \operatorname{Tr}(e'_v) = d \sum_{l=1}^L \frac{m_l}{d} \operatorname{rank}(e_v^l) = du'.$$

Hence, $u' \leq \xi_f(G)^{-1}$ and it follows that $U \leq \xi_f(G)^{-1}$ so that the proof is complete. \square

The following is the analogue of the inequality $\xi_f(G) \leq \chi_q(G)$ established in [26].

Theorem 5.12. *Let G be a graph. Then $\xi_{\text{tr}}(G) \leq \chi_{\text{qc}}(G)$.*

Proof. Given any c -realisation of G , Proposition 5.9 shows that there exists a c -realisation $((E_{v,i})_{i=1}^c, (F_{w,j})_{j=1}^c, \mathcal{H}, \eta)$ of G such that $\langle E_{v,i}\eta, \eta \rangle = c^{-1}$ for all $v \in V$ and all $i = 1, \dots, c$. By Proposition 5.8, $E_{v,1}E_{w,1} = 0$ when (v, w) is an edge of G . Thus, $c^{-1} \leq \xi_{\text{tr}}(G)^{-1}$ and the proof is complete. \square

6. GRAPH HOMOMORPHISMS AND PROJECTIVE RANKS

Recall that we set $C_{\text{loc}}(n, c) = \text{Loc}(n, c)$, $C_{\text{q}}(n, c) = Q(n, c)$ and $C_{\text{qa}}(n, c) = Q(n, c)^-$. Then for $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$ and a graph G on n vertices, we have that $\chi_x(G) \leq c$ if and only if there exists $A \in C_x(n, c)$ such that $L_{G,c}(A) = 0$.

The condition $L_{G,c}((p(i, j|v, w))_{v,i,w,j}) = 0$ can more compactly be written as

$$p(i = j|v = w) = 1 \text{ and } p(i = j|v \sim w) = 0,$$

where $p(i = j|v \sim w) = 0$ means that $p(i = j|v, w) = 0$ whenever $(v, w) \in E(G)$. If we write $p(i, j|v, w) = \langle E_{v,i}F_{w,j}\eta, \eta \rangle$, where $(E_{v,i})_{i=1}^c$ and $(F_{w,j})_{j=1}^c$ are mutually commuting PVM's on a Hilbert space \mathcal{H} , $v, w \in V$, and $\eta \in \mathcal{H}$ is a unit vector, then we have that

$$p_A(i|v) := \sum_{j=1}^c p(i, j|v, w) = \langle E_{v,i}\eta, \eta \rangle$$

does not depend on w and j ; a similar statement holds for $p_B(j|w)$.

Remark 6.1. In the notation introduced above, Proposition 5.9 shows that, for any $x \in \{\text{loc}, \text{q}, \text{qa}, \text{c}\}$, if G is a graph on n vertices and the correlation $(p(i, j|v, w))_{v,i,w,j} \in C_x(n, c)$ satisfies $p(i = j|v = w) = 1$ and $p(i = j|v \sim w) = 0$, then there is a correlation $(p'(i, j|v, w))_{v,i,w,j} \in C_x(n, c)$ additionally satisfying $p'_A(i|v) = p'_B(j|w) = c^{-1}$ for every v, w, i, j .

We recall the following characterization of points in $C_{\text{loc}}(n, c)$:

(16)

$$C_{\text{loc}}(n, c) = \text{Loc}(n, c)$$

$$(17) \quad = \left\{ (p(i, j|v, w))_{v,i,w,j} : p(i, j|v, w) = \sum_k \lambda_k \delta(i = f_k(v)) \delta(j = g_k(w)), \right.$$

$$(18) \quad \left. \text{for some } \lambda_k > 0 \text{ with } \sum_k \lambda_k = 1 \text{ and some } f_k, g_k : V \rightarrow \{1, \dots, c\} \right\}.$$

(Here, the δ function evaluates to 1 when its condition argument is true and 0 otherwise, like the Iverson bracket.)

Definition 6.2. Let $G = (V(G), E(G))$ and $H = (V(H), E(H))$ be graphs on n and m vertices, respectively. For $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$ write $G \xrightarrow{x} H$ if there is a correlation $(p(i, j|v, w))_{v, i, w, j} \in C_x(n, m)$ with $v, w \in V(G)$ and $i, j \in V(H)$ such that

$$\begin{aligned} p(i = j|v = w) &= 1 \\ p(i \sim_H j|v \sim_G w) &= 1, \end{aligned}$$

where $p(i \sim_H j|v, w) := \sum_{(i, j) \in E(H)} p(i, j|v, w)$ and $p(i \sim_H j|v \sim_G w) = 1$ means that $p(i \sim_H j|v, w) = 1$ whenever $(v, w) \in E(G)$.

We will say that such a $(p(i, j|v, w))_{v, i, w, j}$ is an x -homomorphism from G to H .

Stated briefly, the above conditions are the requirement that p be synchronous and that whenever the inputs v and w are adjacent in G , then, with probability 1, the output pair (i, j) is adjacent in H .

We will sometimes write $G \rightarrow H$ for $G \xrightarrow{\text{loc}} H$, since it can be shown that this corresponds to the classical definition of a graph homomorphism. The homomorphism variant $G \xrightarrow{\text{q}} H$ has been extensively studied in [25] and [26]. The following is immediate from the definitions of [24] and [3]:

We let K_c denotes the complete graph on c vertices, i.e., $(i, j) \in E(K_c)$ for all $i \neq j$.

Proposition 6.3. Let G be a graph. For $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$, we have that $\chi_x(G) = \min\{c : G \xrightarrow{x} K_c\}$.

Let us denote by \overline{G} the complementary graph of G , that is, the graph whose vertex set coincides with that of G and for which (v, w) is an edge precisely when (v, w) is not an edge of G (here it is assumed that $v \neq w$). Proposition 6.3 motivates us to define, for $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$,

$$\alpha_x(\overline{G}) = \omega_x(G) = \max\{c : K_c \xrightarrow{x} G\}.$$

The parameters $\omega_x(G)$ are *quantum clique numbers* of G and are complementary to the corresponding chromatic numbers $\chi_x(G)$. They will not be used later on in this paper. Note, however, that $\omega_{\text{loc}}(G)$ (resp. $\alpha_{\text{loc}}(G)$) coincides with the classical clique number $\omega(G)$ (resp. independence number $\alpha(G)$) of G .

Definition 6.4. Let G be a graph on n vertices. For $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$, let $\xi_x(G)$ be the infimum of the positive real numbers t such that there exists $(p(a, b|v, w))_{v, a, w, b} \in C_x(n, 2)$ satisfying

$$\begin{aligned} p(a = b|v = w) &= 1 \\ p(a = 1, b = 1|v \sim w) &= 0 \\ p(a = 1|v) &= t^{-1}. \end{aligned}$$

Note that it makes sense, in the above definition, to use only v in the third condition since $C_x(n, 2)$ is non-signaling.

Suppose that

$$p' = (p'(i, j|v, w))_{v, i, w, j} \in C_{\text{qc}}(n, c) \quad \text{and} \quad p'' = (p''(a, b|i, j))_{i, a, j, b} \in C_{\text{qc}}(c, l).$$

We let $p''p' = ((p''p')(a, b|v, w))_{v, a, w, b}$ be the matrix whose entries are given by

$$(p''p')(a, b|v, w) = \sum_{i, j=1}^c p''(a, b|i, j)p'(i, j|v, w).$$

Thus, if p' (resp. p'') is considered as an element of M_{c^2, n^2} (resp. M_{l^2, c^2}), whose rows are indexed by the pairs (a, b) (resp. (i, j)) and whose columns – by the pairs (i, j) (resp. (v, w)), then $p''p'$ is the matrix product of p'' and p' .

Lemma 6.5. *Let $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$.*

- (i) *If $p' \in C_x(n, c)$ and $p'' \in C_x(c, l)$ then $p''p' \in C_x(n, l)$;*
- (ii) *If $p' \in C_x(n, c)$ and $p'' \in C_{\text{loc}}(c, l)$ then $p''p' \in C_x(n, l)$.*

Proof. (i) Assume first that $x = \text{qc}$. Suppose that \mathcal{H}' (resp. \mathcal{H}'') is a Hilbert space, $\eta \in \mathcal{H}'$ (resp. $\eta \in \mathcal{H}''$) is a unit vector and $(E'_{v,i})_{i=1}^c$ and $(F'_{w,j})_{j=1}^c$ (resp. $(E''_{i,a})_{a=1}^l$ and $(F''_{j,b})_{b=1}^l$) are mutually commuting PVM's such that

$$p'(i, j|v, w) = \langle E'_{v,i} F'_{w,j} \eta', \eta' \rangle \quad (\text{resp. } p''(a, b|i, j) = \langle E''_{i,a} F''_{j,b} \eta'', \eta'' \rangle),$$

for all v, w, i, j, a, b . Let $\mathcal{H} = \mathcal{H}'' \otimes \mathcal{H}'$, $\eta = \eta'' \otimes \eta'$,

$$E_{v,a} = \sum_{i=1}^c E''_{i,a} \otimes E'_{v,i} \quad \text{and} \quad F_{w,b} = \sum_{j=1}^c F''_{j,b} \otimes F'_{w,j}.$$

It is clear that $(E_{v,a})_{a=1}^l$ and $(F_{w,b})_{b=1}^l$ are mutually commuting POVM's for all v and w . Moreover,

$$\begin{aligned} \langle E_{v,a} F_{w,b} \eta, \eta \rangle &= \sum_{i, j=1}^c \langle (E''_{i,a} \otimes E'_{v,i})(F''_{j,b} \otimes F'_{w,j})(\eta'' \otimes \eta'), (\eta'' \otimes \eta') \rangle \\ &= \sum_{i, j=1}^c \langle E''_{i,a} F''_{j,b} \eta'', \eta'' \rangle \langle E'_{v,i} F'_{w,j} \eta', \eta' \rangle \\ &= \sum_{i, j=1}^c p''(a, b|i, j)p'(i, j|v, w) = (p''p')(a, b|v, w). \end{aligned}$$

It follows that $p''p' \in C_{\text{qc}}(n, l)$.

The arguments given above also apply in the case $x = \text{q}$. The claim concerning $x = \text{qa}$ follows from the fact that $C_{\text{qa}}(n, c) = \overline{C_{\text{q}}(n, c)}$ for all n and c . The case $x = \text{loc}$ follows from the observation preceding Proposition 2.3.

- (ii) follows from (i) and the fact that $C_{\text{loc}}(c, l) \subseteq C_x(c, l)$. \square

Theorem 6.6. *For $x \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$, we have that $\xi_x(G) \leq \chi_x(G)$. Moreover, if $G \xrightarrow{x} H$ then $\xi_x(G) \leq \xi_x(H)$.*

Proof. Let $(p(i, j|v, w))_{v, i, w, j}$ be an x -homomorphism from G to K_c with $c = \chi_x(G)$. By Proposition 5.9, we may assume that $p(i|v) = \frac{1}{c}$ for all i and all v . Let $p'_A(a|i)$ (resp. $p'_B(b|j)$) be the probability distribution given by $p'_A(1|1) = 1, p'_A(0|1) = 0$ (resp. $p'_B(1|1) = 1, p'_B(0|1) = 0$) and $p'_A(1|i) = 0, p'_A(0|i) = 1$ (resp. $p'_B(1|j) = 0, p'_B(0|j) = 1$) if $i \neq 1$ (resp. $j \neq 1$). Set

$$p'(a, b|i, j) = p'_A(a|i)p'_B(b|j), \quad a, b = 0, 1, i, j = 1, \dots, c.$$

It is clear that $p' \in C_x(c, 2)$. By Lemma 6.5, $p'p \in C_x(n, 2)$. It remains to check that $p'p$ satisfies the conditions of Definition 6.4.

Suppose that $a \neq b$. If $i = j$ then $p'(a, b|i, j) = 0$, while if $i \neq j$ then $p(i, j|v, v) = 0$. It follows that $(p'p)(a, b|v, v) = 0$ for all v . Suppose that $v \sim w$. Then $p(i, i|v, w) = 0$ for all i , while $p'(1, 1|i, j) = 0$ if $i \neq j$. It follows that $(p'p)(1, 1|v, w) = 0$. Finally, for fixed v , we have

$$\begin{aligned} (p'p)(1|v) &= \sum_{i, j=1}^c p'(1, 0|i, j)p(i, j|v, v) + p'(1, 1|i, j)p(i, j|v, v) \\ &= \sum_{i, j=1}^c p'_A(1|i)p'_B(0|j)p(i, j|v, v) + p'_A(1|i)p'_B(1|j)p(i, j|v, v) \\ &= \sum_{j=1}^c p'_B(0|j)p(1, j|v, v) + p'_B(1|j)p(1, j|v, v) \\ &= \sum_{j=1}^c p(1, j|v, v) = \frac{1}{c}. \end{aligned}$$

We now show the monotonicity of ξ_x . Suppose that $G \xrightarrow{x} H$ and let

$$(p(i, j|v, w))_{v, i, w, j} \in C_x(|V(G)|, |V(H)|)$$

be as in Definition 6.2. Let also $(p'(a, b|i, j))_{i, a, j, b} \in C_x(|V(H)|, 2)$ satisfy the three equations of Definition 6.4 for the graph H . Suppose that $v \in V(G)$ and $a \neq b$. Then, if $i \neq j$ we have that $p(i, j|v, v) = 0$, while $p'(a, b|i, i) = 0$. Thus, $(p'p)(a, b|v, v) = 0$. Suppose that $(v, w) \in E(G)$. If $(i, j) \notin E(H)$ then $p(i, j|v, w) = 0$, while if $(i, j) \in E(H)$ then $p'(1, 1|i, j) = 0$; thus, $(p'p)(1, 1|v, w) = 0$. Finally,

$$\begin{aligned} (p'p)(1|v) &= \sum_{i, j=1}^c p'(1, 0|i, j)p(i, j|v, v) + p'(1, 1|i, j)p(i, j|v, v) \\ &= \sum_{i=1}^c p'(1, 0|i, i)p(i, i|v, v) + p'(1, 1|i, i)p(i, i|v, v) \\ &= \sum_{i=1}^c p'(1, 1|i, i)p(i, i|v, v) = \frac{1}{t} \sum_{i=1}^c p(i, i|v, v) = \frac{1}{t}, \end{aligned}$$

for all $v \in V(G)$. By Lemma 6.5, $p'p \in C_x(|V(G)|, 2)$. Thus, $\xi_x(G) \leq \xi_x(H)$. \square

Lemma 6.7. *Suppose that $p(i, j|x, y) \in C_q(n, 2)$ (respectively, $C_{qc}(n, 2)$) satisfies $p(i = j|x = y) = 1$. Then there exist a finite dimensional C^* -algebra \mathcal{A} (resp. a C^* -algebra \mathcal{A}), a tracial state $s : \mathcal{A} \rightarrow \mathbb{C}$ and projections $E_{v,i} \in \mathcal{A}$, $v \in V$, $i = 1, 2$, such that*

- (1) $\sum_i E_{v,i} = I$, $v \in V$;
- (2) $p(i, j|v, w) = s(E_{v,i}E_{w,j})$ for all $v, w \in V$ and all $i, j = 1, 2$;
- (3) $E_{v,i}E_{w,j} = 0$ if and only if $p(i, j|v, w) = 0$.

Proof. The existence of the C^* -algebra, tracial state and corresponding operators follow from the fact that the state is synchronous, Corollary 5.6 and Proposition 5.8. \square

A graph G is said to have an a/b -coloring provided that to each vertex we can assign a b element subset of $\{1, \dots, a\}$ such that whenever two vertices are adjacent, their corresponding subsets are disjoint. The *fractional chromatic number* of G is then defined by $\chi_f(G) = \inf\{a/b \mid G \text{ has an } a/b\text{-coloring}\}$. Alternatively, there is a family of graphs known as the *Kneser graphs* $K(a, b)$, where each vertex corresponds to a b element subset of an a element set with vertices adjacent when the sets are disjoint, and $\chi_f(G) = \inf\{a/b : G \rightarrow K(a, b)\}$. For more discussion of these ideas and proofs see [13].

Theorem 6.8. *We have that*

- (1) $\xi_{loc}(G)$ is equal to the fractional chromatic number $\chi_f(G)$;
- (2) $\xi_q(G)$ is equal to the projective rank, $\xi_f(G)$;
- (3) $\xi_{qc}(G)$ is equal to the tracial rank, $\xi_{tr}(G)$.

Proof. To prove $\xi_{loc}(G) = \chi_f(G)$, colour the graph G with subsets $S_v \subseteq \{1, \dots, p\}$ of size $|S_v| = q$ where $p/q = \chi_f(G)$ (this is possible since $\chi_f(G)$ can be interpreted in terms of homomorphisms to Kneser graphs). Consider the following protocol in $C_{loc}(n, 2)$: Alice and Bob receive vertices v and w . They use shared randomness to choose $k \in \{1, \dots, p\}$. Alice outputs 1 if $k \in S_v$ while Bob outputs 1 if $k \in S_w$. The corresponding correlation satisfies the conditions of Definition 6.4 with $t = p/q$, so $\xi_{loc}(G) \leq \chi_f(G)$.

Conversely, suppose that $p(a, b|x, y) \in C_{loc}(n, 2)$ satisfies the conditions of Definition 6.4 for some t . By (16), we have $p(a, b|v, w) = \sum_k \lambda_k \delta(a = f_k(v))\delta(b = g_k(w))$ with $\lambda_k > 0$ and $\sum_k \lambda_k = 1$. The condition $p(a = b|v = w) = 1$ requires $f_k = g_k$ for all k . The condition $p(a = 1, b = 1|v \sim w) = 0$ guarantees that $f_k(v)f_k(w) = 0$ for all k when $v \sim w$; consequently $V_k := \{v \in V(G) : f_k(v) = 1\}$ is an independent set. Assigning weight $t\lambda_k$ to set V_k gives a fractional colouring of weight t (see Section 7.1 of [13]). Indeed, $\sum_k t\lambda_k = t$ and for each $v \in V(G)$ we have $\sum_{V_k \ni v} t\lambda_k = t \sum_k f_k(v)\lambda_k = tp(a = 1|v) = 1$. So $\chi_f(G) \leq \xi_{loc}(G)$.

To prove $\xi_q(G) = \xi_f(G)$, suppose $p(a, b|v, w) \in C_q(n, 2)$ satisfies the conditions of Definition 6.4. In particular it is synchronous. Let $\{E_{v,i} : i =$

$1, 2, v \in V\} \in M_d$ be the representation guaranteed by 6.7. The operators $E_{v,1}$ then satisfy the conditions in Proposition 5.11 and hence, referring to the proof, $t^{-1} \in \mathcal{U}$. Taking the infimum over all possible t gives $\xi_f(G) \leq \xi_q(G)$.

Conversely, suppose that $(E_v)_{v \in V}$ is a d/r -projective representation of G . Let $\eta = d^{-1/2} \sum_i e_i \otimes e_i$, where the $\{e_i\}_{i=1}^d$ is the standard orthonormal basis and set $E_{v,1} = E_v$, $E_{v,0} = (I - E_v)$, $F_{w,1} = \overline{E}_w$, and $F_{w,0} = (I - \overline{E}_w)$. The probability distribution $p(a, b|v, w) = \langle E_{v,a} \otimes F_{w,b} \eta, \eta \rangle$ is feasible for 6.4 with value $t^{-1} = p(a = 1|v) = \langle E_{v,1} \eta, \eta \rangle = d^{-1} \text{Tr}(E_v) = r/d$. So $\xi_q(G) \leq d/r$. Taking the infimum over possible values of d/r gives $\xi_q(G) \leq \xi_f(G)$.

Finally, we prove $\xi_{qc}(G) = \xi_{tr}(G)$. It follows from Lemma 6.7 that if t is feasible for $\xi_{qc}(G)$ then there exists a C*-algebra and tracial state satisfying the conditions of Definition 5.10. Thus, $\xi_{tr}(G) \leq \xi_{qc}(G)$.

Conversely, assume that we have a C*-algebra \mathcal{A} a tracial state s and projections E_v and a real number $u = t^{-1}$ satisfying the hypotheses of Definition 5.10. We set $E_{v,1} = E_v$ and $E_{v,0} = I - E_v$.

If we set $p(i, j|v, w) = s(E_{v,i} E_{w,j})$ then by Corollary 5.6, we have that $(p(i, j|v, w)) \in C_{qc}^s(n, 2) \subseteq C_{qc}(n, 2)$. Thus t is feasible for $\xi_{qc}(G)$. This shows $\xi_{qc}(G) \leq \xi_{tr}(G)$ and the proof is complete. \square

Theorem 6.9. *If there exists a graph G for which $\xi_f(G)$ is irrational, then the closure conjecture is false, and consequently, Tsirelson's conjecture is false. In fact, if G is a graph on n vertices with $\xi_f(G)$ irrational, then $C_q(n, 2) \neq C_q(n, 2)^-$.*

Proof. Let n be the number of vertices of G . By Theorem 6.8, $\xi_f(G)^{-1} = \xi_q(G)^{-1}$ and this value is characterized as the infimum of the positive real numbers t over the elements $(p(a, b|v, w))_{v,a,w,b} \in C_q(n, 2)$ such that $p(a = 1|v) = t^{-1}$. If this infimum was attained, then there would exist a representation of $(p(a, b|v, w))_{v,a,w,b}$ via a finite dimensional C*-algebra. It follows by the proof of Proposition 5.11 that the infimum is a rational number.

Hence, if the infimum is attained by a point in $C_q(n, 2)$ then it must be rational. Thus, if $\xi_q(G)$ is irrational, then we must have a point in $C_q(n, 2)^-$ that is not in $C_q(n, 2)$. \square

Corollary 6.10. *If there exists a graph G with $\xi_{tr}(G)$ irrational, then Tsirelson's conjecture is false. In fact, if G has n vertices, then $C_{qc}(n, 2) \neq C_q(n, 2)$.*

Proof. If Tsirelson's conjecture is true, then the closure conjecture is true and $\xi_{tr}(G) = \xi_{qc}(G) = \xi_q(G) = \xi_f(G)$, contradicting the previous result. \square

Given that $\xi_{loc}(G) = \chi_f(G)$, $\xi_q(G) = \xi_f(G)$, and $\xi_{qc}(G) = \xi_{tr}(G)$, we will henceforth use the more established notation $\chi_f(G)$, $\xi_f(G)$, $\xi_{tr}(G)$ and will drop the notation $\xi_{loc}(G)$, $\xi_q(G)$, $\xi_{qc}(G)$.

We turn now to a deeper investigation of $\xi_{tr}(G)$. We first show that the equality for a feasible value can be relaxed to an inequality.

Theorem 6.11. *The number $\xi_{\text{tr}}(G)$ is equal to the minimum of the positive real numbers t such that there exist a Hilbert space \mathcal{H} , a unit vector $\eta \in \mathcal{H}$, a (unital) C^* -algebra $\mathcal{A} \subseteq \mathcal{B}(H)$ and projections $E_v \in \mathcal{A}$, $v \in V$, satisfying*

$$(19) \quad \text{the map } X \rightarrow s(X) = \langle X\eta, \eta \rangle \text{ is a tracial state on } \mathcal{A};$$

$$(20) \quad E_v E_w = 0 \text{ if } v \sim w;$$

$$(21) \quad \langle E_v \eta, \eta \rangle \geq t^{-1} \text{ for all } v \in V(G).$$

Proof. Any solution feasible for Definition 5.10 induces a solution feasible for the above conditions, with the same value. Note that, using the GNS construction, we can assume, without loss of generality, that $s(X) = \langle X\eta, \eta \rangle$, $X \in \mathcal{A}$, for some unit vector η .

Conversely, suppose we have a feasible solution to the above conditions. Let $c_v = s(E_v)$, where $c_v \geq t^{-1}$. Set $c = \min\{c_v : v \in V(G)\}$.

Let $\hat{\mathcal{H}} = \mathcal{H} \otimes L^2(0, 1)$, $\hat{\eta} = \eta \otimes \chi_{[0,1]} \in \hat{\mathcal{H}}$ and, for $0 \leq r \leq 1$, let $P_r : L^2(0, 1) \rightarrow L^2(0, 1)$ denote the projection onto the subspace $L^2(0, r)$. Let \mathcal{D} be the multiplication algebra of $L^\infty(0, 1)$ acting on $L^2(0, 1)$, $\hat{E}_v = E_v \otimes P_{c/c_v}$ and $\hat{s}(X) = \langle X\hat{\eta}, \hat{\eta} \rangle$, $X \in \mathcal{A} \otimes \mathcal{D}$.

The state \hat{s} is tracial because it is the tensor product of two tracial states. It is easily verified that this new family of projections \hat{E}_v and state \hat{s} satisfy the conditions of Definition 5.10 with $\hat{s}(E_v) = r^{-1}$ where $r^{-1} = c \geq t^{-1}$, so that $r \leq t$.

Thus, we attain the same infimum if we require equality in (21) for all v . \square

We shall refer to a vector and set of operators satisfying (19)–(21) for some t a *feasible set for $\xi_{\text{tr}}(G)$ with value t* . The following is an adaptation of a proof from [5].

Theorem 6.12. $\xi_{\text{tr}}(G * H) = \xi_{\text{tr}}(G[H]) = \xi_{\text{tr}}(G)\xi_{\text{tr}}(H)$, where $G * H$ is the disjointive product (co-normal product, OR product) and $G[H]$ is the lexicographical product.

Proof. The inequality $\xi_{\text{tr}}(G[H]) \leq \xi_{\text{tr}}(G * H)$ follows from the inclusion $G[H] \subseteq G * H$.

To prove the inequality, $\xi_{\text{tr}}(G * H) \leq \xi_{\text{tr}}(G)\xi_{\text{tr}}(H)$, let η and E_v form a set feasible for $\xi_{\text{tr}}(G)$ with value t and let η' and E'_v form a set feasible for $\xi_{\text{tr}}(H)$ with value t' . Then $\eta \otimes \eta'$ and $E_v \otimes E'_v$ form a set feasible for $\xi_{\text{tr}}(G * H)$ with value tt' .

$\xi_{\text{tr}}(G)\xi_{\text{tr}}(H) \leq \xi_{\text{tr}}(G[H])$: Let η and $E_{g,h}$ be a feasible set for $\xi_{\text{tr}}(G[H])$ with value t , where $g \in V(G)$ and $h \in V(H)$. For $g \in V(G)$ define \tilde{E}_g to be the projection onto the span of the ranges of $\{E_{g,h}\}_{h \in H}$.

If $g \sim g'$ in G then $(g, h) \sim (g', h')$ for every $h, h' \in V(H)$ and hence $E_{g,h}E_{g',h'} = 0$. From this it follows that $g \sim g'$ implies that $\tilde{E}_g\tilde{E}_{g'} = 0$. Recall that \tilde{E}_g is the strong limit of $(\sum_h E_{g,h})^{1/n}$ and hence $X \rightarrow s(X) = \langle X\eta, \eta \rangle$ is a tracial state on the C^* -algebra generated by the \tilde{E}_g (indeed, the

latter C*-algebra is a subalgebra of the von Neumann algebra generated by the set $\{E_{g,h} : g \in G, h \in H\}$. Thus, η and \tilde{E}_g satisfy all the conditions to be a feasible set for G . Set $c_g = s(\tilde{E}_g)$ and let $r^{-1} = \min\{c_g : g \in V(G)\} = c_f$ for some $f \in V(G)$ so that we have a feasible set for G with value r .

Let $\tilde{\eta} = \sqrt{r}\tilde{E}_f\eta$ and $\tilde{s}(X) = \langle \tilde{\eta}, X\tilde{\eta} \rangle = rs(\tilde{E}_fX\tilde{E}_f)$. Then \tilde{s} is a state, tracial on the algebra generated by $\{E_{f,h} : h \in V(H)\}$. Since $\tilde{s}(E_{f,h}) = rs(E_{f,h}) \geq rt^{-1} = (r^{-1}t)^{-1}$, we see that $\tilde{\eta}$ and $\{\tilde{E}_{f,h}\}_{h \in V(H)}$ is a feasible set for $\xi_{\text{tr}}(H)$ with value $r^{-1}t$.

Thus,

$$\xi_{\text{tr}}(G)\xi_{\text{tr}}(H) \leq r \cdot (r^{-1}t) = t,$$

and since t was an arbitrary feasible value for $G[H]$, we have

$$\xi_{\text{tr}}(G)\xi_{\text{tr}}(H) \leq \xi_{\text{tr}}(G[H]).$$

□

7. A SDP LOWER BOUND FOR THE COMMUTING QUANTUM CHROMATIC NUMBER

We now explore a quantity which can be seen either as a semidefinite relaxation of the quantity $\xi_{\text{tr}}(G)$ or as a strengthening of the quantity $\bar{\vartheta}^+(G)$. As before, we assume that $|V(G)| = n$. For a matrix $Y \in M_{n+1}(\mathbb{R})$, we index the first row and column by 0, and the rest – by the elements of the set $V(G)$. Since we are discussing a semidefinite programming problem in this section, we shall use the more familiar $Y \succeq 0$ to indicate that the matrix Y is positive semidefinite.

Definition 7.1. *Define*

$$\begin{aligned} \xi_{\text{SDP}}(G) = \min \left\{ Y_{00} \mid \exists Y \in M_{n+1}(\mathbb{R}), Y \succeq 0, Y_{vw} \geq 0, \right. \\ Y_{0v} = Y_{vv} = 1 \text{ for } v \in V(G), \\ Y_{vw} = 0 \text{ for } v \sim w, \\ \sum_{v \in S} Y_{vw} \leq 1 \text{ for } S \text{ a clique of } G \text{ and } w \in V(G), \\ \left. Y_{00} + \sum_{v \in S} \sum_{w \in T} Y_{vw} \geq |S| + |T| \text{ for } S, T \text{ cliques of } G \right\}. \end{aligned} \tag{22}$$

Remark 7.2. If in the above definition we assumed instead that Y is a complex matrix that satisfied the remaining conditions, then since $Y = Y^*$ we see that $\frac{Y+Y^t}{2}$ is a real matrix that also satisfies these equations and has the same value for the $(0,0)$ -entry. Thus, we obtain the same value for $\xi_{\text{SDP}}(G)$ if we require $Y \in M_{n+1}(\mathbb{R})$ or $Y \in M_{n+1}(\mathbb{C})$.

Theorem 7.3. *We have that $\bar{\vartheta}^+(G) \leq \xi_{\text{SDP}}(G) \leq \xi_{\text{tr}}(G)$.*

Proof. We first show that $\bar{\vartheta}^+(G) \leq \xi_{\text{SDP}}(G)$. Let $\mathbf{e} \in \mathbb{R}^{|V(G)|}$ be the vector all of whose entries are equal to 1, and J be the matrix all of whose entries are equal to 1. Let Y satisfy conditions (22). Then Y has the block form

$$Y = \begin{pmatrix} Y_{00} & \mathbf{e}^T \\ \mathbf{e} & M \end{pmatrix}.$$

By Cholesky's Lemma, $Y \succeq 0$ if and only if $M \succeq Y_{00}^{-1}J$. Define $Z = Y_{00}M - J$; then $Z \succeq 0$, $Z_{vw} \geq -1$ for all $v, w \in V(G)$, $Z_{vw} = -1$ whenever $v \sim w$, and $Z_{vv} = Y_{00} - 1$ for all $v \in V(G)$. So Z is feasible for $\bar{\vartheta}^+(G)$ with value Y_{00} (see [5] for the definition of $\bar{\vartheta}^+(G)$).

We next show that $\xi_{\text{SDP}}(G) \leq \xi_{\text{tr}}(G)$. Let η and E_v , $v \in V(G)$ be as in Theorem 6.11, with $t = \xi_{\text{tr}}(G)$. Note that by the proof of Theorem 6.11 we can assume equality in (21). Set $\eta_v = E_v\eta$, $v \in V(G)$, and $Y \in M_{n+1}(\mathbb{R})$ as $Y_{vw} = t\langle \eta_v, \eta_w \rangle$, where we take $\eta_0 = \eta$. We have that $Y_{vw} = ts(E_v E_w)$, $Y_{0v} = ts(E_v) = 1$, and $Y_{00} = ts(I) = t$.

Conditions (19)–(21), along with the fact that $E_v^2 = E_v$, give

$$\begin{aligned} Y_{vv} &= ts(E_v E_v) = Y_{0v} = 1 \\ Y_{vw} &= ts(E_v E_w) = 0 \text{ for } v \sim w. \end{aligned}$$

Consider a clique S of G . We have that $E_v \perp E_w$ when $v \sim w$, so $I - \sum_{v \in S} E_v$ is a projection. For S a clique of G and $w \in V(G)$,

$$\begin{aligned} 1 - \sum_{v \in S} Y_{vw} &= ts(E_w) - \sum_{v \in S} ts(E_v E_w) \\ &= ts\left(\left(I - \sum_{v \in S} E_v\right)E_w\right) \\ &= ts\left(E_w\left(I - \sum_{v \in S} E_v\right)E_w\right) \geq 0, \end{aligned}$$

where in the last step we used that s is a tracial state. Similarly, for S, T cliques of G ,

$$\begin{aligned} Y_{00} + \sum_{v \in S} \sum_{w \in T} Y_{vw} - |S| - |T| &= ts\left(\left(I - \sum_{v \in S} E_v\right)\left(I - \sum_{w \in T} E_w\right)\right) \\ &= ts\left(\left(I - \sum_{v \in S} E_v\right)\left(I - \sum_{w \in T} E_w\right)\left(I - \sum_{v \in S} E_v\right)\right) \geq 0. \end{aligned}$$

So Y is a feasible solution to (22) with value t . \square

We can now compute the tracial rank of an odd cycle.

Theorem 7.4. $\xi_{\text{tr}}(C_{2k+1}) = \frac{2k+1}{k}$.

Proof. The following is inspired by a proof from [25]. Let Y be feasible for (22). Considering cliques $S = \{1, 2\}$ and $T = \{2, 3\}$, we have

$$(23) \quad Y_{00} + \sum_{v \in \{1,2\}} \sum_{w \in \{2,3\}} Y_{vw} \geq 4 \implies Y_{13} \geq 3 - Y_{00}.$$

Let $n = 2k + 1$. We equip the set $\{1, \dots, n\}$ with addition modulo n . For any $a, b \in \{1, \dots, n\}$, we have

$$\begin{aligned} - \sum_{v \in \{a, a+1\}} Y_{vb} &\geq -1 \implies -Y_{a,b} - Y_{(a+1),b} \geq -1 \\ Y_{00} + \sum_{v \in \{(a+1), (a+2)\}} \sum_{w \in \{b\}} Y_{vw} &\geq 3 \implies Y_{(a+1),b} + Y_{(a+2),b} \geq 3 - Y_{00}. \end{aligned}$$

Adding these inequalities gives

$$Y_{(a+2),b} - Y_{a,b} \geq 2 - Y_{00}.$$

Adding together c instances of this inequality gives $Y_{(a+2c),b} - Y_{a,b} \geq c(2 - Y_{00})$. Taking $a = 3, b = 1, c = k - 1$ gives $Y_{n1} - Y_{31} \geq (k - 1)(2 - Y_{00})$. But $Y_{n1} = 0$ since $1 \sim n$. Adding (23) gives $0 \geq (k - 1)(2 - Y_{00}) + (3 - Y_{00}) = (2k + 1) - kY_{00}$. So $Y_{00} \geq (2k + 1)/k$.

Hence,

$$(2k + 1)/k \leq \xi_{\text{SDP}}(C_k) \leq \xi_{\text{tr}}(C_k) \leq \xi_{\text{loc}}(C_k) = \chi_{\text{f}}(C_k).$$

And $\chi_{\text{f}}(C_k) = (2k + 1)/k$ [25] so the result follows. \square

Note that the preceding proof did not make use of $Y \succeq 0$. So in fact it would even follow from an LP relaxation of the SDP.

Corollary 7.5. *There is a graph G for which $\lceil \bar{\vartheta}^+(G) \rceil < \chi_{\text{qc}}(G)$.*

Proof. By [5, Theorem 17],

$$\lceil \bar{\vartheta}^+(C_5 * K_3) \rceil = \lceil \bar{\vartheta}^+(C_5) \bar{\vartheta}^+(K_3) \rceil = \lceil 3\sqrt{5} \rceil = 7,$$

while

$$7 < \frac{15}{2} = \xi_{\text{tr}}(K_3) \xi_{\text{tr}}(C_5) = \xi_{\text{tr}}(C_5 * K_3) \leq \chi_{\text{qc}}(C_5 * K_3) \leq \chi(C_5 * K_3) = 8.$$

\square

For all graphs G we have

$$\xi_{\text{SDP}}(G) \leq \xi_{\text{tr}}(G) \leq \xi_{\text{f}}(G) \leq \chi_{\text{f}}(G).$$

In the proof of Theorem 7.4 we see that for odd cycles $\xi_{\text{SDP}}(G) = \chi_{\text{f}}(G)$, so this chain of inequalities collapses. Numerical results show that, in fact, $\xi_{\text{SDP}}(G) = \chi_{\text{f}}(G)$ for all graphs on 9 vertices or less. So now we know the value of ξ_{f} on all of these graphs (it is equal to χ_{f}) whereas before the only nontrivial graphs for which this quantity was known were the Kneser graphs and odd cycles.

ACKNOWLEDGEMENT

The authors would like to thank the referee for a diligent reading that improved the exposition considerably.

REFERENCES

- [1] D. Avis, J. Hagesawa, Y. Kikuchi, and Y. Sasaki, A quantum protocol to win the graph coloring game on all hadamard graphs, *IEICE Trans. Fundam. Electron. Commun. Comput. Sci.*, E89-A(5):1378-1381, 2006. arxiv:quant-ph/0509047v4, doi:10.1093/ietfec/e89-a.5.1378.
- [2] F. Boca, Free products of completely positive maps and spectral sets, *J. Funct. Anal.* 97 (1991), 251-263.
- [3] P.J. Cameron, M.W. Newman, A. Montanaro, S. Severini and A. Winter, On the quantum chromatic number of a graph, *The electronic journal of combinatorics* 14 (2007), R81, arxiv:quant-ph/0608016.
- [4] M. D. Choi and E. Effros, Injectivity and operator spaces, *J. Funct. Anal.* 24 (1977), 156-209.
- [5] T. Cubitt, L. Mančinska, D. Roberson, S. Severini, D. Stahlke, and A. Winter, Bounds on entanglement assisted source-channel coding via the Lovász ϑ number and its variants, *IEEE Trans. Inf. Theory* 60 (2013), arXiv:1310.7120v1.
- [6] R. Duan, S. Severini and A. Winter, Zero-error communication via quantum channels, non-commutative graphs and a quantum Lovász θ function, *IEEE Trans. Inf. Theory* PP:99 (2012), arXiv:1002.2514v2.
- [7] D. Farenick, A. S. Kavruk, V. I. Paulsen and I. G. Todorov, Operator systems from discrete groups, *Comm. Math. Physics* 329 (2014), no. 1, 207-238.
- [8] D. Farenick, A. S. Kavruk, V. I. Paulsen and I. G. Todorov, Characterisations of the weak expectation property, *preprint*, arXiv:1307.1055, 2013.
- [9] D. Farenick and V.I. Paulsen, Operator system quotients of matrix algebras and their tensor products, *Math. Scand.* 111 (2012), 210-243.
- [10] T. Fritz, Operator system structures on the unital direct sum of C^* -algebras, *Rocky Mountain J. Math.* 44 (2014), no. 3, 913-936.
- [11] T. Fritz, Tsirelson's problem and Kirchberg's conjecture, *Rev. Math. Phys.* 24 (2012), no. 5, 1250012, 67 pp.
- [12] V. Galliard and S. Wolf, "Pseudo-telepathy, entanglement, and graph colorings," in *Proc. IEEE International Symposium on Information Theory (ISIT), 2002*, 2002, p. 101.
- [13] C. Godsil and G. Royle, Algebraic graph theory, *Springer-Verlag, New York*, 2001.
- [14] Z. Ji, Binary constraint system games and locally commutative reductions, *preprint (arXiv: 1310.3794)*, 2013.
- [15] M. Junge, M. Navascues, C. Palazuelos, D. Perez-Garcia, V. B. Scholtz and R. F. Werner, Connes' embedding problem and Tsirelson's problem, *J. Math. Physics* 52, 012102 (2011).
- [16] R. V. Kadison and J. R. Ringrose, Fundamentals of the theory of operator algebras II, *American Mathematical Society*, Providence, 1997.
- [17] A. S. Kavruk, Nuclearity related properties in operator systems, *J. Operator Th.* 71 (2014), no. 1, 95-156.
- [18] A. S. Kavruk, V. I. Paulsen, I. G. Todorov, and M. Tomforde, Tensor products of operator systems, *J. Funct. Anal.* 261 (2011), 267-299.
- [19] A. S. Kavruk, V. I. Paulsen, I. G. Todorov, and M. Tomforde, Quotients, exactness, and nuclearity in the operator system category, *Adv. Math.* 235 (2013), 321-360.
- [20] L. Lovász, On the Shannon Capacity of a Graph, *IEEE Transactions on Information Theory*, Vol. II-25, no 1, January 1979, 1-7.

- [21] M. Navascués, S. Pironio, and A. Acín, A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations, *New Journal of Physics*, 10(7): 073013, 2008.
- [22] N. Ozawa, About the Connes’ embedding problem—algebraic approaches, *Japan. J. Math.* 8 (2013), no. 1, 147-183.
- [23] V. I. Paulsen, Completely Bounded Maps and Operator Algebras, *Cambridge University Press*, 2002.
- [24] V. I. Paulsen and I. G. Todorov, Quantum chromatic numbers via operator systems, *Quarterly J. Math.* 66 (2015), no. 2, 677-692.
- [25] David E Roberson, *Variations on a Theme: Graph Homomorphisms*, PhD thesis, University of Waterloo, 2013.
- [26] David E. Roberson and Laura Mančinska, Graph homomorphisms for quantum players, *preprint*, *arXiv:1212.1724*, 2012.
- [27] G. Scarpa and S. Severini, Kochen-Specker sets and the rank-1 quantum chromatic number, *IEEE Trans. Inf. Theory*, 58 (2012), no. 4, 2524-2529.
- [28] V. B. Scholz and R. F. Werner, Tsirelson’s Problem, *preprint*, *arXiv:0812.4305v1*, 2008.
- [29] D. Spielman, *Spectral Graph Theory*, online lecture notes, <http://www.cs.yale.edu/homes/spielman/561/>.
- [30] M. Szegedy, “A note on the ϑ number of Lovász and the generalized Delsarte bound,” in *Proc. 35th Annual Symposium on Foundations of Computer Science, 1994*, 1994, pp. 36–39.
- [31] B. S. Tsirelson, Quantum generalizations of Bell’s inequality, *Lett. Math. Phys.*, 4 (1980), no. 4, 93-100.
- [32] B. S. Tsirelson, Some results and problems on quantum Bell-type inequalities, *Hadronic J. Suppl.*, 8 (1993), no. 4, 329-345.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HOUSTON, HOUSTON, TEXAS 77204-3476, U.S.A.

E-mail address: `vern@math.uh.edu`

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON WC1E 6BT, UNITED KINGDOM

E-mail address: `s.severini@ucl.ac.uk`

DEPARTMENT OF PHYSICS, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PENNSYLVANIA 15213, USA

E-mail address: `dan@stahlke.org`

PURE MATHEMATICS RESEARCH CENTRE, QUEEN’S UNIVERSITY BELFAST, BELFAST BT7 1NN, UNITED KINGDOM

E-mail address: `i.todorov@qub.ac.uk`

ICREA AND FÍSICA TEÒRICA: INFORMACIÓ I FENOMENS QUÀNTICS, UNIVERSITAT AUTÒNOMA DE BARCELONA, ES-08193 BELLATERRA (BARCELONA), SPAIN

E-mail address: `andreas.winter@uab.cat`