



**QUEEN'S
UNIVERSITY
BELFAST**

Integrating geophysical data and upscaling techniques to support regional groundwater flow modelling: A practical example of the Lagan Valley, Northern Ireland

Dickson, N., Comte, J-C., McKinley, J., & Offerdinger, U. (2014). *Integrating geophysical data and upscaling techniques to support regional groundwater flow modelling: A practical example of the Lagan Valley, Northern Ireland*. Poster session presented at European Geosciences Union General Assembly 2014, Vienna, Austria. <http://meetingorganizer.copernicus.org/EGU2014/EGU2014-12147.pdf>

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

CC Attribution 3.0 License.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.



Integrating geophysical data and upscaling techniques to support regional groundwater flow modelling: A practical example of the Lagan Valley, Northern Ireland.

Neil Dickson (1), Jean-Christophe Comte (1), Jennifer McKinley (2), and Ulrich Ofterdinger (1)

(1) School of Planning, Architecture and Civil Engineering, Queen's University Belfast, United Kingdom (ndickson03@qub.ac.uk), (2) School of Geography, Archaeology and Palaeoecology, Queen's University Belfast, United Kingdom

When studying heterogeneous aquifer systems, especially at regional scale, a degree of generalization is anticipated. This can be due to sparse sampling regimes, complex depositional environments or lack of accessibility to measure the subsurface. This can lead to an inaccurate conceptualization which can be detrimental when applied to groundwater flow models. It is important that numerical models are based on observed and accurate geological information and do not rely on the distribution of artificial aquifer properties. This can still be problematic as data will be modelled at a different scale to which it was collected. It is proposed here that integrating geophysics and upscaling techniques can assist in a more realistic and deterministic groundwater flow model. In this study, the sedimentary aquifer of the Lagan Valley in Northern Ireland is chosen due to intruding sub-vertical dolerite dykes. These dykes are of a lower permeability than the sandstone aquifer. The use of airborne magnetics allows the delineation of heterogeneities, confirmed by field analysis. Permeability measured at the field scale is then upscaled to different levels using a correlation with the geophysical data, creating equivalent parameters that can be directly imported into numerical groundwater flow models. These parameters include directional equivalent permeabilities and anisotropy. Several stages of upscaling are modelled in finite element. Initial modelling is providing promising results, especially at the intermediate scale, suggesting an accurate distribution of aquifer properties. This deterministic based methodology is being expanded to include stochastic methods of obtaining heterogeneity location based on airborne geophysical data. This is through the Direct Sample method of Multiple-Point Statistics (MPS). This method uses the magnetics as a training image to computationally determine a probabilistic occurrence of heterogeneity. There is also a need to apply the method to alternate geological contexts where the heterogeneity is of a higher permeability than the host rock.