



**QUEEN'S
UNIVERSITY
BELFAST**

Volunteering and mortality risk: A partner-controlled quasi-experimental design

O'Reilly, D., Rosato, M., Moriarty, J., & Leavey, G. (2017). Volunteering and mortality risk: A partner-controlled quasi-experimental design. *International Journal of Epidemiology*. <https://doi.org/10.1093/ije/dyx037>

Published in:
International Journal of Epidemiology

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© The Author 2017; all rights reserved. Published by Oxford University Press on behalf of the International Epidemiological Association. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access
This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

Volunteering and mortality risk: A partner-controlled quasi-experimental design.

Authors:

Dermot O'Reilly¹; Michael Rosato ²; John Moriarty³; Gerard Leavey²

Affiliations:

1. Centre for Public Health; Queen's University Belfast; Belfast
2. Bamford Centre for Mental health and Wellbeing; Ulster University; Northern Ireland
3. Centre for Evidence and Social Innovation, Queen's University Belfast

Correspondence to
Dr Dermot O'Reilly
Centre for Public Health;
Queen's University Belfast;
Belfast BT12 6BJ,
Northern Ireland
Tel : 028 90632738
Email: d.oreilly@qub.ac.uk

Word count :

Abstract: 233

Paper: 2790

Abstract

Background

The consensus that volunteering is associated with a lower mortality risk is derived from a body of observational studies and therefore vulnerable to uncontrolled or residual confounding. This potential limitation is likely to be particularly problematic for volunteers who, by definition, are self-selected and known to be significantly different from non-volunteers across a range of factors associated with better survival.

Methods

This is a census-based record linkage study of 308,733 married couples aged 25 and over, including 100,571 volunteers, with mortality follow-up for thirty-three months. We used a standard Cox model to examine if mortality risk in the partners of volunteers was influenced by partner volunteering status - something expected if the effects of volunteering on mortality risk were due to shared household or behavioural characteristics.

Results

Volunteers were general more affluent, better educated and more religious than their non-volunteering peers; they also had a lower mortality risk ($HR_{adj}=0.78$: 95%CI=0.71, 0.85 for males and $HR_{adj}=0.77$: 95%CI=0.68, 0.88 for females). However, amongst cohort members who were not volunteers, having a partner who was a volunteer was not associated with a mortality advantage ($HR_{adj}=1.01$: 95%CI=0.92, 1.11 for men and $HR_{adj}=1.00$: 95%CI=0.88, 1.13 women)

Conclusions

This study provides further evidence that the lower mortality associated with volunteering is unlikely to be due to health selection or to residual confounding arising from unmeasured selection effects within households. It therefore increases the plausibility of a direct causal effect.

Key words: Volunteering, mortality risk, quasi-experimental study

Key messages

- Most of the information suggesting that volunteering benefits the volunteer as well as the wider society comes from observational studies and is therefore vulnerable to uncontrolled or residual confounding
- This census-based study uses the known similarities between married co-habiting partners to provide better adjustment for unmeasured or poorly controlled potential confounders
- It shows a clear mortality advantage for volunteers but not their non-volunteering partners providing the strongest evidence yet for a direct causal effect of volunteering on the volunteer.

Background

There is now a general consensus that volunteering carries social and health benefits, not just for recipients and the wider society but also for the volunteers themselves. Many governments are now actively seeking ways of increasing the opportunities for volunteering as a mechanism for civic engagement, especially amongst older members of society. In a series of reviews and systematic reviews¹⁻⁶ this consensus has recently been extended to include an associated reduced mortality risk. One recent meta-analysis of fourteen studies by Okun *et al*⁷ showed that organisational or formal volunteering reduced the mortality risk of people aged fifty-five or more by 47% (95%CI=38-55%), and by 24% (16-31%) after adjustment for mediating factors, and concluded that *'it is no longer a question of whether volunteering is predictive of reduced mortality: rather... that the volunteering-mortality association is reliable and that the magnitude of the relationship is sizable'*. Others have qualified this - Anderson *et al*⁸, in a narrative review, suggested that while benefits (including reduced mortality) were evident at moderate levels of volunteering these may be less apparent at high-intensity levels; and Jenkinson *et al*⁴ noted that most mortality-related studies have been US-based where there is both a strong tradition of volunteering and wide disparities in health.

However, most studies reporting the salutogenic effects of volunteering, and all of the mortality studies⁸⁻¹⁹, are based on observational study designs and therefore potentially subject to confounding due to factors that are unmeasured or difficult to adjust for using standard analytic approaches. Therefore, despite consensus about its benefits, there is still uncertainty as to whether the lower mortality risk associated with volunteering is a result of the activity of volunteering per se or a consequence of the characteristics of those who volunteer. This may be important for purported health benefits, as volunteers are, by definition, self-selected and it is recognised that volunteers differ from those who don't across a range of factors related to mortality risk: for example they have higher levels of social and material resources^{20, 21}; tend to be more affluent and better educated; have better

health (especially physical health); and better social integration and more religious involvement ²²⁻²⁴.

The ideal solution - randomisation - is impractical where mortality is the outcome and while there have been some attempts to trial the effects of volunteering ²⁵⁻²⁸ none have included mortality as an outcome. However, other methodologies such as propensity score ²⁹ matching or the use of instrumental variables ³⁰ are increasingly advocated to better adjust for other potential confounders and enhance causal inference from observational studies. Another approach is to undertake comparisons within a family, effective examples of which include studies of early life factors where sibling exposures and outcomes ³¹⁻³⁴ have been matched to detect and control for familial confounding, and they have produced greater clarity about the role of maternal smoking or body mass index and hyperactivity disorder in offspring ³⁵⁻³⁷ or use of psychotropic medication in later life³⁸ and about the role of education and cardiovascular disease ³⁹.

The current study uses married partners rather than siblings as controls to provide the additional adjustment for the social, environmental and lifestyle covariates that may be associated with both volunteering and mortality risk. This study utilises the fact that not only do married and cohabiting couples share to a great extent the same physical, social, and socio-economic environment but are often similar across a range of other physical and behavioural attributes. This can arise from the combined effects of social homogamy or phenotypic assortative mating whereby people tend to marry people similar to themselves in terms of race, level of education, social class, behaviours etc, or through cohabitation or social interaction effects ie from a continued exposure to a largely shared physical and social environment. Collectively this means that there is a well-recognised concordance between spouses in behaviours such as smoking, diabetes and obesity and other cardiovascular risk factors ⁴⁰⁻⁴¹, alcohol dependency and other health behaviours ^{42, 43}, mental health ⁴⁴, and work-related disability ⁴⁵. We hypothesise that if the lower mortality risk associated with

volunteering is due to unadjusted or poorly controlled confounders present at the household level then this lower mortality risk will be also be evident amongst the non-volunteering partners of these volunteers. If on the other hand volunteering has real protective effects then the mortality advantage should not necessarily be evident for their partner.

Methods

The Northern Ireland Mortality Study (NIMS) is a record-linkage study comprising the census returns for the whole enumerated population and subsequently registered deaths. The details of both NIMS and linkage processes are described elsewhere⁴⁶. For this study the population-at-risk were those enumerated in the Northern Ireland Census (March 2011), aged twenty-five and over and not living in institutional care, with mortality follow-up from the Census until December 2013 (a total of thirty-three months).

All personal characteristics were drawn from the census and selected on the basis of their known association with either volunteering or mortality risk: these include age (in ten year bands to seventy-five and over); gender; and marital status (married, never married, and – as a single group - those widowed, separated or divorced). Religious affiliation was also included as religiosity has been associated with both volunteering and the effects of volunteering, and previous analyses have suggested higher levels of religiosity amongst more conservative Christians: here six religious groups, including no affiliation, were classified (see table 1). In this context the more conservative Christians included smaller Protestant denominations such as Pentecostal or Evangelical groups. Socioeconomic status was assessed using (i) household car availability (two or more cars, one only, no access); (ii) educational attainment (third-level, intermediate, no qualifications); (iii) economic activity; and finally (iv) a combination of housing tenure and the rateable value of the property. Rateable value had been derived as part of an exercise by central government in 2005 to determine the level of local residential tax levels payable for each household, and this data was combined with housing tenure to produce an eight-fold classification of tenure/capital value: private renting; social renting; and, for owner-occupiers, five categories ranging from less than £75,000 to over £200,000 (see table 1), with an additional category for homes as yet unvalued.

Volunteer status:

Although researchers and organisations use a wide array of definitions of volunteering (see ref 47 for overview), most agree that it incorporates three essential components - while the activity is voluntary, unpaid, and benefits recipient individuals or communities, it excludes help to close family members (caregiving). Some researchers also differentiate between formal volunteering, which is structured through an organisation and informal volunteering which happens outside the auspices of a formal organisation. For this study volunteer status was based on a single census question: "*In the past year, have you helped with or carried out any voluntary work without pay?*", with responses 'yes' or 'no'. No further detail was sought in terms of the hours spent volunteering or whether this was in a formal or informal capacity.

Health status:

The health status of cohort members at baseline was also based on responses to a range of census questions: the first asked "*how is your health in general*", with five responses ranging from very good to very bad; a second asked if people had a health problem or disability which limited day-to-day activity "*a little*", or "*a lot*" (and had lasted, or was expected to last, at least twelve months). A further question asked about specific chronic conditions "*Do you have any of the following conditions which have lasted, or are expected to last, at least 12 months?*" – from which we selected four that covered a wide range of physical and mental health problems: (i) "*a mobility or dexterity difficulty (a condition that substantially limits one or more basic physical activities such as walking, climbing stairs lifting or carrying)*"; (ii) "*an emotional, psychological or mental health condition (such as depression or schizophrenia)*"; (iii) "*long-term pain or discomfort*"; and finally, (iv) "*shortness of breath or difficulty breathing (such as asthma)*".

The outcome for analysis was risk of all-cause mortality during follow-up. The resulting linked data were anonymised, held in a safe setting by the Northern Ireland Statistics and Research Agency (NISRA) and made available to the research team for the purpose of this

study. The use of the NIMS for research was approved by the Office for Research Ethics Committees Northern Ireland (ORECNI).

Data manipulation and analysis strategy

Of the 1,123,205 people aged twenty-five and over at the census and not living in institutionalised care, we identified 308,733 married couples living in the same household with complete data on all variables examined. Descriptive statistics illustrated the distributions of: (a) individual-level socio-demographic and self-reported health characteristics, by gender and volunteer-status; and (b) household-level characteristics, by number of volunteers in the household. Sex-specific Cox proportional hazards models were used to examine the all-cause mortality risk associated with volunteering and the mortality associated with spouse-partners.

Results

Of the 617,466 individuals identified in the study 16.3% identified as volunteers, representing almost a quarter (24.3%) of the 308,733 households. Overall there were 48,357 male and 52,214 female volunteers (15.7% and 16.9% of their respective populations). Table 1(a) shows the socio-demographic characteristics associated with volunteering: most prevalent in middle-age, and more common amongst people from more conservative religious faiths; those better educated, employed and in better health (recording lower levels of both mental and physical chronic ill-health). Male and female volunteers have similar demographic profiles except for age, where a higher proportion of older men than older women volunteer. Table 1(b) shows the distribution of volunteers in a household by indicators of relative affluence (housing tenure and car availability), and shows clear differences between those households with a volunteer present and those without, and also that those households with two volunteers are slightly more affluent than those with one only. Almost 30% of households with two or more cars, and 37% of the most expensive houses had at least one volunteer.

In 34% of volunteer households both partners were volunteers - 53% of male and 49% of female volunteers also had a partner who was a volunteer. Even after adjusting for age, religion and socio-economic status, having a partner who was a volunteer increased the likelihood of being a volunteer by almost eightfold (full models available on request). There was a reasonable correspondence in the individual characteristics of couples across a range of variables, with 83% sharing the same religious affiliation, 59% the same level of educational attainment and 58% the same level of general health. In models adjusted for age, religious affiliation, volunteer status, educational attainment, car availability and housing tenure and value having one partner with chronic poor mental health increased the risk of poor mental health in the other partner by four-fold (OR=4.17: 95%CIs=3.99, 4.36 for men and OR=4.18: 95%CIs=4.00, 4.37 for women).

During the thirty-three months of follow-up there were 12,260 deaths, 6.9% of which were to volunteers. In models adjusted for all the demographic and socioeconomic variables listed in

Table 1, the mortality risk associated with being a volunteer was HR=0.65: 95%CI=0.62, 0.69 for males and HR=0.57: 95%CI=0.53, 0.61 for females. With further adjustment for baseline health status this mortality risk attenuated to HR=0.79: 95%CI=0.71, 0.85 for males and HR=0.77: 95%CI=0.68, 0.88 for females. Table 2 shows the mortality variations for all four volunteer-non-volunteer combinations for both sexes. Though the first series of analyses where the comparison is between those non-volunteer subjects who do and do not have a volunteering partner are the most germane to this paper. In models adjusted for age non-volunteering men and women with partners who were volunteers are about 15% less likely to die during follow-up than their peers with non-volunteering partners. This does not change with adjustment for variations in religious affiliation but the difference disappears entirely with further adjustment for socio-economic. Further adjustment for baseline health status produces little further change. In the fully adjusted models, there are no additional mortality benefit for volunteers in having a partner who is also a volunteer.

Discussion

This study shows the following: it confirms the established characteristics associated with volunteering – when compared to non-volunteers they are more affluent, better educated, from more conservative religious affiliations, and physically and mentally healthier. It also confirms that even after adjustment for a range of socio-economic and baseline health factors, volunteers have an approximate 25% lower mortality risk than their non-volunteering peers and that this is true for both men and women. However, the novel finding here is the clear demonstration that amongst co-habiting married couples where one partner is a volunteer the effects of volunteering are seen only in the volunteer and not in their non-volunteering partner: evidence that the known associations between volunteering and lower mortality risk are unlikely to be due to unmeasured or poorly adjusted-for confounders.

Although this study design allows a robust examination of the mortality risk associated with volunteering its strength rests on the assumptions that (a) spousal controls provide a good adjustment for unmeasured potential confounders and (b) that the effects of volunteering are only likely to be experienced by the volunteer. While it is evident that comparison between spouse-pairs provides excellent adjustment for most household-level factors, including socio-economic status, it is possible that some aspects of wealth or income may not be equally distributed within a household. It is also likely that between-partner comparisons provides some adjustment for other potential social, behavioural or attitudinal confounders, though it is acknowledged that this is likely to be less successful than adjustment for shared factors such as physical environment and socio-economic status. A large body of existing evidence attests to the general concordance between spouses in lifestyles and behaviours and in levels of health status⁴⁰⁻⁴⁵, and this study also demonstrates a reasonably high degree of similarity between spouses across an array of social and health factors including religious affiliation, educational attainment, general and mental health. It is therefore unlikely that the mortality advantage associated with volunteering is due to residual confounding as this would be expected to present as a somewhat lower mortality risk amongst the non-

volunteering partner of a volunteer. It is however acknowledged that this study design cannot adjust for other possibly important unshared factors or intrinsic attributes that might confound the relationship between volunteering and mortality such as the personality trait of conscientiousness which is known to be related to both the propensity to volunteer⁴⁸ and to mortality risk⁴⁹.

The second assumption, that any benefits accruing from volunteering are predominantly experienced by the volunteer, is in keeping with the large body of research related to the health benefits associated with volunteering. While we accept that there may be some minor advantage to the non-volunteering partner in terms of wider social interaction or better access to health and other information it is evident that the suggested salutogenic mechanisms emphasise the primacy of the effects on the volunteer. Social integration and role theories, underpinned by putative bio-psychological mechanisms⁵⁰, suggests that volunteering leads to improvements in mental and physical health by providing a sense of meaning and purpose in life⁵¹; through facilitating social integration and interaction⁵²; and by affecting personal self-control promoting and bolstering self-esteem⁵³; increased self-efficacy and competence⁵⁴; and distraction from personal troubles⁵⁵. Volunteering may also be associated with increased levels of physical activity.

This study has significant strengths and limitations. It is a large and representative study with volunteering defined at baseline and full ascertainment of mortality records through official registrations. That the study was based on census returns avoids the recognised selection bias of volunteers into social surveys⁵⁶. The census also provided adjustment for other socio-demographic, socio-economic and health factors known to be associated with both volunteering and mortality risk. The census, however, also presents some limitations, as it has to trade-off a population-wide coverage for quality and extent of data capture. The responses to the census question on volunteering only allowed a dichotomous classification with no additional information available as to the type, duration or intensity of the activity, or

whether this was in a formal or informal capacity. However, the prevalence of approximately 16% recorded here is in keeping with previously reported European and UK levels^{57,58}, and their socio-demographic profile matches that of these and other studies world-wide. Furthermore, the lower mortality risk for those identified as volunteering in the current study is also very similar to that reported by Jenkinson *et al*⁴ and Okun *et al*⁷ in separate meta-analyses on the effects of volunteering.

In conclusion, while this study adds to the evidence for an independent effect, additional work is required to understand how the benefits of volunteering relate to its type, quality and context, its relationship with other pro-social activity, and how the effects of these activities vary across individual characteristics before it can be recommended as a public health intervention^{4,6}.

Acknowledgement

The help provided by the staff of the Northern Ireland Mortality Study (NIMS) and the NILS Research Support Unit is acknowledged. The NIMS is funded by the Health and Social Care Research and Development Division of the Public Health Agency (HSC R&D Division) and NISRA. The NILS-RSU is funded by the ESRC and the Northern Ireland Government. The authors alone are responsible for the interpretation of the data and any views or opinions presented are solely those of the author and do not necessarily represent those of NISRA/NILS.

References

1. Cassidy R, Kinsman E, Fisher C, Bambra C. (2008), Volunteering and Health: What Impact Does It Really Have? Final report to Volunteering England. London, UK: Volunteering England; 2008
2. Von Bonsdorff MB, Rantanen T. Benefits of formal voluntary work among older people – a review. *Aging Clin Exp Res* 2010.
3. Anderson N D, Damianakis T, Kröger E, Wagner L M, Dawson D R, Binns M A, Bernstein S, Caspi E, Cook S L, The BRAVO Team. The benefits associated with volunteering among seniors: A critical review and recommendations for future research. *Psychol Bull.* 2014; 140:1505-33
4. Jenkinson CE, Dickens AP, Jones KS, Thompson–Coon J, Taylor RS, Rogers M, Bambra CL, Lang I, Richards SH. Is volunteering a public health intervention? A systematic review and meta-analysis of the health and survival of volunteers. *BMC Public Health*, 2013; 13: 773-773
5. Wilson J, Musick M. The Effects of Volunteering on the Volunteer, *Law and Contemporary Problems* 1999: 62 141-168
6. Morrow-Howell N. Volunteering in later life: Research frontiers. *J Gerontol B Psychol Sci Soc Sci* 2010; 65B: 461-469
7. Okun MA, Yeung EW, Brown S. Volunteering by older adults and risk of mortality: A meta-analysis. *Psychol Aging.* 2013; 28: 564–577.
8. Sabin E. Social relationships and mortality amongst the elderly. *J Applied Gerontol* 1993; 12: 44-60
9. Ayalon L. Volunteering as a predictor of all-cause mortality: what aspects of volunteering really matter? *Int Psychogeriatr* 2008, **20**:1000-1013
10. Harris AHS, Thoresen CE: Volunteering is associated with delayed mortality in older people: analysis of the longitudinal study of aging. *J Health Psychol* 2005, 10:739-752.
11. Okun MA, August KJ, Rook KS, Newsom JT: Does volunteering moderate the relation between functional limitations and mortality? *Soc Sci Med* 2010, **71**:1662-1668.
12. Oman D, Thoresen CE, McMahon K: Volunteerism and mortality among the community-dwelling elderly. *J Health Psychol* 1999, **4**:301-316.
13. Musick MA, Herzog A, House JS: Volunteering and mortality among older adults: findings from a national sample. *J Gerontol B Psychol Sci Soc Sci* 1999, **54B**:S173-S180.
14. Konrath S, Fuhrel-Forbis A, Lou A, Brown S: Motives for volunteering are associated with mortality risk in older adults. *Health Psychol* 2012, **31**:87-96.
15. Menec VH: The relation between everyday activities and successful aging: a 6 year longitudinal study. *J Gerontol B Psychol Sci Soc Sci* 2003, **58B**:S74-S82.
16. Hsu HC. Does social participation by the elderly reduce mortality and cognitive impairment? *Aging Mental Health* 2007; 11: 699-707
17. Ming-Ching L, Herzog AR. Individual consequences of volunteer and paid work in old age: health and mortality. *J Health Soc Behaviour* 2002; 43: 490-509
18. Lum T, Lightfoot E. The effects of volunteering on the physical and mental health of older people. *Res Aging* 2005; 27: 31-55
19. Lee S, Steinman M, Tan E. Volunteering, driving status and mortality in US retirees. *J Am Geriat Soc* 2011; 59: 274-280
20. Martinson M, Minkler M. Civic engagement and older adults: A critical perspective. *The Gerontologist* 2006; 46: 318-324
21. Minkler M, Holstein MB. From civil rights to civic engagement? Concerns of two older critical gerontologists about a “new social movement” and what it portends. *J Aging Studies* 2008; 22: 196-204.
22. McMunn A, Nazroo J, Wahrendorf M, Breeze E, Zaninotto P: Participation in socially-productive activities, reciprocity and wellbeing in later life: baseline results in England. *Ageing Soc* 2009, 29: 765–782.

23. Tang F. What resources are needed for volunteerism? A life course perspective. *Journal of Applied Gerontology* 2006; 25: 375-390.
24. Wilson J, Musick M. Who cares? Toward an integrated theory of volunteer work. *American Sociological Review* 1997;62:694-713.
25. Tan EJ, Xue QL, Li T, Carlson MC, Fried LP: Volunteering: a physical activity intervention for older adults - the experience corps program in Baltimore. *J Urban Health* 2006, 83:954–969.
26. Yuen HK, Huang P, Burik JK, Smith TG: Impact of participation in volunteer activities for residents living in long-term-care facilities. *Am J Occup Ther* 2008, 62:71–76.
27. George DR, Singer ME: Intergenerational volunteering and quality of life for persons with mild to moderate dementia: results from a 5-month intervention study in the United States. *Am J Geriatr Psychiatry* 2011, 19:392–396.
28. Pettigrew S, Jongenelis M, Newton R, Warburton J, Jackson B. Research protocol for a randomised controlled trial of the health effects of volunteering for seniors. *Health Qual Life Outcomes*. 2015; 13: 74.
29. Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects *Biometrika* 1983; 70: 41-55
30. Palmer TM, Sterne JAC, Harbord RM, Sheehan NA, Tobias JH, Timpson NJ, Davey Smith G, Sterne JA. Instrumental variable estimation of causal risk ratios and causal odds ratios in Mendelian randomization analyses. *Am J Epidemiol*. 2011; 173: 1392-1403
31. D'Onofrio BM, Lahey BB, Turkheimer E, Lichtenstein P. Critical Need for Family-Based, Quasi-Experimental Designs in Integrating Genetic and Social Science Research. *AJPH*. 2013; 103: S46-S55.
32. Susser E, Eide MG, Begg M. Invited Commentary: The Use of Sibship Studies to Detect Familial Confounding. *Am. J. Epidemiol*. 2010; 172: 537-539.
33. Donovan SJ, Susser E. Commentary: Advent of sibling designs *Int. J. Epidemiol*. 2011; 40: 345-349
34. Keyes KM, Davey Smith G, Susser E. Commentary: Smoking in pregnancy and offspring. *Int. J. Epidemiol*. 2014, 1381–1388
35. Obel C, Olsen J, Henriksen TB, Rodriguez A, Järvelin MR, Moilanen I, Parner E, Markussen Linnert K, Taanila A, Ebeling H, Heiervang E, Gissler M. Is maternal smoking during pregnancy a risk factor for Hyperkinetic disorder?-findings from a sibling design. *Int. J. Epidemiol*. 2011; 40: 338-345.
36. Chen Q, Sjölander A, Långström N, Rodriguez A, Serlachius E, D'Onofrio BM, Lichtenstein P, Larsson H. Maternal pre-pregnancy body mass index and offspring attention deficit hyperactivity disorder: a population-based cohort study using a sibling-comparison design. *Int. J. Epidemiol*. 2013; 1–8 doi:10.1093/ije/dyt152
37. Skoglund C, Chen Q, D'Onofrio BM, Lichtenstein P, Larsson H. Familial confounding of the association between maternal smoking during pregnancy and ADHD in offspring *J Child Psychol Psychiatry*. 2014; 55: 61-68.
38. Söderström L, Perez-Vicente R, Juárez S, Merlo J Questioning the Causal Link between Maternal Smoking during Pregnancy and Offspring Use of Psychotropic Medication: A Sibling Design Analysis. *PLoS ONE* 2013; 8: e63420. doi:10.1371/journal.pone.0063420
39. Madsen M, Andersen PK, Gerster M, Andersen AM, Christensen K, Osler M. Are the educational differences in incidence of cardiovascular disease explained by underlying familial factors? A twin study *Soc Sci Med* 2014;118: 182-90
40. Di Castelnuovo, A, Quacquarello G, Donati MB, de Gaetano G, Iacoviello L. Spousal Concordance for major coronary risk factors: A systematic review and meta-analysis. *Am. J. Epidemiol*. 2009; 169: 1-8
41. Cobb LK, McAdams-DeMarco MA, Gudzone KA, Anderson CA, Demerath E, Woodward M, Selvin E, Coresh J. Changes in Body Mass Index and Obesity Risk in Married Couples Over 25 Years: The ARIC Cohort Study *Am J Epidemiol* 2015 pii: kwv112

42. Grant JD, Heath AC, Bucholz KK, et al. Spousal concordance for alcohol dependence: evidence for assortative mating or spousal interaction effects? *Alcohol Clin Exp Res.* 2007; 31: 717-728.
43. Falba TA, Sindelar JL. Spousal concordance in health behavior change. *Health Serv Res.* 2008; 43: 96-116.
44. Butterworth P, Rodgers B. Concordance in the mental health of spouses: analysis of a large national household panel survey. *Psychol Med.* 2006; 36:685-697.
45. Vie G, Romundstad P, Krokstad S, Johnsen R, Bjørngaard J. Mortality and work disability in a cohort of Norwegian couples--the HUNT Study *Eur J Public Health* 2015 25: 807-814
46. O'Reilly D, Rosato M, Johnston F, Catney G, Brolly M. Cohort description: The Northern Ireland Longitudinal Study (NILS): *Int J Epidemiol* 2012; 41: 634-641
47. Volunteering England Information Sheet © Volunteering England 2008
48. Mike A, Jackson JJ, Oltmanns TF. The conscientious retiree: The relationship between conscientiousness, retirement, and volunteering. *J Res Personality.* 2014; 52: 68-77.
49. Jokela M, Batty GD, Nyberg ST, Virtanen M, Nabi H, Singh-Manoux A, Kivimäki M. Personality and all-cause mortality: individual participant meta-analysis of 3,947 deaths in 76,150 adults. *Am J Epidemiol.* 2013; 178: 667–675.
50. Kim S, Ferraro KF. Do Productive Activities Reduce Inflammation in Later Life? Multiple Roles, Frequency of Activities, and C - reactive protein. *The Gerontologist* 2014; 54: 830-839.
51. Parament KI. *The psychology of religion and coping: Theory, research, practice.* New York: Guilford 1997.
52. Bergman L, Syme SL. Social networks, host resistance and mortality: a nine-year follow-up study of Alameda County residents. *Am J Epid* 1979; 109: 186-204
53. Li Y, Ferraro KF. Volunteering and Depression in Later Life: Social Benefit or Selection Processes? *J Health Social Behavior* 2005; 46: 68-84
54. Bandura A. *Self-efficacy: The exercise of control.* Ney York: W.H.Freeman 1997
55. Matthews G, Wells A. Attentional processes, dysfunctional coping and clinical intervention. In NSE Moshe Zeidner (ed) *Handbook of coping: Theory, research, applications.* New York: Wiley 1996
56. Abraham KG, Helms S, Presser S. How social processes distort measurement: The impact of survey nonresponse on estimates of volunteer work in the United States. *American Journal of Sociology* 2009; 114: 1129-1165.
57. Haski-Leventhal D. Elderly volunteering and well-being: A cross-European comparison based on SHARE data *Int J Vol NonProfit Org* 2009; 20: 388-404
58. McMunn, A., Nazroo, J., Wahrendorf, M., Breeze, E., & Zaninotto, P. Participation in socially-productive activities, reciprocity and well-being in later life: Baseline results in England. *Ageing & Society*, 2009; 29: 765–782.

Table 1(a): Comparison of individual characteristics of volunteers and non-volunteers in married households: Data represent percentages within volunteer strata.

	Male		Female	
	Non-volunteer	Volunteer	Non-volunteer	Volunteer
	N= 260,376	N= 48,357	N= 256,519	N= 52,214
Age(years)				
25-34	10.1	8.3	13.9	11.1
35-44	21.4	21.2	22.4	24.5
45-54	23.7	28.1	23.7	28.5
55-64	20.7	22.5	19.9	21.1
65-74	15.5	14.6	14.1	12.1
75+	8.6	5.4	6.0	2.6
Religion				
Roman Catholic	38.6	34.7	41.2	34.7
Presbyterian	23.9	24.3	23.5	25.6
Church of Ireland	16.2	14.9	16.6	16.1
Methodist	3.6	3.8	3.8	4.1
Other Christian	6.6	12.6	6.8	12.0
Other religion	0.9	1.1	0.9	1.0
Not stated	10.2	8.7	7.1	6.5
Educational attainment				
Degree	27.7	51.2	31.3	52.6
Intermediate	39.8	38.0	37.7	38.0
No degree	32.4	10.8	31.0	9.3
Economic Activity				
Employed full-time	57.9	66.7	29.3	33.5
Employed part-time	6.7	7.9	26.3	33.2
Unemployed	2.8	2.1	1.4	1.7
Retired	23.8	19.8	24.6	19.8
Homemaker/other	3.2	2.0	11.9	9.8
Permanently sick	5.5	1.5	6.4	1.9
Limiting long term				
None	74.2	84.7	74.3	85.6
Limiting a little	10.9	10.3	11.0	9.9
Limiting a lot	14.9	5.1	14.7	4.5
General health				
Very good	34.8	45.3	35.8	46.6
Good	38.1	40.8	37.2	40.0
Fair	19.7	12.1	19.4	11.9
Bad	5.8	1.5	6.0	1.4
Very bad	1.5	0.2	1.5	0.2
Chronic conditions				
Mental ill-health	4.7	2.6	6.8	3.8
Mobility problems	15.2	6.9	15.7	7.3
Chronic pain	14.1	8.6	15.6	9.4
Breathing difficulty	9.5	6.3	9.3	6.0

Table 1(b): Characteristics of couple household according to number of volunteers

	No volunteers	One volunteer	Two volunteers
Number of households	N= 233,864	N= 49, 167	N= 25,702
Tenure/ property value			
Owner occupier: £200k	11.9	20.1	25.9
£150-199k	14.9	20.0	22.4
£100-149k	26.9	26.6	23.9
£75-99.9k	14.8	10.8	7.9
<£75k	9.1	5.4	3.7
Owner occupier: no rateable value	6.9	8.0	8.5
Private renting	1.2	1.1	1.8
Social renting	14.2	7.9	5.8
Household car access			
Two or more	57.5	71.4	77.9
One	36.5	26.1	20.8
None	6.0	2.5	1.3

Table 2: Mortality risk stratified by sex and own volunteering status according to risk on volunteering status of partner. Data represents the number of deaths in each category and the hazard ratios (and 95% Confidence Intervals) from separate Cox proportional hazard models.

			Adj age	+ religion	+ SES	+ health
Males	Partner	Non-volunteer	1.00	1.00	1.00	1.00
		(7,434)	Volunteer	0.85 (0.78 ,0.93)	0.85 (0.78 ,0.93)	0.99 (0.91 ,1.09)
	Volunteer	Non-volunteer	1.00	1.00	1.00	1.00
		(560)	Volunteer	0.87 (0.73 ,1.02)	0.87 (0.74 ,1.03)	0.97 (0.82 ,1.15)
Female	Partner	Non-volunteer	1.00	1.00	1.00	1.00
		(4,035)	Volunteer	0.84 (0.75 ,0.96)	0.85 (0.75 ,0.96)	1.00 (0.88 ,1.13)
	Volunteer	Non-volunteer	1.00	1.00	1.00	1.00
		(282)	Volunteer	0.81 (0.64 ,1.02)	0.82 (0.65 ,1.04)	0.90 (0.71 ,1.14)