



**QUEEN'S
UNIVERSITY
BELFAST**

Radiobiological Implications of Fukushima Nuclear Accident for Personalized Medical Approach

Fukunaga, H., Yokoya, A., Taki, Y., & Prise, K. M. (2017). Radiobiological Implications of Fukushima Nuclear Accident for Personalized Medical Approach. *The Tohoku Journal of Experimental Medicine*, 242(1), 77-81. <https://doi.org/10.1620/tjem.242.77>

Published in:
The Tohoku Journal of Experimental Medicine

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© 2017 Tohoku University Medical Press.
This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access
This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

Radiobiological implications of Fukushima nuclear accident for personalized medical approach

Hisanori Fukunaga^{1,2,3}, Akinari Yokoya², Yasuyuki Taki³ and Kevin M. Prise¹

¹ Centre for Cancer Research and Cell Biology, Queen's University Belfast, Belfast, UK

² Tokai Quantum Beam Science Center, National Institutes for Quantum and Radiological Science and Technology, Tokai, Ibaraki, Japan

³ Institute of Development, Aging and Cancer, Tohoku University, Sendai, Japan

Correspondence:

Hisanori Fukunaga, M.D.

Centre for Cancer Research and Cell Biology, Queen's University Belfast, 97 Lisburn Road, Belfast BT9 7AE, UK

Tel: +44 28 9097 2760

E-mail: hfukunaga01@qub.ac.uk

Running title:

Radiobiological implications of Fukushima nuclear disaster

Abstract:

On March 11, 2011, a devastating earthquake and subsequent tsunami caused serious damage to areas of the Pacific coast in Fukushima prefecture and prompted fears among the residents about a possible meltdown of the Fukushima Daiichi Nuclear Power Plant reactors. As of 2017, over six years have passed since the Fukushima nuclear crisis and yet the full ramifications of the biological exposures to this accidental release of radioactive substances remain unclear. Furthermore, although several genetic studies have determined that the variation in radiation sensitivity among different individuals is wider than expected, personalized medical approaches for Fukushima victims have seemed to be insufficient. In this commentary, we discuss radiobiological issues arising from low-dose radiation exposure, from the cell-based to the population level. We also introduce the scientific utility of the Integrative Japanese Genome Variation Database (iJGVD), an online database released by the Tohoku Medical Megabank Organization, Tohoku University that covered the whole genome sequences of 2049 healthy individuals in the northeastern part of Japan in 2016. Here we propose a personalized radiation risk assessment and medical approach, which considers the genetic variation of radiation sensitivity among individuals, for next-step developments in radiological protection.

Keywords:

Fukushima Daiichi Nuclear Power Plant; low-dose radiation; precision medicine; radiation sensitivity; radiation-induced bystander effect

Commentary:

In March of 2011, the northeastern part of Japan was devastated by the triple disasters of an earthquake, a tsunami, and the Fukushima Daiichi Nuclear Power Plant (FNPP) accident (Shibahara 2011), leaving local residents fearful about the long-term health risk of low-dose radiation because of radioactive substances released from the FNPP (Ishigaki et al. 2013; Fukunaga and Kumakawa 2015). As of 2017, unfortunately, the ramifications of biological exposure to low dose radiation remains unclear (Hosoda et al. 2016).

Previous epidemiological data indicate a clear increase in the long-term risk of cancer from exposure to radiation at doses exceeding 100 mSv. For this reason, following Japan's worst ever radiation accident at FNPP, radiation doses have been measured in the Fukushima residents, using both personal dosimeters for external exposure and whole-body counters for internal exposure (Yasumura et al. 2012). Several studies have indicated that the radiation exposure levels of Fukushima residents are much lower than predicted (Hayano et al. 2013; Kamiya et al. 2016). Furthermore, Harada *et al.* showed that the mean annual radiation dose rate in 2012 associated with the accident was 0.89–2.51 mSv/y and the mean dose rate estimates for 2022 are comparable with the variations of the average 2 mSv/y background radiation due to exposure to natural radionuclides in Japan (Harada et al. 2014). These findings suggest that the extra lifetime integrated dose after 2012 will increase the estimated lifetime risk of cancer incidence by a factor of 1.03 to 1.05 at most, which is unlikely to be epidemiologically detectable (Harada et al. 2014).

In 2013, the World Health Organization (WHO) reported that the estimates of radiation-induced health risks in Japan for the lifetime risk for some cancers may be somewhat elevated above baseline rates in certain age and sex groups among residents of the areas most affected by the FNPP accident (WHO 2013). Also, the United Nations

Scientific Committee on the Effects of Atomic Radiation (UNSCEAR) published a report suggesting that a general radiation-related increase in the incidence of health effects among the exposed population would be indiscernible over the baseline level; however, it could not rule out the possibility of excess cases of disease due to irradiation, nor to the possibility of detection of a biomarker for certain types of cancer in certain subgroups being identified in the future that could be associated with radiation exposure (UNSCEAR 2014; UNSCEAR 2015). In addition, the dose-response relation for cancer at low doses is assumed, for purposes of radiological protection, to be linear and without a threshold, but this has not been definitively confirmed for all exposure scenarios. An appropriate dose-response relation for the effects of low doses of radiation therefore remains to be established (Kamiya et al. 2015).

The knowledge accumulated concerning the risks of low-dose radiation exposure to humans appears to be insufficient at this time because of important radiological issues arising at low doses due to inhomogeneous dose distribution in living systems. In general, the energy deposition of radiation is localized along its track, resulting in a heterogeneous distribution of exposed or unexposed cells in an organ. A Monte Carlo track simulation study reported that roughly 50% of the cell nuclei in a cell population receive a “hit” from a radiation track produced by an acute exposure of 270 μGy of ^{137}Cs γ -rays (Watanabe 2012). This situation differs substantially from that in higher dose regions when analyzed by the concept of “absorbed dose.” Instead, an “elemental dose,” given by a single radiation track to a target volume, as developed by microdosimetry, should be used in the analysis of low-dose effects (ICRU 1983). This dose was estimated as 0.87 mGy for ^{60}Co γ -rays by Booz and Feinendegen in 1998 (Booz and Feinendegen 1988). In this low-dose situation, two groups exist in a cell population: one consists of cells that received the

elemental dose from a single track, and the other consists of cells that did not receive any dose. Although several papers reveal the clinical utility of inhomogeneous dose distribution in tissue-based level such as micro-slit-beam radiation therapy using synchrotron-generated X-ray beams (Mukumoto et al. 2017), to our knowledge, the radiation risk assessment of such inhomogeneous dose distribution in a living system has not yet been determined. Thus, two main factors are recognized as governing the biological influences of low-dose irradiation. The first is the radiation-induced bystander effect (RIBE) at the cell or tissue levels (Nagasawa and Little 1992; Prise and O'Sullivan 2009; Blyth and Sykes 2011), and the second is radiation sensitivity at the individual or population levels (Fukunaga et al. 2016; Fukunaga and Yokoya 2016).

A simple definition of RIBE is that “a cell responds to the fact that its neighbor(s) have been irradiated” (Fig. 1), and a key characteristic of RIBE, in contrast to direct irradiation effects, is the dose-response relationship. Instead of an increased response with an increasing radiation dose, the RIBE becomes saturated at relatively low doses (typically less than 1 Gy) (Prise and O'Sullivan 2009). Since the first report of RIBE by Nagasawa and Little (Nagasawa and Little 1992), low-dose radiation research, pursued in a variety of experimental systems that have focused on cultured cells, has suggested that DNA damage and genetic instability due to RIBE are responsible for some of the effects of low-dose radiation on cells (Prise and O'Sullivan 2009; Blyth and Sykes 2011). However, the underlying biological mechanism has yet to be clarified, thereby maintaining the assessment of radiation risks at low doses in a very controversial status. To our knowledge at this time, the data accumulated concerning the cancer risks of RIBE at individual levels appears to be insufficient (Blyth and Sykes 2011). The radiation exposure levels of local residents have truly been extremely low after the FNPP accident;

however, the future effects on people's health remain in doubt because of the lack of knowledge of the underlying mechanism of the biological effects of low-dose radiation, including RIBE.

Severe reactions to radiation therapy have been reported in clinical practice in individuals suffering from DNA damage response (DDR) defective disorders, such as ataxia-telangiectasia (AT), Nijmegen breakage syndrome, Fanconi anemia, and DNA ligase IV deficiency (Pollard and Gatti 2009). These patients, and other heterozygous carriers, can be associated with radiation hypersensitivity, as well as with cancer predisposition, although the underlying mechanism of radiation-induced carcinogenesis in these individuals remains to be determined (Jongmans and Hall 1999). Additionally, several single nucleotide polymorphisms (SNPs) have been reported as potential biomarkers for predicting the development of radiation pneumonitis (RP), one of the adverse effects of radiation therapy. These SNPs in genes related to DDR, cell signaling, and inflammation have a statistically confirmed association with RP, suggesting their direct or indirect involvement in radiation sensitivity (Zhang et al. 2010; Yin et al. 2011; Yin et al. 2012; Edvardsen et al. 2013; Tang et al. 2016). In 2016, we reported the estimated prevalence of heterozygous carriers of major DNA repair disorders in Fukushima, such as AT, Werner syndrome, and hereditary breast ovarian cancer syndrome (Fukunaga and Yokoya 2016).

We used the Integrative Japanese Genome Variation Database (iJGVD, available at <https://ijgvd.megabank.tohoku.ac.jp/>) in 2016 to investigate the variations in these radiosensitivity-related gene in a large population. This is an online database released by the Tohoku Medical Megabank Organization, Tohoku University that covered the whole genome sequences of 2049 healthy individuals in the part of northeastern Japan that was

seriously damaged in the 2011 disaster (Nagasaki et al. 2015; Yamaguchi-Kabata et al. 2015). We found that a large number of individuals have SNPs associated with the risk of developing RP, indicating a greater than expected genetic diversity of radiation sensitivity (Fukunaga et al. 2016). We also investigated the frequency of mutations in the *WRN* gene among residents in the northeastern part of Japan to determine the presence of Werner syndrome (WS) patients and/or *WRN* heterozygous carriers. This syndrome is a rare genetic disorder that causes deficiency of the WRN protein and presents with features suggestive of accelerated aging and radiation hypersensitivity (UNSCEAR 2011). Genetic instability has been observed in WS patients and in *WRN* heterozygotes, suggesting an association between *WRN* gene mutations and radiation hypersensitivity, as well as cancer predisposition, in these persons (Moser et al. 2000). To our knowledge, our result is the first evidence of a direct link between the presence of heterozygous carriers with genetic instability and radiation hypersensitivity in the northeastern part of Japan after the 2011 FNPP accident (data not shown). Consideration of these patients and heterozygous carriers with other radiation sensitivity disorders would give a much wider than expected individual variation in radiation sensitivity.

The target of current medical approaches has shown a gradual change from the “average person” to “each person.” However, in the matter of radiation protection, this approach has not been sufficiently developed (Fukunaga et al. 2016). Using the iJGVD and other databases, we are now investigating the heterozygous carriers of these disorders in the northeastern part of Japan including Fukushima, however because of the sample size we cannot significantly detect such high-risk individuals, with the exception of the *WRN* heterozygous mutation carriers. Other diseases including Cockayne’s syndrome are extremely rare, therefore heterozygous carriers could not be significantly detected

because of methodological limitations. It is true that the iJGVD has great potential for future innovations from the view point of precision medicine, nevertheless, the present sample size is too small and needs to be urgently expanded. As previously mentioned, several genomic studies have recently determined a wider than expected variation in radiation sensitivity among individuals, indicating an inadequacy in the conventional approach to radiological protection. As shown in Fig. 2, we need a personalized radiation risk assessment and care approach that considers the genetic variation of radiation sensitivity in individuals to replace the current dose-dependent risk assessment approach. Appropriate actions are expected in Fukushima health facilities for the coming era of precision medicine.

Acknowledgements:

The authors thank Dr. Yumi Yamaguchi-Kabata and Prof. Masao Nagasaki, Tohoku Medical Megabank Organization, Tohoku University, and Dr. Masataka Taguri, Graduate School of Medicine, Yokohama City University for their helpful comments. This work was supported by Marubun Research Promotion Foundation. The funding organization had no role in the design and conduct of the study; in the collection, management, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

Conflict of Interest:

The authors declare no conflict of interest.

References

- Blyth, B.J. & Sykes, P.J. (2011) Radiation-induced bystander effects: what are they, and how relevant are they to human radiation exposures? *Radiat. Res.*, 176, 139-157.
- Booz, J. & Feinendegen, L.E. (1988) A microdosimetric understanding of low-dose radiation effects. *Int. J. Radiat. Biol. Relat. Stud. Phys. Chem. Med.*, 53, 13-21.
- Edvardsen, H., Landmark-Høyvik, H., Reinertsen, K.V., Zhao, X., Grenaker-Alnæs, G.I., Nebdal, D., Syvänen, A.C., Rødningen, O., Alsner, J., Overgaard, J., Borresen-Dale, A.L., Fosså, S.D. & Kristensen, V.N., (2013) SNP in TXNRD2 Associated With Radiation-Induced Fibrosis: A Study of Genetic Variation in Reactive Oxygen Species Metabolism and Signaling. *Int. J. Radiat. Oncol. Biol. Phys.*, 86, 791-799.
- Fukunaga, H. & Kumakawa, H. (2015) Mental Health Crisis in Northeast Fukushima after the 2011 Earthquake, Tsunami and Nuclear Disaster. *Tohoku J. Exp. Med.*, 237, 41-43.
- Fukunaga, H. & Yokoya, A. (2016) Low-dose radiation risk and individual variation in radiation sensitivity in Fukushima. *J. Radiat. Res.*, 57, 98-100.
- Fukunaga, H., Yokoya, A. & Taki, Y. (2016) Now Is the Time to Consider Personalized Effective Dose. *Int. J. Radiat. Oncol. Biol. Phys.*, 96, 479-480.
- Harada K.H., Niisoe, T., Imanaka, M., Takahashi, T., Amako, K., Fujii, Y., Kanameishi, M., Ohse, K., Nakai, Y., Nishikawa, T., Saito, Y., Sakamoto, H., Ueyama, K., Hisaki, K., Ohara, E. et al. (2014) Radiation dose rates now and in the future for residents neighboring restricted areas of the Fukushima Daiichi Nuclear Power Plant. *Proc. Natl. Acad. Sci. U. S. A.* 111, E914-E923.
- Hayano, R.S., Tsubokura, M., Miyazaki, M., Satou, H., Sato, K., Masaki, S. & Sakuma, Y. (2013) Internal radiocesium contamination of adults and children in Fukushima 7

- to 20 months after the Fukushima NPP accident as measured by extensive whole-body-counter surveys. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.*, 89, 157-163.
- Hosoda, M., Tokonami, S., Omori, Y., Ishikawa, T. & Iwaoka, K. (2016) A comparison of the dose from natural radionuclides and artificial radionuclides after the Fukushima nuclear accident. *J. Radiat. Res.*, 57, 422-430.
- International Commission on Radiation Units and Measurements (ICRU). (1983) *Microdosimetry (Report 36)*, International Commission on Radiation Units and Measurements, Inc., Bethesda, MD.
- Ishigaki, A., Higashi, H., Sakamoto, T. & Shibahara, S. (2013) The Great East-Japan Earthquake and devastating tsunami: an update and lessons from the past Great Earthquakes in Japan since 1923. *Tohoku J. Exp. Med.*, 229, 287-299.
- Jongmans, W. & Hall, J. (1999) Cellular responses to radiation and risk of breast cancer. *Eur. J. Cancer*, 35, 540-548.
- Kamiya, K., Ishikawa, T., Yasumura, S., Sakai, A., Ohira, T., Takahashi, H., Ohtsuru, A., Suzuki, S., Hosoya, M., Maeda, M., Yabe, H., Fujimori, K., Yamashita, S., Ohto, H. & Abe, M. (2016) EXTERNAL AND INTERNAL EXPOSURE TO FUKUSHIMA RESIDENTS. *Radiat. Prot. Dosimetry*, 171, 7-13.
- Kamiya, K., Ozasa, K., Akiba, S., Niwa, O., Kodama, K., Takamura, N., Zaharieva, E.K., Kimura, Y. & Wakeford, R. (2015) Long-term effects of radiation exposure on health. *Lancet*, 386, 469-478.
- Moser, M.J., Bigbee, L.W., Grant, S.G., Emond, M.J., Langlois, R.G., Jensen, R.H., Oshima, J. & Monnat, R.J. (2000) Genetic instability and hematologic disease risk in Werner syndrome patients and heterozygotes. *Cancer Res.*, 60, 2492-2496.

- Mukumoto, N., Nakayama, M., Akasaka, H., Shimizu, Y., Osuga, S., Miyawaki, D., Yoshida, K., Ejima, Y., Miura, Y., Umetani, K., Kondoh, T. & Sasaki, R. (2017) Sparing of tissue by using micro-slit-beam radiation therapy reduces neurotoxicity compared with broad-beam radiation therapy. *J. Radiat. Res.*, 58, 17-23.
- Nagasaki, M., Yasuda, J., Katsuoka, F., Nariai, N., Kojima, K., Kawai, Y., Yamaguchi-Kabata, Y., Yokozawa, J., Danjoh, I., Saito, S., Sato, Y., Mimori, T., Tsuda, K., Saito, R., Pan, X. et al. (2015) Rare variant discovery by deep whole-genome sequencing of 1,070 Japanese individuals. *Nat. Commun.*, 6, 8018.
- Nagasawa, H. & Little, J.B. (1992) Induction of Sister Chromatid Exchanges by Extremely Low Doses of α -Particles. *Cancer Res.*, 52, 6394-6396.
- Pollard, J.M. & Gatti, R.A. (2009) Clinical radiation sensitivity with DNA repair disorders: an overview. *Int. J. Radiat. Oncol. Biol. Phys.*, 74, 1323-1331.
- Prise, K.M. & O'Sullivan, J.M. (2009) Radiation-induced bystander signalling in cancer therapy. *Nat. Rev. Cancer*, 9, 351-360.
- Shibahara, S. (2011) The 2011 Tohoku Earthquake and Devastating Tsunami. *Tohoku J. Exp. Med.*, 223, 305-307.
- Tang, Y., Liu, B., Li, J., Wu, H., Yang, J., Zhou, X., Yi, M., Li, Q., Yu, S. & Yuan X. (2016) Genetic variants in PI3K/AKT pathway are associated with severe radiation pneumonitis in lung cancer patients treated with radiation therapy. *Cancer Med.*, 5, 24-32.
- United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR). (2011) *SOURCES AND EFFECTS OF IONIZING RADIATION. UNSCEAR 2008 Report*, United Nations, New York, NY.

United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR).

(2014) *SOURCES, EFFECTS AND RISKS OF IONIZING RADIATION. UNSCEAR 2013 Report*, United Nations, New York, NY.

United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR).

(2015) *DEVELOPMENTS SINCE THE 2013 UNSCEAR REPORT ON THE LEVELS AND EFFECTS OF RADIATION EXPOSURE DUE TO THE NUCLEAR ACCIDENT FOLLOWING THE GREAT EAST-JAPAN EARTHQUAKE AND TSUNAMI. A 2015 white paper to guide the Scientific Committee's future programme of work*. United Nations, New York, NY.

Watanabe, R. (2012) Microscopic energy distribution of low-dose radiation. *Radiat.*

Biol. Res. Commun., 47, 335-346. (in Japanese)

World Health Organization (WHO). (2013) *Health risk assessment from the nuclear accident after the 2011 Great East Japan earthquake and tsunami, based on a preliminary dose estimation*, WHO Press, Geneva.

Yamaguchi-Kabata, Y., Nariai, N., Kawai, Y., Sato, Y., Kojima, K., Tateno, M., Katsuoka, F., Yasuda, J., Yamamoto, M. & Nagasaki, M. (2015) iJGVD: an integrative Japanese genome variation database based on whole-genome sequencing. *Hum. Genome Var.*, 2, 15050.

Yasumura, S., Hosoya, M., Yamashita, S., Kamiya, K., Abe, M., Akashi, M., Kodama, K., Ozasa, K. & Fukushima Health Management Survey Group. (2012) Study protocol for the Fukushima Health Management Survey. *J. Epidemiol.*, 22, 375-383.

Yin, M., Liao, Z., Liu, Z., Wang, L-E., Gomez, D., Komaki, R. & Wei, Q. (2011) Functional Polymorphisms of Base Excision Repair Genes XRCC1 and APEX1 Predict Risk of Radiation Pneumonitis in Patients With Non-Small Cell Lung

Cancer Treated With Definitive Radiation Therapy. *Int. J. Radiat. Oncol. Biol.*

Phys., 81, e67-e73.

Yin, M., Liao, Z., Liu, Z., Wang, L-E., O'Reilly, M., Gomez, D., Li, M., Komaki, R. &

Wei, Q. (2012) Genetic variants of the nonhomologous end joining gene LIG4 and

severe radiation pneumonitis in nonsmall cell lung cancer patients treated with

definitive radiotherapy. *Cancer*, 118, 528-535.

Zhang, L., Yang, M., Bi, N., Fang, M., Sun, T., Ji, W., Tan, W., Zhao, L., Yu, D., Lin,

D. & Wang, L. (2010) ATM Polymorphisms Are Associated With Risk of

Radiation-Induced Pneumonitis. *Int. J. Radiat. Oncol. Biol. Phys.*, 77, 1360-1368.

Figure Legends

Fig. 1. Radiation-induced bystander effect.

Cells that did not directly receive radiation doses but did receive signals from nearby or neighboring irradiated cells behave as though they have been exposed, showing genomic instability and other abnormalities. The effect is mediated either through gap junctions or via soluble factors released by irradiated cells.

Fig. 2. Personalized radiation risk assessment and care system in the large-scale nuclear accident management.

There are two radiation risk assessment and care systems after the nuclear accident: the dose-dependent (A) and the personalized (B). High, moderate, and low radiation-induced health risk person are shown in red, yellow, and green, respectively.