



**QUEEN'S
UNIVERSITY
BELFAST**

Analysis of Strongyle Egg Shedding Consistency in Horses and Factors That Affect It

Lester, H. E., Morgan, E. R., Hodgkinson, J. E., & Matthews, J. B. (2018). Analysis of Strongyle Egg Shedding Consistency in Horses and Factors That Affect It. *Journal of Equine Veterinary Science*, 60, 113-119. <https://doi.org/10.1016/j.jevs.2017.04.006>

Published in:
Journal of Equine Veterinary Science

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2017 Elsevier Inc.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>, which permits distribution and reproduction for noncommercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

1 Analysis of strongyle egg shedding consistency in horses and
2 factors that affect it

3

4 H.E. Lester^{a,c*}, E.R. Morgan^b, J.E. Hodgkinson^c, J.B. Matthews^a

5 ^a*Moredun Research Institute, EH26 0PZ, UK*

6 ^b*School of Veterinary Sciences, University of Bristol, BS40 5DU, UK.*

7 ^c*Institute of Infection and Global Health, University of Liverpool, L3 5RF, UK*

8

9 *Corresponding author.

10 *E-mail address:* lester.hannah@yahoo.com (H.E. Lester).

11

12 **Abstract**

13 Strongyle egg shedding consistency in horses and factors affecting consistency were
14 investigated. Faecal samples were collected from 26 equine populations over one
15 grazing season. Samples were collected on four 'screening' occasions (S1–S4) and
16 FEC performed using a sensitive method (to 1 eggs per gram (epg) egg detection
17 limit). On each screening occasion, FEC were assigned an egg shedding category: 1
18 (<50 epg) to 7 (>500 epg); and a treatment category: <200 epg (no treatment) or ≥
19 200 epg (treatment). Rank changes in shedding and treatment categories between S1
20 and subsequent screening occasions were calculated. Factors affecting the likelihood
21 of an individual changing shedding or treatment category were assessed using
22 multivariable logistic regression of FEC data from horses that had not received
23 anthelmintic during the study. In total, 573 horses were sampled at S1, 468 at S2, 417
24 at S3 and 83 at S4. Results showed that between S1 and S4, 73.5% (61/83) horses
25 remained in the same egg shedding category and 94.0% (78/83) in the same treatment
26 category. For horses that did not receive anthelmintic (n=304), 90.4% (225/304)
27 remained in the same egg shedding category. Horses under 5 years-old were more
28 likely to change egg shedding (odds ratio, OR 3.3, 95% CI: 1.22-8.46) and treatment
29 (OR 2.8, 95% CI: 1.1-6.3) categories compared to older horses. These results suggest
30 a high level of consistency in strongyle egg shedding in individuals within one
31 season, and withholding anthelmintics from horses with negative or low (i.e. <50 epg)
32 FEC does not appear to be associated with significant increases in egg shedding.

33

34

35 *Keywords:* Horse, Cyathostomins, Strongyle egg shedding, Targeted anthelmintic
36 treatment

37 1. Introduction

38 Anthelmintic resistant nematodes pose a threat to equine health and welfare.
39 Targeting anthelmintic treatments to individual horses based on levels of nematode
40 egg shedding is proposed to achieve control, whilst reducing anthelmintic use and
41 selection for resistance. Helminth parasites are typically overdispersed in their hosts:
42 i.e. relatively few individuals within populations are infected with the majority of the
43 associated parasite population [1-4]. Strongyle nematodes in the large intestine are the
44 most important equine parasites and have a high prevalence [5]. Overdispersion has
45 been demonstrated for strongyle faecal egg counts (FEC) in horses, with most
46 individuals shedding relatively low numbers of eggs [6-8]. This overdispersion
47 underpins the rationale behind targeted anthelmintic treatment strategies [5, 9-10],
48 whereby only horses identified as shedding above a specific threshold of eggs
49 (typically ≥ 200 eggs per gram (epg)) are recommended for treatment. Several studies
50 have shown consistency in strongyle egg shedding patterns in individuals over time
51 [7, 11-14], which could further enhance efficient targeting of anthelmintic treatments
52 and reduce the required frequency of FEC testing. The aim here was to investigate
53 strongyle egg shedding consistency in populations of horses in the UK that followed a
54 targeted anthelmintic treatment programme, with a focus on horses identified as low
55 strongyle egg shedders (< 50 epg), to examine whether non-treatment of these
56 individuals was associated with increases in egg shedding levels over time. This study
57 also analysed factors associated with shedding consistency in individuals within these
58 populations. A **longitudinal cohort** study design was used.

59

60 2. Materials and methods

61 2.1. Study population

62 During 2010–2012, 573 horses from 26 equine holdings (yards) in Scotland and
63 England were recruited (see *supplementary data*). The number of individual's resident
64 on each yard ranged from 7-72 (median, 20 horses). Most horses (i.e. those on 19
65 yards) received moxidectin prior to the start of the study, although all had received a
66 macrocyclic lactone (ML) within the six months prior to the start of the study
67 (n=573). On three yards, information on which anthelmintic was last administered
68 was not supplied. Most horses were resident at livery yards (n=23 yards). The
69 remainder comprised of a non-Thoroughbred stud farm, a sport horse yard and a
70 rescue/welfare sanctuary. In total, 573 horses were screened at S1, 468 horses at S2
71 and 417 at S3. A total of 83 horses on three yards were screened at S4. Each horse
72 was assigned an age category. The age distribution was: foal (<2 years; n=26),
73 youngster (≥ 2 and <5 years; n=68), adult (≥ 5 and < 18 years; n=418) and geriatric
74 (≥ 18 years; n=61).

75

76 All horses included in this study had access to grazing, were at pasture during the
77 study and had been treated with a ML anthelmintic within the previous 6 months.
78 Horses were at pasture for a minimum of eight hours per day, and were grazed on the
79 same pastures for the duration of the trial (i.e. returned to the same pasture after
80 sampling and treatment). Yards with a minimum of 10 horses were included and all
81 horses were subject to the same anthelmintic treatment regimen (Section 2.2). Each
82 yard was supplied with a questionnaire to provide information on the demographics
83 (i.e. yard function, number of horses etc.), anthelmintic usage (frequency of treatment,
84 last product used, type of deworming programme followed) and general management

85 practices. Yard managers were asked to supply the age of the horses included in the
86 study, but not specific details on breed and sex. On all yards, the manager was the
87 point of contact and was responsible for completion of the questionnaire, coordinating
88 sample collection and postage, and administering anthelmintic treatments.

89

90 2.2. Targeted treatment protocol

91 Horses that were previously treated with anthelmintic were sampled after a minimum
92 period of 18 weeks had elapsed after administration of moxidectin, or 14 weeks for
93 ivermectin. The first sample (Screen 1, S1, **n=573**) was collected between February
94 and March. Horses with FEC ≥ 200 epg were treated with pyrantel embonate
95 following the manufacturer's instructions (Strongid-P^{TMa} at a dose rate of 19mg/kg
96 bodyweight). All horses were FEC screened 8-10 weeks later: this was based on a 6-
97 week strongyle egg reappearance period (ERP) for pyrantel embonate [15, 16], plus
98 an additional two weeks (Screen 2, S2; May/June, **n=468**). Horses with FEC ≥ 200
99 epg at S2 were treated with ivermectin (Eqvalan[®] oral paste for horses^b; 0.2mg/kg).
100 All horses were screened 10–12 weeks later based on an ivermectin strongyle ERP of
101 8-10 weeks [15], plus two additional weeks (Screen 3, S3; August/September,
102 **n=417**). At this point, horses with a FEC of ≥ 200 epg were treated with moxidectin
103 (Equest^{®c}; 0.4 mg/kg). On some yards a fourth screen (S4; October/December, **n=83**)
104 was performed on horses that had not received anthelmintic treatment following S3.

105

106 2.3. Sample collection and faecal egg count methodology

107 Samples were collected from freshly passed faeces and placed into individually
108 labeled zip-lock bags. Yard managers were provided with instructions on how to
109 collect, store and post the samples, and were asked to collect at least three boli from

110 freshly voided faeces and to place these into the bag, expelling air before sealing. The
111 samples were sent immediately to the Moredun Research Institute and stored at
112 approximately 4°C. All samples were processed within 4 days of excretion to reduce
113 the effect of egg degradation [17]. A modification of the salt flotation method (1.2
114 specific gravity) with a detection limit down to 1 epg was used [18]. All samples were
115 analysed in duplicate by taking two 10 ml aliquots from a 100 ml dilution of a well-
116 mixed 10 g sub-sample and an average taken to estimate the epg count.

117

118 *2.4. Data analysis*

119 *2.4.1. Egg shedding and anthelmintic treatment categories*

120 Each FEC data point on each screening occasion (S1, S2, S3, S4) was assigned a
121 shedding category: 1 (0–49 epg); 2 (50–99 epg); 3 (100–199 epg); 4 (200–299 epg); 5
122 (300–399 epg); 6 (400–499 epg) and 7 (>500 epg). Each FEC data point was also
123 assigned a treatment category: < 200 epg (0, no treatment) or \geq 200 epg (1,
124 treatment).

125

126 *2.4.2. Egg shedding consistency*

127 To assess shedding consistency, the rank change in shedding category between S1 and
128 S2, S1 and S3 and S1 and S4 was calculated. For example, if the category was ranked
129 as 1 for S1 and 6 for S2, the rank change would be 5. Conversely, if the category was
130 measured as 6 for S1 and 3 for S3, the rank change would be -3. If the category
131 remained the same, rank change assigned was 0.

132

133 2.4.3. *Anthelmintic treatment consistency*

134 Treatment category consistency was determined across S1 and S2, S1 and S3 and S1
135 and S4. If an individual's FEC changed from < 200 epg to ≥ 200 epg between two
136 sampling points, the rank assigned was 1. If it changed from ≥ 200 epg to < 200 epg,
137 it was assigned as -1, and if it remained above or below the 200 epg threshold, was
138 assigned 0.

139

140 2.4.4. *Egg shedding and treatment consistency analysis*

141 For each sampling occasion, the number and percentage of horses in each shedding
142 and treatment category were calculated. Further, the number and percentage of horses
143 that were in each rank change category of shedding (-6 to 6) and treatment (-1, 0 or 1)
144 between S1 and each subsequent screening occasion was calculated. To test if there
145 was a significant difference ($p < 0.05$) in the proportion of horses falling into each
146 shedding and treatment category between screening occasions, a binomial test was
147 performed using the prop.test function in RStudio. All analyses were performed in
148 RStudio, version 2.15.1 (The R Foundation for Statistical Computing, 2012). The
149 analysis that was applied to the anthelmintic-treated horses was applied to horses that
150 had not received anthelmintic during the study to assess if no treatment was
151 associated with lower shedding consistency and rising FEC over time. All horses
152 included in this analysis had $FEC < 200$ epg at S1.

153

154 2.4.5. *Factors affecting egg shedding and anthelmintic treatment consistency*

155 The effect of age, last anthelmintic administered and number of weeks after the
156 expected ERP of each anthelmintic on the likelihood of a horse changing shedding or

157 treatment category between S1 and S3 was assessed using multivariable logistic
158 regression. Factors included in the initial model are shown in Table 1. Regression
159 analysis was performed using the GLM function in R (RStudio, version 2.15.1 (The R
160 Foundation for Statistical Computing, 2012)), specifying the family as binomial,
161 linked to logit transformation, g , where P is the probability of a horse changing
162 shedding or treatment category, β_i is the model (slope) coefficient and X_i is the
163 explanatory variable (Equation 1).

$$164 \quad g = \ln[P/(1-P)] = \beta_0 + \beta_1 X_0 + \dots + \beta_k X_k$$

165 Equation 1

166 The probability of a change in egg shedding and treatment category was estimated
167 using Equation 2.

$$168 \quad P = \exp[\beta_0 + \beta_1 X_0 + \dots + \beta_k X_k] / (1 + \exp[\beta_0 + \beta_1 X_0 + \dots + \beta_k X_k])$$

169 Equation 2

170 Regression models were initially populated with all potential explanatory variables
171 (age category, last treatment, time in weeks since expected ERP elapsed, Table 1),
172 then variables with the highest, non-significant p-values removed in a stepwise
173 process until a model with only significant terms remained. The effect of removing
174 factors from the model was evaluated using log-likelihood ratio tests (LRT) [19, 20].
175 P -values (Wald) of ≤ 0.05 indicated factors that had a significant influence on
176 changing shedding or treatment category in the final model. The Hosmer-Lemeshow
177 test [21] was used to assess overall model fit using the ‘ResourceSelection’ package
178 [22].

179

180 3. Results

181

182 3.1. Consistency of strongyle egg shedding and treatment with anthelmintic

183 At S1, 70.0% (401/573) of horses were shedding less than 50 epg (Figure 1a). This
184 level of shedding was not significantly different at S2 (65.0%, 304/468) and S3
185 (66.2%, 276/417), with the proportions in shedding categories 2 and 3 also not
186 significantly different compared to S1. The percentage of horses in category 7 was
187 low compared to the percentage of horses in category 1: for example, 5.8% (33/573)
188 at S1 and 0.0% (0/83) at S4. On each occasion, the percentage of horses in the 'no
189 treatment' category ranged from 83.6% (479/573) at S1 to 91.6% (76/83) at S4
190 (Figure 1b).

191

192 3.2. Egg shedding and treatment consistency

193 Most horses remained in the same egg shedding and treatment category over time
194 (Figure 2a). From S1 to S2, 61.5% horses (288/468) remained in the same shedding
195 category; from S1-S3, 58.3% (243/417), and from S1-S4, 73.5% (61/83). Between S1
196 and S2, 16.6% (77/468) horses moved to a lower shedding category and 22.0%
197 (103/468) to a higher shedding category. Between S1 and S3, 23.7% (99/417) moved
198 into a lower shedding category and 18.0% (75/417) moved into a higher shedding
199 category. Between S1 and S4, 14.5% (12/83) horses moved into a lower shedding
200 category and 8.3% (10/83) individuals moved to a higher shedding category.
201 Between S1 and S2, S1 and S3 and S1 and S4, 81.8% (383/468), 82.7% (345/417)
202 and 94.0% (78/83) horses remained in the same treatment category, respectively
203 (Figure 2b). The egg shedding consistency in a sub-group that did not receive
204 anthelmintic during the entire study ($n=304$) was investigated. The proportion of

205 horses shedding < 50 epg at S1 was 97.3% (296/304), at S2, 84.9% (258/304) and at
206 S3, 85.5% (213/249). The change in proportion of horses shedding <50 epg was
207 consistent throughout the study ($p>0.05$). Between S1 and S2, 92.1% (280/304)
208 horses remained in the same shedding category, 0.3% (1/304) were in a lower
209 shedding category and 7.9% (23/304) in a higher shedding category. Between S1 and
210 S3, the percentage of horses that remained in the same shedding category was 90.4%
211 (225/304) and the percentage of horses that were in a higher shedding category was
212 9.6% (24/304).

213

214 *3.3. Factors associated with changing strongyle egg shedding category or treatment* 215 *category over time*

216 Only horses (n=346) for which information on age and last anthelmintic treatment
217 were available were included in these analyses. Factors affecting the likelihood of a
218 horse changing egg shedding category were investigated (Model 1). In the final
219 model, age (young horses ≥ 2 - <5 years) and last treatment with moxidectin were
220 identified as significant explanatory variables (Table 2). Young horses (≥ 2 - < 5
221 years) were more likely to change strongyle egg shedding category compared to foals
222 (<2 years), adult horses (≥ 5 - <18 years) and geriatric horses (≥ 18 years, OR=3.3,
223 95% CI=1.22 – 8.46, $p=0.02$), while horses that had received moxidectin at the last
224 anthelmintic treatment were less likely to change strongyle egg shedding category
225 compared to those that received ivermectin or pyrantel (OR=0.15, 95% CI=0.05-0.17,
226 $p<0.0001$). Factors affecting the likelihood of a horse changing treatment category
227 between S1 and S3 were also investigated (Model 2). In the final model (Table 2),
228 young horses (≥ 2 - < 5 years) were more likely to change treatment category
229 compared to foals (<2 years), adult horses (≥ 5 - <18 years) and geriatric horses (≥ 18

230 years, OR=2.8, 95% CI = 1.1-6.3, $p=0.03$) and horses that received moxidectin as
231 their last treatment were significantly less likely to change treatment group compared
232 to horses that received ivermectin or pyrantel (OR=0.15, 95% CI=0.1-0.4, $p < 0.0001$).

233

234 **4. Discussion**

235 Knowledge that horses shed strongyle eggs at consistent levels over time can help
236 underpin evidence-based targeted treatment control programmes [23]. Currently,
237 FEC-directed targeted treatment programmes are recommended for adult horses,
238 whereby individuals are screened for strongyle egg shedding every 4-6 weeks during
239 the grazing season, and those excreting ≥ 200 epg treated with anthelmintic [10]. Here,
240 strongyle egg shedding in horses, analysed over three to four sampling occasions
241 within a grazing season, was found to be consistent. These results agree with those of
242 previous studies which sampled horses over longer periods (1–3 years) [12-14] and
243 shorter periods [11]. Each of these studies reported a high level of shedding
244 consistency in horses sampled from the general population. One study reported that, if
245 the first two FEC were 0 epg, there was an 82% probability that the third FEC would
246 be 0 epg and a 91% chance that it would be < 200 epg. Additionally, these authors
247 found that if the first two counts were < 200 epg, there was an 84% chance that the
248 next FEC would be < 200 epg, and if the first two FEC were ≥ 200 epg, there was a
249 59% probability that the next FEC would be ≥ 200 epg [12]. This latter study
250 demonstrated that egg shedding was consistent over a longer sampling period
251 compared to that of the current one, particularly in horses that were measured as
252 shedding 0 epg at the start. In a later survey [13], which followed a similar sampling
253 time frame to ours, horses were treated with anthelmintic when the FEC was
254 measured as ≥ 250 epg and only data from horses that did not receive anthelmintic

255 were analysed. The authors reported that if the first two FEC were 0 epg, there was a
256 62% probability that the maximum FEC of the next seven samples would be 0 epg,
257 and if the first two FEC were 0 epg, there was an 88% probability that the maximum
258 FEC of the next seven counts would be <200 epg. The authors concluded that, for
259 individual horses, the magnitude of the initial FEC was significantly positively
260 correlated to the maximal FEC of the subsequent eight counts [13]. In a further study,
261 the repeatability of strongyle egg counts was assessed in naturally infected horses
262 [14]. In that study, samples were collected over nine consecutive months, and
263 analysed using a McMaster method with an egg count detection limit of 20 epg. These
264 authors defined repeatability as the variance between horses divided by the total
265 variance, meaning that a value of 0 indicates no consistency in FEC and a value of 1
266 indicates perfect consistency. Using raw egg counts (i.e. the number of actual eggs
267 counted before applying the multiplication factor), the within-horse repeatability was
268 0.52 in all horses and 0.53 when horses that had received treatment were excluded
269 from the analysis [14]. Despite each of the studies described following different
270 counting methodology and distinctive statistical analyses, all demonstrate that
271 strongyle egg shedding in horses was consistent over short and longer sampling time-
272 frames, especially in low egg shedding categories, indicating that such horses tend to
273 excrete low numbers of eggs in the absence of anthelmintic treatment, over a
274 prolonged period. In contrast to these reports, one study examined strongyle egg
275 shedding consistency in ponies managed for conservation purposes, which remained
276 largely untreated with anthelmintic over several years, and found that shedding
277 consistency at individual level was generally weak [7]. In that study, FEC data were
278 analysed using general additive mixed models to estimate repeatability of FEC at
279 individual level and to test for differences in mean FEC amongst populations and age

280 classes. Climate and season were found to exert a significant effect on FECs
281 measured in individuals in populations that did not receive anthelmintic and a strong
282 interaction was identified between age and climate. The lack of individual consistency
283 observed by Wood et al. (2013) compared with other studies [11-14] could be due to
284 the length of time over which the data were collected, the FEC method used, the
285 absence of anthelmintic treatments, the type of statistical analysis undertaken, or the
286 nature of and level of exposure to parasite infection.

287 Past studies investigating strongyle egg shedding consistency focused on estimates of
288 strongyle FEC using McMaster methods with egg detection limits (dl) between 20
289 and 50 epg [7, 11-14]. In the current study, a more sensitive count method was used.
290 FEC generated using McMaster methods tend to generate higher epg estimates and
291 greater variance. In addition, using a FEC method with a higher egg dl (i.e.
292 multiplication factor), the methodology will be less sensitive to relatively small
293 changes in egg abundance, and larger multiplication factors will artificially inflate
294 variance [6, 24]. This may potentially lead to lower observed consistency between
295 egg counts from the same individual over time because of the greater degree of
296 artefactual variation in FEC. On the other hand, the higher number of false negative
297 FEC arising from the limited dl of traditional McMaster methods could artificially
298 increase apparent consistency of 0 measured epg over time in low-shedding horses. In
299 the present study, measures were taken to ensure samples were collected, stored and
300 processed in a way to minimise egg degradation [17], to ensure that a representative
301 sample was collected to reduce egg clumping in faeces [10], and to minimise the
302 effects of rounding error by using a very sensitive FEC method. However, the way in
303 which the samples were collected and handled on each yard on each sampling
304 occasion could have impacted the consistency of the results.

305 Here, egg shedding consistency was higher in adult and geriatric horses compared to
306 youngsters. This lower level of consistency in FEC shedding in younger horses could
307 be related to lower immunity [25] compared with older horses, such that variation in
308 parasite challenge is less buffered and feeds through to downstream variation in FEC.

309 In the current study, horses that were treated with moxidectin prior to the start of the
310 study were less likely to change shedding or treatment category compared to those
311 that had been treated with ivermectin or pyrantel. A likely reason for this is that
312 shedding would be lower in horses previously treated with moxidectin due to its
313 persistent effect against parasites *in vivo* [26]. Moxidectin has an elimination half-life
314 of 23.11 days compared to ivermectin (4.25 days) and pyrantel (13.43 hours) [26].
315 This means that parasites are exposed to active anthelmintic for longer periods; hence
316 the greater strongyle ERP observed after moxidectin treatment [26]. Moxidectin
317 exhibits higher larvicidal activity compared to the other two anthelmintics,
318 particularly against mucosal larvae [26-28]. This will also affect the strongyle ERP
319 observed after moxidectin administration compared to other equine anthelmintics [26,
320 29].

321 Recruitment of yards was non-random, being through veterinary practices and the
322 BHS website. Nevertheless, there was heterogeneity in the yards recruited, which is
323 likely to have provided a fairly representative sample of the UK equine population,
324 outside breeding establishments, which have a younger age profile [6]. As such,
325 further research is required to assess consistency of strongyle egg shedding in
326 populations on breeding farms.

327 It should be noted that the FEC data collected here were obtained over a single
328 grazing season, and the effect of season and climate were not accounted for, both of
329 which can impact on strongyle egg shedding patterns [7]. Furthermore, the effects of

330 management practices such as removal of faeces from pasture were not investigated,
331 which has been shown to significantly reduce FEC in donkeys that grazed pasture
332 where dung was removed twice-weekly by reducing larval populations on pasture
333 [30]. Long-term studies investigating strongyle egg shedding patterns taking account
334 of season, climate and management practices are warranted as these factors impact the
335 intensity of larval contamination on pasture, which will in turn affect the egg
336 shedding intensity downstream. This will help to better inform FEC-directed targeted
337 anthelmintic treatment programmes, and to understand the appropriate frequency of
338 FEC testing.

339 In conclusion, egg shedding and treatment status at individual level were found to be
340 relatively consistent, especially in adult horses, regardless of whether or not they had
341 been treated with anthelmintic, indicating that horses with negative or low FEC (<50
342 epg) initially were significantly more likely to have low FEC on subsequent
343 occasions, and horses with a high FEC (≥ 200 epg) were significantly more likely to
344 have a high FEC on subsequent occasions. Further, an adult horse not requiring
345 anthelmintic treatment initially (based on a 200 epg threshold), would be less likely to
346 require treatment on subsequent occasions in that season. These findings would
347 suggest that the majority of adult horses are able to regulate their strongyle burden,
348 leading to a maintained low FEC status. There is a lack of published information on
349 long term patterns of strongyle egg shedding in equine populations, especially in
350 horses managed under FEC-directed anthelmintic treatment programmes; this should
351 be assessed in future, particularly in horses that are not receiving regular anthelmintic
352 treatments.

353

354 **Acknowledgments**

355 The authors acknowledge funding from the Elise Pilkington Trust for this study. They
356 are indebted to the yard owners/managers and/or attending veterinary surgeons for the
357 supply of equine faecal samples and to Sheena Tarrant, Emma Wood and Rachel
358 Cookson for help with the FEC analysis.

359

360 **Manufacturer's details**

361 ^aStrongid-P™ paste, Elanco Animal Health, Basingstoke, Hampshire, UK.

362 ^bEqvalan® oral paste for horses, Merial Animal Health, Harlow, Essex, UK.

363 ^cEquest® Oral Gel, Zoetis UK Limited, Tadworth, Surrey, UK.

364

365 **References**

366 [1] Crofton, H.D. (1971) A model of host–parasite relationships. *Parasitol.* **63**, 343-
367 364.

368 [2] Anderson, R.M. and May RM. (1978) Regulation and Stability of Host-Parasite
369 Population Interactions: I. Regulatory Processes. *J. Anim. Ecol.* **47**, 219-247.

370 [3] Shaw, D.J. and Dobson, A.P. (1995) Patterns of macroparasite abundance and
371 aggregation in wildlife populations: a quantitative review. *Parasitol.* **111**,
372 S111-S133.

373 [4] Calabrese, J.M., Brunner, J.L. and Ostfeld, R.S. (2011) Partitioning the
374 aggregation of parasites on hosts into intrinsic and extrinsic components via an
375 extended Poisson-gamma mixture model. *PloS one.* 6:e29215.

376 [5] Kaplan, R.M. and Nielsen, M.K. (2010) An evidence-based approach to equine
377 parasite control: It ain't the 60s anymore. *Equine Vet. Edu.* **22**, 306-316.

- 378 [6] Relf, V.E., Morgan, E.R., Hodgkinson, J.E. and Matthews, J.B. (2013) Helminth
379 egg excretion with regard to age, gender and management practices on UK
380 Thoroughbred studs. *Parasitol.***140**, 641-652.
- 381 [7] Wood, E.L.D., Matthews, J.B., Stephenson, S., Slote, M. and Nussey, D.H.
382 (2013) Variation in fecal egg counts in horses managed for conservation
383 purposes: individual egg shedding consistency, age effects and seasonal
384 variation. *Parasitol.* **140**, 115-128.
- 385 [8] Lester, H.E., Spanton, J., Stratford, C.H., Bartley, D.J., Morgan, E.R.,
386 Hoddgkinson, J.E., Coumbe, K., Mair, T., Swan, B., Lemon, G., Cookson, R.
387 and Matthews, J.B. (2013). Anthelmintic efficacy against cyathostomins in
388 horses in Southern England. *Vet. Parasitol.* **197**,189-196.
- 389 [9] Gomez, H.H. and Georgi, J.R. (1991) Equine helminth infections: control by
390 selective chemotherapy. *Equine Vet. J.* **23**, 198-200.
- 391 [10] Lester, H.E. and Matthews, J.B. (2014) Faecal worm egg count analysis for
392 targeting anthelmintic treatment in horses: Points to consider. *Equine Vet. J.*
393 **46**, 139-145.
- 394 [11] Dopfer, D., Kerssens, C., Meijer, Y., Boersema, J.H. and Eysker, M. (2004).
395 Shedding consistency of strongyle-type eggs in Dutch boarding horses. *Vet.*
396 *Parasitol.* **124**, 249 - 258.
- 397 [12] Nielsen, M., Haaning, N. and Olsen, S. (2006) Strongyle egg shedding
398 consistency in horses on farms using selective therapy in Denmark. *Vet.*
399 *Parasitol.* **135**, 333 - 335.
- 400 [13] Becher, A.M., Mahling, M., Nielsen, M.K. and Pfister, K. (2010) Selective
401 anthelmintic therapy of horses in the Federal states of Bavaria (Germany) and

- 402 Salzburg (Austria): An investigation into strongyle egg shedding consistency.
403 Vet. Parasitol. **171**, 116-122.
- 404 [14] Scheuerle, M.C., Stear, M.J., Honeder, A., Becher, A.M., Pfister, K. (2016)
405 Repeatability of strongyle egg counts in naturally infected horses. Vet.
406 Parasitol. **228**, 103-107.
- 407 [15] Borgsteede, F.H., Boersma, J.H., Gaasenbeek, C.P. and van der Burg, W.P.
408 (1993) The reappearance of eggs in faeces of horses after treatment with
409 ivermectin. Vet. Q. **15**, 24-26.
- 410 [16] Mercier, P., Chick, B., Alves-Branco, F. and White, C.R. (2001) Comparative
411 efficacy, persistent effect, and treatment intervals of anthelmintic pastes in
412 naturally infected horses. Vet. Parasitol. **99**, 29-39.
- 413 [17] Nielsen, M.K., Vidyashankar, A.N., Andersen, U.V., Delisis, K., Pilegaard, K.,
414 Kaplan, R.M (2010) Effects of faecal collection and storage factors on
415 strongylid egg counts in horses. Vet. Parasitol. **20**, 56-61.
- 416 [18] Christie, M. and Jackson, F. (1982) Specific identification of strongyle eggs in
417 small samples of sheep faeces. Res.Vet. Sci. **32**,113-117.
- 418 [19] Zuur, A., Ieno, E.N., Walker, N., Saveliever, A.A. and Smith, G.M. (2009).
419 Mixed Effects Models and Extensions in Ecology with R, Springer.
- 420 [20] Crawley, M.J. (2012) Regression. In: The R Book, John Wiley & Sons, Ltd.
421 pp. 449-497.
- 422 [21] Hosmer, D.W. and Lemeshow, S. (2005) Assessing the Fit of the Model. In:
423 Applied Logistic Regression, John Wiley & Sons, Inc. pp. 143-202.
- 424 [22] Lele, S.R. (2009) A New Method for Estimation of Resource Selection
425 Probability Function. J. Wildlife Man. **73**, 122-127.

- 426 [23] Nielsen, M.K., Reinemeyer, C.R., Donecker, J.M., Leathwick, D.M.,
427 Marchiondo, A.A. and Kaplan, R.M. (2014) Anthelmintic resistance in equine
428 parasites—Current evidence and knowledge gaps. *Vet. Parasitol.* **204**, 55-63.
- 429 [24] Torgerson, P.R., Paul, M., Lewis, M.I. (2012) The contribution of simple
430 random sampling to observed variations in faecal egg counts. *Vet Parasitol.*
431 **188**, 397-401.
- 432 [25] Herd, R.P. and Gabel, A.A. (1990) Reduced efficacy of anthelmintics in young
433 compared with adult horses. *Equine Vet. J.* **22**, 164-169.
- 434 [26] Cobb, R. and Boeckh, A. (2009). Moxidectin: a review of chemistry,
435 pharmacokinetics and use in horses. *Parasites & Vectors* **2**, S5.
- 436 [27] Xiao, L., Herd, R.P., Majewski, G.A. (1994) Comparative efficacy of
437 moxidectin and ivermectin against hypobiotic and encysted cyathostomes and
438 other equine parasites. *Vet Parasitol.* **53**, 83-90.
- 439 [28] Schumacher, J. and Taintor, J. (2008) A review of the use of moxidectin in
440 horses. *Equine Vet. Edu.* **20**, 546-551.
- 441 [29] Bairden, K., Davies, H.S., Gibson, N.R. (2006) Efficacy of moxidectin 2 per
442 cent oral gel against cyathostomins, particularly third-stage inhibited larvae, in
443 horses. *Vet Rec.* **158**, 766-768.
- 444 [30] Corbett, C., Love, S., Moore, A., Burden, F., Matthews, J., Denwood, M.
445 (2014) The effectiveness of faecal removal methods of pasture management
446 to control the cyathostomin burden of donkeys. *Parasites & Vectors.* **7**:48.
447

448 **Tables**449 **Table 1.** Variables included in the initial logistic regression model

Variable	Responses
Age category	Foal (<2 years), youngster (≥ 2 and <5 years), adult (≥ 5 and < 18 years) and geriatric (≥ 18 years).
Last anthelmintic class administered	IVM, MOX, PYR
Time since expected *ERP elapsed	Weeks

Notes: *Egg reappearance period (ERP); ivermectin (IVM); moxidectin (MOX), pyrantel (PYR)

450

451

452

453

454

455 **Table 2.** Factors significantly affecting the odds of a horse changing egg shedding
 456 category (Model 1) and treatment category (Model 2) between sampling occasions S1
 457 and S3 as assessed by logistic regression. For each significant variable, the logit
 458 coefficient, the standard error (SE), the odds ratio (OR) and associated 95%
 459 confidence intervals (CI), and the significance ($p < 0.05$) are presented.

Model (Fit ¹)	Significant variable	Factor	Logit coefficient	(SE)	OR (95% CI)	<i>p</i> (Wald)
1 (0.11)			1.44	0.27	Na	<0.0001
	Age	Youngsters (2-5 years)	1.20	0.55	3.3 (1.22 – 8.46)	0.02 (LRT)
	Last treatment	MOX	-1.90	0.31	0.15 (0.05-0.17)	<0.0001
2 (0.38)			1.48	0.22	Na	<0.0001
	Age	Youngsters (2-5 years)	1.04	0.47	2.8 (1.1 - 6.3)	0.028 (LRT)
	Last treatment	MOX	-1.89	0.61	0.15 (0.1 – 0.4)	<0.0001

Notes: ¹ Model fit was assessed using the Hosmer and Lemeshow test. **MOX** = moxidectin, **SE**=standard error, **OR** = odds ratio, **Na**= not applicable; **LRT** = likelihood ratio test.

460

461 **Figure Captions**

462

463 **Figure 1.** The percentage of horses that fell into each strongyle egg shedding category (1-7)
464 on each screening occasion (S1-S4), with category 1 (<49 eggs per gram (epg) represented by
465 the darkest shade of grey and subsequent categories in lighter shades (A). The percentage of
466 horses that were either shedding <200 epg (Category 0, dark grey) or \geq 200 epg (Category 2,
467 light grey) as measured by faecal egg count (FEC) on each screening occasion (B). The width
468 of the bars is proportional to the number of observations per screening occasion. No statistical
469 differences ($p<0.05$) in proportions between screening occasions as determined by the
470 binomial test were observed.

471

472 **Figure 2.** The rank change in egg shedding category and treatment category between
473 sampling occasions (S1 and S2, S1 and S3 and S1 and S4). The percentage of horses either
474 remained in the same egg shedding category (0), increased egg shedding category (1 to 6) or
475 decreased egg shedding category (-1 to -6), with a rank change in -6 categories represented by
476 the darkest shade of grey and subsequent ascending categories in lighter shades (A). The
477 percentage of horses that were either remained in the same treatment category (0, mid-grey),
478 increased in treatment category (1, light grey) or decreased in treatment category (-1, dark
479 grey) (B) as measured by faecal egg count (FEC) on each screening occasion. The width of
480 the bars is proportional to the number of observations per screening occasion. There were no
481 statistical differences ($p<0.05$) in proportions between screening occasions as determined by
482 the binomial test.

483