



**QUEEN'S  
UNIVERSITY  
BELFAST**

## **Adiposity Associated Plasma Linoleic Acid is Related to Demographic, Metabolic Health and Haplotypes of FADS1/2 Genes in Irish Adults**

Li, K., Brennan, L., Bloomfield, J. F., Duff, D. J., McNulty, B. A., Flynn, A., Walton, J., Gibney, M. J., & Nugent, A. P. (2018). Adiposity Associated Plasma Linoleic Acid is Related to Demographic, Metabolic Health and Haplotypes of FADS1/2 Genes in Irish Adults. *Molecular Nutrition & Food Research*, [1700785]. <https://doi.org/10.1002/mnfr.201700785>

**Published in:**  
Molecular Nutrition & Food Research

**Document Version:**  
Peer reviewed version

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

**Publisher rights**  
© 2018 WILEY-VCH Verlag GmbH & Co.  
This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

**General rights**  
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**  
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

1 **Adiposity associated plasma linoleic acid is related to demographic, metabolic health and**  
2 **haplotypes of FADS1/2 genes in Irish adults.**

3 **Name of Authors:** Kaifeng Li<sup>1</sup>, Lorraine Brennan<sup>1</sup>, Jack F. Bloomfield<sup>2</sup>, Dan J. Duff<sup>2</sup>, Breige A.  
4 McNulty<sup>1</sup>, Albert Flynn<sup>3</sup>, Janette Walton<sup>3,4</sup>, Michael J. Gibney<sup>1</sup>, Anne P. Nugent<sup>1,5\*</sup>.

5 <sup>1</sup> Institute of Food and Health, School of Agriculture and Food Science, University College Dublin  
6 (UCD), Belfield Dublin, Republic of Ireland; <sup>2</sup> Chemical Analysis Laboratories, Sandycove,  
7 Dublin, Republic of Ireland, <sup>3</sup> School of Food and Nutritional Sciences, University College Cork,  
8 Cork, Ireland. <sup>4</sup> School of Biological Sciences, Cork Institute of Technology, Cork, Ireland.  
9 <sup>5</sup> School of Biological Sciences, Institute for Global Food Security, Queens University Belfast,  
10 Northern Ireland.

11 **Address Correspondence:** A.P. Nugent, Tel: +353 87 9864415

12 Email: [anne.nugent@ucd.ie](mailto:anne.nugent@ucd.ie), Address: Institute of Food and Health, University College Dublin  
13 (UCD), Belfield, Dublin 4, Republic of Ireland.

14 **Abbreviations:** AgRP, agouti-related protein; ARA, arachidonic acid; CVD, cardiovascular  
15 disease; CHD, coronary heart disease; D5D:  $\Delta$ -5-desaturase; D6D,  $\Delta$ -6-desaturase; DGLA,  
16 dihomog- $\gamma$ -linoleic acid; DNL, *de-novo* lipogenesis; FADS, fatty acid desaturase; GLA,  $\gamma$ -linoleic  
17 acid; HOMA-IR, homeostatic model assessment-insulin resistance; hs-CRP, high-sensitivity-C-  
18 reactive protein; JNK, c-Jun amino-terminal kinases; LA, linoleic acid; LD, linkage disequilibrium;  
19 MetS, metabolic syndrome; NANS, National Adult Nutrition Survey; PAI-1, plasminogen  
20 activator inhibitor-1; QUICKI, quantitative sensitivity check index; RCT, randomized control  
21 trials; SNP, single nucleotide polymorphism; TAG, triacylglycerol; TE, total energy; TLR4, toll-  
22 like receptor 4; TNF, tumor necrosis factor.

23 **Keywords:** FADS1, FADS2, genotype, linoleic acid,

24 **Abstract**

25 **Scope:** This study examined to what extent plasma linoleic acid (LA) is modified by adiposity,  
26 and explored any association between plasma LA, demographics, dietary intakes, markers of  
27 metabolic health and haplotypes of the fatty acid desaturase (FADS) 1/2 genes.

28 **Methods and results:** 820 participants with fasting blood samples from Irish National Adult  
29 Nutrition Survey were studied. Plasma fatty acids were determined using GC-MS. 15 SNPs of  
30 FADS 1/2 genes were genotyped. Plasma LA decreased while  $\gamma$ -linoleic acid and dihomo- $\gamma$ -  
31 linoleic acid increased in overweight/obese participants ( $P \leq 0.002$ ). Participants in the highest  
32 quartile of plasma LA showed decreased plasma markers of *de novo* lipogenesis, insulin resistance  
33 and of inflammation (TNF- $\alpha$ , PAI-1) ( $P \leq 0.005$ ). Adiposity (waist circumference and body fat) was  
34 strongly inversely associated with plasma LA accounting for 11.8% of variance observed, which  
35 was followed by FADS1/2 haplotypes (3.9 %), quantity and quality of carbohydrate intakes  
36 (3.8 %), dietary PUFA intakes (3.7 %), systolic blood pressure (3.6 %) and age (3.2 %).

37 **Conclusion:** Plasma LA was inversely associated with adiposity, followed by haplotypes of  
38 FADS1/2 genes, carbohydrate intakes and dietary PUFA intakes. The association observed  
39 between plasma LA and adiposity may be linked to decreased *de novo* lipogenesis, insulin  
40 resistance and inflammation.

41

## 42 **Introduction**

43 The essential fatty acid, dietary linoleic acid (LA) has traditionally been considered as protective  
44 against cardiovascular disease (CVD) [1] and coronary heart disease (CHD) [2, 3]. Randomized  
45 controlled trials suggest that diets high in PUFA may result in lower fat accumulation relative to  
46 other fatty acids [4, 5]. A number of observational studies have also attempted to relate circulating  
47 or plasma LA to health outcomes. In healthy adults, two recent studies which completed pattern  
48 analysis of a suite of plasma and erythrocyte fatty acids identified patterns rich in LA, both of  
49 which were associated with healthier phenotypes [6, 7]. The Epic Interact Study reported a strong  
50 inverse association between plasma linoleic acid and type 2 diabetes incidence [8]. Other studies  
51 in older populations or those at risk of CVD have suggested that circulating LA is inversely related  
52 to adiposity [9, 10], central adiposity [5, 11] and risk of the metabolic syndrome and type 2 diabetes,  
53 but not CHD [12, 13]. In a middle-aged community cohort, half of whom were obese, erythrocyte  
54 LA was inversely related with insulin resistance measured by HOMA score and IL-6 [11].  
55 However any relationship between plasma LA and health may be influenced by body weight status,  
56 with initial reports suggesting that circulating LA is positively related to circulating concentrations  
57 of soluble intercellular adhesion molecule-1 in (ICAM, a marker of endothelial activation and  
58 predictor of future CVD events) obese participants, the opposite being observed for normal weight  
59 adults [6]. Beyond this, little is known in terms of to what extent circulating LA level is associated  
60 with adiposity and how circulating LA is related to other markers of inflammation and metabolic  
61 health, particularly in younger, healthy cohorts.

62 The enzymes  $\Delta$ -5 and  $\Delta$ -6 desaturase, encoded by fatty acid desaturase (FADS) 1 and 2 genes, are  
63 the rate limiting enzymes regulating synthesis of long-chain n-6 polyunsaturated fatty acids (PUFA)  
64 such as the conversion of LA to  $\gamma$ -linoleic acid (GLA, 18:3n-6), dihomo- $\gamma$ -linoleic acid (DGLA,

65 20:3n-6) and arachidonic acid (ARA, 20:4n-6). Using genotype and haplotype analysis, circulating  
66 LA levels were associated with FADS 1 and 2 genes [14-16], with carriers of minor alleles  
67 demonstrating increased circulating LA [17]. Generally, the interaction between adiposity, dietary  
68 intakes, circulating LA and haplotypes of FADS1/2 genes and how any interaction may affect  
69 markers of inflammation and risk of metabolic syndrome (MetS) remain largely unknown. Against  
70 this backdrop, the objective of present study was to explore, in a cohort of healthy Irish adults, the  
71 association between plasma LA and adiposity, demographics, dietary intakes and haplotypes of  
72 FADS1/2 genes.

73

#### 74 **Participants and methods**

##### 75 Design and population

76 This paper uses data from the cross-sectional national food survey of Irish adults: the National  
77 Adult Nutrition Survey (NANS) [18]. The NANS (May 2008-April 2010) examined the habitual  
78 food and beverage intake of a representative sample (n=1500, 740 males and 760 females) of Irish  
79 adults aged 18-90 years who were free-living and who were not pregnant or breast feeding. Ethical  
80 approval was obtained from University College Cork Clinical Research Ethics Committee of the  
81 Cork Teaching Hospitals and the Human Ethics Research Committee of University College Dublin  
82 (ECM 3 (p) 4 September 2008). All procedures in this study were in accordance with the guidelines  
83 laid down in the Declaration of Helsinki, with written informed consent obtained from all  
84 participants.

85 As no national identification system exists in the Republic of Ireland, hence participants were  
86 randomly selected from the Data Ireland (National Postal Service) database of free-living adults

87 in Ireland in 20 geographical clusters nationwide, with clusters selected to provide proportional  
88 representation across the urban-rural continuum and a minimum of 100 participants in the least  
89 populated age and sex subgroups. A sample of 1500 free-living adults, representing the Irish  
90 population of over 4 million people participated in the dietary survey with analyses of the profile  
91 of NANS showing it to be representative of adults in Ireland with respect to age, gender, social  
92 class and urban/rural location compared to the most recent Irish census [18-20]. The overall  
93 response rate was 60%. In the present paper, only participants for whom fatty acid profiles were  
94 available and valid reporters are included, leaving a final sample size of 820 (which remained  
95 representative of the entire cohort). Demographics of participants who provided fasting blood  
96 samples were reported elsewhere [20].

97

#### 98 Dietary assessment

99 Food and beverage consumption was assessed using a 4-day semi-weighed food record. Initial  
100 nutrient analysis was carried out using the Weighed Intake Software Program (Tinuviel Software,  
101 Anglesey UK), which uses data from McCance & Widdowson's The Composition of Foods, fifth  
102 and sixth editions plus all nine supplemental volumes as described elsewhere [18] to generate  
103 nutrient intake data. Adjustments were made to the food composition database to take account of  
104 recipes, nutritional supplements, commonly consumed generic Irish food and new foods on the  
105 market. All entries were double-checked by the researcher and a random sample was checked by  
106 another researcher in-house to minimize the chance of error.

107

108

109 Plasma fatty acids analysis

110 The procedures used to determine plasma fatty acids profiles in the fasted samples are described  
111 elsewhere [6]. In brief, plasma lipids were extracted using the method of Bligh & Dyer [21].  
112 Butylated hydroxytoluene (BHT) and heptadecanoic acid (2 mg/mL in methanol) were used as  
113 antioxidant and internal standard, respectively. Extracted plasma lipids were transesterified using  
114 boron tri-fluoride-methanol solution (14%), and the resulting fatty acid methyl esters (FAME)  
115 were extracted twice using hexane and water [22]. FAMES were separated and identified by gas  
116 chromatography coupled with mass spectrometry (GC-MS) HP 5890 Series II, HP 5971A (Hewlett  
117 Packard) using an Omegawax 250 fused silica capillary column (30m×0.25mm×0.25µm film  
118 thickness, (Supelco). 38 known peaks were identified (including 17:0 as internal standard) using  
119 GC-MS. Laboratory CV of LA in fasting blood samples was 14.4%. The demographics of  
120 participants who provided a blood sample (76% of total sample), and those in the entire sample  
121 have been reported elsewhere, with no significant differences identified [19].

122

123 Assessment of biochemical profiles and metabolic risk factors

124 Collection of anthropometric measurements (including weighed body weight, BMI, waist  
125 circumference, body fat and blood pressure) and biochemistry parameters assessment has been  
126 described elsewhere [23, 24]. In brief, anthropometric measurements were carried out in duplicate  
127 by trained researchers using standardized methods. % body fat and weight were collected using  
128 Tanita ® Model BC-420 MA body composition analyzers (Tanita Ltd. GB) with a % CV of 0.73%  
129 and 0.02% respectively. Height was measured using a Leicester portable height measure (Seca,  
130 Birmingham, UK). All scales were calibrated and placed on a hard surface prior to use with  
131 participants asked to remove any heavy clothing, accessories or shoes as appropriate. Hip and waist

132 circumference were measured using non-stretch measuring tape to the nearest 0.1 cm with % CV  
133 of 0.17% and 0.13% respectively. Weight class was created as normal weight (BMI < 25 kg/m<sup>2</sup>),  
134 overweight (BMI 25-30 kg/m<sup>2</sup>) and obese (BMI > 30 kg/m<sup>2</sup>). Blood samples were collected by  
135 venepuncture by trained phlebotomists and stored appropriately until they reached the lab within  
136 5 hours of collection for processing and further storage at -80°C until further analysis. Biochemical  
137 values were assessed using a clinical bioanalyzer (Randox Daytona, Randox Laboratories) and  
138 cytokines and hormones using a biochip array system (Evidence Investigator, Randox Laboratories)  
139 except adiponectin which was measured by ELISA (ALPCO Diagnostics Kit). All samples were  
140 run in duplicate with standard quality control procedures followed in both analyses to ensure data  
141 integrity. Homeostatic model assessment-IR (HOMA-IR) score was calculated using the formula  
142 (fasting insulin μU/mL x fasting glucose mmol/L)/22.5 [25]. Quantitative sensitivity check index  
143 (QUICKI) score was calculated as follows: 1/(log(fasting insulin μU/mL) + log(fasting glucose  
144 mg/dL) [26]. Diagnosis of MetS was defined according to the National Cholesterol Education  
145 Programme's Adult Treatment Panel III criteria for MetS 2001 [27] including fasting blood  
146 glucose concentrations of 5.5-7.0 mmol/L, serum triacylglycerol (TAG) concentrations of ≥1.7  
147 mmol/L, HDL < 1.0 mmol/L (men), < 1.3 mmol/L (women), blood pressure ≥ 130/85 mmHg and  
148 waist circumference > 102 cm (men) & > 88 cm (women). Physical activity was estimated using a  
149 tri axial accelerometer (Actigraph GT1M, ActiGraph, LLC) for the four days of the survey period  
150 [28].

151

## 152 Genotyping

153 15 single nucleotide polymorphisms (SNPs) were selected based on previous studies which have  
154 been shown to be associated with circulating LA [14, 16, 29] and human health [30]. SNP



155 genotyping was then conducted by the LGC group ([www.lgcgroup.com](http://www.lgcgroup.com)) using their proprietary  
156 KASPar polymerase chain reaction technique [31].

157

158 Statistical analysis

159 Plasma n-6 PUFA (18:2n-6, 18:3n-6, 20:2n-6, 20:3n-6, 20:4n-6, 22:4n-6) profiles were compared  
160 across normal weight, over-weight and obese participants. Quartiles of plasma LA, GLA and  
161 DGLA were obtained using SPSS version 20.0 (IBM Inc. Chicago, USA). General linear contrast  
162 model adjusted for age, body fat, blood pressure, waist circumference, waist hip ratio, smoking  
163 status, working energy expenditure, and BMI was applied across plasma LA quartiles for  
164 comparison of percentage of plasma fatty acid, nutrient density (calculated as % total energy (TE)  
165 or  $\mu\text{g}$  or  $\text{mg}/10\text{MJ}$ ), and circulating concentrations of biochemical biomarkers. Differences in  
166 gender, social class, smoking habits, contraceptive use and metabolic risk factors across quartiles  
167 were compared using Chi-square test.  $\Delta$ -5-desaturase (D5D) and  $\Delta$ -6-desaturase (D6D) were  
168 estimated using fatty acid product/precursor ratios as following:  $\text{D6D}=\text{C18:3n-6}/\text{C18:2n-6}$ , and  
169  $\text{D5D}=\text{C20:4n-6}/\text{C20:3n-6}$  [32]. Contraceptive use in females was not controlled for in subsequent  
170 analysis due to the low proportion (3.5 %) of oral contraceptive users in NANS with no difference  
171 in proportions across LA quartiles ( $P=0.094$ ). One-way ANOVA was also applied across fatty acid  
172 clusters for demographic variables including age, BMI, waist circumference, body fat physical  
173 activity, waist hip ratio, and blood pressure. Skewed data were log transformed to obtain  
174 normalized distribution. Where appropriate in this exploratory analysis, Bonferroni correction was  
175 applied by adjusting the p-values by the number of traits in each table. p-values that exceed 1.0  
176 after correction for multiple testing have been marked down to 1.000.

177

178 Linkage disequilibrium (LD) was assessed using Haploview 4.2 [33], and haplotype blocks were  
179 obtained using the Gabriel's definition [34]. Haplotype frequency was estimated using Haplo.Stat  
180 1.7.6 package in R 3.2 (R Foundation for Statistical Computing, Vienna, Austria; ISBN 3-900051-  
181 07-0; Internet: <http://www.R-project.org>). Multiple linear regression with stepwise backward  
182 selection was used to explore possible determinants of plasma LA status. Demographic and dietary  
183 intake variables were included in the initial model and haplotypes were included in the final model  
184 regardless of the significance. Partial  $R^2$  was used to calculate the explained variation of each  
185 determinants in the final model. The association between plasma LA and haplotypes were  
186 estimated using HapStat 3.0 [35].

187

## 188 **Results**

189 A complete fasting plasma fatty acid dataset of 820 participants (mean age  $41.9 \pm 16.7$  years, with  
190 412 males and 408 females) were included in the analysis. Initial examination showed that plasma  
191 LA and estimated  $\Delta$ -5-desaturase activity decreased while GLA, DGLA, and  $\Delta$ -6-desaturase  
192 activity increased across the body weight categories in Irish adults ( $P \leq 0.003$ ) (Table 1).  
193 Demographics and markers of metabolic health were compared across the quartiles of plasma LA  
194 (Table 2 and Table 3), GLA and DGLA (Table S1) to explore any association between plasma LA,  
195 GLA, DGLA and health parameters. As quartiles of plasma LA showed greater separation in terms  
196 of demographics and markers of metabolic health (as described below) therefore subsequent  
197 analysis focused on plasma LA only.

198

199 Demographics and dietary intakes

200 As shown in Table 2, participants in the highest quartile of plasma LA displayed a healthier  
201 phenotype. Compared with participants in quartile 1 (Q1), participants in quartile 4 (Q4) were  
202 younger (50.7 years vs. 36.5 years,  $P < 1.9 \times 10^{-19}$ ), with lower adiposity (including BMI, body fat,  
203 waist circumference and waist hip ratio,  $P < 3.1 \times 10^{-6}$ ), blood pressure (131.9/81.4 mmHg vs.  
204 118.2/75.3 mmHg,  $P < 2.2 \times 10^{-7}$ ), metabolic risk and a lower proportion were overweight (45.3 %  
205 vs. 32.1 %) or obese (34.7 % vs. 14.0 %) ( $P \leq 2.7 \times 10^{-7}$ ). Higher work activity energy expenditure  
206 was observed for Q4 (47.8 MET hrs/wk vs. 66.1 MET hrs/wk,  $P < 0.026$ ) and a greater proportion  
207 of participants that never smoked (41.9 % vs. 62.7 %,  $P < 0.001$ ). Dietary intakes across the quartiles  
208 of plasma LA are shown in Table S2. Across plasma LA quartiles, intake of starch, PUFA, n-6  
209 PUFA and dietary fibre significantly increased ( $P < 0.003$ ).

210

#### 211 Markers of metabolic health and plasma fatty acids

212 The clinical chemistry biomarkers and plasma fatty acid levels across quartiles are shown in Table  
213 3. After adjustment for confounders, insulin, C-peptide, TAG, HOMA-IR, PAI-1, TNF- $\alpha$ , hs-CRP  
214 decreased, while QUICKI and adiponectin increased across plasma LA quartiles ( $P \leq 0.005$ ).  
215 Regarding plasma fatty acids related to the n-6 PUFA pathway, 20:2n-6, 18:3n-6, 20:3n-6 and  
216 D6D decreased across plasma LA quartiles ( $P \leq 0.019$ ), whereas total n-6 PUFA and n-6/n-3 ratio  
217 increased ( $P \leq 1.4 \times 10^{-138}$ ). Plasma total SFA and total MUFA decreased ( $P \leq 1.1 \times 10^{-21}$ ), as did  
218 plasma fatty acids related to *de novo* lipogenesis (DNL, including 14:0, 14:1n-5, 16:0, 16:1n-7 and  
219 18:1n-9) were also significantly decreased ( $P \leq 4.7 \times 10^{-6}$ ) across plasma LA quartiles.

220

221

222 Effects of haplotypes on plasma n-6 PUFA profiles

223 The degree of LD between the analyzed genetic polymorphisms are shown in Figure S1.  
224 Haplotypes were statistically reconstructed for three windows (Table 4). The first window  
225 contained all 15 SNPs from rs174537 to rs482548. Seven haplotypes were identified in this  
226 window with a frequency over 1 %, of which haplo 7 was the most common (53.9 %) haplotype  
227 with no minor allele. Compared with haplo 7, haplo 1, 3, 4, 5 were positively associated with  
228 plasma LA ( $P \leq 0.02$ ), with all of the haplotypes except haplo 2 negatively associated with D5D  
229 and D6D ( $P \leq 6.44 \times 10^{-3}$ ) (Table 5). The second window contained 8 SNPs from rs174537 to  
230 rs99780, and four haplotypes were reconstructed with frequency above 1 %. Compared with the  
231 most common haplotype (haplo 4 with frequency of 65.6 %) in the second window, all the other  
232 haplotypes were positively associated with plasma LA while negatively associated with D5D and  
233 D6D ( $P \leq 1.67 \times 10^{-3}$ ). The third window contained 3 SNPs from rs2524299 to rs174583, and three  
234 haplotypes were reconstructed with a frequency above 1 %. Compared with the most common  
235 haplotype (haplo 3 with frequency of 65.5 %) in the third window, all of the other haplotypes were  
236 positively associated with plasma LA while negatively associated with D5D and D6D ( $P \leq$   
237  $1.64 \times 10^{-3}$ ). Haplotypes in the second and third windows explained the variance of plasma LA by  
238 3.5 %, respectively (Table S2 and Table S4). We also explored the interaction of dietary PUFA  
239 intakes and haplotypes but no significant interaction was found after adjustment for confounders  
240 ( $P \geq 0.057$ ).

241

242 Possible determinants of plasma LA

243 Multiple linear regression of possible determinants of plasma LA are shown in Table 6. Of the  
244 variables tested, adiposity measures (waist circumference and body fat) were most strongly

245 inversely associated with plasma LA accounting for 11.8 % of the total variance, followed by 15-  
246 SNP haplotypes (3.9 %), quantity and quality of carbohydrate intakes (3.8 %), dietary PUFA  
247 intakes (3.7 %), systolic blood pressure (3.6 %) and age (3.2 %).

248

## 249 **Discussion**

250 In this study, higher plasma LA was related to a healthier phenotype and metabolic profile in Irish  
251 adults. We observed an inverse relationship between plasma LA and the adiposity measures  
252 studied, with waist circumference and body fat most strongly associated with plasma LA, followed  
253 by dietary PUFA and haplotypes of FADS genes. The beneficial effects of plasma LA may be  
254 associated with decreased biomarkers of DNL, insulin resistance and inflammation, all of which  
255 have previously been linked with the pathogenesis of obesity [36, 37].

256 A growing body of evidence suggests that LA may influence health outcomes. An inverse  
257 relationship between serum LA and all-cause mortality (but not CVD risk) has been described in  
258 a cohort of healthy older adults [38]. Further observational studies report lower plasma  
259 phospholipid LA and higher GLA and DGLA levels in overweight/obese adolescents [9, 39] and  
260 higher proportions of serum and cholesterol ester LA being associated with lower markers of  
261 abdominal obesity and BMI and reduced T2D incidence [8, 40-42]. Altered desaturase activity is  
262 documented in overweight and obese participants and those with T2D, with reports of elevated  
263 D6D and decreased D5D [8, 9, 39, 41]. Growing evidence suggests that such benefits may arise  
264 from effects of LA on body fat accumulation. Animal trials and human RCTs indicate that  
265 compared to SFA, high LA intakes result in improvements in body composition outcomes [4, 5,  
266 43]. There is some evidence of similar benefits in populations with regular dietary LA intakes  
267 whereby serum or erythrocyte LA was inversely related to body fat, specifically trunk fat in a

268 healthy elderly cohort [5] and in a middle aged relatively healthy cohort [11]. Of interest, such  
269 benefits were not observed for erythrocyte n-3 PUFA or for the MUFA, OA [11]. In the present  
270 study, we confirm these fatty acid profiles of elevated LA and altered patterns of D5D and D6D  
271 activity according to body weight status and without significant alterations in n-3 PUFA status, in  
272 this nationally representative cohort of healthy adults aged 18-90 y. Our study adds to evidence  
273 which is suggestive of either a protective effect of circulating LA on obesity and/or, of altered fatty  
274 acid metabolism in the presence of excess body weight. However, desaturase activities based on  
275 ratio of circulating fatty acids are indirect measure of enzyme activities which may also be affected  
276 by lifestyle. Further, our results are presented as percentage, rather than absolute, values which  
277 may imply co-dependence on all other fatty acids. Further research is required to explore the  
278 specific health implications of plasma LA as well as any interactions between dietary PUFA  
279 intakes and circulating fatty acid profiles.

280 Previous studies have demonstrated a similar inverse association between circulating LA and  
281 biomarkers such as insulin, glucose, TAG, hs-CRP, with higher adiponectin concentrations and  
282 with reduced risk of insulin resistance, metabolic syndrome, cardio-metabolic risk score and T2D  
283 [5, 8-10, 44-46]. In addition, our results show decreased inflammation markers such as PAI-1 and  
284 TNF- $\alpha$  across plasma LA quartiles, elevated levels of which have been linked to obesity [47].  
285 Nevertheless, this decrease in markers of metabolic health in the present study remained significant  
286 even when controlling for BMI and waist circumference. A negative association between plasma  
287 LA and inflammation markers e.g. IL-6 and CRP has been reported elsewhere [11, 48]. Such  
288 associations may be explained by a recent study, showing that LA may suppress the expression of  
289 toll-like receptor 4 (TLR4), phosphorylation levels of c-Jun amino-terminal kinases (JNK) and  
290 agouti-related protein (AgRP) [49]. AgRP promotes food intake[50], and an absence of JNK1 may

291 lead to decreased adiposity and improved insulin sensitivity [51]. Such potential mechanisms may  
292 help explain the decreased adiposity and markers of inflammation observed in participants in the  
293 highest quartile of plasma LA (Q4). Moreover, bioactive metabolites of LA such as nitro-linoleic  
294 acid may be anti-inflammatory by reducing the activation and gene transcription of nuclear factor-  
295  $\kappa$ B while stimulating activation of peroxisome proliferator-activated receptor  $\gamma$  [52]. Further  
296 research is required in this area.

297 It is possible that the decreased plasma LA observed in overweight/obese participants in our study  
298 may be linked to DNL. DNL is a, typically inhibited, pathway of converting excess carbohydrate  
299 to esterified fatty acids which are subsequently stored as TAG in the liver and adipose tissue [53].  
300 Increased DNL has been observed in the obese [36, 54], in non-alcoholic fatty liver disease [55]  
301 and under conditions of insulin resistance, including inflammation [53]. Further, DNL stimulated  
302 hepatic TAG are reported to be enriched with palmitic acid while depleted in LA [56]. In our study,  
303 in addition to decreased levels of plasma LA observed in obese participants, all adults in the lowest  
304 quartile of plasma LA had a fatty acid profile suggestive of DNL (i.e. reduced LA, increased 14:0,  
305 16:0, 16:1n-7) and with patterns of biochemical markers suggestive of insulin resistance and low-  
306 grade inflammation. Dietary intakes may also influence DNL, whereby high carbohydrate, low  
307 fat diets may promote DNL, particularly when dietary sugar is substituted for starch [57]. In  
308 contrast high dietary fibre intakes can yield less short-chain fatty acids, which may suppress DNL  
309 and improve insulin sensitivity [58, 59]. In our study, participants had a relatively high fat diet  
310 (mean daily intake of 33.8 %TE) and moderate carbohydrate intakes (mean daily intake of  
311 42.6 %TE). Nevertheless, participants with the lowest circulating plasma concentrations (quartile  
312 1) had lower intakes of starch and dietary fibre, while regression analysis identified quantity and  
313 quality of carbohydrate intake as the third largest determinant of plasma LA (3.8%). Collectively,

314 in this cohort of generally healthy adults there exists a group with lipid and metabolic profiles  
315 suggestive of possible future risk and who can be easily identified by nature of circulating plasma  
316 LA concentrations. This is supportive of the hypothesis which aims to clarify whether endogenous  
317 metabolites or lipids constitute simple and effective biomarkers of metabolic diseases such as  
318 NAFLD, type 2 diabetes and metabolic syndrome [6, 55].

319 Polymorphism and haplotype of FADS1/2 genes have been reported to be linked to long-chain  
320 PUFA concentrations/metabolism such as for LA, ARA, D5D and D6D [14, 30, 60, 61]. To our  
321 knowledge, only one study reported variation of haplotypes of FADS genes in relation to serum  
322 phospholipid, which was higher compared with the present study (9.2 % vs. 3.9 %) [14]. This may  
323 due to different SNPs, lipid fractions and study population included. Potential dietary PUFA  
324 intakes  $\times$  haplotypes interactions were also explored in the present study, however, the P values  
325 were above the significant threshold ( $P \geq 0.057$ ) after adjustment for confounders [62]. The impact  
326 of dietary intakes  $\times$  haplotypes/genotype interactions on circulating n-6 PUFA profiles have been  
327 poorly explored and merits future study [63]. Haplotype analysis of FADS genes also revealed  
328 significant association between haplotypes and concentration of hs-CRP and TAG [15, 16].  
329 Similarly, results of genome-wide association studies reported that carriers of the minor allele of  
330 rs174537 and rs174547 showed lower total cholesterol HDL-C and LDL-C [64, 65]. However, no  
331 significant association was identified in our study between haplotypes and inflammation markers,  
332 suggesting that haplotype may not be involved in the association between plasma LA and  
333 inflammation.

334 The major strength of the present study is that we included a comprehensive group of inflammation  
335 markers, genotype information as well as the combination of dietary and plasma fatty acid data  
336 which clearly showed the beneficial effects of a higher plasma LA. Limitations of the study include



337 the observational study design that is unable to reveal the causality of the relations. In this national  
338 food consumption survey (NANS), body composition measurements were collected by trained  
339 field workers with body fat assessed using a Tanita® body fat analyser rather than gold standard  
340 methods such as magnetic resonance imaging, computed tomography or DEXA scans. This was  
341 due to the practical simplicity of using Tanita® as part of a national food consumption survey. It  
342 has been reported that Tanita® is not associated with a clinically significant decrement in  
343 performance relative to traditional impedance devices [66] although it is recognized that this  
344 technique may slightly overestimate fat mass relative to DEXA [67] and MRI [68]. A further  
345 limitation of this study was the use of total plasma fatty acids rather than lipid fractions given that  
346 different associations have been reported between phospholipid and cholesterol esters [69]. Finally,  
347 we acknowledge that alterations in proportions of circulating fatty acids with age are well  
348 described, with reductions in erythrocyte concentrations of linoleic acid reported to be in the order  
349 of 2 percentage points per decade [70]. In our study, those in the lowest quartile of plasma LA  
350 were older than those in the highest quartile. Hence, it is possible that age-related influences on  
351 body fat accumulation and on metabolic health may have influenced outcome. However, we  
352 controlled for age in all relevant analysis to attempt mitigate any such risk.

353 In conclusion, our results showed that plasma LA was inversely associated with adiposity,  
354 followed by haplotype of FADS genes and dietary intakes. A higher plasma LA profile was related  
355 to a healthier phenotype and markers of metabolic health even when controlling for BMI and waist  
356 circumference. Future work is needed to better understand the mechanism of obesity modified  
357 plasma LA profile on human health.

358

359

360 **Acknowledgements**

361 This research was funded by the Irish Department of Agriculture, Food and the Marine and the  
362 Health Research Board under their joint Food for Health Research Initiative (2007-12) (grant  
363 number FHRIUCC2). Mr. Li is in receipt of a PhD student from the China Scholarship Council  
364 2012-2016.

365

366 **Author's contribution**

367 *1) designed research: A.P. Nugent, L. Brennan, M.J. Gibney, B.A. McNulty, A. Flynn, J.*  
368 *Walton, and; 2) conducted research: K. Li, F.J. Bloomfield, D. J. Duff, A.P. Nugent and B.A.*  
369 *McNulty ; 3) analysed data or performed statistical analysis: K. Li, A.P. Nugent, L. Brennan; 4)*  
370 *wrote paper: K. Li, A.P. Nugent; 5) had primary responsibility for final content: K. Li, L.*  
371 *Brennan and A.P. Nugent*

372

373 **Disclaimers: None of the authors declare a conflict of interest**

374

375

376

377 **References**

- 378 [1] Harris, W. S., Mozaffarian, D., Rimm, E., Kris-Etherton, P., Rudel, L. L., Appel, L. J., Engler,  
379 M. M., Engler, M. B., Sacks, F., Omega-6 fatty acids and risk for cardiovascular disease: A science  
380 advisory from the American Heart Association nutrition subcommittee of the council on nutrition,  
381 physical activity, and metabolism; council on cardiovascular nursing; and council on epidemiology  
382 and prevention. *Circulation* 2009, *119*, 902-907.
- 383 [2] Farvid, M. S., Ding, M., Pan, A., Sun, Q., Chiuve, S. E., Steffen, L. M., Willett, W. C., Hu, F.  
384 B., Dietary linoleic acid and risk of coronary heart disease: a systematic review and meta-analysis  
385 of prospective cohort studies. *Circulation* 2014, *130*, 1568-1578.
- 386 [3] Oh, K., Hu, F. B., Manson, J. E., Stampfer, M. J., Willett, W. C., Dietary fat intake and risk of  
387 coronary heart disease in women: 20 years of follow-up of the nurses' health study. *Am J Epidemiol*  
388 2005, *161*, 672-679.
- 389 [4] Bjermo, H., Iggman, D., Kullberg, J., Dahlman, I., Johansson, L., Persson, L., Berglund, J.,  
390 Pulkki, K., Basu, S., Uusitupa, M., Rudling, M., Arner, P., Cederholm, T., Ahlstrom, H., Riserus,  
391 U., Effects of n-6 PUFAs compared with SFAs on liver fat, lipoproteins, and inflammation in  
392 abdominal obesity: a randomized controlled trial. *Am J Clin Nutr* 2012, *95*, 1003-1012.
- 393 [5] Rosqvist, F., Bjermo, H., Kullberg, J., Johansson, L., Michaëlsson, K., Ahlström, H., Lind, L.,  
394 Risérus, U., Fatty acid composition in serum cholesterol esters and phospholipids is linked to  
395 visceral and subcutaneous adipose tissue content in elderly individuals: a cross-sectional study.  
396 *Lipids Health Dis* 2017, *16*, 68.
- 397 [6] Li, K., Brennan, L., McNulty, B. A., Bloomfield, J. F., Duff, D. J., Devlin, N. F. C., Gibney,  
398 M. J., Flynn, A., Walton, J., Nugent, A. P., Plasma fatty acid patterns reflect dietary habits and  
399 metabolic health: a cross-sectional study. *Mol Nutr Food Res* 2016, *60*, 2043-52.

400 [7] Bigornia, S. J., Lichtenstein, A. H., Harris, W. S., Tucker, K. L., Associations of erythrocyte  
401 fatty acid patterns with insulin resistance. *Am J Clin Nutr* 2016, *103*, 902-909.

402 [8] Forouhi, N. G., Imamura, F., Sharp, S. J., Koulman, A., Schulze, M. B., Zheng, J., Ye, Z.,  
403 Sluijs, I., Guevara, M., Huerta, J. M., Kröger, J., Wang, L. Y., Summerhill, K., Griffin, J. L.,  
404 Feskens, E. J. M., Affret, A., Amiano, P., Boeing, H., Dow, C., Fagherazzi, G., Franks, P. W.,  
405 Gonzalez, C., Kaaks, R., Key, T. J., Khaw, K. T., Kühn, T., Mortensen, L. M., Nilsson, P. M.,  
406 Overvad, K., Pala, V., Palli, D., Panico, S., Quirós, J. R., Rodriguez-Barranco, M., Rolandsson,  
407 O., Sacerdote, C., Scalbert, A., Slimani, N., Spijkerman, A. M. W., Tjønneland, A., Tormo, M.-J.,  
408 Tumino, R., van der A, D. L., van der Schouw, Y. T., Langenberg, C., Riboli, E., Wareham, N. J.,  
409 Association of Plasma Phospholipid n-3 and n-6 Polyunsaturated Fatty Acids with Type 2 Diabetes:  
410 The EPIC-InterAct Case-Cohort Study. *PLoS Med* 2016, *13*, e1002094.

411 [9] Steffen, L. M., Vessby, B., Jacobs, D. R., Jr., Steinberger, J., Moran, A., Hong, C. P., Sinaiko,  
412 A. R., Serum phospholipid and cholesteryl ester fatty acids and estimated desaturase activities are  
413 related to overweight and cardiovascular risk factors in adolescents. *Int J Obes* 2008, *32*, 1297-  
414 1304.

415 [10] Steffen, B. T., Steffen, L. M., Tracy, R., Siscovick, D., Hanson, N. Q., Nettleton, J., Tsai, M.  
416 Y., Obesity modifies the association between plasma phospholipid polyunsaturated fatty acids and  
417 markers of inflammation: the Multi-Ethnic Study of Atherosclerosis. *Int J Obes* 2012, *36*, 797-804.

418 [11] Belury, M. A., Cole, R. M., Bailey, B. E., Ke, J. Y., Andridge, R. R., Kiecolt-Glaser, J. K.,  
419 Erythrocyte linoleic acid, but not oleic acid, is associated with improvements in body composition  
420 in men and women. *Mol Nutr Food Res* 2016, *60*, 1206-1212.

421 [12] Matthan, N. R., Ooi, E. M., Van Horn, L., Neuhouser, M. L., Woodman, R., Lichtenstein, A.  
422 H., Plasma phospholipid fatty acid biomarkers of dietary fat quality and endogenous metabolism

423 predict coronary heart disease risk: a nested case-control study within the Women's Health  
424 Initiative observational study. *J Am Heart Assoc* 2014, 3.

425 [13] Chowdhury, R., Warnakula, S., Kunutsor, S., Crowe, F., Ward, H. A., Johnson, L., Franco,  
426 O. H., Butterworth, A. S., Forouhi, N. G., Thompson, S. G., Khaw, K. T., Mozaffarian, D., Danesh,  
427 J., Di Angelantonio, E., Association of dietary, circulating, and supplement fatty acids with  
428 coronary risk: A systematic review and meta-analysis. *Ann Int Med* 2014, 160, 398-406.

429 [14] Schaeffer, L., Gohlke, H., Müller, M., Heid, I. M., Palmer, L. J., Kompauer, I., Demmelmair,  
430 H., Illig, T., Koletzko, B., Heinrich, J., Common genetic variants of the FADS1 FADS2 gene  
431 cluster and their reconstructed haplotypes are associated with the fatty acid composition in  
432 phospholipids. *Hum Mol Genet* 2006, 15, 1745-1756.

433 [15] Martinelli, N., Girelli, D., Malerba, G., Guarini, P., Illig, T., Trabetti, E., Sandri, M., Friso,  
434 S., Pizzolo, F., Schaeffer, L., Heinrich, J., Pignatti, P. F., Corrocher, R., Olivieri, O., FADS  
435 genotypes and desaturase activity estimated by the ratio of arachidonic acid to linoleic acid are  
436 associated with inflammation and coronary artery disease. *Am J Clin Nutr* 2008, 88, 941-949.

437 [16] Bokor, S., Dumont, J., Spinneker, A., Gonzalez-Gross, M., Nova, E., Widhalm, K.,  
438 Moschonis, G., Stehle, P., Amouyel, P., De Henauw, S., Molnàr, D., Moreno, L. A., Meirhaeghe,  
439 A., Dallongeville, J., Group, o. b. o. t. H. S., Single nucleotide polymorphisms in the FADS gene  
440 cluster are associated with delta-5 and delta-6 desaturase activities estimated by serum fatty acid  
441 ratios. *J Lipid Res* 2010, 51, 2325-2333.

442 [17] O'Neill, C. M., Minihane, A.-M., The impact of fatty acid desaturase genotype on fatty acid  
443 status and cardiovascular health in adults. *Proc Nutr Soc* 2016, 1-12.

444 [18] Li, K., B. A. M., Ann M. Tiernery, Niamh F.C. Devlin, Triona Joyce, Joao C. Leite, Albert  
445 Flynn, Janette Walton, Lorraine Brennan, Michael J. Gibney, Anne P. Nugent, Dietary fat intakes  
446 in Irish adults in 2011: how much has changed in 10 years? *Br J Nutr* 2016, *115*, 1798-809.

447 [19] Hopkins, S. M., Gibney, M. J., Nugent, A. P., McNulty, H., Molloy, A. M., Scott, J. M., Flynn,  
448 A., Strain, J., Ward, M., Walton, J., McNulty, B. A., Impact of voluntary fortification and  
449 supplement use on dietary intakes and biomarker status of folate and vitamin B-12 in Irish adults.  
450 *Am J Clin Nutr* 2015, *101*, 1163-72.

451 [20] Cashman, K. D., Muldowney, S., McNulty, B., Nugent, A., Fitzgerald, A. P., Kiely, M.,  
452 Walton, J., Gibney, M. J., Flynn, A., Vitamin D status of Irish adults: Findings from the National  
453 Adult Nutrition Survey. *Br J Nutr* 2013, *109*, 1248-1256.

454 [21] Bligh, E. G., Dyer, W. J., A rapid method of total lipid extraction and purification. *Canadian*  
455 *J Biochem and Physiol* 1959, *37*, 911-917.

456 [22] Eder, K., Gas chromatographic analysis of fatty acid methyl esters. *J Chromatography B:*  
457 *Biomed Applications* 1995, *671*, 113-131.

458 [23] Boylan, E. A., McNulty, B. A., Walton, J., Flynn, A., Nugent, A. P., Gibney, M. J., The  
459 prevalence and trends in overweight and obesity in Irish adults between 1990 and 2011. *Public*  
460 *Health Nutr* 2014, *17*, 2389-97.

461 [24] O'Donovan, C. B., Walsh, M. C., Nugent, A. P., McNulty, B., Walton, J., Flynn, A., Gibney,  
462 M. J., Gibney, E. R., Brennan, L., Use of metabotyping for the delivery of personalised nutrition.  
463 *Mol Nutr Food Res* 2015, *59*, 377-385.

464 [25] Matthews, D. R., Hosker, J. P., Rudenski, A. S., Naylor, B. A., Treacher, D. F., Turner, R. C.,  
465 Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma  
466 glucose and insulin concentrations in man. *Diabetologia* 1985, *28*, 412-419.

467 [26] Katz, A., Nambi, S. S., Mather, K., Baron, A. D., Follmann, D. A., Sullivan, G., Quon, M. J.,  
468 Quantitative insulin sensitivity check index: A simple, accurate method for assessing insulin  
469 sensitivity in humans. *J Clin Endocrinol Metab* 2000, 85, 2402-2410.

470 [27] Expert Panel on Detection, E., and Treatment of High Blood Cholesterol in Adults, Executive  
471 summary of the third report of the national cholesterol education program (ncep) expert panel on  
472 detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel iii).  
473 *JAMA* 2001, 285, 2486-2497.

474 [28] Irish Universities Nutrition Alliance. National Adult Nutrition Survey  
475 Summary Report. Cork: Irish Universities Nutrition Alliance. 2011. Internet:  
476 <http://www.iuna.net/?p=106> 2012. (accessed 16/08/2017).

477 [29] Martinelli, N., Girelli, D., Malerba, G., Guarini, P., Illig, T., Trabetti, E., Sandri, M., Friso,  
478 S., Pizzolo, F., Schaeffer, L., Heinrich, J., Pignatti, P. F., Corrocher, R., Olivieri, O., FADS  
479 genotypes and desaturase activity estimated by the ratio of arachidonic acid to linoleic acid are  
480 associated with inflammation and coronary artery disease. *Am J Clin Nutr* 2008, 88, 941-949.

481 [30] Merino, D. M., Ma, D. W., Mutch, D. M., Genetic variation in lipid desaturases and its impact  
482 on the development of human disease. *Lipids Health Dis* 2010, 9, 63.

483 [31] Ryan, M. F., O'Grada, C. M., Morris, C., Segurado, R., Walsh, M. C., Gibney, E. R., Brennan,  
484 L., Roche, H. M., Gibney, M. J., Within-person variation in the postprandial lipemic response of  
485 healthy adults. *Am J Clin Nutr* 2013, 97, 261-267.

486 [32] Nakamura, M. T., Nara, T. Y., Structure, function, and dietary regulation of delta6, delta5,  
487 and delta9 desaturases. *Ann Re. Nutr*, 2004, 24, 345-76

488 [33] Barrett, J. C., Fry, B., Maller, J., Daly, M. J., Haploview: analysis and visualization of LD  
489 and haplotype maps. *Bioinformatics* 2005, 21, 263-265.

490 [34] Gabriel, S. B., Schaffner, S. F., Nguyen, H., Moore, J. M., Roy, J., Blumenstiel, B., Higgins,  
491 J., DeFelice, M., Lochner, A., Faggart, M., Liu-Cordero, S. N., Rotimi, C., Adeyemo, A., Cooper,  
492 R., Ward, R., Lander, E. S., Daly, M. J., Altshuler, D., The structure of haplotype blocks in the  
493 human genome. *Science* 2002, 296, 2225-2229.

494 [35] Lin, D., Zeng, D., Millikan, R., Maximum likelihood estimation of haplotype effects and  
495 haplotype-environment interactions in association studies. *Genet Epidemiol* 2005, 29, 299.

496 [36] Kusunoki, J., Kanatani, A., Moller, D. E., Modulation of fatty acid metabolism as a potential  
497 approach to the treatment of obesity and the metabolic syndrome. *Endocrine* 2006, 29, 91-100.

498 [37] Winer, Daniel A., Luck, H., Tsai, S., Winer, S., The Intestinal Immune System in Obesity and  
499 Insulin Resistance. *Cell Metab* 2016, 23, 413-426.

500 [38] Marklund, M., Leander, K., Vikström, M., Laguzzi, F., Gigante, B., Sjögren, P., Cederholm,  
501 T., de Faire, U., Hellénus, M.-L., Risérus, U., Polyunsaturated Fat Intake Estimated by Circulating  
502 Biomarkers and Risk of Cardiovascular Disease and All-Cause Mortality in a Population-Based  
503 Cohort of 60-Year-Old Men and Women. *Circulation* 2015, 132, 586-94.

504 [39] Klein-Platat, C., Draï, J., Oujaa, M., Schlienger, J. L., Simon, C., Plasma fatty acid  
505 composition is associated with the metabolic syndrome and low-grade inflammation in overweight  
506 adolescents. *Am J Clin Nutr* 2005, 82, 1178-1184.

507 [40] Warensjö, E., Sundstrom, J., Lind, L., Vessby, B., Factor analysis of fatty acids in serum  
508 lipids as a measure of dietary fat quality in relation to the metabolic syndrome in men. *Am J Clin*  
509 *Nutr* 2006, 84, 442-448.

510 [41] Alsharari, Z. D., Risérus, U., Leander, K., Sjögren, P., Carlsson, A. C., Vikström, M., Laguzzi,  
511 F., Gigante, B., Cederholm, T., De Faire, U., Hellénus, M.-L., Marklund, M., Serum Fatty Acids,



512 Desaturase Activities and Abdominal Obesity – A Population-Based Study of 60-Year Old Men  
513 and Women. *PloS One* 2017, *12*, e0170684.

514 [42] Zheng, Z.J., Folsom, A. R., Ma, J., Arnett, D. K., McGovern, P. G., Eckfeldt, J. H., Plasma  
515 fatty acid composition and 6-year incidence of hypertension in middle-aged adults: the  
516 Atherosclerosis Risk in Communities (ARIC) Study. *Am J Epidemiol* 1999, *150*, 492-500.

517 [43] Norris, L. E., Collene, A. L., Asp, M. L., Hsu, J. C., Liu, L. F., Richardson, J. R., Li, D., Bell,  
518 D., Osei, K., Jackson, R. D., Comparison of dietary conjugated linoleic acid with safflower oil on  
519 body composition in obese postmenopausal women with type 2 diabetes mellitus. *Am J Clin Nutr*  
520 2009, *90*, 468-76.

521 [44] Mayneris-Perxachs, J., Guerendiain, M., Castellote, A. I., Estruch, R., Covas, M. I., Fitó, M.,  
522 Salas-Salvadó, J., Martínez-González, M. A., Aros, F., Lamuela-Raventós, R. M., López-Sabater,  
523 M. C., Plasma fatty acid composition, estimated desaturase activities, and their relation with the  
524 metabolic syndrome in a population at high risk of cardiovascular disease. *Clin Nutr* 2014, *33*, 90-  
525 97.

526 [45] Venalainen, T., Agren, J., Schwab, U., de Mello, V. D., Eloranta, A. M., Laaksonen, D. E.,  
527 Lindi, V., Lakka, T. A., Cross-sectional associations of plasma fatty acid composition and  
528 estimated desaturase and elongase activities with cardiometabolic risk in Finnish children-The  
529 PANIC study. *J Clin Lipidol* 2016, *10*, 82-91.

530 [46] Enzenbach, C., Kroger, J., Zietemann, V., Jansen, E. H., Fritsche, A., Doring, F., Boeing, H.,  
531 Schulze, M. B., Erythrocyte membrane phospholipid polyunsaturated fatty acids are related to  
532 plasma C-reactive protein and adiponectin in middle-aged German women and men. *Eur J Nutr*  
533 2011, *50*, 625-636.

534 [47] Jean-Philippe Bastard, M. M., Claire Lagathu, Min Ji Kim, Martine Caron, Hubert Vidal,  
535 Jacqueline Capeau, Bruno Feve, Recent advances in the relationship between obesity,  
536 inflammation, and insulin resistance. *Eur Cytokine Netw* 2006, 17, 4-12.

537 [48] Muka, T., Kiefte-de Jong, J. C., Hofman, A., Dehghan, A., Rivadeneira, F., Franco, O. H.,  
538 Polyunsaturated Fatty Acids and Serum C-Reactive ProteinThe Rotterdam Study. *Am J Epidemiol*  
539 2015, 181, 846-856.

540 [49] Wang, S., Xiang, N., Yang, L., Zhu, C., Zhu, X., Wang, L., Gao, P., Xi, Q., Zhang, Y., Shu,  
541 G., Jiang, Q., Linoleic acid and stearic acid elicit opposite effects on AgRP expression and  
542 secretion via TLR4-dependent signaling pathways in immortalized hypothalamic N38 cells.  
543 *Biochem Biophys Res Comm* 2016, 471, 566-571.

544 [50] Hirosumi, J., Tuncman, G., Chang, L., Gorgun, C. Z., Uysal, K. T., Maeda, K., Karin, M.,  
545 Hotamisligil, G. S., A central role for JNK in obesity and insulin resistance. *Nature* 2002, 420,  
546 333-336.

547 [51] Schwartz, M. W., Morton, G. J., Obesity: Keeping hunger at bay. *Nature* 2002, 418, 595-597.

548 [52] Delmastro-Greenwood, M., Freeman, B. A., Wendell, S. G., Redox-dependent anti-  
549 inflammatory signaling actions of unsaturated fatty acids. *Annu Rev Physiol* 2014, 76, 79-105.

550 [53] Ameer, F., Scandiuzzi, L., Hasnain, S., Kalbacher, H., Zaidi, N., De novo lipogenesis in health  
551 and disease. *Metab.: Clin Exp* 2014, 63, 895-902.

552 [54] Diraison, F., Dusserre, E., Vidal, H., Sothier, M., Beylot, M., Increased hepatic lipogenesis  
553 but decreased expression of lipogenic gene in adipose tissue in human obesity. *Am J Physiol.*  
554 *Endocrinol Metab* 2002, 282, E46-51.

555 [55] Sanders, F. W., Griffin, J. L., De novo lipogenesis in the liver in health and disease: more  
556 than just a shunting yard for glucose. *Biol Rev* 2016, 91, 452-468.

557 [56] Chong, M. F., Hodson, L., Bickerton, A. S., Roberts, R., Neville, M., Karpe, F., Frayn, K. N.,  
558 Fielding, B. A., Parallel activation of de novo lipogenesis and stearoyl-CoA desaturase activity  
559 after 3 d of high-carbohydrate feeding. *Am J Clin Nutr* 2008, *87*, 817-823.

560 [57] Hudgins, L. C., Seidman, C. E., Diakun, J., Hirsch, J., Human fatty acid synthesis is reduced  
561 after the substitution of dietary starch for sugar. *Am J Clin Nutr* 1998, *67*, 631-639.

562 [58] Weitkunat, K., Schumann, S., Nickel, D., Kappo, K. A., Petzke, K. J., Kipp, A. P., Blaut, M.,  
563 Klaus, S., Importance of propionate for the repression of hepatic lipogenesis and improvement of  
564 insulin sensitivity in high-fat diet-induced obesity. *Mol Nutr Food Res* 2016.

565 [59] Singh, V., Yeoh, B. S., Ntambi, J. M., Vijay-Kumar, M., in: Ntambi, M. J. (Ed.), *Hepatic De*  
566 *Novo Lipogenesis and Regulation of Metabolism*, Springer International Publishing, Cham 2016,  
567 pp. 189-209.

568 [60] Lattka, E., Illig, T., Heinrich, J., Koletzko, B., Do FADS genotypes enhance our knowledge  
569 about fatty acid related phenotypes? *Clin Nutr* 2010, *29*, 277-287.

570 [61] Ameer, A., Enroth, S., Johansson, A., Zaboli, G., Igl, W., Johansson, A. C., Rivas, M. A.,  
571 Daly, M. J., Schmitz, G., Hicks, A. A., Meitinger, T., Feuk, L., van Duijn, C., Oostra, B.,  
572 Pramstaller, P. P., Rudan, I., Wright, A. F., Wilson, J. F., Campbell, H., Gyllensten, U., Genetic  
573 adaptation of fatty-acid metabolism: a human-specific haplotype increasing the biosynthesis of  
574 long-chain omega-3 and omega-6 fatty acids. *Am J Hum Genet* 2012, *90*, 809-820.

575 [62] Keller, M. C., Gene × Environment Interaction Studies Have Not Properly Controlled for  
576 Potential Confounders: The Problem and the (Simple) Solution. *Biol Psy* 2014, *75*, 18-24.

577 [63] Guan, W., Steffen, B. T., Lemaitre, R. N., Wu, J. H. Y., Tanaka, T., Manichaikul, A., Foy,  
578 M., Rich, S. S., Wang, L., Nettleton, J. A., Tang, W., Gu, X., Bandinelli, S., King, I. B., McKnight,  
579 B., Psaty, B. M., Siscovick, D., Djousse, L., Ida Chen, Y.-D., Ferrucci, L., Fornage, M.,

580 Mozafarrian, D., Tsai, M. Y., Steffen, L. M., Genome-Wide Association Study of Plasma N6  
581 Polyunsaturated Fatty Acids Within the Cohorts for Heart and Aging Research in Genomic  
582 Epidemiology Consortium. *Circ: Cardiovas Genet* 2014, 7, 321-331.

583 [64] Tanaka, T., Shen, J., Abecasis, G. R., Kisialiou, A., Ordovas, J. M., Guralnik, J. M., Singleton,  
584 A., Bandinelli, S., Cherubini, A., Arnett, D., Tsai, M. Y., Ferrucci, L., Genome-wide association  
585 study of plasma polyunsaturated fatty acids in the InCHIANTI study. *PLoS genetics* 2009, 5.

586 [65] Kathiresan, S., Willer, C. J., Peloso, G. M., Demissie, S., Musunuru, K., Schadt, E. E., Kaplan,  
587 L., Bennett, D., Li, Y., Tanaka, T., Voight, B. F., Bonnycastle, L. L., Jackson, A. U., Crawford,  
588 G., Surti, A., Guiducci, C., Burt, N. P., Parish, S., Clarke, R., Zelenika, D., Kubalanza, K. A.,  
589 Morken, M. A., Scott, L. J., Stringham, H. M., Galan, P., Swift, A. J., Kuusisto, J., Bergman, R.  
590 N., Sundvall, J., Laakso, M., Ferrucci, L., Scheet, P., Sanna, S., Uda, M., Yang, Q., Lunetta, K. L.,  
591 Dupuis, J., de Bakker, P. I. W., O'Donnell, C. J., Chambers, J. C., Kooner, J. S., Hercberg, S.,  
592 Meneton, P., Lakatta, E. G., Scuteri, A., Schlessinger, D., Tuomilehto, J., Collins, F. S., Groop, L.,  
593 Altshuler, D., Collins, R., Lathrop, G. M., Melander, O., Salomaa, V., Peltonen, L., Orho-  
594 Melander, M., Ordovas, J. M., Boehnke, M., Abecasis, G. R., Mohlke, K. L., Cupples, L. A.,  
595 Common variants at 30 loci contribute to polygenic dyslipidemia. *Nat Genet* 2009, 41, 56-65.

596 [66] Jebb, S. A., Cole, T. J., Doman, D., Murgatroyd, P. R., Prentice, A. M., Evaluation of the  
597 novel Tanita body-fat analyser to measure body composition by comparison with a four-  
598 compartment model. *Br J Nutr* 2000, 83, 115-122.

599 [67] Lloret Linares, C., Ciangura, C., Bouillot, J.-L., Coupaye, M., Declèves, X., Poitou, C.,  
600 Basdevant, A., Oppert, J.-M., Validity of Leg-to-Leg Bioelectrical Impedance Analysis to  
601 Estimate Body Fat in Obesity. *Obesity Surg* 2011, 21, 917-923.

602 [68] Browning, L. M., Mugridge, O., Chatfield, M., Dixon, A., Aitken, S., Joubert, I., Prentice, A.  
603 M., Jebb, S. A., Validity of a new abdominal bioelectrical impedance device to measure abdominal  
604 and visceral fat: comparison with MRI. *Obesity* 2010, *18*, 2385-2391.

605 [69] Fekete, K., Gyorei, E., Lohner, S., Verduci, E., Agostoni, C., Decsi, T., Long-chain  
606 polyunsaturated fatty acid status in obesity: a systematic review and meta-analysis. *Obesity Rev*  
607 2015, *16*, 488-497.

608 [70] Harris, W. S., Pottala, J. V., Varvel, S. A., Borowski, J. J., Ward, J. N., McConnell, J. P.,  
609 Erythrocyte omega-3 fatty acids increase and linoleic acid decreases with age: observations from  
610 160,000 patients. *Prostaglandins Leukot Essent Fatty Acids* 2013, *88*, 257-263.

611

612

613

**Table 1** Profile of plasma n-6 fatty acids between normal weight, overweight and obese Irish adults (n=781)

	Normal weight (n=294, % males=40.5)		Overweight (n=301, % males=58.8)		Obese (n=186, % males=52.2)		P-trend
	Mean	SD	Mean	SD	Mean	SD	
18:2n-6	27.30	3.39	25.65	3.52	24.69	3.85	$2.9 \times 10^{-7}$
18:3n-6	0.52	0.18	0.58	0.18	0.59	0.20	0.003
20:2n-6	0.37	0.19	0.38	0.16	0.39	0.22	1.000
20:3n-6	2.02	0.47	2.02	0.47	2.11	0.47	0.003
20:4n-6	7.80	1.99	7.73	1.75	7.41	1.74	0.312
22:4n-6	0.24	0.09	0.23	0.08	0.25	0.09	1.000
D5D	4.08	1.53	4.10	1.95	3.71	1.32	$8.6 \times 10^{-4}$
D6D	0.020	0.01	0.024	0.01	0.025	0.01	$1.6 \times 10^{-5}$

Participants who are under-weight (n=5) and participants with missing BMI value were excluded from this analysis (n=34). D5D: delta-5-desaturase (20:4n6/20:3n6); D6D: delta-6-desaturase (18:3n6/18:2n6).

General linear contrast model adjusted for age and gender was used, P-values were corrected by multiple number of traits. P-values that exceed 1.0 after correction for multiple testing have been marked down to 1.000.

614

615

616

**Table 2.** Demographics across quartiles of plasma linoleic acid in Irish adults (n=820).

	Quartiles of plasma linoleic acid								P-trend
	1 (n=205)		2 (n=205)		3 (n=205)		4 (n=205)		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
	% LA=21.1		% LA=25.0		% LA=27.4		% LA=30.5		
Age (years)	50.67	17.00	42.31	16.89	38.04	15.06	36.46	14.02	1.9×10 <sup>-19</sup>
BMI (kg/m <sup>2</sup> )	29.02	5.75	27.07	4.99	26.66	4.60	25.30	3.73	2.2×10 <sup>-12</sup>
Body fat (%)	31.46	10.12	28.66	9.16	28.68	8.66	26.32	8.74	3.1×10 <sup>-6</sup>
Waist circumference (cm)	97.83	15.27	91.05	13.76	90.00	12.95	85.56	11.54	1.2×10 <sup>-15</sup>
Waist hip ratio	0.92	0.09	0.87	0.09	0.86	0.08	0.85	0.07	3.5×10 <sup>-13</sup>
Systolic blood pressure (mmHg)	131.93	18.12	123.86	16.03	122.22	17.38	118.23	15.12	1.7×10 <sup>-13</sup>
Diastolic blood pressure (mmHg)	81.44	11.13	78.05	10.43	76.86	10.69	75.34	9.97	2.2×10 <sup>-7</sup>
Physical activity (MET hrs/wk)									
Work activities	47.77	60.31	58.88	52.81	62.70	58.90	66.08	70.89	0.026
Recreational activities	27.55	41.01	37.64	51.90	38.37	56.67	34.12	39.37	1.000
Gender	M=118,F=87		M=105,F=100		M=93,F=112		M=96,F=109		0.061
Oral contraceptive users (%)	2.9		2.4		6.3		2.4		0.094
Metabolic risk factor (%)									3.7×10 <sup>-20</sup>
No Risk Factors	18.0		39.0		46.3		60.5		
1-2 Risk Factors	53.2		48.3		47.3		27.8		
> 3 Risk Factors	22.0		8.8		4.9		4.9		
Smoking status (%)									0.001
Current smoker	22.2		18.6		21.5		18.6		
Former smoker	36.0		27.9		21.5		18.6		
Never smoke	41.9		53.4		57.1		62.7		
Social class (%)									0.153
Professional/managerial/technical	48.2		46.2		42.8		43.2		
Non-manual skilled	12.8		17.6		19.9		18.1		
Manual skilled	21.0		13.6		11.9		14.1		
Semi-skilled/unskilled	17.9		22.6		25.4		24.6		
Weight class									2.7×10 <sup>-7</sup>
Normal weight	20.0		37.8		38.6		53.9		
Overweight	45.3		37.8		39.1		32.1		
Obese	34.7		24.5		22.3		14.0		

Comparisons of age, BMI, body fat, waist circumference, physical activity and blood pressure across quartiles were assessed using Linear contrast analysis; comparisons of gender, smoking status, social class, weight class and metabolic risk factor were based on chi square statistic. P-values were corrected by multiple number of traits. P-values that exceed 1.0 after correction for multiple testing have been marked down to 1.000. M: males; F: females.

**Table 3.** Concentration of biochemistry biomarkers across quartiles of plasma linoleic acid in Irish adults (n=820).

	Quartiles of plasma linoleic acid								P-trend
	1 (n=205)		2 (n=205)		3 (n=205)		4 (n=205)		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
<b>Metabolism</b>									
Glucose (mmol/l)	5.71	1.21	5.23	1.28	5.11	0.66	5.03	0.86	0.058
Insulin ( $\mu$ IU/ml)	12.77	10.80	8.73	6.15	8.97	8.08	7.48	5.28	$5.7 \times 10^{-5}$
C-Peptide (ng/ml)	3.00	2.51	2.08	2.41	1.93	1.97	1.48	1.40	$7.0 \times 10^{-5}$
Triglyceride (mmol/l)	1.83	0.88	1.28	0.58	1.03	0.38	0.86	0.33	$2.1 \times 10^{-35}$
Total Cholesterol (mmol/l)	5.02	1.10	5.02	0.93	4.96	1.01	4.75	0.94	1.000
LDL Cholesterol (mmol/l)	2.72	0.93	2.90	0.79	2.89	0.92	2.76	0.80	0.160
HDL Cholesterol (mmol/l)	1.47	0.41	1.54	0.41	1.60	0.43	1.59	0.42	1.000
HOMA-IR	3.43	3.76	2.09	1.74	2.16	2.59	1.71	1.33	$9.1 \times 10^{-6}$
QUICKI	0.34	0.04	0.36	0.04	0.36	0.04	0.37	0.04	$1.9 \times 10^{-5}$
<b>Immune function and metabolic health</b>									
PAI-1 (ng/ml)	25.46	9.16	24.00	9.68	22.11	8.60	20.29	7.08	$1.2 \times 10^{-4}$
TNF- $\alpha$ (pg/ml)	7.60	2.70	6.99	2.60	6.66	2.48	6.28	1.55	0.005
High Sensitivity CRP (mg/l)	3.49	3.27	2.84	3.00	2.35	2.40	2.08	2.09	0.003
Adiponectin ( $\mu$ g/ml)	5.16	2.69	5.78	3.28	6.52	3.28	6.35	3.13	0.002
Leptin Soluble Receptor (ng/ml)	27.16	6.76	27.19	7.24	27.48	7.21	27.49	6.27	1.000
Resistin(ng/ml)	5.92	2.50	6.30	2.88	6.01	2.64	6.04	2.72	1.000
<b>Plasma fatty acids (%) and desaturase</b>									
Total n-6 PUFA	32.1	2.99	36.1	2.02	38.38	1.93	41.04	2.32	$1.4 \times 10^{-138}$
18:2n-6	21.07	2.03	25.02	0.78	27.36	6.17	30.50	1.86	$4.0 \times 10^{-252}$
20:2n-6	0.44	0.25	0.37	0.15	0.35	0.17	0.35	0.13	$1.9 \times 10^{-7}$
18:3n-6	0.62	0.20	0.58	0.17	0.54	0.18	0.51	0.17	$8.8 \times 10^{-5}$
20:3n-6	2.05	0.49	2.13	0.49	2.02	0.46	1.97	0.42	0.019
20:4n-6	7.62	2.18	7.71	1.72	7.89	1.79	7.49	1.64	1.000
22:4n-6	0.25	0.09	0.24	0.08	0.23	0.09	0.23	0.08	0.156
EPA	1.56	1.00	1.43	0.78	1.32	0.80	1.18	0.70	1.000
DHA	2.42	0.97	2.47	0.93	2.44	0.93	2.25	0.85	1.000
D5D	3.95	1.68	3.82	1.30	4.18	2.09	4.01	1.40	1.000
D6D	0.03	0.01	0.02	0.01	0.02	0.01	0.02	0.01	$4.7 \times 10^{-35}$
n-6/n-3 ratio	5.96	1.64	6.72	1.67	7.59	2.50	8.43	2.26	$1.2 \times 10^{-17}$
Total SFA	37.16	3.50	35.77	3.07	34.65	2.77	33.63	2.81	$6.0 \times 10^{-32}$
Total MUFA	24.94	3.63	22.41	3.60	21.42	3.14	20.07	3.13	$1.1 \times 10^{-21}$
14:0	1.47	0.44	1.19	0.31	1.01	0.32	0.93	0.26	$7.8 \times 10^{-30}$
16:0	24.48	1.96	23.20	1.45	22.50	1.35	21.56	1.18	$9.1 \times 10^{-53}$
18:0	9.86	2.61	9.97	2.62	9.82	2.59	9.80	2.49	0.288



14:1n-5	0.14	0.07	0.10	0.05	0.08	0.05	0.06	0.04	$4.0 \times 10^{-30}$
16:1n-7	3.43	0.94	2.71	0.61	2.30	0.48	1.88	0.49	$7.5 \times 10^{-71}$
18:1n-9	20.38	3.52	18.60	3.49	18.08	3.11	17.20	3.07	$4.7 \times 10^{-6}$

Comparisons of concentration of biochemical biomarkers and plasma fatty acids (log transformed for skewed variable) across quartiles were assessed using General Linear Contrast Model adjusted for age, body fat, blood pressure, waist circumference, waist hip ratio, smoking status, working energy expenditure and BMI. P-values were corrected by multiple number of traits. P-values that exceed 1.0 after correction for multiple testing have been marked down to 1.000. QUICKI: quantitative insulin sensitivity check index; PAI-1: plasminogen activator inhibitor-1; hs-CRP: high-sensitivity-C-reactive protein; D5D: delta-5-desaturase (20:4n6/20:3n6); D6D: delta-6-desaturase (18:3n6/18:2n6).

619

620

**Table 4.** Haplotype characteristics of 15-locus, 8-locus and 3-locus haplotypes (n=790).

Haplotypes		Alleles <sup>a</sup>	Number of minor allele	Frequency
<u>15-locus haplotype (rs174537 to rs482548)</u>				
Haplo1	TTGTCTTTCGAATGC	222222221211221	11	17.0%
Haplo2	GCCCTCCCCAACCT	111111111111112	1	11.1%
Haplo3	TTGTCTCTCGTGTCC	22222121222211	11	2.3%
Haplo4	TTGTCTCTTCTGTCC	22222122122211	11	7.1%
Haplo5	TTCCTTCTTGAATGC	221112122211221	8	3.0%
Haplo6	TTGTCTCTTGAATGC	22222122211221	11	1.6%
Haplo7	GCCCTCCCCAACCC	111111111111111	0	53.9%
<u>8-locus haplotype (rs174537 to rs99780)</u>				
Haplo1	TTGTCTTT	22222222	8	19.2%
Haplo2	TTGTCTCT	22222212	7	11.4%
Haplo3	TTCCTTCT	22111212	4	3.0%
Haplo4	GCCCTCCC	11111111	0	65.6%
<u>3-locus haplotype (rs2524299 to rs174583)</u>				
Haplo1	TGT	222	3	10.9%
Haplo2	AAT	112	1	22.8%
Haplo3	AAC	111	0	65.5%

<sup>a</sup>1, major allele; 2, minor allele.

621

622

623

624

625

626

627

628

**Table 5.** Estimates of haplotypes effects on plasma linoleic acid,  $\Delta$ -5 desaturase and  $\Delta$ -6 desaturase (n=790).

	Plasma LA		$\Delta$ -5 desaturase		$\Delta$ -6 desaturase	
	Estimates $\pm$ SE	P value	Estimates $\pm$ SE	P value	Estimates $\pm$ SE	P value
<u>15-locus haplotype</u>						
Haplo1	0.82 $\pm$ 0.24	7.1 $\times 10^{-4}$	-0.99 $\pm$ 0.11	1.81 $\times 10^{-19}$	-0.004 $\pm$ 0.001	4.4 $\times 10^{-9}$
Haplo2	0.25 $\pm$ 0.28	0.37	-0.10 $\pm$ 0.13	0.446	-0.001 $\pm$ 0.001	0.211
Haplo3	1.31 $\pm$ 0.53	0.01	-0.86 $\pm$ 0.25	6.37 $\times 10^{-4}$	-0.006 $\pm$ 0.001	5.5 $\times 10^{-5}$
Haplo4	0.77 $\pm$ 0.34	0.02	-0.81 $\pm$ 0.15	1.21 $\times 10^{-7}$	-0.005 $\pm$ 0.001	1.1 $\times 10^{-9}$
Haplo5	1.74 $\pm$ 0.51	6.9 $\times 10^{-4}$	-0.34 $\pm$ 0.23	0.146	-0.006 $\pm$ 0.001	4.9 $\times 10^{-6}$
Haplo6	0.94 $\pm$ 0.70	0.18	-0.87 $\pm$ 0.32	6.44 $\times 10^{-3}$	-0.006 $\pm$ 0.002	4.9 $\times 10^{-4}$
<u>8-locus haplotype</u>						
Haplo1	0.78 $\pm$ 0.22	4.06 $\times 10^{-4}$	-1.02 $\pm$ 1.00	2.45 $\times 10^{-24}$	-0.003 $\pm$ 0.001	8.9 $\times 10^{-10}$
Haplo2	0.91 $\pm$ 0.27	6.86 $\times 10^{-4}$	-0.85 $\pm$ 0.01	4.27 $\times 10^{-12}$	-0.005 $\pm$ 0.001	6.4 $\times 10^{-15}$
Haplo3	1.59 $\pm$ 0.50	1.67 $\times 10^{-3}$	-0.32 $\pm$ 0.50	1.67 $\times 10^{-3}$	-0.006 $\pm$ 0.002	8.8 $\times 10^{-6}$
<u>3-locus haplotype</u>						
Haplo1	0.86 $\pm$ 0.27	1.64 $\times 10^{-3}$	-0.85 $\pm$ 0.13	2.34 $\times 10^{-11}$	-0.004 $\pm$ 0.001	2.4 $\times 10^{-12}$
Haplo2	0.90 $\pm$ 0.20	1.03 $\times 10^{-5}$	-0.88 $\pm$ 0.09	5.34 $\times 10^{-21}$	-0.004 $\pm$ 0.002	4.7 $\times 10^{-14}$

The most common haplotype was used as reference.

Maximum likelihood estimation adjusted for age, BMI, body fat, waist circumference, working energy expenditure, waist hip ratio, blood pressure and smoking habits was used.

629

630

631

**Table 6.** Multiple linear regression analysis of possible plasma linoleic acid determinates with 15-locus haplotype (n=790).

	$\beta$ Coefficient	Standard error	P	Variance explained
Age (years)	-0.047	0.010	<0.001	3.2%
Waist circumference (cm)	-0.036	0.012	0.002	8.1%
PUFA intakes (%TE)	0.325	0.065	<0.001	3.7%
Fibre intakes (g/10MJ)	0.035	0.019	0.066	0.5%
Systolic blood pressure (mmHg)	-0.028	0.010	0.002	3.6%
Body fat (%)	-0.040	0.017	0.020	3.7%
Sugar intakes (%TE)	-0.090	0.032	0.005	1.3%
Carbohydrate intakes (%TE)	0.098	0.027	<0.001	2.0%
15-locus Haplotype <sup>a</sup>				3.9%
Haplo1	0.796	0.253	0.002	
Haplo2	0.455	0.305	0.136	
Haplo3	1.843	0.604	0.002	
Haplo4	1.011	0.345	0.003	
Haplo5	1.177	0.569	0.039	
Haplo6	0.893	0.817	0.275	

<sup>a</sup> Most common haplotype was used as a reference