



**QUEEN'S
UNIVERSITY
BELFAST**

Up and away: ontogenic transference as a pathway for aerial dispersal of microplastics

Al-Jaibachi, R., Cuthbert, R. N., & Callaghan, A. (2018). Up and away: ontogenic transference as a pathway for aerial dispersal of microplastics. *Biology Letters*, 14(9). <https://doi.org/10.1098/rsbl.2018.0479>

Published in:
Biology Letters

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2018 The Authors. Published by the Royal Society. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Up and away: ontogenic transference as a pathway for aerial dispersal of microplastics

Rana Al-Jaibachi¹, Ross N. Cuthbert^{1,2}, Amanda Callaghan¹

¹*Ecology and Evolutionary Biology, School of Biological Sciences, University of Reading, Harborne Building, Reading RG6 6AS, England*

²*Institute for Global Food Security, School of Biological Sciences, Medical Biology Centre, Queen's University Belfast, Belfast BT9 7BL, Northern Ireland*

**Corresponding author: e-mail, a.callaghan@reading.ac.uk*

Abstract

Microplastics (MPs) are ubiquitous pollutants found in marine, freshwater and terrestrial ecosystems. With so many MPs in aquatic systems it is inevitable that they will be ingested by aquatic organisms, and be transferred up through the food chain. However, to date, no study has considered whether MPs can be transmitted by means of ontogenic transference i.e. between life stages that utilise different habitats. Here, we determine whether fluorescent polystyrene beads could transfer between *Culex* mosquito life stages and, particularly, could move into the flying adult stage. We show for the first time that MPs can be transferred ontogenically from a feeding (larva) into a non-feeding (pupa) life stage and subsequently into the adult terrestrial life stage. However, transference is dependent on particle size, with smaller 2µm MPs transferring readily into pupae and adult stages, whilst 15µm MPs transferred at a significantly reduced rate. Microplastics appear to accumulate in the Malpighian tubule renal excretion system. The transfer of MPs to the adults represents a potential aerial pathway to contamination of new environments. Thus, any organism that feeds on terrestrial life phases of freshwater insects could be impacted by MPs found in aquatic ecosystems.

Keywords

Food chain: ontology; life stage; Malpighian tubules, microplastics; *Culex pipiens*

Introduction

Microplastics (MPs) are ubiquitous pollutants found in marine, freshwater and terrestrial ecosystems [1–3]. There is little doubt that plastic and MP pollution is a major environmental concern globally. Despite this, there is relatively little research into the impact of MPs on freshwater ecosystems, with most research concentrating on marine systems and organisms [2]. MPs have been defined as plastic particles smaller than 5mm in size [4,5]. However, this simple description covers a wide range of types, with polypropylene, polyethylene and polystyrene MPs entering the environment in different shapes and sizes, including fibres, pellets and cosmetic beads [6,7]. MPs are categorised based on their origin as primary or secondary types, depending on whether they were released into the environment as MPs (primary) or have degraded to that size in the environment (secondary) [8,9]. Microplastics pass through terrestrial environments in wastewater [2,10]. Rivers can subsequently deliver MPs into lakes, where they can be found in high concentrations [11–13].

Microplastics are ingested by aquatic organisms, and can be transferred through the food chain in both freshwater and marine environments [14–18]. However, to date no study has considered whether MPs can be transmitted by means of ontogenic transference i.e. between life stages that utilise different habitats. Freshwater environments are inhabited by insects that spend their juvenile stages in water but their adult stages in the terrestrial environment. Such insects include mayflies, dragonflies, midges and mosquitoes, most of which are eaten by terrestrial vertebrates. This raises the potential for MPs to enter terrestrial ecosystems from freshwater habitats aerially *via* transference to adult invertebrate life stages. Here, we thus determine whether 2 and 15µm fluorescent polystyrene beads could transfer between insect life stages and, particularly, could move into the flying adult stage. Fluorescent beads were selected to enable MPs to be easily detected in the non-feeding stages and also to allow an investigation of location within the body during metamorphosis. The *Culex pipiens* mosquito complex was

selected as a model for this study given their worldwide distribution and broad habitat preference [19]. Mosquitoes develop through four feeding larval instars and a non-feeding pupal stage, and finally emerge into a flying adult.

Materials and methods

For additional details of all methods and analyses, see the electronic supplementary material.

Two types of MPs were used: a 2µm fluorescent yellow-green carboxylate-modified polystyrene (density 1.050g/cm³, excitation 470nm; emission 505nm, Sigma-Aldrich, UK) and a 15.45±1.1µm fluorescent dragon green polystyrene (density 1.06 g/cm³ (5x10⁶ particles/ml, excitation 480nm; emission 520nm, Bangs Laboratories Inc., USA). Four treatments were used; a control with no microplastics, a treatment of 8x10⁵ 2µm particles/ml, a treatment of 8x10² 15µm particles/ml, and a 1:1 mixture of both treatments. Each replicate (five per treatment) contained ten 3rd instar *C. pipiens* larvae in a 50ml glass beaker filled with 50ml of tap water. The control and all treatments contained 100mg of pelleted guinea pig food. Treatments were assigned randomly to a position on the laboratory bench to reduce experimental error.

One random individual was removed from each beaker when every mosquito had moulted into the 4th instar, and again when they pupated or emerged as adults. All samples were then placed in separate 1.5ml Eppendorf tubes and stored at -20 °C prior to examination. Microplastics were extracted from mosquitoes by homogenization and filtration. The filter membrane was examined using an epi-fluorescent microscope (Zeiss Axioskop) under a 20x lens to count the number of fluorescent MPs. Adults were further dissected under a binocular stereo microscope (0.7X-4.5X) to extract the gut and quantify the numbers of MPs under the epi-fluorescent microscope [20].

All data were analyzed using the statistical software R v3.4.2 [21]. Microplastic counts were analysed using generalized linear models (GLMs) assuming a quasi-Poisson distribution. Uptake of microplastics was examined with respect to ‘particle size’, ‘treatment’ and ‘life stage’. We performed model simplification via stepwise removal of non-significant effects. Tukey tests were used post hoc for multiple comparisons.

Results

No MPs were found in control groups of any mosquito life stage. However, when exposed to MPs, particles were transferred from larvae to pupae, and subsequently to adults (Table S1). Significantly more 2µm particles were found in mosquito life stages than 15µm particles overall ($F_{1, 58}=303.98$, $P<0.001$). However, densities of MPs were significantly different between life stages ($F_{2, 56}=160.42$, $P<0.001$), with MP numbers significantly falling as mosquitoes moved between successive ontogenic levels (all $P<0.001$). Microplastics uptake was also significantly greater overall in mixed exposure treatments ($F_{1, 55}=6.00$, $P=0.02$). Although 2µm particles were transferred to adults in all instances, we found no transference of 15µm particles following single treatment exposures. However, in the mixed MPs treatment, transference to adults of both 2µm and 15µm particles was evidenced (Figure 1). Microplastic ingestion was confirmed by fluorescent microscopy where the beads were detected in the adult abdomen, specifically inside the Malpighian tubules (Figure 2).

Discussion

Here, we show for the first time that MPs can be transferred ontogenically from a feeding (larval) into a non-feeding (pupal) life stage and subsequently into the flying (adult) life stage. Transference through to adults was found in both MP sizes, although the larger 15µm MPs were not ingested as readily as the 2µm MPs. Dissection of mosquito adults showed that 2µm

MPs accumulated in the renal excretion system of Malpighian tubules which, unlike the gut, pass from larvae to adult stages without visible reorganization [22]. This has been demonstrated previously to provide a physical transport system between stages during metamorphosis for *Pseudomonas* bacteria and seems to be important for ontogenic transmission from larvae to adults [23].

Few 15µm MPs were transferred into adults suggesting that MP size is an important factor in ontogenic transfer which could be related to the transfer and accumulation of MPs in the Malpighian tubes. Although the translocation mechanism of MPs to the Malpighian tubules is unclear in mosquitoes, analysis of fish, fiddler crab and marine mussels has demonstrated that MPs can be translocated from gastrointestinal tracts into other tissues in a wide range of phyla [24, 25,26]. Malpighian tubules have an entry point to the gut between the mid- and hindgut of mosquitoes, but the flow of fluid is from the Malpighian tubules to the hindgut [27]. Diptera are known to produce structures called concretions in the Malpighian tubules which have been shown to sequester heavy metals [28]. However, it is unlikely that this pathway would operate with a solid MP.

Our results have important implications since any aquatic life stage that is able to consume MPs and transfer them to their terrestrial life stage is a potential vector of MPs onto novel aerial and terrestrial habitats. Ingestion of MP-contaminated organisms by terrestrial organisms is not new [29]. Indeed, the widespread distribution of MPs in marine environments has meant that animals such as fish and shellfish sold for human consumption are contaminated with a range of plastics with a consequent transference of MPs between trophic levels [24]. Unlike MP fibres, which are common in the air and atmosphere, there has been no evidence for MPs being transported into the air [24]. We have demonstrated that species with both aquatic and terrestrial stages can harbour MPs through their life history, with aerial and terrestrial animals accordingly open to MP exposure.

References

1. Sighicelli M, Pietrelli L, Lecce F, Iannilli V, Falconieri M, Coscia L, Di Vito S, Nuglio S, Zampetti G. 2018 Microplastic pollution in the surface waters of Italian Subalpine Lakes. *Environ. Pollut.* **236**, 645–651. (doi:10.1016/j.envpol.2018.02.008)
2. Wagner M, Lambert S. 2018 *Freshwater Microplastics*. Cham: Springer International Publishing. (doi:10.1007/978-3-319-61615-5)
3. Mason SA, Welch V, Neratko J. 2018 Synthetic polymer contamination in bottled water. *Fredonia State Univ. New York* , 1–17.
4. Eriksen M, Lebreton LCM, Carson HS, Thiel M, Moore CJ, Borerro JC, Galgani F, Ryan PG, Reisser J. 2014 Plastic Pollution in the World's Oceans: More than 5 Trillion Plastic Pieces Weighing over 250,000 Tons Afloat at Sea. *PLoS One* **9**, e111913. (doi:10.1371/journal.pone.0111913)
5. Imhof HK, Ivleva NP, Schmid J, Niessner R, Laforsch C. 2013 Contamination of beach sediments of a subalpine lake with microplastic particles. *Curr. Biol.* **23**, 1–15. (doi:10.1016/j.cub.2013.09.001)
6. Andrady AL, Neal MA. 2009 Applications and societal benefits of plastics. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **364**, 1977–1984. (doi:10.1098/rstb.2008.0304)
7. Rocha-Santos T, Duarte AC. 2014 A critical overview of the analytical approaches to the occurrence, the fate and the behavior of microplastics in the environment. *Trends Anal. Chem.* **65**, 47–53. (doi:10.1016/j.trac.2014.10.011)
8. Moore CJ. 2008 Synthetic polymers in the marine environment: A rapidly increasing, long-term threat. *Environ. Res.* **108**, 131–139. (doi:10.1016/j.envres.2008.07.025)
9. Barnes DKA, Galgani F, Thompson RC, Barlaz M. 2009 Accumulation and fragmentation of plastic debris in global environments. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **364**, 1985–98. (doi:10.1098/rstb.2008.0205)
10. Mason SA, Garneau D, Sutton R, Chu Y, Ehmann K, Barnes J, Fink P, Papazissimos D,

178 Rogers DL. 2016 Microplastic pollution is widely detected in US municipal wastewater
 179 treatment plant effluent. *Environ. Pollut.* **218**, 1045–1054.
 180 (doi:10.1016/j.envpol.2016.08.056)

181 11. Eriksen M, Mason S, Wilson S, Box C, Zellers A, Edwards W, Farley H, Amato S. 2013
 182 Microplastic pollution in the surface waters of the Laurentian Great Lakes. *Mar. Pollut.*
 183 *Bull.* **77**, 177–182. (doi:10.1016/j.marpolbul.2013.10.007)

184 12. Fischer EK, Paglialonga L, Czech E, Tamminga M. 2016 Microplastic pollution in lakes
 185 and lake shoreline sediments - A case study on Lake Bolsena and Lake Chiusi (central
 186 Italy). *Environ. Pollut.* **213**, 648–657. (doi:10.1016/j.envpol.2016.03.012)

187 13. Su L, Xue Y, Li L, Yang D, Kolandhasamy P, Li D, Shi H. 2016 Microplastics in Taihu
 188 Lake, China. *Environ. Pollut.* **216**, 711–719. (doi:10.1016/j.envpol.2016.06.036)

189 14. Aljaibachi R, Callaghan A. 2018 Impact of polystyrene microplastics on *Daphnia*
 190 *magna* mortality and reproduction in relation to food availability. *PeerJ* **6**, e4601.
 191 (doi:10.7717/peerj.4601)

192 15. Cole M, Lindeque P, Fileman E, Halsband C, Goodhead R, Moger J, Galloway TS. 2013
 193 Microplastic ingestion by zooplankton. *Environ. Sci. Technol.* **47**, 6646–6655.
 194 (doi:10.1021/es400663f)

195 16. Scherer C, Brennholt N, Reifferscheid G, Wagner M. 2017 Feeding type and
 196 development drive the ingestion of microplastics by freshwater invertebrates. *Sci. Rep.*
 197 **7**, 17006. (doi:10.1038/s41598-017-17191-7)

198 17. Sussarellu R *et al.* 2016 Oyster reproduction is affected by exposure to polystyrene
 199 microplastics. *Proc. Natl. Acad. Sci.* **113**, 2430–2435. (doi:10.1073/pnas.1519019113)

200 18. Messinetti S, Mercurio S, Parolini M, Sugni M, Pennati R. 2018 Effects of polystyrene
 201 microplastics on early stages of two marine invertebrates with different feeding
 202 strategies. *Environ. Pollut.* **237**, 1080–1087. (doi:10.1016/j.envpol.2017.11.030)

- 203 19. Dow JA, Maddrell SH, Görtz A, Skaer NJ, Brogan S, Kaiser K. 1994 The malpighian
204 tubules of *Drosophila melanogaster*: a novel phenotype for studies of fluid secretion
205 and its control. *J. Exp. Biol.* **197**, 421–8.
- 206 20. Coleman J, Juhn J, James AAA. 2007 Dissection of Midgut and Salivary Glands from
207 *Ae. aegypti* Mosquitoes. *J. Vis. Exp.* , 2007. (doi:10.3791/228)
- 208 21. R Development Core Team. 2017 R: A Language and Environment for Statistical
209 Computing. *R Found. Stat. Comput. Vienna Austria* (doi:10.1038/sj.hdy.6800737)
- 210 22. Clements AN. 1992 *The biology of mosquitoes. Volume 1: Development, nutrition and*
211 *reproduction*. London: Chapman and Hall.
- 212 23. Chavshin AR, Oshaghi MA, Vatandoost H, Yakhchali B, Zarenejad F, Terenius O. 2015
213 Malpighian tubules are important determinants of *Pseudomonas* transstadial
214 transmission and longtime persistence in *Anopheles stephensi*. *Parasites and Vectors* **8**,
215 1–7. (doi:10.1186/s13071-015-0635-6)
- 216 24. Von Moos N, Burkhardt-Holm P, Köhler A. 2012 Uptake and Effects of Microplastics
217 on Cells and Tissue of the Blue Mussel *Mytilus edulis* L. after an Experimental
218 Exposure. *Environ. Sci. Technol.* **46**, 11327–11335.
- 219 25. Brennecke D, Ferreira EC, Costa TMM, Appel D, da Gama BAP, Lenz M. 2015
220 Ingested microplastics (>100µm) are translocated to organs of the tropical fiddler crab
221 *Uca rapax*. *Mar. Pollut. Bull.* **96**, 491–495. (doi:10.1016/j.marpolbul.2015.05.001)
- 222 26. Avio CG, Gorbi S, Regoli F. 2015 Experimental development of a new protocol for
223 extraction and characterization of microplastics in fish tissues: First observations in
224 commercial species from Adriatic Sea. *Mar. Environ. Res.* **111**, 18–26.
225 (doi:10.1016/j.marenvres.2015.06.014)
- 226 27. Piermarini PM. 2016 *Renal Excretory Processes in Mosquitoes*. 1st edn. Elsevier Ltd.
227 (doi:10.1016/bs.aiip.2016.04.003)

28. Leonard EM, Pierce LM, Gillis PL, Wood CM, O'Donnell MJ. 2009 Cadmium transport by the gut and Malpighian tubules of *Chironomus riparius*. *Aquat. Toxicol.* **92**, 179–186. (doi:10.1016/j.aquatox.2009.01.011)
29. Huerta Lwanga E, Gertsen H, Gooren H, Peters P, Salánki T, Van Der Ploeg M, Besseling E, Koelmans AA, Geissen V. 2016 Microplastics in the Terrestrial Ecosystem: Implications for *Lumbricus terrestris* (Oligochaeta, Lumbricidae). *Environ. Sci. Technol.* (doi:10.1021/acs.est.5b05478)
30. Dris R, Gasperi J, Mirande C, Mandin C, Guerrouache M, Langlois V, Tassin B. 2017 A first overview of textile fibers, including microplastics, in indoor and outdoor environments. *Environ. Pollut.* (doi:10.1016/j.envpol.2016.12.013)

Figure legends

Figure 1: Mean (log count) uptake of microplastics following single 2µm/15µm and mixed 2µm/15µm exposures. Means are ±SE (*n*=5 per experimental group).

Figure 2. Epi-fluorescent microscope images showing fluorescent microplastic particles within (A) the abdomen of an adult mosquito before dissection, and (B) the abdominal Malpighian tubules following dissection.

Ethics

Ethics committee approval was not required.

Data accessibility

Data files are available in online supplementary material.

Author contribution

A.C. and R.A. designed the study, R.A. collected data and R.A. and R.N.C. analysed data and drafted the manuscript. AC revised and approved the manuscript.

Competing interests

255 We declare we have no competing interests.

256 **Funding**

257 A.C. is funded by the University of Reading. R.A. is self-funded and R.N.C. is funded
258 through the Department for the Economy, Northern Ireland.

259 **Acknowledgements**

260 We thank Natali Ortiz-Perea for assisting with mosquito colony rearing.