



**QUEEN'S
UNIVERSITY
BELFAST**

Norms of basic elementary operators on algebras of regular operators

Wickstead, A. W. (2015). Norms of basic elementary operators on algebras of regular operators. *Proceedings of the American Mathematical Society*. <https://doi.org/10.1090/proc/12664>

Published in:

Proceedings of the American Mathematical Society

Document Version:

Peer reviewed version

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2015 AMS

This is an open access article published under a Creative Commons Attribution-NonCommercial-NoDerivs License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

First published in the Proceedings of the American Mathematical Society, 2015, published by the American Mathematical Society.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

NORMS OF BASIC ELEMENTARY OPERATORS ON ALGEBRAS OF REGULAR OPERATORS

A.W. WICKSTEAD

(Communicated by)

ABSTRACT. We show that if E is an atomic Banach lattice with an order continuous norm, $A, B \in \mathcal{L}^r(E)$ and $M_{A,B}$ is the operator on $\mathcal{L}^r(E)$ defined by $M_{A,B}(T) = ATB$ then $\|M_{A,B}\|_r = \|A\|_r\|B\|_r$ but that there is no real $\alpha > 0$ such that $\|M_{A,B}\| \geq \alpha\|A\|_r\|B\|_r$.

1. INTRODUCTION.

An *elementary* operator on a Banach algebra \mathcal{A} is an operator of the form $T \mapsto \sum_{k=1}^n A_k T B_k$, where each $A_k, B_k \in \mathcal{A}$. If $n = 1$ such an elementary operator is called *basic*. There is a substantial literature about elementary operators, much of it in the setting of C^* -algebras with there also being significant results about the norm in the case of the algebra, $\mathcal{L}(E)$, of all bounded operators on a Banach space E . The elementary starting point for such results in that case is the fact, which is a simple consequence of the Hahn-Banach theorem, that the norm of a basic elementary operator $T \mapsto ATB$ is precisely $\|A\|\|B\|$. This statement refers, of course, to the norm of $T \mapsto ATB$ as an element of the space of bounded operators on $\mathcal{L}(E)$. In this note, we look at the corresponding question for the space of *regular* operators, $\mathcal{L}^r(E)$ on a Dedekind complete Banach lattice E . This is both a Banach lattice (for the usual order for operators between Banach lattices) and a Banach algebra under composition provided it is equipped with the *regular* norm $\|T\|_r = \|\|T\|\|$. As $T \mapsto ATB$ defines elements of both $\mathcal{L}(\mathcal{L}^r(E))$ and $\mathcal{L}^r(\mathcal{L}^r(E))$, this means that we can ask about both its regular and operator norms. We will see that, at least in the slightly specialized setting in which we are able to obtain satisfactory results, there are rather different answers in the two cases. This is in contrast to the only related result in the literature, Proposition 1 of [6], that this author knows of.

Our interest in this problem is not purely for the sake of generalization. For $A_k, B_k \in Z(E)$, the *centre* of E (the order ideal generated by the identity on E), operators of the form $T \mapsto \sum_{k=1}^n A_k T B_k$ have already featured in the literature in connection with descriptions of the centre of $\mathcal{L}^r(E)$. The interested reader is referred to [1], [2] and [9] for details, albeit in slightly more general setting.

We refer the reader to [5] for the standard terminology in the field of Banach lattices and linear operators on them.

2010 *Mathematics Subject Classification*. Primary: 47B48, 47B60.

Key words and phrases. Regular operators, basic elementary operators, Banach lattices.

©XXXX American Mathematical Society

The author would like to thank Chen Jinxi for pointing out an error in the proof of Theorem 3.1 in a previous version of this paper. He would also like to thank Anton Schep for providing a correction for that error and for pointing out that our Theorem 3.1 is essentially stated in the remark following Hilfsatz 3 in [7].

2. NOTATION AND HISTORY.

We will always assume that E is a Dedekind complete Banach lattice, so that $\mathcal{L}^r(E)$ is also a Dedekind complete Banach lattice (and Banach algebra) under the regular norm. If $A, B \in \mathcal{L}^r(E)$ then we will denote by $M_{A,B}$ the operator $T \mapsto ATB$ which is easily seen to map $\mathcal{L}^r(E)$ into itself and to be an element of both $\mathcal{L}(\mathcal{L}^r(E))$ and of $\mathcal{L}^r(\mathcal{L}^r(E))$. *A priori* there is no reason to expect the two norms $\|M_{A,B}\|$ and $\|M_{A,B}\|_r$ to be the same, but what historical results exist might lead one to believe in such a possibility.

In [6], Schep studied the *left regular representation* of $\mathcal{L}^r(E)$ which, in our notation, is the mapping $A \mapsto M_{A,I}$ where I denotes the identity operator on E . In Proposition 1, he establishes that $\|M_{A,I}\| = \|M_{A,I}\|_r = \|A\|_r$. After that result he also remarks that $\|M_{I,B}\| = \|M_{I,B}\|_r = \|B\|_r$. As part of his Proposition 1, he also establishes the rather significant fact that $|M_{A,I}| = M_{|A|,I}$. Rather earlier, in Satz 3.1 (a) of [8], Synnatzschke proved the slightly more general result that $|M_{A,B}| = M_{|A|,B}$ whenever B is positive. In the remarks following Proposition 1, Schep leaves open the possibility that $|M_{I,B}| = M_{I,|B|}$. We must leave open the same question, but do wish to point out that this is obviously true when E is reflexive. This is because the mapping $T \mapsto T^* : \mathcal{L}^r(E) \rightarrow \mathcal{L}^r(E^*)$ is a linear order isomorphism and $M_{I,B}(T)^* = M_{B^*,I^*}(T^*)$. We feel that it should be true that $|M_{I,B}| = M_{I,|B|}$ at least when E has an order continuous norm. Satz 3.1 (b) of [8] is relevant here.

3. THE REGULAR NORM.

Our results for the regular norm of $M_{A,B}$ are restricted to the setting that E is atomic with an order continuous norm. Such Banach lattices are certainly Dedekind complete. The reader will not go too far astray in thinking of E as being something like ℓ_p for $1 \leq p < \infty$.

One reason that we are able to obtain results for such spaces is the observation that if e is an atom of E and $T \in \mathcal{L}^r(E)$ then $|T|(e) = |T(e)|$. This follows from the well known formula $|T|(e) = \bigvee\{T(x) : |x| \leq e\}$. As e is an atom, the x 's here are of the form λe for $\lambda \in [-1, 1]$ so that $|T|(e) = \bigvee\{|\lambda T(e)| : \lambda \in [-1, 1]\} = |T(e)|$. If $f \in E^*$ and $x \in E$, then the operator $T = f \otimes x : e \mapsto f(e)x$ is certainly a member of $\mathcal{L}^r(E)$. Furthermore, $M_{A,B}(T)(e) = A(T(B(e))) = A(f(B(e))x) = f(B(e))A(x)$, for any $e \in E$.

Theorem 3.1. *If E is an atomic Banach lattice with an order continuous norm and $A, B \in \mathcal{L}^r(E)$ then $|M_{A,B}| = M_{|A|,|B|}$.*

Proof. Suppose that $T \in \mathcal{L}^r(E)_+$. As products of positive operators are positive, decomposing A and B into their positive and negative parts we have

$$\begin{aligned} M_{A,B}(T) &= ATB = (A_+ - A_-)T(B_+ - B_-) \\ &= A_+TB_+ - A_+TB_- - A_-TB_+ + A_-TB_- \\ &\leq A_+TB_+ + A_+TB_- + A_-TB_+ + A_-TB_- \\ &= (A_+ + A_-)T(B_+ + B_-) \\ &= |A|T|B| = M_{|A|,|B|}(T), \end{aligned}$$

which shows that $|M_{A,B}| \leq M_{|A|,|B|}$.

We establish the reverse inequality first for the particular case that $T = f \otimes x$ where $f \in E_+^*$ and $x \in E_+$. As non-negative linear combinations of the atoms in E are norm dense in the positive cone of E , it suffices to prove that $|M_{A,B}|(f \otimes x)(e) \geq M_{|A|,|B|}(f \otimes x)(e)$ for any atom $e \in E_+$. Note that

$$\begin{aligned} |M_{A,B}|(f \otimes x) &= \bigvee \{M_{A,B}(S) : |S| \leq f \otimes x\} \\ &\geq \bigvee \{M_{A,B}(g \otimes y) : |y| \leq x, |g| \leq f\}, \end{aligned}$$

so that

$$\begin{aligned} |M_{A,B}|(f \otimes x)(e) &\geq \bigvee \{|M_{A,B}(g \otimes y)|(e) : |y| \leq x, |g| \leq f\} \\ &= \bigvee \{|M_{A,B}(g \otimes y)(e)| : |y| \leq x, |g| \leq f\} \\ &= \bigvee \{|g(Be)Ay| : |y| \leq x, |g| \leq f\} \\ &= \bigvee \{|g(Be)||Ay| : |y| \leq x, |g| \leq f\} \\ &= \bigvee \{|g(Be)| : |g| \leq f\} \bigvee \{|Ay| : |y| \leq x\} \\ &= f(|Be|)|A|(x) \\ &= f(|Be|)|A|(x) \quad [\text{As } e \text{ is an atom.}] \\ &= M_{|A|,|B|}(f \otimes x)(e). \end{aligned}$$

We have now established that $|M_{A,B}|(f \otimes x) = (M_{|A|,|B|}(f \otimes x))$, if $f \in E_+^*$ and $x \in E_+$. By linearity this will also hold for any $f \in E^*$ and $x \in E$ and therefore $|M_{A,B}|(T) = M_{|A|,|B|}(T)$ for any finite rank operator T on E .

Now take any $T \in \mathcal{L}^r(E)$, F a finite set of atoms in E and P_F the band projection onto the linear span of these atoms. As E is atomic with an order continuous norm, $P_F(x) \rightarrow x$ in norm as the sets F increase. In particular, this means that the finite rank operators $P_F \circ T$ converge upwards to T for the strong operator topology. It follows easily that $M_{|A|,|B|}(P_F \circ T)$ converges strongly to $M_{|A|,|B|}(T)$. On the other hand if $T \geq 0$ and $x \in E_+$ then we have

$$0 \leq |M_{A,B}|(T - P_F \circ T)(x) \leq M_{|A|,|B|}(T - P_F \circ T)(x) \rightarrow 0$$

and hence $|M_{A,B}|(T - P_F \circ T)(x) \rightarrow 0$ from which it follows that $|M_{A,B}|(P_F \circ T) \rightarrow |M_{A,B}|(T)$ strongly for positive T and hence for all regular T . From these observations it is immediate that $|M_{A,B}|(T) = M_{|A|,|B|}(T)$ for all $T \in \mathcal{L}^r(E)$ and hence that $|M_{A,B}| = M_{|A|,|B|}$. \square

Corollary 3.2. *If E is an atomic Banach lattice with an order continuous norm and $A, B \in \mathcal{L}^r(E)$ then $\|M_{A,B}\|_r = \|A\|_r \|B\|_r$.*

Proof. We certainly always have $\|M_{A,B}\|_r \leq \|A\|_r \|B\|_r$. If A and B are positive then, as in the Banach space case, we have $\|M_{A,B}\| = \|A\| \|B\|$. To see this take $\epsilon > 0$ and pick $x \in E_+$ with $\|x\| = 1$ and $\|A(x)\| \geq \|A\| - \epsilon$, where we may choose $x \geq 0$ as A is positive. Similarly pick $y \in E_+$ with $\|y\| = 1$ and $\|B(y)\| \geq \|B\| - \epsilon$. Let $f \in E_+^*$ with $\|f\| = 1$ and $f(B(y)) = \|B(y)\|$. Take $T = f \otimes x$ which is positive with $\|T\| = \|T\|_r = \|f\| \times \|x\| = 1$. As

$$\|M_{A,B}(T)(y)\| = |f(B(y))| \|A(x)\| \geq (\|A\| - \epsilon)(\|B\| - \epsilon)$$

we have $\|M_{A,B}\| \geq \|A\| \|B\| = \|A\|_r \|B\|_r$. From the preceding Theorem, in general we have $\|M_{A,B}\|_r = \| |M_{A,B}| \| = \|M_{|A|,|B|}\| = \| |A| \| \| |B| \| = \|A\|_r \|B\|_r$. \square

4. THE OPERATOR NORM

In this section, not only do we establish that we need not have $\|M_{A,B}\| = \|A\|_r \|B\|_r$ (even in the setting of Corollary 3.2), but that there need not even be a constant $\alpha > 0$ such that $\|M_{A,B}\| \geq \alpha \|A\|_r \|B\|_r$ for all $A, B \in \mathcal{L}^r(E)$. Our example will be on the space ℓ_2 and will be built from finite dimensional examples. We need to estimate the operator norms, $\|A\|$, and regular norms, $\|A\|_r$, of certain square matrices $A = (a_{ij})_{i,j=1}^m$ when considered as acting as linear operators on an m -dimensional Hilbert space ℓ_2^m . We write $A_i = \sum_{j=1}^m a_{ij}$ for the sum of the i 'th row of A and then set $c_j = \sum_{i=1}^m a_{ij} A_i$.

Proposition 4.1. *If A is a $m \times m$ matrix then*

- (1) $\frac{1}{m} \sum_{j=1}^m c_j = \frac{1}{m} \sum_{i=1}^m A_i^2 = \frac{1}{m} \|(A_i)_{i=1}^m\|_2^2 \leq \|A\|^2$
- (2) *If A has non-negative entries then also $\|A\|^2 \leq \max\{c_j : 1 \leq j \leq m\}$.*

Proof. For (1) set $x = (\frac{1}{\sqrt{m}})_1^m$ so that $\|x\|_2 = 1$ and $Ax = \frac{1}{\sqrt{m}}(A_1, A_2, \dots, A_m)$ so that $\|Ax\|_2^2 = \frac{1}{m} \sum_{i=1}^m A_i^2$, showing that $\|A\|^2 \geq \frac{1}{m} \sum_{i=1}^m A_i^2$. Note also that

$$\begin{aligned} \sum_{j=1}^m c_j &= \sum_{j=1}^m \left(\sum_{i=1}^m a_{ij} A_i \right) \\ &= \sum_{i=1}^m \left(\sum_{j=1}^m a_{ij} \right) A_i = \sum_{i=1}^m A_i^2. \end{aligned}$$

To establish (2) we need to estimate $\|Ax\|_2$ for $x \in \ell_2^m$ with $\|x\|_2 \leq 1$ and (as A has non-negative entries) all $x_i \geq 0$.

$$\begin{aligned} \|Ax\|_2^2 &= \sum_{i=1}^m \left(\sum_{j=1}^m a_{ij}x_j \right)^2 = \sum_{i=1}^m \left[\sum_{j=1}^m a_{ij}^2x_j^2 + 2 \sum_{1 \leq j < k \leq m} a_{ij}a_{ik}x_jx_k \right] \\ &\leq \sum_{i=1}^m \left[\sum_{j=1}^m a_{ij}^2x_j^2 + \sum_{1 \leq j < k \leq m} a_{ij}a_{ik}(x_j^2 + x_k^2) \right] = \sum_{i=1}^m \left[\sum_{j=1}^m \left(\sum_{k=1}^m a_{ij}a_{ik}x_j^2 \right) \right] \\ &= \sum_{i=1}^m \left[\sum_{j=1}^m a_{ij} \left(\sum_{k=1}^m a_{ik} \right) x_j^2 \right] = \sum_{i=1}^m \left[\sum_{j=1}^m a_{ij}A_i x_j^2 \right] = \sum_{j=1}^m \left[\sum_{i=1}^m a_{ij}A_i \right] x_j^2 \\ &= \sum_{j=1}^m c_j x_j^2 \leq \max\{c_j : 1 \leq j \leq m\} \sum_{j=1}^m x_j^2 \leq \max\{c_j : 1 \leq j \leq m\} \end{aligned}$$

so that $\|A\|^2 \leq \max\{c_j : 1 \leq j \leq m\}$. \square

Corollary 4.2. *If A is a $m \times m$ matrix of non-negative reals and each row of A is constant then*

$$\|A\|^2 = \frac{1}{m} \sum_{i=1}^m A_i^2 = \frac{1}{m} \|(A_i)_{i=1}^m\|_2$$

Proof. This follows immediately as all the c_j are the same. \square

Corollary 4.3. *If A is any real $m \times m$ matrix with each row constant then*

$$\|A\|_r^2 = \frac{1}{n} \sum_{i=1}^m A_i^2.$$

Proof. If $B = |A|$ and B_i is the sum of the i 'th row of B then $B_i = |A_i|$ so that $B_i^2 = A_i^2$. B has constant rows so that

$$\|A\|_r = \|B\| = \frac{1}{m} \sum_{i=1}^m B_i^2 = \frac{1}{m} \sum_{i=1}^m A_i^2 = \frac{1}{m} \|(A_i)_{i=1}^m\|_2^2.$$

\square

Define a sequence of matrices inductively by $K_0 = (1)$ and

$$K_{n+1} = \begin{pmatrix} K_n & K_n \\ -K_n & K_n \end{pmatrix}.$$

These matrices were introduced to the study of positive operators by Krengel in [3] and [4]. The matrix $2^{-n/2}K_n$ is an orthogonal matrix so that as an operator on $\ell_2^{2^n}$ it is an isometry. It follows that $\|K_n x\|_2 = 2^{n/2}\|x\|_2$ for all $x \in \ell_2^{2^n}$ and in particular $\|K_n\| = 2^{n/2}$ whilst it is easily checked, and well known, that $\|K_n\|_r = \| |K_n| \| = 2^n$. In particular, we will make use of the fact that $|K_n|$ is the $2^n \times 2^n$ matrix with all its entries equal to 1.

Our intention is to estimate the norm of the operators $M_{K_n, |K_n|} \in \mathcal{L}(\mathcal{L}^r(\ell_2^{2^n}))$ defined by $M_{K_n, |K_n|}(T) = K_n T |K_n|$.

Example 4.4. Although $\|K_n\|_r = \| |K_n| \| = 2^n$, $\|M_{K_n, |K_n|}\| \leq 2^{3n/2}$.

Proof. Take an arbitrary real $2^n \times 2^n$ matrix T . All the entries in the i 'th row of $T|K_n|$ are equal to T_i , the sum of the i 'th row in T . The columns in $K_n T|K_n|$ are therefore all equal to the product of K_n with the column vector $(T_i)_{i=1}^{2^n}$ so that the column vectors of row sums for $K_n T|K_n|$ is precisely $2^n K_n (T_i)_{i=1}^{2^n}$. We may thus apply Corollary 4.3 to see that

$$\begin{aligned} \|K_n T|K_n|\|_r^2 &= 2^{-n} \times 2^{2n} \|K_n (T_i)_{i=1}^{2^n}\|_2^2 \\ &= 2^n \|K_n (T_i)_{i=1}^{2^n}\|_2^2 \\ &= 2^n (2^{n/2} \|(T_i)_{i=1}^{2^n}\|_2)^2 \\ &= 2^{2n} \|(T_i)_{i=1}^{2^n}\|_2^2 \end{aligned}$$

whilst by Proposition 4.1, $\|T\|_r^2 \geq \|T\|^2 \geq 2^{-n} \|(T_i)_{i=1}^{2^n}\|_2^2$, so that $\|M_{K_n, |K_n|}(T)\|_r^2 = \|K_n T|K_n|\|_r^2 \leq 2^{3n} \|T\|_r^2$ and $\|M_{K_n, |K_n|}\|_r \leq 2^{3n/2}$. \square

Embedding each of the examples on $\ell_2^{2^n}$ into $\mathcal{L}^r(\ell_2)$ shows that for regular operators A, B on ℓ_2 there is no real $\alpha > 0$ such that $\|M_{A, B}\| \geq \alpha \|A\|_r \|B\|_r$.

REFERENCES

- [1] G. J. H. M. Buskes, P. G. Dodds, B. de Pagter, and A. R. Schep, *Up-down theorems in the centre of $\mathcal{L}_b(E, F)$* , *Nederl. Akad. Wetensch. Indag. Math.* **48** (1986), no. 1, 1–9. MR834315 (87f:47052)
- [2] Brian Jefferies and Susumu Okada, *An operator bound related to regular operators*, *Arch. Math. (Basel)* **66** (1996), no. 3, 219–227, DOI 10.1007/BF01195708. MR1373249 (97a:47042)
- [3] Ulrich Krengel, *Über den Absolutbetrag stetiger linearer Operatoren und seine Anwendung auf ergodische Zerlegungen*, *Math. Scand.* **13** (1963), 151–187 (German). MR0176034 (31 #310)
- [4] ———, *Remark on the modulus of compact operators*, *Bull. Amer. Math. Soc.* **72** (1966), 132–133. MR0190752 (32 #8162)
- [5] Peter Meyer-Nieberg, *Banach lattices*, Universitext, Springer-Verlag, Berlin, 1991. MR1128093 (93f:46025)
- [6] Anton R. Schep, *A remark on the uniform zero-two law for positive contractions*, *Arch. Math. (Basel)* **53** (1989), no. 5, 493–496, DOI 10.1007/BF01324724. MR1019163 (90j:47046)
- [7] J. Synnatzschke, *Der Operator ${}_B T_C : A \rightarrow BAC$ als vollstetiger Fastintegraloperator*, *Wiss. Z. Karl-Marx-Univ. Leipzig Math.-Natur. Reihe* **24** (1975), no. 5, 529–534. Beiträge zur Theorie nichtnormierbarer topologischer Algebren. MR0428103 (55 #1132)
- [8] ———, *Über einige verbandstheoretische Eigenschaften der Multiplikation von Operatoren in Vektorverbänden*, *Math. Nachr.* **95** (1980), 273–292, DOI 10.1002/mana.19800950125 (German). MR592901 (82b:47048)
- [9] A. W. Wickstead, *The centre of spaces of regular operators*, *Math. Z.* **241** (2002), no. 1, 165–179, DOI 10.1007/s002090200411. MR1930989 (2004b:47069)

PURE MATHEMATICS RESEARCH CENTRE, QUEEN'S UNIVERSITY BELFAST, BELFAST BT7 1NN, NORTHERN IRELAND.

E-mail address: A.Wickstead@qub.ac.uk