



**QUEEN'S  
UNIVERSITY  
BELFAST**

## Direct Healthcare Costs of Sedentary Behaviour in the UK

Heron, L., O'Neill, C., McAneney, H., Kee, F., & Tully, M. (2019). Direct Healthcare Costs of Sedentary Behaviour in the UK. *Journal of Epidemiology & Community Health*, 73(7), 625-629. <https://doi.org/10.1136/jech-2018-211758>

**Published in:**  
Journal of Epidemiology & Community Health

**Document Version:**  
Peer reviewed version

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

### **Publisher rights**

Copyright 2019 BMJ.

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

### **General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

### **Open Access**

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

1 Title: The Direct Healthcare Costs of Sedentary Behaviour in the UK

2 Authors:

3 Leonie Heron<sup>1</sup>, Professor Ciaran O'Neill<sup>1</sup>, Dr Helen McAneney<sup>1</sup>, Professor Frank Kee<sup>1</sup>,  
4 Professor Mark A Tully<sup>2</sup>

5 <sup>1</sup>Centre for Public Health, Queen's University Belfast, Institute of Clinical Sciences, Block A,  
6 Royal Victoria Hospital, Belfast, BT12 6BA

7 <sup>2</sup>School of Health Sciences, Ulster University, Shore Road, Newtownabbey, Co. Antrim,  
8 BT37 0QB

9

10 Corresponding author:

11 Professor Mark Tully

12 Room 01D116, School of Health Sciences, Ulster University, Shore Road, Newtownabbey,  
13 Co. Antrim, BT37 0QB

14 Email: [m.tully@ulster.ac.uk](mailto:m.tully@ulster.ac.uk)

15 Telephone: +44 28 9036 6977

16

17 Word count, excluding title page, abstract, references, figures and tables: 2728

18 2877

19

### **What is already known on this subject?**

Recent evidence indicates that prolonged sedentary behaviour increases the risk of several non-communicable diseases. However, a large proportion of the UK population spend their lives in sedentary jobs and leisure activities, and official physical activity recommendations regarding sedentariness are vague.

### **What does this study add?**

Diseases associated with prolonged sedentary behaviour cost the NHS £0.8 billion in the 2016-17 financial year. This estimate can inform decision-makers who are prioritising resources in healthcare and make a financial case for reducing sedentary behaviour in the UK.

1

## 2 **ABSTRACT**

3 **Background:** Growing evidence indicates that prolonged sedentary behaviour increases the  
4 risk of several chronic health conditions and all-cause mortality. Sedentary behaviour is  
5 prevalent among adults in the United Kingdom (UK). Quantifying the costs associated with  
6 sedentary behaviour is an important step in the development of public health policy.

7 **Methods:** National Health Service (NHS) costs associated with prolonged sedentary  
8 behaviour ( $\geq 6$  hours/day) were estimated over a one-year period in 2016-17 costs. We  
9 calculated a population attributable fraction (PAF) for five health outcomes (type 2 diabetes,  
10 cardiovascular disease [CVD], colon cancer, endometrial cancer, and lung cancer).  
11 Adjustments were made for potential double counting due to co-morbidities. We also  
12 calculated the avoidable deaths due to prolonged sedentary behaviour using the PAF for all-  
13 cause mortality.

14 **Results:** The total NHS costs attributable to prolonged sedentary behaviour in the UK in  
15 2016-17 were £0.8 billion, which included expenditure on CVD (£424 million), type 2  
16 diabetes (£281 million), colon cancer (£30 million), lung cancer (£19 million), and  
17 endometrial cancer (£7 million). After adjustment for potential double-counting, the  
18 estimated total was £0.7 billion. If prolonged sedentary behaviour was eliminated, 48,024 UK  
19 deaths might have been avoided in 2016.

1 **Conclusions:** In this conservative estimate of the direct healthcare costs in the UK, prolonged  
2 sedentary behaviour causes a considerable burden to the NHS. This estimate may be used by  
3 decision makers when prioritising healthcare resources and investing in preventative public  
4 health programmes.

5 **Keywords:** public health; sedentary behaviour; sitting time; cost analysis; health expenditure;  
6 healthcare cost; physical activity.

7

## 1 INTRODUCTION

2 Adults in the United Kingdom (UK) have become increasingly sedentary as modern  
3 technology has changed everyday life.[1] Sedentary behaviour is distinct from physical  
4 inactivity and refers to sitting or lying while expending low amounts of energy ( $\leq 1.5$   
5 metabolic equivalents [METs]).[2] National guidelines recommend minimising time spent  
6 sedentary[3] without specifying how many hours/day of sitting might be harmful. A recent  
7 meta-analysis reported that spending 6-8 hours/day sedentary increases future risk of all-  
8 cause and cardiovascular disease (CVD).[4] This study defined sedentary behaviour as  
9 spending at least six hours of waking time sedentary. Thirty percent of adults in the UK are  
10 sedentary for a least six hours/day during the week, which rises to 37% at the weekend.[5]  
11 Consequently, many individuals in the UK are at greater risk of chronic disease.

12 Sedentary behaviour is an established risk factor for several non-communicable diseases.  
13 Strong evidence suggests that high levels of sitting time lead to increased risk of CVD, type 2  
14 diabetes, and all-cause mortality (risk of mortality from all causes, not only those mentioned  
15 here).[6] Additionally, moderate evidence indicates an increased risk of colon, endometrial,  
16 and lung cancer.[6] These diseases all contribute considerably to morbidity and mortality in  
17 the UK. Thus, addressing the problem of sedentary behaviour could potentially reduce the  
18 burden of disease.

19 Awareness of the economic burden of sedentary behaviour could inform and motivate  
20 policymakers to address this risk factor. Estimates of the cost impacts allow decision makers  
21 to prioritise funding and make an economic argument for investment in prevention. Estimates  
22 for the financial impact of many lifestyle risk factors in the UK are available, such as obesity,  
23 smoking, and physical inactivity,[7, 8] however none exist thus far for sedentary  
24 behaviour. As a result, this study aims to estimate the direct healthcare costs of prolonged  
25 sedentary behaviour in the UK.

## 26 METHODS

27 Costs were estimated from a healthcare payer perspective (UK National Health Service  
28 [NHS]) using a prevalence-based and population attributable fraction- (PAF) approach,  
29 following methodology employed by Ding et al.[7]

### 30 **Quantifying the increased risk to health due to sedentary behaviour**

1 We selected the most suitable meta-analyses cited in a recent report of the relationship  
2 between sedentary behaviour and health[6] in order to extract the relative risks (RRs).  
3 Appropriate studies employed a prospective design, non-diseased participants at baseline, and  
4 adjusted for levels of physical activity in their statistical model. Furthermore, the researchers  
5 had investigated the association by comparing the most sedentary individuals with the least  
6 sedentary, and we preferred studies which had used sedentary time as an exposure. Two  
7 studies were appropriate for the outcome of CVD[19,21]: we chose the more recent meta-  
8 analysis by Pandey et al. as it had included three additional applicable studies. After  
9 examining data from the primary studies, we excluded those that did not meet the exact  
10 criteria above and repooled the risk estimate using Review Manager (RevMan version 5.3).

### 11 **Estimating the extent of sedentary behaviour in the UK population**

12 The Health Survey for England (HSE) 2012[5] reported that 30% of adults in England spent  
13 at least six hours/day sedentary on weekdays, and 37% of adults at the weekend. We used  
14 these figures to estimate the percentage of UK adults who are sedentary on any given day of  
15 the week.

16 The PAF formula we have used requires the prevalence of sedentary behaviour at baseline in  
17 those who went on to become cases (i.e., experiencing the adverse outcome). This  
18 information is not readily available. Therefore, we calculated prevalence “adjustment  
19 factors”[9] using data from cohort studies. We searched for cohort studies on Pubmed that  
20 fitted the same criteria mentioned in the previous section and had specifically measured and  
21 reported sedentary behaviour for the total population and for cases only at baseline. We  
22 preferred European-based studies and larger studies with longer follow-up times to give more  
23 reliable adjustment factors. The proportion of cases in the highest reported category of  
24 sedentary behaviour was divided by the proportion of people at baseline in the highest  
25 category to produce an adjustment factor. For example, Stamatakis et al.[10] reported that  
26 34.1% of all study participants and 38.3% of diabetes cases were sedentary at baseline. The  
27 adjustment factor was 1.12 (38.3/34.1). We then multiplied the adjustment factor by the  
28 prevalence of sedentary behaviour in the general population in order to estimate the  
29 additional prevalence among cases.

30 Table 1. Prevalence adjustment factors calculated from longitudinal study data

Disease	Study	Country	Prevalence of prolonged sedentary behaviour <sup>1</sup> at baseline	Prevalence of prolonged sedentary behaviour <sup>1</sup> in cases	Adjustment factor
Type 2 diabetes	Stamatakis et al., 2017[10]	UK	0.34	0.38	1.12
CVD incidence	Bjork Petersen et al., 2014[35]	Denmark	0.13	0.16	1.23
All-cause mortality	van der Ploeg et al., 2012[36]	Australia	0.06	0.12	1.87
Lung cancer	Ukawa et al., 2013[37]	Japan	0.25	0.28	1.10
Colon cancer	Simons et al., 2013[38]	Netherlands	0.26	0.32	1.22
Endometrial cancer	Gierach et al., 2009[39]	USA	0.08	0.10	1.20

CVD = cardiovascular disease. <sup>1</sup>Prolonged sedentary behaviour indicates spending at least six hours sedentary during waking hours.

1

## 2 Calculating PAFs for each health outcome

3 The PAF estimate the contribution of a risk factor to the total burden of a disease in a given  
4 population. Here, PAFs estimate the reduction in disease that would occur if prolonged  
5 sedentary behaviour was eliminated. The following formula from Rockhill, Newman and  
6 Weinberg[11] was used:

$$7 \quad PAF(\%) = \frac{p_1(RR_{adj} - 1)}{RR_{adj}} \times 100$$

8 where  $p_1$  is the prevalence of sedentary behaviour among cases and  $RR_{adj}$  is the pooled  
9 adjusted RR, comparing the most sedentary individuals with the least sedentary.

10 It integrates the pooled adjusted RR ( $RR_{adj}$ ) estimates and the proportion of sedentary  
11 individuals who became cases ( $p_1$ ). It is appropriate to use when confounding is present.[11]  
12 We calculated Wald intervals for each of the PAFs using Monte Carlo simulation methods  
13 (250,000 simulations) on Microsoft Excel (2016).[12] These techniques accounted for  
14 random error and uncertainty in confounding from the pooled RR estimates and the  
15 prevalence of sedentary behaviour (see supplementary file 1 for further details).

## 16 Estimating NHS expenditure for each disease

1 Healthcare budgets for specific disease groupings was available for the NHS in England for  
2 the nearest financial year 2012-13,[13] Wales for 2016-17,[14] and Scotland 2011-12.[15]  
3 Costs were standardised to the year 2017 by adjusting costs for inflation using the hospital  
4 and community health services (HCHS) index, a weighted average of annual increases in pay  
5 and prices in healthcare services.[16] Healthcare budget data for Northern Ireland was  
6 unavailable, thus we estimated costs for this region based on the number of diagnoses  
7 compared to the rest of UK. Further details are reported in supplementary file 2. All costs are  
8 in pounds sterling (GBP).

### 9 **Calculating costs attributable to sedentary behaviour**

10 We multiplied the adjusted PAFs and their 95% CIs by the total disease expenditure to  
11 estimate the NHS costs attributable to sedentary behaviour in the UK. Since the timeframe  
12 for this analysis is one year, discounting was unnecessary.

13 Thirty percent of Europeans with type 2 diabetes are also affected by CVD.[17] Therefore,  
14 30% of the type 2 diabetes expenditure attributable to sedentary behaviour was subtracted  
15 from the total costs to adjust for double-counting caused by this co-morbidity. This is  
16 consistent with the approach used by Ding et al.[7]

### 17 **Estimating the avoidable deaths due to sedentary behaviour**

18 In addition, we multiplied the PAF for all-cause mortality by the total number of UK deaths  
19 in 2016 to estimate the number of deaths that would have been avoided if prolonged  
20 sedentary behaviour was completely eliminated. As complete elimination is unrealistic, the  
21 number of avoidable deaths was also estimated for 10%, 30%, and 50% potential reductions  
22 in the proportion of sedentary individuals (i.e., sedentary  $\geq 6$  hours/day).

## 23 **RESULTS**

24 The health outcomes that we considered most relevant for this analysis were type 2 diabetes,  
25 CVD, and all-cause mortality (strong evidence), and endometrial, colon, and lung cancers  
26 (moderate evidence).[6] Pooled analyses of crude or age-adjusted estimates were not  
27 available in the literature. The PAF formula given required a pooled risk estimate and so we  
28 extracted RRs from the least adjusted models and pooled them to give an unadjusted RR  
29 estimate. Most models were age-adjusted only, however several of the least-adjusted models  
30 were already adjusted for more variables. Crucially, none of the models had adjusted for  
31 physical activity level, an important confounder in the association between sitting time and



1 health.[18] Table 2 presents the prevalence of sedentary behaviour for each health outcome  
 2 with the associated RRs and PAFs.

3 Table 2. Estimates of prevalence, relative risk of disease, and population attributable fractions  
 4 for sedentary behaviour in the UK.

<b>Strength of evidence<sup>1</sup></b>	<b>Health outcome</b>	<b>Proportion of prolonged sedentary adults in cases(%)<sup>2</sup></b>	<b>RR (95% CI)</b>	<b>PAF (95% CI)</b>
Strong	Type 2 diabetes	34%	1.88 (1.62, 2.17)	16.9% (14.0%, 19.6%)
	CVD incidence	38%	1.14 (1.09, 1.19)	4.9% (4.2%, 5.5%)
	All-cause mortality	57%	1.25 (1.16, 1.34)	11.6% (10.3%, 12.9%)
Moderate	Lung cancer	34%	1.27 (1.06, 1.52)	7.5% (3.9%, 11.0%)
	Colon cancer	37%	1.30 (1.12, 1.49)	9.0% (7.3%, 10.7%)
	Endometrial cancer <sup>5</sup>	40%	1.28 (1.08, 1.53)	8.0% (6.0%, 10.0%)

RR = relative risk; CI = confidence interval; PAF = population attributable fraction; CVD = cardiovascular disease.

<sup>1</sup>Strength of evidence as reported by 2018 Physical Activity Guidelines Advisory Committee (2018).

<sup>2</sup>Estimated from weekday and weekend proportions available from Health Survey for England.[5]

5 We re-pooled the adjusted RR for the association between sedentary behaviour and type 2  
 6 diabetes presented by Biswas et al.[19] to exclude a cross-sectional study.[20] The updated  
 7 pooled RR estimate was 1.88 (95% CI 1.62, 2.17). Based on the PAF calculations, 16.9%  
 8 (14.0%, 19.6%) of cases of type 2 diabetes were associated with sedentary behaviour. Pandey  
 9 et al.[21] reported an adjusted RR of 1.14 (95% CI 1.09, 1.19) for the association between  
 10 CVD and sedentary behaviour. Just under five per cent (4.9% [4.2%, 5.5%]) of CVD could  
 11 be attributable to sedentary behaviour. The adjusted RR for the association between sedentary  
 12 behaviour and all-cause mortality[21] was reanalysed in order to exclude four studies. The  
 13 studies were inappropriate for the following reasons: their baseline populations were not free  
 14 of disease;[22] they reported a per-hour association,[23] rather than comparing individuals in  
 15 the most and least sedentary categories; their design was cross-sectional;[20] or they reported  
 16 an inapplicable association (one study reported the association between those who were

1 ‘consistently nonsedentary’ vs. ‘consistently sedentary’).[24] The sedentary time definition  
 2 that they used, reported ranges, and estimated median sedentary time are reported in  
 3 supplementary file 3.

4 The new pooled RR estimate was 1.25 (95% CI 1.16, 1.33) and the corresponding PAF for  
 5 this association was 11.6% (10.3%, 12.9%). Shen et al.[25] investigated the risk of cancer  
 6 associated with higher sedentary behaviour. They reported adjusted RRs for lung cancer (1.27  
 7 [95% CI 1.06, 1.52]), colon cancer (1.30 [95% CI 1.12, 1.49]), and endometrial cancer (1.28  
 8 [95% CI 1.08, 1.53]). The PAF calculations showed that 7.5% (3.9%, 11.0%) of lung cancer;  
 9 9.0% (7.3%, 10.7%) of colon cancer; and 8.0% (6.0%, 10.0%) of endometrial cancer could be  
 10 attributable to sedentary behaviour.

11 Thus, if sedentary behaviour was eliminated in the UK, 48,024 deaths in 2016 might have  
 12 been avoided. More realistically, if levels of sedentary behaviour were 10%, 30%, or 50%  
 13 lower in 2016, we might have avoided 4,802, 12,006, or 24,012 deaths respectfully.

14 It is also important to note that the total budgets adjusted for inflation to 2016/17 costs were  
 15 considerably lower than reported total budgets for 2016/17 for England[26] and Scotland.[27]  
 16 Individual healthcare budgets were not available for these years and so costs had to be  
 17 inflated. Table 3 provides the NHS costs attributable to sedentary behaviour and 95% CIs.  
 18 CVD is associated with the greatest cost attributable to sedentary behaviour of £424 million  
 19 (£367, £480 million), followed by £281 million (£233, £327 million) for type 2 diabetes.  
 20 Costs for specific cancers attributable for sedentary behaviour were much lower; £19 million  
 21 (£10, £28 million) for lung cancer, £30 (£24, £35 million) for colon cancer, and £7 million  
 22 (£5, £9 million) for endometrial cancer. Together, the total costs attributable to sedentary  
 23 behaviour are £762 million (£639, £879 million). Total UK NHS health expenditure is  
 24 estimated to be £65.7 billion for 2016/17, indicating that sedentary behaviour accounted for  
 25 1.2% of total expenditure.

26 Table 3. Costs Attributable to Diseases Associated with Sedentary Behaviour

Disease	Costs Attributable to Sedentary Behaviour by UK region (£million, 2016-17)				Total UK NHS costs attributable to sedentary behaviour (£million, 2016-17 [95% CI])
	England	Scotland	Wales	NI	
Type 2 diabetes	£242.54	£14.86	£16.90	£7.03	£281.34 (£233.46, £326.85)
CVD	£348.95	£40.75	£22.80	£11.88	£424.38 (£366.61, £480.09)

Lung cancer	£13.54	£3.78	£1.32	£0.52	£19.16 (£9.92, £27.98)
Colon cancer	£22.74	£3.99	£2.90	£0.80	£29.64 (£23.96, £35.12)
Endometrial cancer	£5.72	£0.74	£0.63	£0.20	£7.29 (£5.44, £9.07)
<b>Total costs</b>	<b>£633.49</b>	<b>£64.13</b>	<b>£44.55</b>	<b>£20.44</b>	<b>£761.80 (£639.40, £879.11)</b>

NHS = National Health Service; CI = confidence interval; CVD = cardiovascular disease; NI = Northern Ireland.

1

2 After adjustment for double-counting, the NHS costs attributable to sedentary behaviour is  
3 £677 million. An alternative method[7] was also used as a sensitivity analysis. A meta-  
4 analysis reported the RR of having CVD as being 206% higher for people with type 2  
5 diabetes compared to those without type 2 diabetes.[28] Based on the prevalence of CVD in  
6 the general population (4.28%, as reported by the British Heart Foundation),[29] we estimate  
7 that 8.82% of people with type 2 diabetes have CVD. After subtracting 8.82% of type 2  
8 diabetes expenditure, the total costs attributable to sedentary behaviour were £737 million.

9 After an additional sensitivity analysis which excluded diseases for which only moderate  
10 evidence of an association was available, the total costs attributable to sedentary behaviour  
11 were £706 million (£600, £807 million), i.e., approximately eight per cent lower. The small  
12 change is due to the much lower incidence and prevalence of the individual cancers in  
13 comparison to CVD and type 2 diabetes expenditure.

14

## 15 **DISCUSSION**

16 This cost-of-illness analysis found that prolonged sedentary behaviour costs the UK NHS  
17 £0.8 billion in the financial year 2016-17. After adjustments for double-counting, this  
18 estimate was slightly reduced to £0.7 billion. The results suggested that 11.6% of all-cause  
19 mortality was associated with sedentary behaviour. Therefore, 48,024 deaths might have been  
20 avoided in 2016 if sedentary behaviour was eliminated in the UK.

21 The total costs presented are likely to be a conservative estimate of the true burden of  
22 sedentary behaviour. There are reported links between sedentary behaviour and several other  
23 cancers, musculoskeletal disorders, and mental health disorders.[30–32] However, the  
24 evidence remains limited, hence they were excluded from this study. Moreover, the analysis

1 used a PAF-approach which typically produces lower estimates than alternative econometric  
2 approaches.[33]

3 CVD, type 2 diabetes, and colon, endometrial and lung cancers are all linked to sedentary  
4 behaviour (PAFs ranged from 4.9%-16.9%). Patterson et al.[4] also calculated PAFs for  
5 sedentary behaviour in a recent meta-analysis, where the exposure was TV viewing time and  
6 the methodology (using a Monte-Carlo micro-simulation) was somewhat different. Thus, it is  
7 difficult to compare these estimates. Nonetheless, it is interesting to note that the PAFs for  
8 type 2 diabetes, CVD and all-cause mortality are of the same order of magnitude (i.e., type 2  
9 diabetes > all-cause mortality > CVD). The PAFs for CVD are similar (5% [95% CI: 1%,  
10 8%] from Patterson vs. 4.9% (95% CI: 1.8%, 7.9%) in the present study). This indicates that  
11 although the studies differ in their definition of sedentary behaviour and in the methods used,  
12 there is considerable agreement in the observed pattern of the relationships.

13 This study had several strengths. We have calculated PAFs for sedentary behaviour in the UK  
14 using the best data available, and we have included all conditions reported as having  
15 moderate to strong evidence of an association.[6] The analysis followed several suggestions  
16 from a checklist for reporting estimates of the economic costs of risk factors by Ding et  
17 al.[33] Importantly, all extracted RRs had been adjusted for physical activity. We provided  
18 uncertainty limits in the form of 95% CIs for the PAFs and the subsequent cost estimates.  
19 Finally, we subtracted a proportion of costs to account for the strong likelihood of double-  
20 counting due to co-morbidities.

21 However, the study was limited by the evidence available for sedentary behaviour and health  
22 outcomes. We included a non-European study[37] in order to estimate the prevalence of  
23 sedentary behaviour in lung cancer cases, which may not fully reflect a UK population.  
24 Individual studies included in the meta-analyses which were used in this analysis varied in  
25 their choice of cut-off values for each category, definition of sedentary behaviour, and in the  
26 questionnaire used. Crucially, six hours/day was the minimum median time spent in  
27 sedentary behaviour in the highest categories (supplementary file 3). Nevertheless,  
28 theoretically the definition used for the prevalence of sedentary behaviour should match the  
29 RR when calculating the PAF. We believe that since the minimum median sedentary time in  
30 the most sedentary class is 6 hours, and our definition of sedentary behaviour is spending at  
31 least six hours sedentary, that the RRs reported are reasonable estimations. Therefore, the  
32 PAFs are also reasonable estimations. We were further limited by self-reported data for

1 sedentary behaviour, which may have either underestimate or overestimate sedentary  
2 behaviour[34] and could subsequently bias the results in either direction.

3 Future research is still needed to elucidate the complex relationship between sedentary  
4 behaviour and health, and which of these are truly independent of physical activity.[23]  
5 Ideally, prospective studies could use a combined method of both accelerometry and  
6 behaviour logs, repeated over time, when measuring this behaviour. Consensus on how many  
7 hours/day of sedentary behaviour is harmful would be helpful in research, in line with the  
8 more specific guidelines for physical activity.[3]

9 Indirect costs that incorporate the financial burden on society, such as productivity losses to  
10 the workforce, can be very high. Physical inactivity was responsible for an estimated \$0.5  
11 billion (international dollars) outside of the healthcare setting in 2013 in the UK.[7] There are  
12 no known estimates for the wider societal costs of sedentary behaviour. Economic estimates  
13 will need to be updated as further evidence on sedentary behaviour emerges.

14 There are several barriers that cause a gap between evidence and practice. Evidence may be  
15 non-existent or arrive too late for policymakers. They may prefer uncomplicated papers and a  
16 wide range of evidence to inform their decisions.[40] We have been explicit about the  
17 strengths and weakness of this straightforward cost estimation for the benefit of other  
18 academics and policymakers. We hope that these results can be easily understood and  
19 synthesized with other evidence on sedentary behaviour. An economic case could be made  
20 for investment in reducing the prevalence of sedentary behaviour in the UK. These cost  
21 estimates can be compared with those of other risk factors in order to inform decision-making  
22 and prioritise preventative health programmes. Many individuals in the UK spend their  
23 leisure time in sedentary behaviour, but the workplace represents a significant proportion of  
24 unavoidable daily sitting time for many people. Measures should be taken to reduce sedentary  
25 behaviour with the aim of improving population health and reducing the financial burden to  
26 the health service.

27 This analysis presents the first estimate of direct healthcare costs due to prolonged sedentary  
28 behaviour in the UK. After adjustment for co-morbidities, diseases associated with prolonged  
29 sedentary behaviour cost the NHS £0.7 billion in 2016-17 costs. Furthermore, 48,024 deaths  
30 could have potentially been avoided in 2016 if prolonged sedentary behaviour in the UK was  
31 eliminated. It is hoped that these estimates will help policymakers prioritise resources to  
32 address a major public health issue.

## 1 **Contributors**

2 CO, FK, LH, and MT developed the research question. LH did the analysis, with  
3 methodological guidance from CO. HM provided statistical expertise. LH drafted the  
4 manuscript. All authors read and approved the final manuscript.

## 5 **Funding**

6 Leonie Heron is funded by a Department for the Economy (Northern Ireland) studentship.

7 The Corresponding Author has the right to grant on behalf of all authors and does grant on  
8 behalf of all authors, an exclusive licence (or non exclusive for government employees) on a  
9 worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be  
10 published in JECH and any other BMJ PGL products and sublicences such use and exploit all  
11 subsidiary rights, as set out in our licence  
12 (<http://group.bmj.com/products/journals/instructions-for-authors/licence-forms>).

13 Competing Interest: None declared.

## 14 **REFERENCES**

- 15 1. Dunstan DW, Healy GN, Sugiyama T, et al. “Too much sitting” and metabolic risk -  
16 Has modern technology caught up with us? *Eur Endocrinol.* 2009;5:29–33.
- 17 2. Tremblay MS, Aubert S, Barnes JD, et al. Sedentary Behavior Research Network  
18 (SBRN) – Terminology Consensus Project process and outcome. *Int J Behav Nutr*  
19 *Phys Act.* 2017;14(1):75.
- 20 3. UK Department of Health. *Physical Activity Guidelines [Internet]*. 2011. Available  
21 from: <https://www.gov.uk/government/publications/uk-physical-activity-guidelines>.  
22 (Accessed Jan 2018).
- 23 4. Patterson R, McNamara E, Tainio M, et al. Sedentary behaviour and risk of all-cause,  
24 cardiovascular and cancer mortality, and incident type 2 diabetes: a systematic review  
25 and dose response meta-analysis. *Eur J Epidemiol.* 2018;1:1–19.
- 26 5. NHS England. *Health Survey England 2012 [Internet]*. 2013. Available from:  
27 [https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-](https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/health-survey-for-england-2012)  
28 [england/health-survey-for-england-2012](https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/health-survey-for-england-2012) (Accessed Jan 2018).
- 29 6. 2018 Physical Activity Guidelines Advisory Committee. *2018 Physical Activity*  
30 *Guidelines Advisory Committee Scientific Report [Internet]*. Washington, DC; 2018.  
31 Available

- 1 from:[https://health.gov/paguidelines/secondedition/report/pdf/PAG\\_Advisory\\_Comm](https://health.gov/paguidelines/secondedition/report/pdf/PAG_Advisory_Comm)  
2 [itee\\_Report.pdf](https://health.gov/paguidelines/secondedition/report/pdf/PAG_Advisory_Comm) (Accessed Jul 2018).
- 3 7. Ding D, Lawson KD, Kolbe-Alexander TL, et al. The economic burden of physical  
4 inactivity: a global analysis of major non-communicable diseases. *Lancet*.  
5 2016;388(10051):1311–24.
- 6 8. Scarborough P, Bhatnagar P, Wickramasinghe KK, et al. The economic burden of ill  
7 health due to diet, physical inactivity, smoking, alcohol and obesity in the UK: An  
8 update to 2006-07 NHS costs. *J Public Health (Oxf)*. 2011;33(4):527-35
- 9 9. Lee I-M, Shiroma EJ, Lobelo F, et al. Impact of Physical Inactivity on the World’s  
10 Major Non-Communicable Diseases. *Lancet*. 2012;380(9838):219–29.
- 11 10. Stamatakis E, Pulsford RM, Brunner EJ, et al. Sitting behaviour is not associated with  
12 incident diabetes over 13 years: The Whitehall II cohort study. *Br J Sports Med*.  
13 2017;51(10):818–23.
- 14 11. Rockhill B, Newman B, Weinberg C. Use and misuse of population attributable  
15 fractions. *Am J Public Health*. 1998;88(1):15–9.
- 16 12. Greenland S. Interval estimation by simulation as an alternative to and extension of  
17 confidence intervals. *Int J Epidemiol*. 2004;33(6):1389–97.
- 18 13. NHS. 2012/13 *Programme Budgeting Data [Internet]*. 2018. Available from:  
19 <https://www.networks.nhs.uk/nhs-networks/health-investment-network/news/2012->  
20 [13-programme-budgeting-data-is-now-available](https://www.networks.nhs.uk/nhs-networks/health-investment-network/news/2012-) (accessed Jul 2018).
- 21 14. Statistics for Wales. *NHS Expenditure Programme Budgets, 2016-17 [Internet]*. 2018.  
22 Available from: <https://gov.wales/docs/statistics/2018/180417-nhs-expenditure->  
23 [programme-budgets-2016-17-en.pdf](https://gov.wales/docs/statistics/2018/180417-nhs-expenditure-) (Accessed Jun 2018).
- 24 15. Scottish Government. *Programme Budgeting in NHS Scotland [Internet]*. 2015.  
25 Available from: <https://www.gov.scot/Publications/2015/08/4735/0> (Accessed Jul  
26 2018).
- 27 16. Curtis, L. & Burns, A. (2017) Unit Costs of Health and Social Care 2017, Personal  
28 Social Services Research Unit, University of Kent, Canterbury.  
29 <https://doi.org/10.22024/UniKent/01.02/65559>.
- 30 17. Einarson TR, Acs A, Ludwig C, et al. Prevalence of cardiovascular disease in type 2  
31 diabetes : a systematic literature review of scientific evidence from across the world in  
32 2007 – 2017. *Cardiovasc Diabetol*. 2018;17(1):83.
- 33 18. Ekelund U, Steene-Johannessen J, Brown WJ, et al. Does physical activity attenuate,

- 1 or even eliminate, the detrimental association of sitting time with mortality? A  
2 harmonised meta-analysis of data from more than 1 million men and women. *Lancet*.  
3 2016;388(10051):1302–10.
- 4 19. Biswas A, Oh PI, Faulkner GE, et al. Sedentary time and its association with risk for  
5 disease incidence, mortality, and hospitalization in adults a systematic review and  
6 meta-analysis. *Ann Intern Med*. 2015;162(2):123–32.
- 7 20. Dunstan DW, Salmon J, Owen N, et al. Associations of TV viewing and physical  
8 activity with the metabolic syndrome in Australian adults. *Diabetologia*.  
9 2005;48(11):2254–61.
- 10 21. Pandey A, Salahuddin U, Garg S, et al. Continuous Dose-Response Association  
11 Between Sedentary Time and Risk for Cardiovascular Disease. *JAMA Cardiol*.  
12 2016;1(5):575-83.
- 13 22. George ES, Rosenkranz RR, Kolt GS. Chronic disease and sitting time in middle-aged  
14 Australian males: Findings from the 45 and Up Study. *Int J Behav Nutr Phys Act*.  
15 2013;10:20.
- 16 23. Wijndaele K, Brage S, Besson H, et al. Television viewing time independently  
17 predicts all-cause and cardiovascular mortality: The EPIC Norfolk study. *Int J*  
18 *Epidemiol*. 2011;40(1):150–9.
- 19 24. León-Muñoz LM, Martínez-Gómez D, Balboa-Castillo T, et al. Continued  
20 sedentariness, change in sitting time, and mortality in older adults. *Med Sci Sports*  
21 *Exerc*. 2013;45(8):1501–7.
- 22 25. Shen D, Mao W, Liu T, et al. Sedentary behavior and incident cancer: A meta-  
23 analysis of prospective studies. *PLoS One*. 2014;9(8):e105709.
- 24 26. Nuffield Trust, The Health Foundation, and The King's Fund (2016) *The Autumn*  
25 *Statement: joint statement on health and social care [Internet]*. Available from:  
26 [https://www.nuffieldtrust.org.uk/resource/the-autumn-statement-joint-statement-on-](https://www.nuffieldtrust.org.uk/resource/the-autumn-statement-joint-statement-on-health-and-social-care)  
27 [health-and-social-care](https://www.nuffieldtrust.org.uk/resource/the-autumn-statement-joint-statement-on-health-and-social-care) (Accessed Jul 2018).
- 28 27. Audit Scotland. *NHS in Scotland 2017 [Internet]*. 2017. Available from:  
29 [http://www.audit-](http://www.audit-scotland.gov.uk/uploads/docs/report/2017/nr_171026_nhs_overview.pdf)  
30 [scotland.gov.uk/uploads/docs/report/2017/nr\\_171026\\_nhs\\_overview.pdf](http://www.audit-scotland.gov.uk/uploads/docs/report/2017/nr_171026_nhs_overview.pdf) (Accessed  
31 Jul 2018).
- 32 28. Sarwar N, Gao P, Seshasai SR, et al. Diabetes mellitus, fasting blood glucose  
33 concentration, and risk of vascular disease : a collaborative meta-analysis of 102



- 1 prospective studies. *Lancet*. 2010;375(9733):2215–22.
- 2 29. British Heart Foundation. *Heart and Circulatory Diseases Statistics 2017 [Internet]*.  
3 2017. Available from: <https://www.bhf.org.uk/what-we-do/our-research/heart->  
4 [statistics/heart-statistics-publications/cardiovascular-disease-statistics-2017](https://www.bhf.org.uk/what-we-do/our-research/heart-statistics/heart-statistics-publications/cardiovascular-disease-statistics-2017) (Accessed  
5 Oct 2018)
- 6 30. Rezende LFM de, Rodrigues Lopes M, Rey-López JP, et al. Sedentary Behavior and  
7 Health Outcomes: An Overview of Systematic Reviews. *PLoS One*.  
8 2014;9(8):e105620.
- 9 31. Werneck AO, Oyeyemi AL, Szwarcwald CL, et al. Associations between TV viewing  
10 and depressive symptoms among 60,202 Brazilian adults: The Brazilian national  
11 health survey. *J Affect Disord*. 2018;236:23-30.
- 12 32. Hoare E, Milton K, Foster C, et al. The associations between sedentary behaviour and  
13 mental health among adolescents: A systematic review. *Int J Behav Nutr Phys Act*.  
14 2016 Oct; 13(1):108.
- 15 33. Ding D, Kolbe-Alexander T, Nguyen B, et al. The economic burden of physical  
16 inactivity: a systematic review and critical appraisal. *Br J Sports Med*. 2017  
17 Oct;51(19):1392–409.
- 18 34. Healy GN, Clark BK, Winkler EAH, et al. Measurement of adults' sedentary time in  
19 population-based studies. *Am J Prev Med*. 2011;41(2):216-227.
- 20 35. Bjork Petersen C, Bauman A, Grønbaek M, et al. Total sitting time and risk of  
21 myocardial infarction, coronary heart disease and all-cause mortality in a prospective  
22 cohort of Danish adults. *Int J Behav Nutr Phys Act*. 2014;11(1):13.
- 23 36. van der Ploeg H, Chey T, Korda R, et al. Sitting time and all cause mortality risk in  
24 222,497 Australian adults. *Arch Intern Med*. 2012;172(6):494-500.
- 25 37. Ukawa S, Tamakoshi A, Wakai K, et al. Prospective cohort study on television  
26 viewing time and incidence of lung cancer: Findings from the Japan Collaborative  
27 Cohort Study. *Cancer Causes Control*. 2013;24(8):1547–53.
- 28 38. Simons CCJM, Hughes LAE, Van Engeland M, et al. Physical activity, Occupational  
29 sitting time, and colorectal cancer risk in the netherlands cohort study. *Am J*  
30 *Epidemiol*. 2013;177(6):514–30.
- 31 39. Gierach GL, Chang S-C, Brinton LA, et al. Physical activity, sedentary behavior, and  
32 endometrial cancer risk in the NIH-AARP Diet and Health Study. *Int J Cancer*.  
33 2009;124(9):2139–47.

1 40. Whitty, C. J. M. What makes an academic paper useful for health policy? *BMC Med.*  
2 2015;13(1):301.

3

4

5

6

7