



**QUEEN'S
UNIVERSITY
BELFAST**

Practitioners' experiences of using blended models within family support: A proof of concept study involving Cognitive-Behavioural Therapy (CBT), Multisystemic Therapy (MST) and Incredible Years (IY) interventions

Walsh, C., & Best, P. (2019). Practitioners' experiences of using blended models within family support: A proof of concept study involving Cognitive-Behavioural Therapy (CBT), Multisystemic Therapy (MST) and Incredible Years (IY) interventions. *Journal of Family Social Work*. <https://doi.org/10.1080/10522158.2019.1616240>

Published in:

Journal of Family Social Work

Document Version:

Peer reviewed version

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2019 Taylor & Francis. This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Practitioners' experiences of using blended models within family support: A proof of concept study involving Cognitive Behavioural Therapy (CBT), Multisystemic Therapy (MST) and Incredible Years interventions

Introduction

Since the emergence of the violent sectarian conflict in Northern Ireland during the 1960's - commonly referred to as the '*Troubles*' and generally regarded as Europe's longest running violent conflict (Lynch & Joyce, 2018), approximately 3,700 people lost their lives, about 40,000 people sustained physical injuries (McAlinden & Dwyer, 2015) and countless others suffered psychological trauma. Some estimates suggest that 80% of the population knew someone who had been killed or injured during the *Troubles* (Breen-Smith, 2012). Crimes against person and property permeated all aspects of life. Routine tasks such as shopping, travelling and socialising were often affected by conflict-related incidents such as police check-points, bomb scares, car hijacking, protests, parades and mass rioting and this had a significant psychosocial impact on the whole population (McAlister, Scraton & Haydon, 2014). In the context of this, evidence has consistently demonstrated that families living in Northern Ireland experience multiple adversities, particularly in those areas that were characterised as having been exposed to conflict and social deprivation (Davidson, Bunting & Webb, 2012; Bunting & Lazenbatt, 2015). Some of these families, were often referred for, or offered additional supports for their family.

Family support is a familiar term that has been used widely over recent years and yet is a contested concept (Cameron et al., 2007; Devaney & Dolan, 2014) which varies according to context, purpose and underpinning values within specific models (Churchill & Sen, 2016). Rather than illustrating a particular type of intervention, 'family support' refers to a range of interventions or strategies that are delivered across many disciplines to address a well-defined problem (McKeown, 2000) within specific family systems. The emergence of the concept as well as significant increase in interventions defined as 'family support' models in many ways reflects social trends fuelled by concerns around welfare, family functioning, mental health, violence and inequality (Bonoli, 2013). Over recent decades there has been an exponential growth in social, psychological and therapeutic specialisms (Ovretviet, Hansson & Brommels, 2010) in response to the increased complexity of needs presented by service users (Lemmens, Molema, Versnel, Baan & De Bruin, et al., 2015). By some accounts, the figures for primary care centres in the US who also offered behavioural interventions rose from 42%

to 75% between 2000 and 2011 (Padwa et al., 2016) suggestive of a rise in crisis experienced by families and members within those families.

Adversity affects many families but a minority find themselves in crisis following complex and sometimes cumulative adversities such as abuse, poverty, incarceration and ill health (Lee et al., 2017). Within a social ecological framework, these issues impact upon both individuals and wider family members. Often one issue exacerbates other pre-existing issues whilst also contributing to new and more complex challenges. Whilst an array of social programmes and human service interventions are available, accessing those who need them and retaining them in support can be difficult.

The emergence of evidence-based practices (EBPs) and implementation science (Fixsen et al., 2005) provided opportunities for human services to understand what works in improving outcomes as well as the mechanisms to successfully (and sustainably) replicate what works (Olsson, 2010; Wiggins et al., 2010). This includes practices to both identify and retain those who are most in need of support so that the potential impact of evidence based practices can be fully achieved.

Depending on the criteria applied and how 'evidence' is defined, there are between 50 and 200 evidence-based programmes (Little, 2010) currently available in the areas of parenting (Furlong et al., 2012), educational attainment (Valdebenito et al., 2018) and violence prevention (Tolan, 2013). In the UK there has been a drive to embed these models into routine practice. Between 2008 and 2011, a UK government sponsored implementation strategy called the Parenting Early Intervention Programme (PEIP) was designed to help facilitate the replication of a range of evidence based parenting models across 150 local government areas in England (Lindsay et al., 2013). And between 2012 and 2017, the Big Lottery, the UK's largest charitable grant making body, funded the '*Realising Ambition*' Programme aimed at improving the evidence base of what works with preventing youth offending and was supported by a financial pot of \$32m (Catch 22, 2014). In Northern Ireland too, there has been an increasing focus on integrating evidence into practice and this was built into the previous programme for government (OFMDFM, 2011). Commitments have been made to deliver a range of measures to tackle poverty and social exclusion through a 'Delivering Social Change (DSC) Framework' itself funded to the tune of \$150M (CYPSP, 2014). The programme was underpinned by a focus on early intervention, the use of best evidence to inform practice. In an era of increased austerity, successfully implementing such interventions within the context of family support also had the additional value of saving money whilst improving outcomes (Odom, 2009).

However, these are often implemented either sequentially (as one treatment proves ineffective) or in parallel but also in isolation from the other with needs assessment, clinical decision making and reviews taking place.

Given that families' needs are often complex and co-occurring, with different family members requiring different types and levels of support simultaneously, single, specific and targeted programmes may not comprehensively address the complex needs experienced by some families (Guastaferrero et al., 2017). In a social, economic and political climate that has demonstrated significant interest and investment in evidence based interventions that seek to improve family outcomes (Batty & Flint, 2012), multiple component interventions that address the distinct needs of family members have been significantly under-evaluated (Aarons, Hulburt & Horwitz, 2011). Few studies have actively sought to explore whether a combination or blending of evidence based models and practices enhance outcomes that families experience, particularly when they experience multiple adversities concurrently. Additionally, there are even fewer studies that have sought to explore the implementation challenges with this type of novel blended approach.

For the purpose of this paper, blended models (Walsh & Doherty, 2016) have been defined as the concurrent implementation of more than one evidence based model or practice facilitated by practitioners within the same programme but to address distinct needs. Using our definition of blended models, family therapy followed by cognitive behavioural therapy (or vice-versa) is not a blended model. Likewise, an individual within a family receiving two interventions, neither of which is integrated or coordinated is not a blended model. In this conceptualisation, a blended model is one in which a family receive more than one intervention which is integrated, coordinated and is also intended to address more than one issue (e.g. CBT for depression and MST for a reduction in violent behaviour).

Given the dearth of primary studies exploring the viability and acceptability of this approach, the aim of this paper is to explore the feasibility of blended approaches within complex family support programmes as a means of addressing these complex and concurrent needs and retaining families in interventions most likely to achieve the best possible outcomes for them. The medical research council provide some guidance around evaluating complex interventions. They suggest that prior to an expensive process evaluation or outcome study, an early stage review could be considered with the aim of answering 'would this be possible to use' (Craig et al., 2008). This proof of concept study aims to highlight the potential feasibility of blending different interventions to meet those concurrent and complex needs that families

experience. Proof of concept (PoC) is a realization of a certain method or idea in order to demonstrate its feasibility or a demonstration in principle with the aim of verifying that some concept or theory has practical potential (Ungar et al., 2017). The studies aim to explore the feasibility of a novel approach (Kush et al. 2007). In health care, PoC studies have been used to understand challenges and reduce inefficiencies in clinical trials serving to prepare the ground for full outcome studies. Within prevention, early intervention and parenting initiatives, PoC studies are increasingly being used to highlight gaps in current practice, the added value of novel approaches and the mechanics of those new practices (Rabinowitz et al., 2013). But given their primary aim is to demonstrate that a programme design has practical potential rather than generate conclusive evidence about their relative efficacy, further studies to determine effectiveness are often required (Cobham et al., 2017).

In this study, a large voluntary organisation implemented three evidence based models and practices within the same service environment.

For the current paper the focus is on the concurrent implementation of three evidence based models and practices implemented concurrently within a family support service: Multisystemic Therapy (MST), Incredible Years (IY), and Cognitive Behaviour Therapy (CBT).

In 2014, as part of a coordinated approach in reducing family adversity and improving outcomes for children and young people in Northern Ireland, the Extern Organization, a large NGO supporting children, young people and families with complex needs, piloted a new model of practice called the Intensive Family Support Service (IFSS). The pilot sought to blend together a range of evidence based, evidence informed and basic needs practices under one roof and to be delivered by a skilled but also multi-disciplinary team. Three evidence based models (Multisystemic Therapy-MST, Cognitive Behavioural Therapy-CBT and Incredible Years) were implemented (see table 1). In addition to the evidence based interventions, a range of evidence informed practices such as person centred counselling and motivational interviewing were implemented.

The aims of the pilot were to engage families who had been known to multiple services but were considered 'hard to reach' either because they had failed to complete interventions or presented with challenges following previous support; to help prevent placement breakdown; to reduce offending and other risk taking behaviours; to improve mental health issues; and to increase academic engagement and attainment.

Table 1:
Overview of EBPs within IFSS

Model or practice	Target group	Desired outcome	N=
Multisystemic Therapy (MST)	Parents of young people 12-18	Reduced anti-social behaviour and improved family functioning	56
Cognitive Behavioural Therapy (CBT)	Young people aged 10-18	Improved mental health symptomatology	32
Incredible Years	Parents of young children	Improved family functioning	12

A range of evidence based practices and programmes were available to families and within the IFSS programme. Three of these are covered in this article: Multisystemic Therapy (MST); Incredible Years; and Cognitive Behavioral Therapy (CBT).

Incredible Years supports parents of younger children to improve parenting practices and family function; MST supports parents to address adolescent risk factors associated with family breakdown; and CBT provides individual therapeutic support to those with mental health difficulties. It has been proposed that the combined delivery of these three models are both complementary and the potential for improved outcomes is potentially enhanced through the novel blending.

MST is an intensive community based model that uses evidence-based interventions to address problem behaviours and attempts to mitigate the risks associated with out of home placement by placing the family at the centre of all elements of the work. Since being developed in the 1980's, MST provision has extended beyond the USA where it was first developed and into 15 counties across more than 200 teams (Olsson, 2010). Nine treatment outcome studies (including 3 controlled trials) have been published and for seven of these follow-up data from 1-4 years have been reported to be effective for those who completed the treatment programmes (Carr, 2005).

In addition to MST, it was decided to replicate Incredible Years, an evidence based programme that involves parenting and teaching practices that have been shown to positively and effectively reduce conduct problems and strengthen children's social and emotional

competence. The model is guided by the behavioural principles of operant conditioning and social learning, and aim to strengthen parent and teacher skills respectively and ultimately improve child outcomes. The National Institute for Health and Clinical Excellence has published guidelines for parenting programmes for the treatment of conduct disorders and when published, Incredible Years was only 1 of 2 models that met the criteria (NICE, 2006). The programme, aimed at children aged 3 to 12 years, is founded on Social Learning Theory (Bandura and consists of at least 12 weekly, two-hour group sessions delivered by skilled practitioners. Overall, the Incredible Years Programme aims to: promote positive parenting and improve parent-child relationships, reduce critical and physical discipline and increase the use of positive strategies and help parents to identify social learning theory principles for managing behaviour and improve home-school relationships (McDaniel et al., 2009). Reviews of the efficacy of a number of parent training prevention programmes and treatment programmes have shown these programmes to be promising for changing maladaptive parent and child behaviour in younger children (Webster-Stratton & Reid, 2017) and a study in Ireland found that benefits were experienced by families both in the short term and longer term (McGilloway et al., 2009). An additional study (McGilloway, 2012), investigating the impact of incredible years on parenting, educational engagement and childhood functioning found significant reductions in childhood conduct disorder, improved parenting skills and improved academic engagement.

Cognitive approaches to therapy first emerged in the work of Alfred Adler, Albert Ellis and Aaron T. Beck (Beck, 1995). Beck et al (1979: 3) note that cognitive therapy is an “active, directive, time-limited and structured approach used to treat a variety of psychiatric disorders”. The therapeutic techniques used within Beckian CBT are designed to identify, test and ultimately correct the dysfunctional schemas that underlie the client’s cognitions. CBT is based on the idea that how we think (cognition), how we feel (emotion) and how we act (behavior) all interact together. Specifically, our thoughts determine our feelings and our behaviour (Beck, 2011). Cognitive behavioural approaches use components of Cognitive Behavioural Therapy and are well established psycho-social therapeutic practices, particularly for ‘at risk’ populations. Largely developed in clinical settings and adapted for a range of populations including those with anxiety, depression, and those who have experienced trauma (Armeliu & Andreassen 2007), more recently the approaches have been integrated into preventative practices with the aim of reducing risk taking and/or offending behaviour (Littell & Forsyth, 2005). The practices have been widely used in youth justice and probation environments as a means of addressing offending at an early stage as well as reducing the likelihood of reoffending by engaging clients or service users in reasoning and/or problem solving skills (Joughin, 2006)

The rationale to include these three models is related to both the function and structure of them. All three are evidence based models widely used. All three are implemented for well-defined reasons, with well-defined desired outcomes. A critical feature is that the target populations are also well defined. For instance, MST would not be implemented to address parenting concerns regarding a toddler, but Incredible Years would. Conversely, Incredible Years would not be implemented to prevent further offending by a 17 year old but MST would.

The aim of this proof of concept study is to compare and contrast the experiences of practitioners implementing both blended and non-blended approaches to family support.

The objectives of this study are:

- To explore the experience and perceptions of practitioners who implemented the blended model.
- To explore the novelty of blended approaches
- To examine the extent to which families were engaged in and were retained in this programme;
- To compare the outcomes for families who received blended and non-blended approaches;

Methods

Data collection

This exploratory study used qualitative methods as a primary data collection tool. This was supplemented by pre-existing quantitative data gathered by the organisation (See below for further details). A mixed methods approach (Bronstein et al., 2013) was employed in the study in which family data (reasons for referral, amount of support received and whether families were engaged in blended or non-blended support), and staff demographic data dovetailed (gender, no. of year's experience, time with the agency, primary discipline), (Bryman, 1998) into semi-structured interviews in order to increase the accuracy of the findings (Creswell, 2009; Moran-Ellis et al., 2013).

The sample were practitioners from the project and included primary data from thirteen individual interviews as well as secondary data from 41 staff.

Practitioner demographics

A total of forty one staff completed the initial questionnaires representing 91% of the total staff team. The returns represented the various disciplines (Youth work n=5; Social work n=15; Psychology n=6; Counselling-n=2; Play and art therapy n=3; Health and social care n=4; Combined social work and psychology n=6) (see table 2).

Table 2:

Practitioner demographics

	M	%	N
Age	25		
Gender	Male	14	6
	Female	86	35
	Total	100	41
Year's experience	3.5		
Months with the organisation	14		
Discipline	Youth work	12	5
	Social work	36	15
	Psychology	14	6
	Counselling	4	2
	Play therapy	7	3
	Social care	9	4
	Combined	14	6

Quantitative data collection tools

The quantitative data included existing information which was designed and collected by the project delivery staff. These were non-standardised measures designed in line with the operational requirements of the project. Data was collected from families who engaged in the pilot was collated and analysed. This included access to referral forms, case notes and

closure reports. Referral forms were received from a range of referral agents on a standardised programme template which outlined reasons for referrals and contact information for each family. Following receipt of this, the family were allocated a project worker whose initial role it was to undertake an assessment for service. This assessment aimed to understand the strengths and needs of each family, what they wished to receive support with and negotiate how best the programme could serve them. Following this assessment, the project worker, the family and the service manager agreed a family support plan and what that plan would consist of. At this stage, objective and measurable desired outcomes were agreed (e.g. X will return to education and attend daily, consistently for four weeks). During period reviews and at the end of each intervention, these overarching goals were reviewed. Following intervention, a determination around the extent to which those goals were met, partially met, or not met were agreed between the family, the project worker, the programme manager and the referral agent.

Not all families received the same type of intensity of support. Some received single interventions and others received blended interventions. Using this assessment data, comparisons were then made between those who received a blended model of support and those who did not.

Following allocation, work which was undertaken with each family was recorded following each session using routinely collected session template. The data included categorical (such as reasons for referral; models engaged in; completion status; outcomes achieved). Additionally, continuous data was collected (hours of support provided).

In addition to family profile data, all staff (n=41) were invited to engage in the research and complete an initial questionnaire. This questionnaire included demographic details (gender & age); and professional details (discipline, level of education, years of experience and time with the agency).

Qualitative data collection tools

All key work staff within the programme (n=41) were invited to engage in the interview process. Fourteen staff consented and were interviewed. A semi-structured interview schedule was developed (Creswell & Clarke, 2007). Each interview lasted between thirty five and forty five minutes. Interviews were facilitated by two trained investigators with experience in undertaking qualitative studies neither of whom were directly associated with the interventions being studied. Notes were taken by one of the investigators verbatim whilst the other conducted the

interview. Following the interview, participants were given an opportunity to review notes taken during the interview and asked to confirm if they accurately reflected what they said. Following each interview, a de-brief was undertaken. Using more than one interviewer enabled greater access to participants and their experiences, greater understanding of the responses, gaining the trust of the participants and establishing a rapport more quickly and the opportunity to take more detailed notes (Fontana & Fay, 2005). Critically this allowed for a more informed reflection of each interview (Matteson & Lincoln, 2009). Each interview followed a standard process with key workers asked to confirm that they voluntarily opted into the study; that they were aware of the purpose; and that they understood how the data collected would be used. Following this confirmation key workers were then asked a series of open questions related to their experiences within the project (their motivation for joining the project; prior experience of evidence based practices; perceived level of support in relation to implementing the work; previous training/discipline/qualifications; challenges with implementing the model; and recommendations moving forward were all explored.

All of the interviews included staff who had delivered a blended approach.

Ethical considerations

Data was collected as part of a service evaluation. As such, no formal ethical approval was sought (NRES, 2009). Despite this, ethical issues were addressed. Participants were provided with information about the study and an opportunity to opt in. The information sheet outlined what would be involved if they chose to be involved and how the data would be used. They were provided with one week between receiving the initial information and being provided with a consent form to engage. No paper copies of personal information was held by the researchers. Confidentiality of participant's personal data was ensured by not collecting individual names on the enrolment or measurement forms. All data will be stored in a pseudonymised manner. During enrolment participants were issued an identification code. Consent forms were held securely on the premises in a locked filing cabinet. For qualitative data, transcripts were anonymised (names, places, services) and no identifying information was used in dissemination.

Data Analysis

The analysis was undertaken sequentially. Secondary analysis was undertaken using existing data routinely collected within the project. This data was analysed prior to the interviews and helped inform that interview schedule that was undertaken with project staff. That quantitative data was analysed using SPSS Version 22 (SPSS, 2013). Independent samples t-tests were used to compare means between variables of interest. Crosstabulations were used to compare categorical data (reasons for referral, engagement in the three evidence based models and outcomes attained) and Chi-square tests of independence were used to explore statistical relationships between them.

Qualitative data was captured during the semi-structured interviews. Notes were taken by the researcher during the interview, reviewed immediately following the interview, typed and uploaded onto the NVivo version 10 qualitative software programme. Each interview was coded within NVivo, nodes were created and a process of thematic analysis was facilitated to interrogate the data in order to explore themes emerging from the interviews. Thematic analysis is a method for identifying, analysing, and reporting patterns (themes) within the data (Braun & Clarke, 2006). Themes that emerged from the informants stories were pieced together to form a comprehensive picture of the collective experience. Given the dearth of empirical or theoretical evidence for this type of approach, themes emerged inductively (Miles & Huberman, 1994). That is to say, the themes were data driven and not confined to pre-specified coding systems.

Results

Quantitative results

Reasons for referral

Families were referred for a number of complex psychosocial issues (see table 3) and all families were being supported with at least 2 of these complex issues simultaneously at the point of referral. In fact, 89% of families were being supported with 4 or more issues.

Table 3

Overview of reasons for referral

Reason for referral	% of total families referred	N=	% of families referred blended	N=	% of families referred-non-blended	N=
<i>Safety, risk and aggression</i>	66	104	84	26	61	78
<i>Education</i>	74	116	74	23	73	93
<i>Mental health</i>	46	73	81	25	38	48
<i>Substance use</i>	25	39	19	6	26	33
<i>Health</i>	31	49	29	9	32	40
<i>Finances</i>	25	39	7	2	29	37
<i>Home conditions</i>	33	52	13	4	38	48
<i>Employment</i>	14	22	0	0	100	22

The intensity with which families were supported varied depending on need at various points in the programme journey. The average number of face to face hours families were afforded was 11.5 per month. However, the range was a huge 56 hours per month and the standard deviation was more than 7 hours per month.

Blended approaches

Not all families received the same intensity or type of support. Some families received a blended approaches and others did not. Families were supported from staff from a range of professional backgrounds. Interestingly, only those from youth work (3%), social work (29%) psychology (10%) or combined social work/psychology (58%) backgrounds implemented a blended model approach.

Families who were referred to the programme for issues around safety, risk and/or aggression (34%) were more likely than other families to undertake risk management work and safety planning. Those families were also more likely to achieve positive outcomes (X^2 2, $n=86=21.29$, $p=.00$, $\phi=.50$). Interestingly 66% ($n=21$) of those who were not referred with safety concerns also undertook risk management work suggesting that more families experienced issues related to risk than statutory services were aware of at the point of referral. Families referred with finance ($n=29$) and/or home conditions issues were more likely to engage in more practical supports (80%, $n=23$).

A range of evidence based practices and programmes were available to families. In total, 68% of families engaged in at least one evidence based programme or practice.

There was a strong relationship between use of evidence based practices and outcomes being met ($X^2, 2, n=158= 26, p=.00, \phi=.41$) suggesting that those with more complex needs were better served by evidence based models or approaches. 70% (n=57) of those who used an evidence based practice also fully met their target goals compared with 38% (n=24) who did not use an EBP fully met their goals. But then families with basic needs issues (such as finances or employment were less likely

38% (N=31) of those who engaged in an EBP had access to more than 1 evidence based practice or intervention. Within this group, families had access to an increased number of hours (m=16.13, sd=6.46) compared to families who did not engage in a blended intervention (m=10.43, sd=6.82). This may partly be because families who engaged in a blended approach were also those who more likely to experience multiple and higher risk issues (See table 4). The more intensive nature of the support was statistically significant ($t(156)=4.21, p<.01$).

Table 4:

Overview of reasons for referral and outcomes by blended/not blended

Population	Reason for referral	%	Outcome (% fully met)	Complete?
Blended	Safety, risk or aggression	84% (n=26)	81% (n=21)	100% (n=31)
	Mental Health	81% (n=25)	80% (n=20)	100% (n=25)
	Employment	0% (n=0)	0% (n=0)	0% (n=0)
	Finances	7% (n=2)	100% (n=2)	100% (n=2)
Non-blended	Safety, risk or aggression	61% (n=78)	41% (n=32)	81% (n=63)
	Mental Health	38% (n=48)	35% (n=17)	83% (n=40)
	Employment	100% (n=22)	46% (n=10)	96% (n=21)
	Finances	29% (n=37)	57% (n=21)	97% (n=36)

And whilst it appears that those who engaged in multiple EBPs were families who were more likely to experience more significant and risky challenges, they were also more likely to

achieve their outcomes. 81% of those who engaged in a blended approach (n=25) had their goals fully met compared with only 48% (n=61) of those who did not engage in a blended approach. Likewise, only 3% (n=1) of those within the blended approach met none of their goals whilst 28% (n=36) of those who did not engage in a blended approach met none of their goals. These were at the point of statistical significance (χ^2 , 2, n=158= 12.22, p=.002). When specific reasons for referral are explored in more detail, it seems that families with complex and risky issues who engaged in blended approaches were more likely to achieve their outcomes compared with non-blended families

In addition to achieving positive outcomes, those who engaged in a blended approach were more likely to fully complete the programme than those who engaged in a non-blended intervention. 100% (n=31) of families who engaged in blended approaches fully completed the programme compared to 87% of those who did not (χ^2 , 1, n=158= 4.35, p=.04).

Qualitative results

Benefits of blended approaches

It was clear from interviews with staff that the complexities families face were not underestimated.

Some of my families are so vulnerable, I would be better moving in with them. I'm not joking. It's not always the case but there are some families I could be with in the morning, on the phone to them in the afternoon and back out with them in the evening. That could be the pattern for weeks depending on what's happening. But then as time goes on, you expect to see intensity reduce and intervention approaches change (Social Worker 2: 1.5 years' experience)

The intensity of support was echoed during all interviews. Practitioners felt that a thorough assessment enabled them to make sense of the variety of issues affecting families and prioritise in some way the challenges being presented. Most staff believed that given the complexities, single responses were often not what was required.

Families don't come to IFSS because they have a few issues. They usually get referred in after years, maybe generations of support from services. They usually have loads of issues and they take time to methodically address all of these. The benefit of IFSS is we can target multiple things at once. We help coordinate services. We are in the family's homes, we see what they see and we can provide a birds' eye perspective on what is going on. That means we can target support where it needs to go to get the biggest bang for our buck! (Youth Worker 1: 3.5 years' experience)

And yet, despite the complexities that practitioners described in great detail, many of the staff reported seeing significant change taking place with their families. Practitioners reported something unique about this type of blended approach. Even those more seasoned practitioners with experience in a variety of other support services indicated that the blended approach was contributing to change that was tangible.

I've been involved in a range of projects and to be honest, I've seen and heard about many of the same families coming through IFSS. There's families that have been through loads of different programmes. But something different has happened here. I know one family who's kids are now off the child protection register. That says loads. It says that they're in a place to manage things themselves after years of support. (Psychologist 3: 2 years' experience)

Having a variety of interventions that could be implemented with different members of a family to address their specific needs was seen not only as a resource for the family but also for the practitioner. So often, practitioners described uncertainty about what to implement, how and when. The combination of evidence based and manualised approaches enabled practitioners to provide support in an evidence based way with the understanding that these interventions have been rigorously tested.

I think because we have some many resources at our disposal, you can draw on one model for this and another for that so you don't need loads of different services as long as you know how to prioritise (Psychologist 1: 3 years' experience).

This model incorporates all family members and so isn't solely behaviour modification work. Because there are so many issues, we have a menu of work that is growing. We can draw on practical expertise and evidence based theoretical interventions sometimes at the same time. (Counsellor 1: 2 years' experience)

Implementation Challenges with blended approaches

Implementation can be defined as a series of measurable steps towards full and competent delivery. Within this process of implementation, different actors play different but complimentary roles. Whilst the focus of service evaluations is often on the competence and delivery of practitioners, during interviews, staff were keenly aware of the role that the organisation had to achieve implementation of this type of approach

The managers here have invested a lot in us as staff. There is loads of training that we need to take part in. you see, if you are delivering Incredible Years, you need to know how to deliver it. So you need trained in it. It's the same with other types of intervention.

In addition to the resource implications (time and financial), practitioners also described the role that the organisation had in other areas of implementation

Because we deliver evidence based models, there is a strong focus on fidelity. You need to deliver the model as it was intended to be delivered. So there's a lot of systems in place that the organisation had to develop and then stay on top of. They monitor how close we are delivering to the model and give us feedback (Social worker 1: 1 years' experience)

Organisational systems and supports appeared to be critical for practitioners to engage in a blended approach. For others however, regardless of the type of support provided and training afforded, some staff struggled to engage in the blended approaches.

They [the organisation] expect us to do everything. I think this is a way of cutting down on costs and just expecting us to do all the work that 3 or 4 organisations would have done before (Youth worker 1: 5 years' experience)

And whilst this was echoed several times, others had a different perspective on reducing the amount of service providers involved with families

My families don't know where they are from day to day, who's coming out and even when they're out, it's a struggle for the families to remember what their role is. They need one service that can do a whole lot of things for them. You can build up a relationship and they know where they're at. (Social worker 3: 7 years' experience)

But despite the opportunities with blended intervention, for some, the combination of interventions and approaches created a more complex working environment. This was particularly the case for staff who had not previously implemented a manualised model and for those who had concerns around prioritising what to implement and when. There are some indications that particular disciplines are more likely to be open to evidence based practice in general and this may in part be explained by their exposure to EBP during initial professional training

Using evidence based models wasn't that new to me. I covered it in uni and then when I was on placement, we got to implement a model called Strengthening Families. (Social worker 1: 1 years' experience)

But whether practitioners felt the approach was useful or not, there was a general consensus that too little information was available to them on the mechanics of implementation. There were real concerns around how decisions were taken and the consistency around those decisions.

I look to my manager for some guidance on what a family should receive. We have so much here but that can be overwhelming (Social worker 3: 7 years' experience)

Indeed, it appeared that the programme lacked a coherence around decision making process, particularly when it came to which family would receive which type of service and which family would engage in which component/s of the blended model. In the absence of a coherent framework, preference filled the vacuum.

There could be better guidance around what options to choose to handle certain difficulties. I think it depends on the staff member and the team they are in at the minute what options are available and what options are suggested (Psychologist 3: 2 years' experience)

Discussion

The viability of the blended model

A central design feature of the IFSS service were the range of supports that families had access to- a menu of strategies, supports and interventions that were blended within a complex arrangement of internal and external relationships to meet the needs of individual families. Interventions that were actioned when specific issues presented. These issues only became visible as the relationship developed. This suggests that some adversities were often unknown to statutory agencies but when made aware, IFSS keyworkers were able to implement appropriate interventions. This adds weight to the benefit families achieved by engaging in an environment where a menu of interventions is available.

There exists a dearth of information related to outcomes of blended models, their frequency and their impact. Whilst other models such as Integrated Care provide some empirical basis for the development of blended programme designs, there are clear distinctions. For instance, whilst integrated approaches seek to improve coordination, this is often across different sites and operating structures. Within the IFSS model, the blended design assimilates different approaches within one site and one operating structure. In addition, integrated approaches involve the joining of disciplines as a conscious and active effort to share best practice. In contrast, whilst the IFSS model did actively seek to recruit staff from across various disciplines, the blended approach requires that staff, from whatever sector, fully implement a range of evidence based practices and interventions.

Findings from the quantitative analysis appears to show a link between the use of a blended approach and improved outcomes for children, young people and families and despite limitations with the measures used, there are indications that this novel design did contribute

towards improved outcomes for families with complex needs and multiple risks factors. As noted, despite the array of support services available to families, not all families received the same level or categories of support. Interestingly, not every family needed an evidence based programme to achieve their desired outcomes. This suggests that not every family would receive an additional benefit from a blended approach but for others, with more complex needs, a blended approach could contribute towards enhanced outcomes. However, further investigation is needed.

The challenges

Given the multiple adversities experienced by families within IFSS, greater understanding is needed around how, why and when to use particular strategies and/or an evidence based practice or programme would greatly benefit staff in their decision making. Decision making processes were overwhelmingly unclear and inconsistent. However, during the interviews it was clear that some key considerations around which intervention or strategy to use included:

- Is there any immediate risk to family, community or worker?
- Based on the up to date assessment of need, what are the immediate issues facing the family?
- Based on this up to date assessment of need, does the intensity of support need to be increased/decreased?
- How do these priorities relate to the short and longer term family goals?
- Is there a need for revision of goals?
- Based on these identified priorities, what model, intervention or strategy is best suited and proven to address this identified issues or need?
- Is there consensus on all the above (family, referral agents, other key organisations, team)?

Conclusion

The implementation of evidence based models has been a growing phenomenon within human services. However, evidence based models have tended to be developed and proven effective for specific populations and for addressing well defined, albeit complex issues. Within some environments, this limits the extent to which specific models can successfully address concurrent issues within complex social systems, such as the family.

This proof of concept study has demonstrated some potential to support families who experience multiple complexities and significant adversity simultaneously. Despite the limitations noted, families with such complex needs experienced positive outcomes. Additionally, practitioners described in detail the acceptability of this type of approach.

Limitations and Future Directions

The concept of blending distinct interventions to improve outcomes for children and families has practical implications for professionals and front line services. Despite the potential of the blended model approach to supporting families with complex and concurrent needs, the authors acknowledge the current limitations.

Measures used within this study were non-standardised.. Whilst they were based upon the mutual concerns of the family, the project worker and the project manager, there is a need to enhance the objectivity of such findings by incorporating validated measures. .

Both in terms of IFSS replication and the evolving study of implementation, future studies could explore in greater detail the mechanism/s successfully driving a multiple implementation site and complexities around such implementation including defining core components across multiple interventions and fidelity issues.

However, given the novelty of the blended model approach, there are few other empirical studies to anchor this article onto. As such, further investigation is needed using larger sample sizes and explore the feasibility of this approach using a wider set of interventions.

References

- Allen, G. (2011) Early Intervention: The Next Steps. An Independent Report to Her Majesty's Government. Cabinet Office. Available from:<http://grahamallenmp.wordpress.com>
- Armeliuss, B., & Andreassen, T. H. (2007). Cognitive-behavioral treatment for antisocial behavior in youth in residential treatment. Cochrane Database of Systematic Reviews. doi:10.1002/14651858.cd005650.pub2
- Barbour, R. (1999) The case for combining qualitative and quantitative approaches in health services research. *Journal of Health Services Research & Policy*, Vol1
- Bazeley, P., Jackson, K (2013) *Qualitative Data Analysis with NVIVO*, 2nd edn. Sage
- Beck, A. T., Rush, A. J., Shaw, B. F., and Emery, G. (1979). *Cognitive therapy of depression*. New York: Guilford.
- Beck, J. S. (1995). *Cognitive therapy: Basics and beyond*. London: Guilford Press

- Beck, J. (2011). *Cognitive Behaviour Therapy: Basics and Beyond* (2nd ed). New York. Guilford Press
- Bunting, L., & Lazenbett, A. (2015) Changing the narrative: lifespan perspectives on multiple adversity. *Qualitative Social Work*, 15,4, 484-500
- Braun, V & Clarke, V. (2006) Using thematic analysis in psychology. *Qualitative Research in Psychology*, 3 (2), pp. 77-101
- Bronstein, L., Kovacs, P. (2013) Writing a Mixed Methods Report in Social Work Research. *Research on Social Work Practice*, 23(3)
- Bryman, A. (2004) *Social Research Methods*. 2nd ed. Oxford: Oxford University Press
- Butler, M., Kane, R.L., McAlpine, D., Kathol, R.G., Fu, S., Hagedorn, H., Wilt, T.J. (2008). *Integration of mental health/substance abuse and primary care*. Evidence Report/Technology Assessment (Full Report), 1-362
- Cabinet Office (2011). *Early Intervention: The Next Steps. An Independent Report to Her Majesty's Government*. Graham Allen MP. London: Stationary Office. <http://media.education.gov.uk/assets/files/pdf/g/graham%20allens%20review%20of%20early%20intervention.pdf>
- Catch 22 (2014) Lessons on replication and evidence-based interventions: Mid-programme report. Catch 22 <http://www.catch-22.org.uk/wp-content/uploads/2014/10/Realising-Ambition-Mid-Programme-Report-Summary.pdf>
- Churchill, H and Sen, R. (2016) Introduction: Intensive Family Support Services: Politics, Policy and Practice Across Contexts. *Social Policy & Society*, 15(02), pp 251-261
- Cobham, V.E., Filus, A., Sanders, M.R. (2017) Working with parents to treat anxiety-disordered children: A proof of concept RCT evaluating Fear-less Triple P. *Behavior and Research Therapy*, 95, pp.128-138
- Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., Petticrew, M. (2006) *Developing and evaluating complex interventions: new guidance*. Medical Research Council. Retrieved from: <https://mrc.ukri.org/documents/pdf/complex-interventions-guidance/>
- Creswell, J & Clark, V. (2007) *Designing and conducting mixed methods research*. Sage Publications: London
- Creswell, J. (2009) *Research design: Qualitative, quantitative, and mixed methods approaches*. 3rd ed. Sage Publications, Thousand Oaks, CA
- CYPSP. (2014). Early Intervention Transformation Programme (Rep.). Retrieved <http://www.cypsp.hscni.net/wp-content/uploads/2015/03/Website-Version-Early-Intervention-Transformation-Programme-EITP-Newsletter-Issue-1-March-2015.pdf>
- Davidson, G., Bunting, L. and Webb, M. A. (2012) *Families experiencing multiple adversities: A review of the international literature*. Belfast, Barnardos.
- Dartington (2013) Realising Ambition: Lessons on replication and evidence-based interventions-one year in. Social Research Unit Publication.
- Denscombe, M. (2007) *The Good Research Guide*. 3rd ed. Open University Press: Berkshire
- Department of Children and Youth Affairs. (2014) *Better Outcomes, Brighter Futures: The National Policy Framework for Children and Young People 2014 – 2020*. Retrieved from: https://www.dcy.gov.ie/documents/cypp_framework/BetterOutcomesBetterFutureReport.pdf
- Devaney, C., Canavan, J., Landy, F., Gillen, A. (2013) *What Works in Family Support?* Dublin: Child & Family Agency.
- Dillane, J., Hill, M., Bannister, J., Scott, S. (2001) Evaluation of the Dundee Families Project, Edinburgh: Scottish Executive, Dundee City Council and NCH Action for Children.
- Dixon, J., Schneider, V., Lloyd, C., Reeves, A., White, C., Tomaszewski, W., Green, R., Ireland, E. (2010). *Monitoring and Evaluation of Family Interventions (information on families supported to March 2010)*. London: Department for Education.
- Fontana, A., & Frey, J. H. (2005). The interview: From neutral stance to political involvement. In N. K. Denzin & Y. S. Lincoln (Eds.), *The Sage handbook of qualitative research* (3rd ed., pp. 695-727). Thousand Oaks, CA: Sage.
- Fixsen, D., Blasé, K., Naoom, S., Wallace, F. (2005) *Implementation Research: A Synthesis of the Literature*. Tampa, FL: University of South Florida. The National Implementation Research Network.
- Furlong, M., McGilloway, S., Bywater, T., Hutchings, J., Donnelly, M., Smith, S. (2012) Behavioural and cognitive-behavioural group-based parenting programmes for early-onset conduct problems in children aged 3 to 12 years. *Campbell Systematic Reviews*, 2012. <https://campbellcollaboration.org/library/group-parenting-programmes-conduct-problems-children-3-to-12.html>

- Gambrill, E. (2006). *Social Work Practice: A Critical Thinker's Guide*. 2nd ed. New York: Oxford University Press.
- Joughin, C. (2006) *Cognitive behaviour therapy can be effective in managing behavioural problems and conduct disorder in pre-adolescence*. What Works for Children group: Evidence Nugget; September 2006.
- Kazak, A.E., Hoagwood, K., Weisz, J.R., Hood, K., Kratochwill, T.R., Vargas, L.A., et al. (2010) A meta-systems approach to evidence-based practice for children and adolescents. *American Psychologists*, 65(2).
- Kelle, U (2001) 'Sociological Explanations Between Macro and Macro and the Integration of Qualitative and Quantitative Methods', *FQS* 2(1). Online, accessed on DATE at URL <http://qualitative-research.net>
- Kush, R., Alschuler, L., Ruggeri, R., Cassells, S., Gupta, N., Bain, L., Claise, K., Shah, M., Meredith Nahm, M. (2007) Implementing Single Source: The STARBRITE Proof-of-Concept Study. *Journal of the American Medical Informatics Association*, 14, 5
- Lee, Y., Anderson, E., Quranta, J., Shim, B. (2017) An interdisciplinary family-centered program: An innovative approach to better meet the multifaceted needs of grandparent-headed families. *Journal of Family Social Work*, In press. [doi.10.1080/10522158.2017.1300111](https://doi.org/10.1080/10522158.2017.1300111)
- Lemmens, L., Molema, C., Versnel, N., Baan, C., De Bruin, S. (2015) Integrated care programs for patients with psychological comorbidity: A systemic review and meta-analysis. *Journal of Psychosomatic Research*, 79, 580-94
- Lindsay, G., Strand, S. (2013). Evaluation of the national roll-out of parenting programmes across England: the parenting early intervention programme (PEIP). *BMC Public Health*, 13, pp.972
- Littell, J.H., Popa, M., Forsythe, B. Multisystemic Therapy for social, emotional, and behavioral problems in youth aged 10-17. *Cochrane Database of Systematic Reviews* 2005, Issue 4
DOI: 10.1002/14651858.CD004
- Littell, J. (2006) Letter to the Editor: The case for Multisystemic Therapy: Evidence or Orthodoxy. *Children and Youth Services Review*, 28, pp. 458-472
- Little, M. (2010) *Proof Positive*. DEMOS: London
- Lonne, B., Parton, N., Thomson, J., Harries, M. (2009) *Reforming Child Protection*. Routledge: Oxon
- McDaniel, B., Braiden, H., Regan, H. (2009) *Incredible Years Parenting Programme*. Northern Ireland: Barnardos
- Lynch, O., Joyce, C. (2018) Functions of collective victimhood Political violence and the case of the Troubles in Northern Ireland. *International Review of Victimology*, 24, 2
- McGilloway, S; Bywater, T; Ni Mhaille, G et al (2009) *Proving the power of positive parenting: A randomised controlled trial to investigate the effectiveness of the Incredible Years BASIC Parent Training Programme in an Irish context (short-term outcomes)*. Archways. Dublin: Archways and NUI Maynooth
- McAlinden, A.M., Dwyer, C. (eds.) (2015) *Criminal Justice in Transition: The Northern Ireland Context*. Oxford: Hart Publishing.
- McAlister, S., Scraton, P., Haydon, D. (2014) Childhood in transition: growing up in 'post conflict Northern Ireland. *Children's Geographies*, 12, 3, 297-311
- McGilloway, S., NiMhaille, G., Fulong, M., Hyland, L., Leckey, Y., Kelly, P., Bywater, T., et al. (2012) *The Incredible Years Ireland Study: Parents, teacher and childhood intervention*. Archways. Dublin: Archways and NUI Maynooth, Dublin.
- McEvoy, Levy, S. (2001) Youth Violence and Conflict Transformation. *Peace Review: A Journal of Social Justice*, 13 (1)
- McKeown, K. (2000) *A Guide to What Works in Family Support Services for Vulnerable Families*. Dublin: Area Development Management Limited.
- Manuel, J., Mullen, E., Fany, L., Bellamy, J., Bledsoe, S. (2009) Preparing Social Work Practitioners to Use Evidence-based Practice: A Comparison of Experiences From an Implementation Project. *Research on Social Work Practice*, 19
- Matteson, S., & Lincoln, Y. (2009) Using multiple interviewers in qualitative research. *Qualitative Inquiry*, 15, 4, 659-674
- Miles, M. B., & Huberman, A. M. (1994). *Qualitative data analysis: An expanded sourcebook*. 2nd ed. Thousand Oaks, CA: Sage.
- Miller, R., Brewer, J. (2003) *The A-Z of Social Research*. Sage: London
- Moran-Ellis, J., Alexander, V., Dickinson, M., Fielding, J., Slaney, J., Thomas, H. (2006) Triangulation and integration: processes, claims and implications. *Qualitative Research*, 6 (1)
- Munro, E (2011). *The Munro Review of Child Protection*. London: Department of Education Cm 8062

NICCY (2016) *Advice to the Northern Ireland Executive on its Draft Programme for Government* Belfast :Framework. Northern Ireland Commissioner for Children and Young People. Retrieved from: www.niccy.org/media/2506/niccy-pfg-advice-220716.docx

National Institute for Health and Clinical Excellence (2006) *Parent-training/education programmes in the management of children with conduct disorders*. London: NICE.

National Research Ethics Service (2009) *Defining Research*. UK: NHS retrieved from http://www.uhbristol.nhs.uk/media/1572809/defining_research_leaflet_1_.pdf

NISRA (2015). *NISRA Statistical Bulletin: Population and Migration Estimates in Northern Ireland* (2014). Northern Ireland: Northern Ireland Statistics and Research Agency. Retrieved from http://www.nisra.gov.uk/archive/demography/population/midyear/MYE14_Bulletin.pdf

Nixon, J., Parr, S., Hunter, C., Myers, S., Sanderson, D. and Whittle, S. (2006) *Anti-Social Behaviour Intensive Family Support Projects: An Evaluation of Six Pioneering Projects*. London: Communities and Local Government.

Nixon, J., Parr, S., Hunter, C., Myers, S., Sanderson, D. and Whittle, S. (2008) *The Longer Term Outcomes for Families Who Had Worked with Intensive Family Support Projects*. London: Communities and Local Government.

Odom, S. (2009) The tie that binds: Evidence-based practice, implementation science, and outcomes for children. *Topics in Early Childhood Special Education* 2009. 29 (1), 53-61

OFMDFM (2011) *Our Children Our Pledge: A strategy for Children and Young People in Northern Ireland*. Office of the First and Deputy First Minister of Northern Ireland: Northern Ireland.

Olsson, T. (2010). MST with Conduct Disordered Youth in Sweden: Costs and Benefits After 2 Years. *Research on Social Work Practice*, 20

Ovretveit, J., Hansson, J., Brommels, M. (2010) An integrated health and social care organisation in Sweden: Creation and structure of a unique local public health and social care system. *Health Policy*, 97

Padwa, H., Teruya, C., Tran, E., Lovinger, K., Antonini, V., Overholt, C., Urada, D. (2016) The implementation of integrated behavioural health protocols in primary care setting in Project Care. *Journal of Substance Abuse*, 62

Palinkas, L., Aarons, G., Chorpita, B., Hoagwood, K., Landsverk, J., Weisz, J. (2009). Cultural Exchange and the Implementation of Evidence-Based Practices: Two Case Studies. *Research on Social Work Practice*, 19

Paulsell, D., Del Grosso, P., Dynarski, M. (2009) *The Atlantic Philanthropies' Disadvantaged Children and Youth Program in Ireland and Northern Ireland: Overview of Program Evaluation Findings*. Dublin: Atlantic Philanthropies Limited

Pawson, H., Davidson, E., Sosenko, F., Flint, J., Nixon, J., Casey, R. and Sanderson, D. (2009) *Evaluation of Intensive Family Support Projects in Scotland*. Edinburgh: Scottish Government.

Rabinowitz, P. M., Kock, R., Kachani, M., Kunkel, R., Thomas, J., Gilbert, J., et al. (2013). Toward Proof of Concept of a One Health Approach to Disease Prediction and Control. *Emerging Infectious Disease Journal* 19 (12).

Rogers, E. (1995). *Diffusion of Innovations*. 3rd ed. New York: The Free Press

Schoenwald, S., Brown, T., Henggeler, S. (2000) Inside Multisystemic Therapy: Therapist, Supervisory and Program Practices. *Journal of Emotional and Behavioral Disorders*, 8

Silverman, D. (ed) (2004) *Qualitative Research, Theory, Method and Practice*. 2nd ed. London: Sage

Sneddon, H (2014) Parenting Programmes Review for Workstream 2-Executive Summary. Northern Ireland: Centre for Effective Services.

Snyder, D., Balderrama-Durbin, C. (2012) Integrative Approaches to Couple Therapy: Implications for Clinical Practice and Research. *Behaviour Therapy*, 43,13-24

Tolan, P. H. (2013). Making and using lists of empirically tested programs: Value for violence interventions for progress and impact. Washington, DC: The National Academies Press.

Ungar, T., Norman, C., Knaak, S. (2017) Think You Can Shrink? A Proof-of-Concept Study for Men's Health Education Through Edutainment. *Journal of Technology in Behavioral Science*, 2(2)

Valdebenito, S., Eisner, M., Farrington, D., Tfofi, M., Sutherland, A. (2018) School-based interventions for reducing disciplinary school exclusion. *Campbell Systematic Review*, 2018, 14 <https://campbellcollaboration.org/library/reducing-school-exclusion-school-based-interventions.html>

Waller, D.(2006) Art Therapy for Children: How It Leads to Change. *Clinical Child Psychology and Psychiatry*. 11 (2).

Walsh, C., & Doherty, F. (2016) *IFSS: A blended model of intervention*. Northern Ireland: Extern

Webster-Stratton, C. and J. Reid, *The Incredible Years Parents, Teachers and Children Training Series: A Multifaceted Treatment Approach for Young Children with Conduct Problems* in Evidence-based

psychotherapies for children and adolescents. 3rd edition. A.E. Kazdin and J.R. Weisz, Editors. 2017, Guildford Publications New York

White, C., Warrener, M., Reeves, A. and La Valle, I. (2008) *Family Intervention Projects: An Evaluation of Their Design, Set-up and Early Outcomes*. Department for Children, Schools and Families/Communities and Local Government: London

Wright, S., Gray, P., Watts, E., McAteer, L., Hazel, N., Liddle, M. and Haines, K. (2010) *Evaluation of Early Intervention Pilot Projects*. Swansea: University of Swansea: