



**QUEEN'S  
UNIVERSITY  
BELFAST**

## **Compositional data-driven alternatives to single geochemical component maps**

van den Boogaart, K. G., McKinley, J., de Caritat, P., Filzmoser, P., Grunsky, E., Hron, K., Reimann, C., & Tolosana-Delgado, R. (2015). Compositional data-driven alternatives to single geochemical component maps. In *Proceedings of the The 17th annual conference of the International Association for Mathematical Geosciences* (pp. 280). IAMG. <http://www.iamg2015.de/>

### **Published in:**

Proceedings of the The 17th annual conference of the International Association for Mathematical Geosciences

### **Document Version:**

Publisher's PDF, also known as Version of record

### **Queen's University Belfast - Research Portal:**

[Link to publication record in Queen's University Belfast Research Portal](#)

### **Publisher rights**

© 2015 The Authors

### **General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

### **Open Access**

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

## Queen's University Belfast - Research Portal

### Compositional data-driven alternatives to single geochemical component maps

van den Boogaart, K. G., McKinley, J., de Caritat, P., Filzmoser, P., Grunsky, E., Hron, K., ... Tolosana-Delgado, R. (2015). Compositional data-driven alternatives to single geochemical component maps. 280. Paper presented at The 17th Annual Conference of the International Association of Mathematical Geosciences (IAMG), Freiberg, Germany. ISBN 978-3-00-050337-5 (DVD)

**Document Version:**

Publisher final version (usually the publisher pdf)

**Link:**

[Link to publication record in Queen's University Belfast Research Portal](#)

**General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

## **Compositional data-driven alternatives to single geochemical component maps**

K. G. VAN DEN BOOGAART<sup>1,8\*</sup>, J. MCKINLEY<sup>2</sup>, P. DE CARITAT<sup>3</sup>, P. FILZMOSER<sup>4</sup>,  
E. GRUNSKY<sup>5</sup>, K. HRON<sup>6</sup>, C. REIMANN<sup>7</sup>, R. TOLOSANA DELGADO<sup>1</sup>

<sup>1</sup>Helmholtz Institute Freiberg for Resource Technology - Helmholtz Center Dresden Rossendorf, Germany  
[boogaart@hzdr.de](mailto:boogaart@hzdr.de)

<sup>2</sup>School of Geography, Archaeology and Palaeoecology - Queens Univ. Belfast, UK

<sup>3</sup>Geoscience Australia, Australia

<sup>4</sup>Dept. of Statistics and Mathematical Methods in Economics - Vienna University of Technology, Austria

<sup>5</sup>Geological Survey of Canada - Natural Resources Canada, Canada

<sup>6</sup>Dept. of Mathematical Analysis and Applications of Mathematics - Palacky University Olomouc, Czech Republic

<sup>7</sup>Geological Survey of Norway, Norway

<sup>8</sup>Institute for Stochastics - TU Bergakademie Freiberg, Germany

\* presenting author

Conventional practice in Regional Geochemistry includes as a final step of any geochemical campaign the generation of a series of maps, to show the spatial distribution of each of the components considered. Such maps, though necessary, do not comply with the compositional, relative nature of the data, which unfortunately make any conclusion based on them sensitive to spurious correlation problems. This is one of the reasons why these maps are never interpreted isolated. This contribution aims at gathering a series of statistical methods to produce individual maps of multiplicative combinations of components (logcontrasts), much in the flavor of equilibrium constants, which are designed on purpose to capture certain aspects of the data.

We distinguish between supervised and unsupervised methods, where the first require an external, non-compositional variable (besides the compositional geochemical information) available in an analogous training set. This external variable can be a quantity (soil density, collocated magnetics, collocated ratio of Th/U spectral gamma counts, proportion of clay particle fraction, etc) or a category (rock type, land use type, etc). In the supervised methods, a regression-like model between the external variable and the geochemical composition is derived in the training set, and then this model is mapped on the whole region. This case is illustrated with the Tellus dataset, covering Northern Ireland at a density of 1 soil sample per 2 square km, where we map the presence of blanket peat and the underlying geology.

The unsupervised methods considered include principal components and principal balances (Pawlowsky-Glahn et al., CoDaWork2013), i.e. logcontrasts of the data that are devised to capture very large variability or else be quasi-constant. Using the Tellus dataset again, it is found that geological features are highlighted by the quasi-constant ratios Hf/Nb and their ratio against SiO<sub>2</sub>; Rb/K<sub>2</sub>O and Zr/Na<sub>2</sub>O and the balance between these two groups of two variables; the balance of Al<sub>2</sub>O<sub>3</sub> and TiO<sub>2</sub> vs. MgO; or the balance of Cr, Ni and Co vs. V and Fe<sub>2</sub>O<sub>3</sub>. The largest variability appears to be related to the presence/absence of peat.