



**QUEEN'S
UNIVERSITY
BELFAST**

Compositional data-driven alternatives to single geochemical component maps

van den Boogaart, K. G., McKinley, J., de Caritat, P., Filzmoser, P., Grunsky, E., Hron, K., Reimann, C., & Tolosana-Delgado, R. (2015). Compositional data-driven alternatives to single geochemical component maps. In *Proceedings of the The 17th annual conference of the International Association for Mathematical Geosciences* (pp. 280). IAMG. <http://www.iamg2015.de/>

Published in:

Proceedings of the The 17th annual conference of the International Association for Mathematical Geosciences

Document Version:

Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2015 The Authors

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Queen's University Belfast - Research Portal

Compositional data-driven alternatives to single geochemical component maps

van den Boogaart, K. G., McKinley, J., de Caritat, P., Filzmoser, P., Grunsky, E., Hron, K., ... Tolosana-Delgado, R. (2015). Compositional data-driven alternatives to single geochemical component maps. 280. Paper presented at The 17th Annual Conference of the International Association of Mathematical Geosciences (IAMG), Freiberg, Germany. ISBN 978-3-00-050337-5 (DVD)

Document Version:

Publisher final version (usually the publisher pdf)

Link:

[Link to publication record in Queen's University Belfast Research Portal](#)

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Compositional data-driven alternatives to single geochemical component maps

K. G. VAN DEN BOOGAART^{1,8*}, J. MCKINLEY², P. DE CARITAT³, P. FILZMOSER⁴,
E. GRUNSKY⁵, K. HRON⁶, C. REIMANN⁷, R. TOLOSANA DELGADO¹

¹Helmholtz Institute Freiberg for Resource Technology - Helmholtz Center Dresden Rossendorf, Germany
boogaart@hzdr.de

²School of Geography, Archaeology and Palaeoecology - Queens Univ. Belfast, UK

³Geoscience Australia, Australia

⁴Dept. of Statistics and Mathematical Methods in Economics - Vienna University of Technology, Austria

⁵Geological Survey of Canada - Natural Resources Canada, Canada

⁶Dept. of Mathematical Analysis and Applications of Mathematics - Palacky University Olomouc, Czech Republic

⁷Geological Survey of Norway, Norway

⁸Institute for Stochastics - TU Bergakademie Freiberg, Germany

* presenting author

Conventional practice in Regional Geochemistry includes as a final step of any geochemical campaign the generation of a series of maps, to show the spatial distribution of each of the components considered. Such maps, though necessary, do not comply with the compositional, relative nature of the data, which unfortunately make any conclusion based on them sensitive to spurious correlation problems. This is one of the reasons why these maps are never interpreted isolated. This contribution aims at gathering a series of statistical methods to produce individual maps of multiplicative combinations of components (logcontrasts), much in the flavor of equilibrium constants, which are designed on purpose to capture certain aspects of the data.

We distinguish between supervised and unsupervised methods, where the first require an external, non-compositional variable (besides the compositional geochemical information) available in an analogous training set. This external variable can be a quantity (soil density, collocated magnetics, collocated ratio of Th/U spectral gamma counts, proportion of clay particle fraction, etc) or a category (rock type, land use type, etc). In the supervised methods, a regression-like model between the external variable and the geochemical composition is derived in the training set, and then this model is mapped on the whole region. This case is illustrated with the Tellus dataset, covering Northern Ireland at a density of 1 soil sample per 2 square km, where we map the presence of blanket peat and the underlying geology.

The unsupervised methods considered include principal components and principal balances (Pawlowsky-Glahn et al., CoDaWork2013), i.e. logcontrasts of the data that are devised to capture very large variability or else be quasi-constant. Using the Tellus dataset again, it is found that geological features are highlighted by the quasi-constant ratios Hf/Nb and their ratio against SiO₂; Rb/K₂O and Zr/Na₂O and the balance between these two groups of two variables; the balance of Al₂O₃ and TiO₂ vs. MgO; or the balance of Cr, Ni and Co vs. V and Fe₂O₃. The largest variability appears to be related to the presence/absence of peat.