



**QUEEN'S
UNIVERSITY
BELFAST**

Characterizing Jordan Homomorphisms

Mathieu, M. (2020). Characterizing Jordan Homomorphisms. *Acta Scientiarum Mathematicarum*, 86(3-4), 697-701. <https://doi.org/10.14232/actasm-020-067-7>

Published in:
Acta Scientiarum Mathematicarum

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© 2020 ACTA.

This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

CHARACTERIZING JORDAN HOMOMORPHISMS

MARTIN MATHIEU

ABSTRACT. It is shown that every bounded, unital linear mapping that preserves elements of square zero from a C^* -algebra of real rank zero and without tracial states into a Banach algebra is a Jordan homomorphism.

Throughout, A and B will be (at least) unital complex Banach algebras. A *Jordan homomorphism* between A and B is a linear mapping $T: A \rightarrow B$ such that $T(x^2) = (Tx)^2$ for all $x \in A$. That is, T is a homomorphism of the associated special Jordan algebras A^+ and B^+ . The quest of characterizing Jordan homomorphisms by spectral conditions is largely motivated by Kaplansky's question from the 1970s whether every unital, bounded, surjective linear mapping T which preserves invertibility between semisimple Banach algebras must be a Jordan epimorphism (i.e., a surjective Jordan homomorphism). All listed conditions are well known to be necessary (where *unital* means $T1 = 1$). While this problem is open in the stated generality, many contributions under additional conditions on the algebras have been obtained (see, e.g., the survey [9]), notably if the domain is a C^* -algebra. Aupetit [3] affirmed Kaplansky's question for von Neumann algebras, and his proof can easily be adapted to the situation where A is a unital C^* -algebra with real rank zero and B is any semisimple unital Banach algebra; cf. [8, Theorem 1.1].

The abundance of projections in C^* -algebras with real rank zero (i.e., every self-adjoint element can be approximated by finite linear combinations of orthogonal projections) is a key ingredient via the following result; see [10, Lemma 2.1] or [7, Lemma 1].

Lemma 1. *Let $T: A \rightarrow B$ be a bounded linear operator from a C^* -algebra A with real rank zero into a Banach algebra B sending projections in A to idempotents in B . Then T is a Jordan homomorphism.*

With few exceptions, see, e.g., [11], the assumption of real rank zero in the domain has remained essential. The connection between idempotents and elements of square zero is provided by the following lemma from [10, Lemma 3.3].

Lemma 2. *Let $T: A \rightarrow B$ be a linear mapping between Banach algebras A and B which preserves elements with square zero. If e, f are orthogonal idempotents in A , then*

$$(Ta)(Tb) + (Tb)(Ta) = 0$$

2020 *Mathematics Subject Classification.* 47B48, 46L05, 46L30, 16W10, 17C65.

Key words and phrases. C^* -algebras, commutators, nilpotents, tracial states, Jordan homomorphisms, spectrally bounded operators.

for all $a \in eAe$, $b \in fAf$ which can be written as finite sums of elements with square zero.

Let A be a unital C^* -algebra and denote by $N^{(2)}$ the linear span of all elements in A with square zero and by $[A, A]$ the linear span of all commutators $[x, y] = xy - yx$, $x, y \in A$. Every square-zero element x is a commutator; e.g., write $x = [v|x|^{1/2}, |x|^{1/2}]$ where $x = v|x|$ is the polar decomposition of x (in a faithful representation of A or the enveloping von Neumann algebra A^{**}). Indeed, $(|x|^{1/2}v|x|^{1/2})(|x|^{1/2}v|x|^{1/2})^* = |x|^{1/2}xv^*|x|^{1/2} = 0$ since $|x|^{1/2}$ belongs to the C^* -subalgebra generated by x^*x and $(x^*x)x = 0$. Consequently, $N^{(2)} \subseteq [A, A]$ in general.

Recall that a *tracial state* on A is a positive linear functional τ of norm 1 such that $\tau(xy) = \tau(yx)$ for all $x, y \in A$. Clearly $[A, A] \subseteq \ker \tau$ for every tracial state τ . Therefore, an obstruction for the equality $A = [A, A]$ is the existence of a tracial state. Pop [13] proved that this is the only obstruction. On the other hand, Robert showed in [14, Theorem 4.2] that $[A, A] = N^{(2)}$ provided A has no characters. Combining these two results we obtain a characterization of the identity $N^{(2)} = A$.

Lemma 3. *Let A be a unital C^* -algebra. Then every element of A is a finite sum of square zero elements if and only if A has no tracial states.*

This is clear as every character is a tracial state.

We now put the results above together to verify a conjecture, in the real rank zero case, which we first stated in [8]; see also [6].

Theorem. *Let A be a unital C^* -algebra with real rank zero and $\dim A \geq 2$. The following conditions are equivalent.*

- (a) *A has no tracial states;*
- (b) *every bounded unital linear mapping from A into a unital Banach algebra preserving square zero elements is a Jordan homomorphism.*

Proof. Evidently we have to exclude the case $\dim A < 2$ since every unital linear mapping is multiplicative in this case.

Suppose τ is a tracial state on A . Since $N^{(2)} \subseteq [A, A] \subseteq \ker \tau$, τ preserves square zero elements. Therefore, assumption (b) entails that τ is a character on A . Hence $A = \ker \tau \oplus \mathbb{C}1$ and $\ker \tau$ is a maximal ideal. Assume first that A is not commutative. Let S be an $*$ -automorphism of A and pick $\gamma \in (0, 1)$. Set

$$T: A \rightarrow A, \quad Tx = \gamma Sx + (1 - \gamma)\tau(x).$$

Then T is a unital bounded linear mapping on A , with $\|T\| = 1$, and, for all $x, y \in A$,

$$(Tx)(Ty) = \gamma^2 (Sx)(Sy) + (1 - \gamma)^2 \tau(x)\tau(y) + \gamma(1 - \gamma)\tau(y)Sx + \gamma(1 - \gamma)\tau(x)Sy.$$

Take $x \in A$ non-zero such that $x^2 = 0$ (such x exists only if A is noncommutative) and put $y = x$. Then $(Tx)^2 = \gamma^2 (Sx)^2 = \gamma^2 S(x^2) = 0$ so T preserves elements of square

zero and hence is a Jordan homomorphism under condition (b). On the other hand, for $y = x^*$,

$$\begin{aligned} (Tx)(Ty) + (Ty)(Tx) &= \gamma^2 ((Sx)(Sy) + (Sy)(Sx)) = \gamma^2 S(xy + yx) \\ &\neq \gamma S(xy + yx) = T(xy + yx), \end{aligned}$$

a contradiction. Assume now that A is commutative. Since $\dim A \geq 2$ there is another character $\tau' \neq \tau$ of A . Take $x, y \in A$ such that $\tau(x) = 0$, $\tau'(x) = 1$, $\tau(y) = 1$ and $\tau'(y) = 0$. Put $\rho = \frac{1}{2}(\tau + \tau')$ which is another tracial state thus a character, by condition (b). Then

$$\rho(xy) = \frac{1}{2}(\tau(x)\tau(y) + \tau'(x)\tau'(y)) = 0 \neq \frac{1}{4} = \rho(x)\rho(y)$$

which is a contradiction.

Consequently, condition (b) rules out the existence of tracial states on A , that is, condition (a) holds.

Conversely assume condition (a). Let $T: A \rightarrow B$ be a bounded linear mapping from A into a unital Banach algebra B with $T1 = 1$ and suppose that $(Tx)^2 = 0$ for every $x \in A$ with $x^2 = 0$. Let p be a non-trivial projection in A . As every bounded trace τ on the hereditary C^* -subalgebra pAp extends to A , pAp has no tracial states. (This is known and follows, e.g., from combining [4, Proposition II.4.2] to extend τ from the (completely) full hereditary C^* -algebra to the closed ideal \overline{ApA} and then onto all of A , a slick proof of the latter is contained in [15, Lemma 3.1].) Lemma 3 thus entails that every element in pAp is a finite sum of square zero elements. By Lemma 2, we conclude that

$$0 = (Tp)(T(1-p)) + (T(1-p))(Tp) = 2Tp - (Tp)^2$$

and thus Tp is an idempotent in B . Now Lemma 1 finishes the proof. \square

The above result extends Theorem 2.4 in [6] for the purely infinite case. Note that every bounded linear mapping which vanishes on elements of square zero necessarily is a trace as $\overline{N^{(2)}} = \overline{[A, A]}$ ([1, Proposition 2.2] or [14, Corollary 2.3]).

A linear mapping $T: A \rightarrow B$ is said to be *spectrally bounded* if $r(Tx) \leq M r(x)$ for some constant $M \geq 0$ and all $x \in A$. Here, $r(\cdot)$ denotes the spectral radius. When B is semisimple and T is surjective, spectral boundedness implies boundedness of T [2, Theorem 5.5.2]. Every Jordan epimorphism T is unital and preserves invertibility [5, Lemma 4.1]; hence, it is spectrally bounded with constant 1.

Suppose that A is a unital C^* -algebra and B is a unital semisimple Banach algebra. It was observed in [10, Lemma 3.1] that surjective spectrally bounded operators preserve nilpotency, we restrict ourselves to elements of square zero here.

Lemma 4. *Let $T: A \rightarrow B$ be spectrally bounded and surjective. For every $x \in A$ with $x^2 = 0$ we have $(Tx)^2 = 0$.*

Corollary. *Let $T: A \rightarrow B$ be a unital surjective spectrally bounded operator from a unital C^* -algebra A with real rank zero and without tracial states onto a unital semisimple Banach algebra B . Then T is a Jordan epimorphism.*

This is now immediate from the theorem and Lemma 4.

The statement of the corollary was obtained under the additional assumption that A is simple in [8, Theorem 3.1]. For properly infinite von Neumann algebras it is in [10, Theorem 3.6] and for A purely infinite of real rank zero in [6, Corollary 2.5] (which extended the simple case treated in [7, Theorem B]).

In the case when the domain C^* -algebra has a tracial state, the above statement fails already in the finite-dimensional situation. In fact, a unital surjective linear mapping $T: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$ is spectrally bounded if and only if it is of the form $Tx = \gamma Sx + (1 - \gamma)\tau(x)$, $x \in M_n(\mathbb{C})$ for a unique non-zero complex number γ and a unique Jordan isomorphism S on $M_n(\mathbb{C})$, where τ denotes the unique tracial state on $M_n(\mathbb{C})$; cf. [9], Example 5.3 and Remark 5.5.

On the other hand the assumption that T is unital can sometimes be relaxed, see for example [9], in particular when T is a *spectral isometry*, that is, $r(Tx) = r(x)$ for all $x \in A$; see [12], Proposition 2.3 and Corollary 2.4. The above corollary also removes an additional assumption on the primitive ideal space of the domain algebra from [11, Theorem 3.6].

REFERENCES

- [1] J. Alaminos, M. Brešar, J. Extremera, Š. Špenko and A. R. Villena, *Commutators and square-zero elements in Banach algebras*, Quart. J. Math. **67** (2016), 1–13.
- [2] B. Aupetit, *A Primer on Spectral Theory*, Springer-Verlag, New York, 1991.
- [3] B. Aupetit, *Spectrum-preserving linear mappings between Banach algebras or Jordan–Banach algebras*, J. London Math. Soc. **62** (2000), 917–924.
- [4] B. Blackadar and D. Handelman, *Dimension functions and traces on C^* -algebras*, J. Funct. Anal. **45** (1982), 297–340.
- [5] P. Civin and B. Yood, *Lie and Jordan structures in Banach algebras*, Pacific J. Math. **15** (1965), 775–797.
- [6] Y.-F. Lin and M. Mathieu, *Jordan isomorphism of purely infinite C^* -algebras*, Quart. J. Math. **58** (2007), 249–253.
- [7] M. Mathieu, *Spectrally bounded operators on simple C^* -algebras*, Proc. Amer. Math. Soc. **132** (2004), 443–446.
- [8] M. Mathieu, *Spectrally bounded operators on simple C^* -algebras II*, Irish Math. Soc. Bull. **54** (2004), 33–40.
- [9] M. Mathieu, *A collection of problems on spectrally bounded operators*, Asian-Eur. J. Math. **2** (2009), 487–501.
- [10] M. Mathieu and G. J. Schick, *Spectrally bounded operators from von Neumann algebras*, J. Operator Theory **49** (2003), 285–293.
- [11] M. Mathieu and A. R. Sourour, *Spectral isometries on non-simple C^* -algebras*, Proc. Amer. Math. Soc. **142** (2014), 129–145.
- [12] M. Mathieu and M. Young, *Spectral isometries into commutative Banach algebras*, Contemp. Math. **645** (2015), 217–222.

- [13] C. Pop, *Finite sums of commutators*, Proc. Amer. Math. Soc. **130** (2002), 3039–3041.
- [14] L. Robert, *On the Lie ideals of C^* -algebras*, J. Operator Theory **75** (2016), 387–408.
- [15] E. P. Scarparo, *Supramenable groups and partial actions*, Ergodic Theory Dynam. Systems **37** (2017), 1592–1606.

MATHEMATICAL SCIENCES RESEARCH CENTRE, QUEEN'S UNIVERSITY BELFAST, BELFAST BT7
1NN, NORTHERN IRELAND

Email address: m.m@qub.ac.uk