



**QUEEN'S  
UNIVERSITY  
BELFAST**

## **Smoked cigarette butt leachate impacts survival and behaviour of freshwater invertebrates**

Green, D. S., Kregting, L., & Boots, B. (2020). Smoked cigarette butt leachate impacts survival and behaviour of freshwater invertebrates. *Environmental Pollution*, 266, [115286]. <https://doi.org/10.1016/j.envpol.2020.115286>

**Published in:**  
Environmental Pollution

**Document Version:**  
Peer reviewed version

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

### **Publisher rights**

© 2020 Elsevier Ltd.

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

### **General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

1 **Please see link for Publisher’s version: <https://doi.org/10.1016/j.envpol.2020.115286>**

2 **Smoked cigarette butt leachate impacts survival and behaviour of freshwater**  
3 **invertebrates**

4 Dannielle Senga Green<sup>1\*</sup>, Louise Kregting<sup>2</sup>, Bas Boots<sup>1</sup>

5 *<sup>1</sup>Applied Ecology Research Group, School of Life Sciences, Anglia Ruskin University,*  
6 *Cambridge, CB1 1PT, United Kingdom.*

7 *<sup>2</sup>Marine Biology Station, Queen’s University Belfast, Portaferry, Newtownards, BT22 1PF,*  
8 *United Kingdom.*

9 \*Corresponding author: [dannielle.green@aru.ac.uk](mailto:dannielle.green@aru.ac.uk).

## 10 **Abstract**

11 Smoked cigarette filters a.k.a. “butts”, composed of plastic (e.g. cellulose acetate) are one of  
12 the world’s most common litter items. In response to concerns about plastic pollution,  
13 biodegradable cellulose filters are being promoted as an environmentally safe alternative,  
14 however, once smoked, both contain toxins which can leach once discarded. The impacts of  
15 biodegradable butts as littered items on the receiving environment, in comparison with  
16 conventional butts has not yet been assessed. A freshwater mesocosm experiment was used to  
17 test the effects of leachate from smoked cellulose acetate versus smoked cellulose filters at a  
18 range of concentrations (0, 0.2, 1 and 5 butts L<sup>-1</sup>) on the mortality and behaviour of four  
19 freshwater invertebrates (*Dreissena polymorpha*, *Polycelis nigra*, *Planorbis planorbis* and  
20 *Bithynia tentaculata*). Leachate derived from 5 butts L<sup>-1</sup> of either type of filter caused 60-100%  
21 mortality to all species within 5 days. Leachate derived from 1 butt L<sup>-1</sup> of either type resulted  
22 in adults being less active than those exposed to no or 0.2 butts L<sup>-1</sup> leachate. Cigarette butts,  
23 therefore, regardless of their perceived degradability can cause mortality and decreased activity  
24 of key freshwater invertebrates and should always be disposed of responsibly.

25 **Key words:** smoking, cigarette butts, leachate, molluscs, platyhelminth.

26 **Capsule:** As litter in enclosed aquatic habitats, conventional and biodegradable cigarette butts  
27 have the same effects causing mortality and behavioural changes to invertebrates.

28

## 29 **1. Introduction**

30 Cigarette butts (used cigarette filters) are the most common form of personal litter worldwide  
31 due to the majority (>75%) of smokers littering them after use (Patel et al. 2013). Each year,  
32 ~6 trillion cigarettes are smoked globally, possibly resulting in an estimated deposition of ~4.5  
33 trillion used cigarette butts in the environment (Novotny and Slaughter 2014). Despite their  
34 prevalence as litter in the environment, the effects of cigarette butts on marine, freshwater and  
35 terrestrial habitats is still vastly understudied. The majority (~90%) of cigarette filters are  
36 composed of cellulose acetate (Pauly et al. 2002), a type of plastic which is not readily  
37 biodegradable, but can break down into smaller pieces and persist as microplastics and  
38 nanoplastics (Chevalier et al. 2018). Cellulose acetate itself can cause environmental impacts  
39 as litter, with some studies finding that even unsmoked plastic filters can cause a detrimental  
40 effect on the receiving ecosystem, for example, decreasing plant growth (Green et al. 2019)  
41 causing mortality to fish (Slaughter et al. 2011) and amphibians (Lawal and Ologundudu 2013).  
42 In response to concerns about plastic, alternative materials, including pure, unbleached  
43 cellulose, are being promoted for use in cigarette filters instead of cellulose acetate plastic.  
44 These alternative filters have been described as “green”, “biodegradable” and “environmentally  
45 friendly” giving the impression that these items would be benign as litter (Amos et al. 2017).  
46 There is, however, no research providing evidence of their level of toxicity as litter items nor  
47 any research comparing their effects with that of the cellulose acetate butts.

48 As litter, cigarette butts present a unique combination of physical and chemical contamination.  
49 Once smoked, cigarette butts contain thousands of chemicals including nicotine, polycyclic  
50 aromatic hydrocarbons and heavy metals which, once entering an aquatic environment, can  
51 leach out into the surrounding water (Moerman and Potts 2011; Roder Green et al. 2014;  
52 Dobaradaran et al. 2019). Such leachates are likely to pose a greater threat to lotic habitats that  
53 can have slow rates of water turnover such as ponds, low energy streams or rockpools than to  
54 habitats where the rate of water replacement is rapid (e.g. the ocean and in fast flowing streams  
55 and rivers). Indeed, leachate from smoked cigarette butts can be lethal for freshwater organisms  
56 such as microalgae, including *Raphidocelis subcapitata* (Bonanomi et al. 2020), water fleas,  
57 including *Ceriodaphnia dubia* (Warne et al. 2002, Micevska et al. 2006), *Daphnia magna*  
58 (Register 2000), fish including *Pimephales promelas* (Slaughter et al. 2011) and amphibians  
59 including *Hymenochirus curtipes* and *Clarias gariepinus* (Lawal and Ologundudu 2013).  
60 Although mortality often occurs at high concentrations of cigarette butt leachate ( $> 1$  butt  $L^{-1}$ ),  
61 sublethal impacts at lower, more environmentally realistic concentrations ( $<0.2$  butts  $L^{-1}$ ) have  
62 been observed, including mutagenic effects (Montalvão et al. 2019), developmental retardation  
63 (Lee and Lee 2015, Parker and Rayburn 2017) and alterations to behaviour (Booth et al. 2015;  
64 Wright et al. 2015). Such sublethal effects are often overlooked by policymakers, but may  
65 invoke important cascading ecological effects (Relyea and Hoverman 2006).

66 To explore toxicological effects of leachate from smoked cigarette butts at incremental  
67 concentrations, four different aquatic invertebrate species were studied in a controlled  
68 environment. The selected organisms included *Dreissena polymorpha* (Pallas 1771) (zebra  
69 mussel), *Polycelis nigra* (Müller 1774) (a flatworm), *Planorbis planorbis* (Linnaeus 1758)  
70 (ramshorn snail) and *Bithynia tentaculata* (Linnaeus 1758) (faucet snail). These were chosen  
71 as model organisms as each are commonly found in pond ecosystems across Europe and the  
72 UK and fulfil a range of ecosystem functions (as e.g. detritivores, grazers, filter feeders,

73 predators and prey organisms). Here, lethal (mortality) and sublethal (behaviour) effects were  
74 measured in response to leachate derived from smoked cigarettes with either conventional  
75 cellulose acetate filters or biodegradable cellulose filters. The hypothesis tested was that  
76 alternative, cellulose cigarette butts would not cause the same lethal and sublethal effects as  
77 conventional, cellulose acetate cigarette butts on the aquatic invertebrates.

78

## 79 **2. Materials and methods**

### 80 *2.1. Preparation of leachate from smoked cigarette filters*

81 Cigarettes were rolled manually using standard cigarette papers to an average ( $\pm$  S.E.) of 0.543  
82  $\pm$  0.002 g per cigarette of a leading brand of tobacco in the UK, with either a cellulose acetate  
83 or a cellulose (unbleached) filter. All cigarettes were smoked using a hand-operated vacuum  
84 pump with silicone tubing attached to the filter of the cigarettes. After lighting, approximately  
85 30 ( $\pm$  1) ml of air was drawn in, simulating a draft and each cigarette was smoked for a total  
86 inhalation volume of  $\sim$ 600 ml per cigarette, thereby emulating a similar total inhalation volume  
87 of cigarettes smoked by humans ( $585 \pm 245$ ) ml; McBride et al. 1984). Cigarettes were smoked  
88 until 2 mm from the edge of the filter and stubbed out in an aluminium tray. Any remaining  
89 tobacco was removed, leaving the filter with the cigarette paper attached. A stock solution of  
90 leachate from each type of filter used (cellulose and cellulose acetate) was prepared separately  
91 by soaking 14 smoked butts in 1 L of fresh, filtered (20  $\mu$ m) rainwater obtained from an  
92 artificial pond in glass volumetric flasks and gently agitating (100 rpm) on an orbital shaker  
93 for 18 h at room temperature ( $\sim$ 18  $^{\circ}$ C). Rainwater was chosen to represent how cigarettes butts  
94 may experience leaching when exposed to precipitation in the environment. Furthermore,  
95 rainwater resembles pond water more closely than media such as distilled water. Rainwater has  
96 been shown to also leach potential contaminants from cigarette butts (e.g. Koutela et al. 2020).

97 2.2. Mesocosm set-up and experimental design

98 The experiment was carried out in a temperature and light controlled facility at the Portaferry  
99 Marine Laboratory with a 12/12 h light/dark cycle. Mesocosms were set up in the laboratory,  
100 using conical glasses (86 mm diameter at top, 65 mm diameter at bottom) that were filled with  
101 rainwater (400 ml), extracted from the same artificial pond as the test organisms, and left to  
102 settle without any added leachate for 24 h before the experimental exposures were initiated.  
103 On day 1 (19<sup>th</sup> March, 2020) of the experiment, treatments were randomly assigned to  
104 mesocosms and corresponding leachate was added by removing the required volume of water  
105 and substituting with 5.7, 28.6 or 142.8 ml of stock leachate representing incremental  
106 concentrations based on 0.2, 1 or 5 smoked butts L<sup>-1</sup> of either cellulose or cellulose acetate  
107 smoked filters. The experimental organisms including *D. polymorpha*, *P. nigra*, *P. planorbis*  
108 and *B. tentaculata* were harvested using a net from an artificial pond (1.4 x 2.1 x 0.9 m). One  
109 individual of each species was added to each mesocosm along with five *B. tentaculata* juveniles  
110 thereby creating representative communities of similar densities to those found in the sampled  
111 pond (Table S1). A treatment with no added leachate served as a control. Therefore, the  
112 experiment consisted of an asymmetric design with 2 fixed factors; “Butts” (2 levels; cellulose  
113 versus cellulose acetate filters) and “Concentration” (3 levels; 0.2, 1 and 5 butts L<sup>-1</sup> added as  
114 leachate). Each treatment was replicated using 5 separate mesocosms (n = 5, N = 35) (Figure  
115 1). Water temperature within the mesocosms had an average pH of 8.13 (± 0.02), salinity <  
116 0.05 ppt and was maintained at 15 (± 0.42) °C throughout the experiment.

117 The experiment was repeatedly sampled every 24 h for a total of 120 h. At each sampling  
118 occasion, mortality was recorded and a number of behavioural observations were recorded into  
119 categories including (i) filtering or (ii) closed for the bivalves, (i) moving, (ii) open (antennae  
120 and foot extended) or (iii) closed (antennae and foot withdrawn into shell) for the gastropods  
121 and (i) moving, (ii) open (body elongated) or (iii) closed (body compressed into a spherical

122 shape) for the flatworms. Observations were made in real time by the same observer each time.  
123 Due to the high mortality rate in the 5 butts L<sup>-1</sup> treatments, behavioural observations were only  
124 recorded for mesocosms exposed to 0, 0.2 or 1 butt L<sup>-1</sup>.

125

### 126 2.3. Statistical analysis

127 Mortality data was categorised into a mortality scale ranging from 0 to 5 with “0” meaning no  
128 mortality at the end of the experiment and “1”, “2”, “3”, “4” and “5” meaning that death  
129 occurred after >120, 96, 72, 48 and 24 hours respectively. In this way, the higher the number,  
130 the more rapidly the animal died representing a more lethal effect. The survival of juvenile *B.*  
131 *tentaculata* was converted to percentage out of 5 which were still alive at each time point.  
132 Mortality and juvenile survival were analysed using asymmetrical ANOVA (see e.g. Green et  
133 al. 2016 for more details) to account for a single set of control units for the two experimental  
134 levels Butt and Concentration. The survival of juvenile *B. tentaculata* was analysed separately  
135 for each time point to avoid complications involved with repeated measures. Univariate data  
136 were screened for normality and homogeneity of variance to check assumptions of ANOVA  
137 and any necessary transformations are where appropriate. Statistical analyses were done using  
138 R V.3.6.2 (R Core Team 2019).

139 To test effects of leachate on the behaviour over the duration of the experiment, the behavioural  
140 data over the course of the 5 days was pooled and analysed mirroring the univariate analysis  
141 except with only 2 levels of leachate concentration (0.1 and 1 butt L<sup>-1</sup>) instead of three due to  
142 the removal of the 5 butts L<sup>-1</sup> treatment. Multivariate ANOVA was done on Bray-Curtis  
143 dissimilarities of untransformed data with 9999 permutations under the reduced model using  
144 Type I SS using the vegan package v2.5-2 (Oksanen et al. 2019). The asymmetric analysis was  
145 done by fitting each main effect (‘Butt’ and ‘Concentration’) in turn with a Type I (sequential)  
146 SS model, swapping the order of the terms and combining the results of these 2 analyses. The

147 multivariate behaviour data were visualised using a non-metric multidimensional scaling  
148 ordination approach reflecting the dissimilarity matrix used for the PERMANOVA with  
149 variables with a Pearson's correlation  $R > 0.6$  overlain as vectors. SIMPER was used to  
150 elucidate which behaviours were driving the significant differences between treatments  
151 (contributing  $>5\%$  to the dissimilarity) found by PERMANOVA analysis. Note that  
152 behavioural data is a sublethal response variable so data from either of the 5 butts  $L^{-1}$  treatments  
153 was omitted since there was a high instance of mortality in these treatments. The nMDS and  
154 the SIMPER analyses were generated using Primer V6.1.13 (PRIMER-e, Plymouth, UK).

155

### 156 **3. Results**

#### 157 *3.1. Effects of leachate from smoked cigarette butts on mortality of aquatic invertebrates.*

158 At 5 butts  $L^{-1}$  most of *D. polymorpha*, *P. planorbis*, *B. tentaculta* and *P. nigra* died after 72  
159 hours of exposure on average (Figure 1), which was significantly (Table 1) different from  
160 mesocosms treated with 1 butt  $L^{-1}$  (Concentration [5 vs 1 butt  $L^{-1}$ ]:  $P < 0.001$ ), 0.2 butt  $L^{-1}$   
161 (Concentration [5 vs 0.2 butt  $L^{-1}$ ]:  $P < 0.001$ ) or mesocosm with no leachate (Concentration [5  
162 butt  $L^{-1}$  vs control]:  $P < 0.001$ ). There was no significant difference between survival of the test  
163 organisms based on leachate derived from cellulose versus cellulose acetate butts (Table 1).

164 Significantly fewer juvenile *B. tentaculata* survived in mesocosms with either 5 butts  $L^{-1}$  of  
165 cellulose acetate or cellulose butts compared with controls with less than 20% surviving even  
166 after just 24 h (Table 1, Concentration [Control vs 5 cellulose butts  $L^{-1}$ ]:  $P < 0.001$ ,  
167 Concentration [control vs 5 cellulose acetate butts  $L^{-1}$ ]:  $P < 0.001$  for each time point). After  
168 48 and 72 h, survival with 1 cellulose acetate butt  $L^{-1}$  was  $\sim 50\%$  which was significantly lower  
169 than in the Controls (Control vs 1 cellulose acetate butt  $L^{-1}$ :  $P < 0.001$  at 48 and 72 h). At the  
170 same time points (48 and 72 h) 1 cellulose butt  $L^{-1}$  did not have a significant effect on survival  
171 (Control vs 1 cellulose butt  $L^{-1}$ :  $P = 0.690$ ). After 120 h, however, there were no differences



172 between cellulose acetate and cellulose butts and 1 butt L<sup>-1</sup> of either type caused survival to  
173 drop to ~30% (Table 1, Figure 2). In addition, by 120 h, survival decreased with increasing  
174 concentration of leachate with 100% survival at 0.2 butts L<sup>-1</sup>, ~30% at 1 butt L<sup>-1</sup> and <5% at 5  
175 butts L<sup>-1</sup> (post-hoc tests for concentration at 120 h; 0.2 vs 1: P < 0.001, 0.2 vs 5: P < 0.001 and  
176 1 vs 5: P < 0.001).

177

### 178 3.2. Sub-lethal effects of leachate from smoked cigarette butts on aquatic invertebrates.

179 Behaviour of the surviving individuals did not significantly differ (Figure 3) regardless of the  
180 source of the leachate (Butt [cellulose vs cellulose acetate], P = 0.458). The concentration of  
181 leachate, however, did significantly alter patterns of behaviour. In particular, the mesocosms  
182 exposed to 1 butt L<sup>-1</sup> exhibited different types of behaviour compared to those in mesocosms  
183 with 0.2 butts L<sup>-1</sup> or no leachate (Concentration [control vs 1 butt L<sup>-1</sup>]: P < 0.003 and [0.2 vs 1  
184 butt L<sup>-1</sup>]: P = 0.002). These differences were mostly due to a greater occurrence of movement  
185 or filtering in the case of *D. polymorpha* (accounting for ~40% of the variation in the  
186 multivariate pattern), and less occurrence of being in a closed state (accounting for ~38% of  
187 the variation in the multivariate pattern), of all four species in mesocosms without leachate or  
188 with 0.2 butts L<sup>-1</sup> leachate compared with those in leachate from 1 butt L<sup>-1</sup> (Figure 3).

189

## 190 4. Discussion

191 Cigarette butt leachate derived from biodegradable (i.e. cellulose) filters was equally as  
192 detrimental to freshwater pond invertebrates as leachate derived from conventional (i.e.  
193 cellulose acetate) filters. Leachate from 5 butts L<sup>-1</sup> derived from either type of butt was lethal  
194 to ~60% of adult *P. nigra*, *P. planorbis* and *B. tentaculata* and to ~40% of adult *D. polymorpha*  
195 within 48 hours. This is similar, albeit less lethal, to the results of Booth et al. (2015) who found  
196 100% mortality of two species of marine gastropod (*Austrocochlea porcata* and *Nerita*

197 *atramentosa*) after 24 hours of continuous exposure to leachate from 5 butts L<sup>-1</sup>, but 100%  
198 mortality of a third species (*Bembecium nanum*) did not occur until 150 hours.

199 In the current study, mortality of adults was low at exposure to leachate from 1 butt L<sup>-1</sup>  
200 equivalent and no animals died during the experimental period in mesocosms with no or just  
201 0.2 butts L<sup>-1</sup> equivalent of leachate. Juvenile *B. tentaculata*, however, were more sensitive to 1  
202 butt L<sup>-1</sup> than their adult counterparts with only ~30% of juveniles surviving after 120 h of  
203 exposure versus ~80% of adults. This is not surprising given that early life stages of  
204 invertebrates are typically more sensitive to toxicants and hence are often prioritised for use in  
205 ecotoxicological studies (Mohammed 2013). For example, early life stage (ELS) tests (such as  
206 OECD 2018) are widely conducted to estimate toxicity for the registration of industrial  
207 chemicals, pesticides, biocides, and pharmaceuticals. Early life stages are also important  
208 ecologically because a reduction in successful recruitment can result in changes to population  
209 dynamics over the longer term and cause shifts in freshwater biodiversity and ecosystem  
210 functioning (Strayer and Malcom 2012).

211 It is important to measure sublethal responses to contaminants as these may be ecologically  
212 important, for example, movement facilitates feeding, predator avoidance, reproduction and  
213 migration and so can link effects on individuals to a population level (Bayley et al. 1997). Even  
214 though there was little mortality of adults at 1 butt L<sup>-1</sup> of leachate, significant alterations to  
215 behaviour did occur whereby the test animals were less active. It is likely that this indicates  
216 that they were under stress and in the longer-term this may have led to mortality (Rubach et al.  
217 2011). In a study by Wright et al (2015), a marine polychaete (*Hediste diversicolor*) was also  
218 found to be less active, decreasing burrowing in response to >2 butts L<sup>-1</sup> leachate. Alteration  
219 to behaviour also occurred in marine gastropods exposed to 1.25 butts L<sup>-1</sup>, but this differed  
220 depending on the species (Booth et al. 2015). Lee and Lee (2015) found contrasting effects at  
221 increasing concentrations of cigarette butt leachate, with significantly increased heart rates and

222 accelerated embryonic development at lower concentrations (0.2 - 2 butts L<sup>-1</sup>), but lower heart  
223 rates and suppressed development at high concentrations (5 - 10 butts L<sup>-1</sup>). In addition,  
224 Montalvão et al. (2019) found that freshwater mussels, *Anadontites trapesialis*, exposed to  
225 leachate from smoked cigarette butts accumulated heavy metals in their tissues and experienced  
226 mutagenetic effects even at low environmentally relevant concentrations (<0.2 butts L<sup>-1</sup>),  
227 although the treatments were pseudo-replicated. Therefore, the response over time to sublethal  
228 toxicity may manifest in factors such as reproduction or growth performance, important for  
229 population sustainability and warrants further investigation.

230 We currently know very little about how the toxicity of cigarette butts may change over time  
231 when in the environment, but recent research indicates that butts continue to exude toxic  
232 chemicals into the air at least 1 week after being extinguished (Gong et al. 2020). Furthermore,  
233 Bonanomi et al. (2020) found that cellulose acetate cigarette butts remained toxic to the  
234 microalga *Raphidocelis subcapitata* after 5 years of degradation in the terrestrial environment.  
235 Whether or not cellulose cigarette butts also remain toxic for this length of time is unknown  
236 but should be a priority of future work in order to ascertain comparative effects of these  
237 different filter materials. International testing standards designed to evaluate the  
238 biodegradability of materials for use in cigarette butts do not test biodegradation after smoking,  
239 therefore are not environmentally realistic and when smoked, cellulose cigarette butts  
240 deposited as litter in the environment can also persist for years (Joly and Coulis 2018).

241

## 242 **Conclusion**

243 Overall, leachate from either type of butt at 5 butts L<sup>-1</sup> caused mortality of most of the  
244 individuals in the experiment. Additionally, at 1 butt L<sup>-1</sup>, both types of butt had a lethal effect  
245 on juvenile snails and reduced the activity levels of all four species of invertebrate. This

246 emphasises that, once smoked, cigarette filters, biodegradable or not, therefore are likely to  
247 have a detrimental effect on the environment due to toxins concentrated from smoking tobacco.  
248 Filters manufactured of cellulose, once smoked, can pose the same ecological threat as  
249 conventional cellulose acetate butts if they become litter in an enclosed water body such as a  
250 lake or pond. Considering their lack of rapid biodegradation in terrestrial habitats and their  
251 toxic effects in freshwater habitats, any shift to cellulose cigarette filters should be  
252 accompanied with the same plans for their appropriate post-use disposal as those made from  
253 cellulose acetate.

254

## 255 **Acknowledgements**

256 We would like to thank Anglia Ruskin University for granting Dr Green a research sabbatical  
257 and Emma Gorman for maintaining laboratory facilities at Portaferry.

258

## 259 **References**

- 260 Amos, C., Allred, A. & Zhang, L. 2017. Do Biodegradable Labels Lead to an Eco-safety  
261 Halo Effect? *J Consum Policy* 40, 279–298.
- 262 Bayley M, Baatrup E, Bjerregaard P. 1997. Woodlouse locomotor behavior in the assessment  
263 of clean and contaminated field sites. *Environ Toxicol Chem.* 16:2309–2314.
- 264 Bonanomi G, Maisto G, De Marco A, Cesarano G, Zotti M, Mazzei P, Libralato G, Staropoli  
265 A, Siciliano A, De Filippis F, La Storia A, Piccolo A, Vinale F, Crasto A, Guida M, Ercolini  
266 D, Incerti G. 2020. The fate of cigarette butts in different environments: Decay rate, chemical  
267 changes and ecotoxicity revealed by a 5-years decomposition experiment. *Environmental*  
268 *Pollution*, Volume 261,114108.

269 Booth DJ, Gribben P, Parkinson K. 2015. Impact of cigarette butt leachate on tidepool snails.  
270 Marine Pollution Bulletin, 15, 95(1):362-4.

271 Chevalier Q, Hadri HE, Petitjean P, Le Coz MB, Reynaud S, Grassl B, Gigault J. 2018.  
272 Nano-litter from cigarette butts: Environmental implications and urgent consideration,  
273 Chemosphere, 194: 125-130.

274 Dobaradaran S, Schmidt TC, Lorenzo-Parodi N, Jochmann MA, Nabipour I, Raeisi A,  
275 Stojanović N, Mahmoodi M. 2019. Cigarette butts: An overlooked source of PAHs in the  
276 environment? Environmental Pollution, 249: 932-939.

277 Gong M, Daniels N, Poppendieck D. 2020. Measurement of chemical emission rates from  
278 cigarette butts into air. Indoor Air, 00: 1– 14.

279 Green DS, Boots B, Carvalho J, Starkey T. 2019. Cigarette butts have adverse effects on  
280 initial growth of perennial ryegrass (gramineae: *Lolium perenne* L.) and white clover  
281 (leguminosae: *Trifolium repens* L.). Ecotoxicology and Environmental Safety, 182: 109418.

282 Green DS, Boots B, Sigwart J, Jiang S, Rocha C. 2016. Effects of conventional and  
283 biodegradable microplastics on a marine ecosystem engineer (*Arenicola marina*) and  
284 sediment nutrient cycling. Environmental Pollution, 208: 426-434.

285 Koutela N, Fernández E, Saru ML, Psillakis E. 2020. A comprehensive study on the leaching  
286 of metals from heated tobacco sticks and cigarettes in water and natural waters. Science of  
287 The Total Environment. 714. 136700.

288 Lawal MS, Ologundudu SO. 2013. Toxicity of cigarette filter leachates on *Hymenochirus*  
289 *curtipes* and *Clarias gariepinus* in Nigeria. J Environ Ext 11:7–14.

290 Lee W, Lee CC. 2015. Developmental toxicity of cigarette butts – An underdeveloped issue.  
291 Ecotoxicology and Environmental Safety, 113: 362-368.

292 Joly FX, Coulis M, 2018. Comparison of cellulose vs. plastic cigarette filter decomposition  
293 under distinct disposal environments. *Waste Management*, 72: 349-353.

294 McBride MJ, Guyatt AR, Kirkham AJ, Cumming G. 1984. Assessment of smoking behaviour  
295 and ventilation with cigarettes of differing nicotine yields. *Clin Sci (Lond)* 67:619–31.

296 Micevska T, Warne MS, Pablo T, Patra R. 2006. Variation in, and causes of, toxicity of CB  
297 to a cladoceran and microtox. *Arch. Environ. Contam. Toxicol.*, 50, pp. 205-212.

298 Moerman JW, Potts GE. 2011. Analysis of metals leached from smoked cigarette litter.  
299 *Tobacco Control*, 20: 30-35.

300 Mohammed A. 2013. Why are Early Life Stages of Aquatic Organisms more Sensitive to  
301 Toxicants than Adults? *New Insights into Toxicity and Drug Testing*. Gowder S (Ed). DOI:  
302 10.5772/55187 Available from: <https://www.intechopen.com/books>.

303 Montalvão MF, Chagas TQ, da Silva Alvarez TG, Mesak C, da Costa Araújo AP, Gomes  
304 AR, Vieira JE, Malafaia G. 2019. How leachates from wasted cigarette butts influence  
305 aquatic life? A case study on freshwater mussel *Anodontites trapesiali*. *Science of The Total*  
306 *Environment*, 689: 381-389.

307 Novotny TE, Slaughter E. 2014. Tobacco product waste: an environmental approach to  
308 reduce tobacco consumption. *Curr. Environ. Health Rpt*, 1: 208-216.

309 OECD (2018), "Fish, Early-Life Stage (FELS) Toxicity Test (OECD TG 210)", in Revised  
310 Guidance Document 150 on Standardised Test Guidelines for Evaluating Chemicals for  
311 Endocrine Disruption, OECD Publishing, Paris, [https://doi.org/10.1787/9789264304741-13-](https://doi.org/10.1787/9789264304741-13-en)  
312 en.

313 Oksanen J, Guillaume Blanchet F, Friendly M, Kindt R, Legendre P, McGlenn

314 D, Minchin P, O'Hara R, Simpson G, Solymos P, Stevens M, Szoecs E, Wagner H. 2019.  
315 Vegan: community Ecology Package. R Package Version 2.5-2. Available: [https://cran.r-](https://cran.r-project.org/web/packages/vegan/vegan.pdf)  
316 [project.org/web/packages/vegan/vegan.pdf](https://cran.r-project.org/web/packages/vegan/vegan.pdf).

317 Patel V, Thomson GW, Wilson N. 2013. Cigarette butt littering in city streets: a new  
318 methodology for studying and results. *Tob. Control*, 22: 59-62

319 Parker TT, Rayburn J. 2017. A comparison of electronic and traditional cigarette butt  
320 leachate on the development of *Xenopus laevis* embryos. *Toxicology Reports* 4: 77-82.

321 Pauly JL, Mepani AB, Lesses JD, Cummings KM, Streck RJ. 2002. Cigarettes with defective  
322 filters marketed for 40 years: what Philip Morris never told smokers. *Tob Control*. 2002  
323 Mar;11 Suppl 1: I51-61.

324 R Core Team. 2019. R: A language and environment for statistical computing. R Foundation  
325 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>

326 Register K. 2000. Cigarette butts as litter-toxic as well as ugly? *Bulletin of the American*  
327 *Littoral Society*, 25: 23-29.

328 Relyea R, Hoverman J. 2006. Assessing the ecology in ecotoxicology: a review and synthesis  
329 in freshwater systems. *Ecology Letters*, 9: 1157-1171.

330 Roder Green AL, Putschew A, Nehls T. 2014. Littered cigarette butts as a source of nicotine  
331 in urban waters. *Journal of Hydrology*, 519: 3466-3474.

332 Rubach MN, Crum SJH, Van den Brink PJ. 2011. Variability in the dynamics of mortality  
333 and immobility responses of freshwater arthropods exposed to chlorpyrifos. *Arch Environ*  
334 *Contam Toxicol*. 60:708–721.

335 Slaughter E, Gersberg EM, Watanabe K, Rudolph J, Stransky C, Novotny TE. 2011. Toxicity  
336 of CB, and their chemical components, to marine and freshwater fish. *Tob. Control*, 20: 25-  
337 29.

338 Strayer DL, Malcom HM. 2012. Causes of recruitment failure in freshwater mussel  
339 populations in southeastern New York. *Ecological Applications*, 22: 1780-1790.

340 Wright SL, Rowe D, Reid MJ, Thomas KV, Galloway TS. 2015. Bioaccumulation and  
341 biological effects of cigarette litter in marine worms. *Scientific Reports*, 5:14119.



342 **Tables and figures**


343 **Table 1.** Results of asymmetrical ANOVA for (a) the lethality of leachate to each species  
 344 throughout the experiment and (b) the survival of juvenile *B. tentaculata* at each time point  
 345 (from 24 to 120 h). *d.f.* = degrees of freedom, *F* = F-ratio and *p* = p-value. Significance at  $\alpha <$   
 346 0.05 and is indicated by values in **bold**.

(a)										
Source of variation	d.f.	<i>D. polymorpha</i>		<i>P. nigra</i>		<i>P. planorbis</i>		<i>B. tentaculata</i>		
		<i>F</i>	<i>p</i>	<i>F</i>	<i>p</i>	<i>F</i>	<i>p</i>	<i>F</i>	<i>p</i>	
Treatment (one way)	6	<b>8.63</b>	<b>&lt;0.001</b>	<b>15.72</b>	<b>&lt;0.001</b>	<b>4.47</b>	<b>0.003</b>	<b>10.64</b>	<b>&lt;0.001</b>	
Control vs others	1	1.93	0.175	3.72	0.064	2.27	0.143	<b>5.31</b>	<b>0.029</b>	
Butt (B)	1	1.27	0.297	2.62	0.090	0.07	0.930	0.72	0.497	
Concentration (C)	2	<b>22.37</b>	<b>&lt;0.001</b>	<b>40.85</b>	<b>&lt;0.001</b>	<b>11.97</b>	<b>&lt;0.001</b>	<b>27.82</b>	<b>&lt;0.001</b>	
B x C	2	1.27	0.297	1.85	0.176	0.24	0.791	0.72	0.497	
Residuals	52									

(b)											
Source of variation	d.f.	24 h		48 h		72 h		96 h		120 h	
		<i>F</i>	<i>p</i>	<i>F</i>	<i>p</i>	<i>F</i>	<i>p</i>	<i>F</i>	<i>p</i>	<i>F</i>	<i>p</i>
Treatment (one way)	6	<b>10.64</b>	<b>&lt;0.001</b>	<b>39.31</b>	<b>&lt;0.001</b>	<b>54.67</b>	<b>&lt;0.001</b>	<b>50.37</b>	<b>&lt;0.001</b>	<b>35.47</b>	<b>&lt;0.001</b>
Control vs others	1	<b>5.31</b>	<b>0.029</b>	<b>27.53</b>	<b>&lt;0.001</b>	<b>44.16</b>	<b>&lt;0.001</b>	<b>48.26</b>	<b>&lt;0.001</b>	<b>54.40</b>	<b>&lt;0.001</b>
Butt (B)	1	0.72	0.497	2.60	0.092	2.71	0.084	1.13	0.338	0.70	0.504
Concentration (C)	2	<b>27.82</b>	<b>&lt;0.001</b>	<b>96.10</b>	<b>&lt;0.001</b>	<b>132.88</b>	<b>&lt;0.001</b>	<b>122.19</b>	<b>&lt;0.001</b>	<b>76.21</b>	<b>&lt;0.001</b>
B x C	2	0.72	0.497	5.47	0.010	6.33	0.005	3.67	0.039	2.29	0.120
Residuals	52										


347



348


349 **Figure 1.** Heatmap showing the lethality of leachate derived from smoked cigarettes butts  
 350 made from either cellulose or cellulose acetate filters on *D. polymorpha*, *P. nigra*, *P. planorbis*  
 351 and *B. tentaculata* for each replicate mesocosm. Mortality scale is shown and is based on the  
 352 time taken for death to occur, i.e. the darker the cell, the higher the mortality in a replicate  
 353 mesocosm, with 0 the least lethal (no deaths within 120 h) and 5 the most lethal (died within 24  
 354 h).

355



356

357 **Figure 2.** Survival (%) out of 5 individual juvenile *B. tentaculata* snails in either rainwater  
 358 without leachate (Control) and leachate from 0.2, 1 or 5 cellulose, or cellulose acetate butts L<sup>-1</sup>  
 359 <sup>1</sup> at 24 (□), 48 (□), 72 (□), 96 (■) and >120 (■) hours of exposure. Data are mean ± SEM, n  
 360 = 5.



361

362 **Figure 3.** Non-metric multidimensional scaling diagram of the behaviour exhibited by all

363 species pooled over the 5 days of the experiment exposed to either no leachate (○) or to leachate

364 from 0.2 (◇) or 1 (◆) cellulose butts L<sup>-1</sup> or to 0.2 (▣) or 1 (■) cellulose acetate butts L<sup>-1</sup>.

365 Vectors are overlain for behaviours classifications correlated to the multivariate pattern at  $r >$

366 0.6. Included are results of the asymmetric PERMANOVA analysis, with associated pseudo-F

367 values and observed p-values based on 9999 permutations of the data.