



**QUEEN'S
UNIVERSITY
BELFAST**

Community structure and environmental factors affecting diatom abundance and diversity in a Mediterranean climate river system

Dalu, T., Cuthbert, R. N., Weyl, O. L. F., & Wasserman, R. J. (2021). Community structure and environmental factors affecting diatom abundance and diversity in a Mediterranean climate river system. *The Science of the Total Environment*, Article 152366. Advance online publication. <https://doi.org/10.1016/j.scitotenv.2021.152366>

Published in:

The Science of the Total Environment

Document Version:

Peer reviewed version

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2021 Elsevier.

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License

(<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

Journal Pre-proof

Community structure and environmental factors affecting diatom abundance and diversity in a Mediterranean climate river system

Tatenda Dalu, Ross N. Cuthbert, Olaf L.F. Weyl, Ryan J. Wasserman



PII: S0048-9697(21)07443-X

DOI: <https://doi.org/10.1016/j.scitotenv.2021.152366>

Reference: STOTEN 152366

To appear in: *Science of the Total Environment*

Received date: 26 October 2021

Revised date: 8 December 2021

Accepted date: 9 December 2021

Please cite this article as: T. Dalu, R.N. Cuthbert, O.L.F. Weyl, et al., Community structure and environmental factors affecting diatom abundance and diversity in a Mediterranean climate river system, *Science of the Total Environment* (2021), <https://doi.org/10.1016/j.scitotenv.2021.152366>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier B.V.

Community structure and environmental factors affecting diatom abundance and diversity in a Mediterranean climate river system

Tatenda Dalu^{a,b,c,*}, Ross N Cuthbert^{d,c,*}, Olaf LF Weyl^{c,+}, Ryan J Wasserman^{e,f}

^a *Department of Ecology and Resource Management, University of Venda, Thohoyandou 0950, South Africa*

^b *Wissenschaftskolleg zu Berlin - Institute for Advanced Study, Berlin 14193, Germany*

^c *South African Institute for Aquatic Biodiversity, Makhanda 6140, South Africa*

^d *GEOMAR Helmholtz-Zentrum für Ozeanforschung Kiel, 24105 Kiel, Germany*

^e *Department of Zoology and Entomology, Rhodes University, Makhanda 6140, South Africa*

^f *School of Science, Monash University Malaysia, Bandar Sunway, Selangor Darul Ehsan, Malaysia*

⁺ *Deceased*

**Corresponding e-mail: rossnoe.cuthbert@gmail.com (RNC); dalutatenda@yahoo.co.uk*

(TD)

Abstract

Mediterranean climate river systems are among the most threatened ecosystems worldwide, due to a long history of anthropogenic impacts and alien invasive species introductions. Many of such rivers naturally exhibit a non-perennial flow regime, with distinct seasonal, inter-annual and spatial heterogeneity. The present study seeks to detect diatom community patterns and to understand the processes that cause these structures in an Austral Mediterranean river system among different months and river sections. In general, most environmental variables showed an increasing trend downstream for both months, with the exception of pH, dissolved oxygen, PO_4^{3-} and substrate embeddedness, which decreased downstream. A total of 110 diatom species between the two study months (October – 106 taxa; January – 78 taxa) were identified, dominated by 50 species with at least >2 % abundance. Diatom community structure differed significantly across river zones, while no significant differences were observed between the study months. A boosted regression trees model showed that B (43.3 %), Cu (20.3 %), Fe (3.4 %) and water depth (3.2 %) were the most significant variables structuring diatoms. Diatom species communities reflected environmental variables (i.e., sediment and water chemistry) in this Mediterranean climate river system, as sediment metals such as B, Cu and Fe were found to be important in structuring diatom communities. Biotic influences from fish communities had little effect on diversity, but shifted diatom community structure. Therefore, the current study highlights how river systems have complex interactions that play an important role in determining diatom species composition.

Keywords: biological invasion, Diatoms, freshwater ecosystem, Krom River, Mediterranean climate, Boosted regression trees

Introduction

Rivers are important ecosystems and are home to diverse and abundant animal and plant species, providing vital ecosystem services (Soria et al., 2017; Messenger et al., 2021). Lotic freshwater systems are either perennial or temporary, depending on groundwater, surface flow and prevailing environmental conditions. Freshwater ecological communities within these systems are thus adapted to the natural variability through short life spans, mechanisms to resist or avoid desiccation, and high colonisation rates (Feio et al., 2014). Rivers are also critical for humans, being a source of freshwater, food and recreational activities, among other ecosystem services (Dunham et al., 2018). Thus, the conflict between human needs and river system function — given their continuing exposure to human impacts (e.g., considerable changes in sediment delivery, declining water quality, biodiversity loss, flow patterns, morphological modifications) — necessitates management adaptations (Dudgeon et al., 2006; Tan et al., 2017).

Mediterranean climate river systems are among the most threatened ecosystems worldwide, due to a long history of anthropogenic impacts and alien invasive species introductions (Colin et al., 2016). Water diversion, flow regulation, increased salinity, pollution and introduced species have impacted Mediterranean climate river ecosystems over time (Feio et al., 2014). Many of such rivers naturally exhibit a non-perennial flow regime, with distinct seasonal, inter-annual and spatial heterogeneity (Skoulikidis et al., 2017). That is because these systems have climatic conditions that are characterised by marked seasonal differences, such as hot-dry summers and cool-wet winters, sometimes compounded by large inter-annual rainfall variability which affects aquatic biota (Boix et al., 2010; Feio et al., 2014). These Mediterranean river communities therefore show strong inter-annual fluctuations in species richness and composition (Feio et al., 2014). Such systems often additionally have a very

complex history and environmental features that have contributed to the high levels of biodiversity and endemism currently hosted (Colin et al., 2016). Thus, organisms in these river systems are often patchily distributed in time and space, rather than homogeneously or randomly, generating considerable spatiotemporal heterogeneity (Bonada et al., 2005; Stomp et al., 2011).

Diatoms are single-celled microscopic organisms belonging to a group of algae, exhibiting a large diversity in terms of taxonomy, morphology and ecology, and are present in most aquatic and humid environments (Round et al., 1990; Liang et al., 2020). Diatoms are cosmopolitan in nature, and are known to be efficient indicators of river ecological quality (Vasselon et al., 2017; Wood et al., 2019; Mangadze et al., 2019a). Bioindicators track environmental variations at differing temporal scales, ranging from weeks (i.e., diatoms, invertebrates) to years (i.e., fish) (Colins, et al., 2016; Mangadze et al., 2019b). Diatom community variability is influenced both by natural and anthropogenic factors and species evolutionary adaptations (Stenger-Kovács et al., 2013). Diatoms are accordingly commonly used as biological indicators to assess habitat degradation in river systems, due to their sensitivity to a wide range of environmental stresses (Liu et al., 2020; Gelis et al., 2021). Their use in habitat assessments is supported by their rapid response to changes in environmental conditions, their substantial abundances, and their short generational times (Wood et al., 2019). Their indicator efficiency also relies on their marked taxonomical diversity and their species-specific ecological preferences to different levels of habitat degradation (Pandey et al., 2018; Tan et al., 2017, 2021). However, knowledge gaps in their responses to key environmental variables remain in the Global South and under particular climatic regimes, such as in changeable Mediterranean river systems, hampering understandings of their broad scale efficacy in habitat assessments. As such, it important to

analyse diatom species composition in response to environmental factors associated with anthropogenic impacts for better management and protection of aquatic environments. For example, light, nutrients and environmental properties, such as pH and ionic composition, are essential abiotic factors in river ecosystems and their extent and/or availability change spatiotemporally, potentially affecting diatom communities (Stenger-Kovács et al., 2013; Mangadze et al., 2016).

Biotic factors may also be pertinent in influencing diatom communities. In river ecosystems, top predators such as fish often influence primary producer (i.e., diatom (phytoplankton)) abundance and diversity through trophic interactions (Vanni and Lynn, 1997; Sommer et al., 2012). Fish-driven trophic cascades can influence primary producers, but with varying strengths (Stevenson et al., 2016; do Nascimento Filho and do Nascimento Moura, 2021) and may arise due to biological invasion (Weyl and Ellender, 2014). For example, diatom community structure is often negatively affected by fish directly or indirectly via reduced abundance and diversity when planktivorous and/or herbivorous fish are abundant. However, diatoms could be positively affected if predatory fish feed on lower-order planktivorous and/or herbivorous fish, reducing their abundance and thus feeding rates on diatoms in consequence (Wasserman et al., 2013). Fish species confer effects through many food web pathways; however, quantifying the importance of these direct and indirect effects is important to understand diatom community structuring and inferences of ecosystem health (Polis et al. 1996; Vanni and Lynn, 1997). Therefore, in addition to abiotic factors, diatom species composition is also affected by upper trophic levels, with studies elucidating diatom community structuring by fish fundamental to our understanding of river ecosystem functioning.

The present study seeks to detect diatom community patterns and to understand the processes that cause these structures in an austral Mediterranean river system among different months and river sections, characterized by variable physico-chemical parameters and dominant fish species. Besides the high conservation value of these systems, biodiversity is related to the ecosystem services that these systems provide to society, as well as their ability to cope with multiple stressors. Thus, we expect the abundance and presence of diatoms to be a result of several multi-scale filters, such as the ecological constraints ranging from the landscape to micro-habitat scale (Bonada et al., 2005).

Materials and methods

Study area

The Krom River system is a tributary of the O'ifants-Doring River system in the Western Cape Province of South Africa. The river starts at an elevation of ~1500 m in the Cederberg Mountains and flows in an easterly direction for approximately 20 km before joining the Matjies River, a tributary of the Doring River (Fig. 1). The upper river section is narrow (width <5 m) and shallow (depth <1 m), comprising of several pools and bedrock steps interspersed by occasional cobble-bed riffles (Marr et al., 2019). The lower river section has deep pools (>3 m), and is isolated during low flow periods (i.e., mid to late summer). The area falls within the Mediterranean climate region, characterised by cold, wet winters and warm, dry summers (Cowling and Holmes, 1992). The mean annual rainfall is estimated at 1000 mm, 500 mm and 250 mm, respectively for the upper, middle and lower reaches. The geology of the river channel is predominantly quartzite sandstones, with minor shale bands of the Nardouw Subgroup, Table Mountain Group and Cape Supergroup (Thamm and Johnson, 2006). The vegetation is dominated by the Cederberg Sandstone Fynbos above the Krom River Cederberg Park, before changing to the northern inland shale band vegetation, just east

of a farm complex, and lastly the arid extreme of Cederberg Sandstone Fynbos (Mucina and Rutherford, 2006).

Sampling was conducted in October 2017 and January 2018 (i.e., two seasons). Twenty-two sites were selected on the Krom River system along a zonation of dominant fish presence into three sections i.e., invasive rainbow trout *Oncorhynchus mykiss* (sites K1–K10), native Clanwilliam rock catfish *Austroglanis gilli* (sites K11–K16) and native sawfin *Cheilobarbus serra* (sites K17–K22) zones. It is important to note that in the different fish zones highlighted above, these fish species were the most dominant between both study months in previous studies (Marr et al., 2019), however, other fish species are also present in each zone, such as the invasive bluegill sunfish *Lepomis macrochirus*. Sites K16 – K18 are located within the Cederberg Park camp area and site K22 at the confluence with the Matjies River (Fig. 1; Marr et al., 2019). Approximately 17 km above the Krom River, Cederberg Park is near-pristine and unaffected by human impacts. The river then flows through highly transformed Krom River Cederberg Park, where major abstraction from the Krom River and water diversion to several small reservoirs for irrigation and recreational purposes occurs. Below the park area, the river becomes more seasonal and flows through the near-pristine Matjies River Nature Reserve before joining the Matjies River system.

Environmental variables sampling and analysis

The key water parameters of conductivity, salinity, pH, temperature and total dissolved solids (TDS) were measured per site and each season using a portable probe PCTestr 35 (Eutech/Oakton Instruments, Singapore). The dissolved oxygen (DO) and turbidity were measured using Per Scientific DO (DO 850045, Taiwan) and turbidity (AL250T-IR, Aqualytic, Germany) meters, respectively. Two 500 mL water samples were collected at each

site and season for analysis of nutrients (i.e., ammonium – NH_4^+ , phosphates – PO_4^{3-}) via in-field laboratory analyses within 8 hours of collection using a HI 83203 multiparameter photometer (Hanna Instruments Inc., Rhode Island). The key habitat characteristics of substrate embeddedness, riparian cover, channel width, water depth and detrital cover, were determined according to Dalu et al. (2017a).

Integrated 1.5 kg sediment samples ($n = 2$) were collected from each of the 22 study sites for each sampling event using acid-washed wooden splints and placed in new, labelled ziplock bags for further laboratory analysis. In the laboratory, all sediment samples were oven-dried at 60 °C for 48–72 hrs before disaggregation and sieving using 0.05 mm mesh size to remove organic matter and large stones. All sediment samples were sent for analysis at BEMLAB, a South African National Accreditation System (SANS) certified laboratory. The following were analysed: (i) cation elements: boron (B), calcium (Ca), potassium (K), magnesium (Mg), sodium (Na), (ii) heavy metal: manganese (Mn), copper (Cu), iron (Fe), zinc (Zn) and nutrients: nitrogen (N) and phosphorus (P) (see Dalu et al. 2020a,b for detailed methods).

Diatom sampling and analysis

Diatom sample collection and analysis were conducted according to Taylor et al. (2005) and Dalu et al. (2020c) by sampling 10 random rocks and/or pebbles per site and season. The collected samples were processed using the hot hydrochloric acid and potassium permanganate method, with diatoms counted (i.e., 300–650 valves per sample) and identified under a phase-contrast Olympus CX light microscope to their lowest taxonomic level using identification guides from Taylor et al. (2007). The following diversity indices were computed: taxa richness, evenness, Shannon-Weiner, and Simpson's indices to assess variations in community structure among sites, zones and seasons.

Data analyses

Prior to multivariate analyses of environmental data, a test for homogeneity of variance and data normality was conducted and the data were found to violate assumptions of a parametric test. Hence, a non-parametric Kruskal-Wallis test was conducted to assess for differences in environmental variables and diversity indices (i.e., taxa richness, evenness, Shannon-Weiner, Simpson's) data between seasons (i.e., October and January) and river zones (i.e., rainbow trout, rock catfish and sawfish zone) using SPSS version 16.0 (SPSS Inc., 2007). Distance-based Permutational Analysis of Variance (PERMANOVA; Anderson, 2001) based on Bray-Curtis and Euclidean distance dissimilarities and 9999 permutations with Monte Carlo tests were performed to analyse differences in diatom communities data among months and river sections using PERMANOVA+ for PRIMER version 6 (Anderson et al., 2008).

Differences in environmental variables and diatom species composition between months and zones were tested using a two-way analysis of similarity (ANOSIM) with 9999 permutations using PRIMER v6 add-on package PERMANOVA+ (Anderson et al., 2008; Clarke, 1993). The ANOSIM statistic R was based on the difference of mean ranks between groups and within groups. It varies between -1 and $+1$, with value 0 indicating random grouping.

We further employed a Boosted Regression Tree (BRT) model to assess the relationships between environmental variables and diatom species richness. The BRT uses two algorithms, namely boosting and regression (Schapire, 2003; Elith et al., 2008). The fitting of multiple trees in BRT overcomes the biggest drawback of single tree models, and the BRT model was carried out in the R statistical package (R Core Team, 2020), using the BRT extension `gbm`

package (i.e., `gbm.step` and `gbm.simplify` function) (Ridgeway, 2006; Elith et al., 2008). A Type I error of 0.05 was used in all analyses.

Results

Environmental variables

In general, most environmental variables showed an increasing trend downstream for both months, with the exception of pH, dissolved oxygen, PO_4^{3-} and substrate embeddedness, which decreased downstream (Table 1). Most of the water and habitat characteristics (i.e., temperature, pH, DO, conductivity, TDS, salinity, turbidity, riparian cover, detrital cover, water depth, substrate embeddedness, NH_4^+) and several sediment variables (i.e., N, Mg, Na, Zn) were found to be significantly different (Kruskal Wallis (K-W), $p < 0.05$) among the river zones, whereas significant seasonal differences (K-W, $p < 0.05$) were observed for water and habitat characteristics (i.e., temperature, pH, conductivity, salinity, turbidity, water depth, riparian cover, detrital cover, PO_4^{3-}) and several sediment variables (i.e., K, Ca, Mn, Cu, B) (Table 2). Sediment P was below detection limit among all sites and seasons.

Diatom community structure and relationship with environmental variables

A total of 110 diatom species between the two study months (October – 106 taxa; January – 78 taxa) were identified, dominated by 30 species with at least >2 % abundance (Tables 3, S1). The most common taxa were *Diatoma vulgare*, *Eunotia formica*, *Eunotia minor*, *Navicula radiosa*, *Pinnularia viridiformis* and *Tabellaria flocculosa*. Based on PERMANOVA analysis, significant differences were observed in diatom community structure across the river zones (Pseudo-F = 3.527, $p = 0.0001$) and months (Pseudo-F = 3.527, $p = 0.0001$), with significant pair-wise differences being observed between the rainbow trout and rock catfish zones ($t = 1.594$, $p = 0.0034$). The sawfish zone had the

highest taxa richness, with diatom diversity indices being relatively high in the rock catfish zone (Table 3). No significant differences (K-W, $p > 0.05$) were observed among the river zones (all diversity indices) and for Simpson's diversity index between seasons (K-W, $H = 2.511$, $p = 0.113$), with significant seasonal differences being observed for taxa richness (K-W, $H = 31.715$, $p < 0.001$), Shannon-Weiner (K-W, $H = 13.067$, $p < 0.001$) and Evenness (K-W, $H = 21.819$, $p < 0.001$) diversity indices.

Based on the ANOSIM results, diatom community structure differed significantly across river zones ($R = 0.245$, $p = 0.010$), while no significant differences were observed between the study months ($R = 0.030$, $p = 0.283$). No significant differences ($p > 0.05$) were, however, observed in diatom community structure based on pair-wise comparisons among the different river zones. The boosted regression trees (BRT) model predictive performance relating diatom species richness to environment variables was as follows: 3780 trees fitted (boosted models); 53.4 and 0.09 total deviance and mean residual of the model, respectively; 7.18 ± 2.23 estimated cross-validated (CV) residual deviance \pm standard error; and CV correlation of 0.94 ± 0.02 . According to the BRT model, all variables were retained as significant terms relating diatom species richness to environment variables, with the first eleven variables (i.e., B (43.3 %), Cu (20.8 %), Fe (3.4 %), water depth (3.2 %), temperature (3.0 %), PO_4^{3-} (2.9 %), Mn (2.6 %), detrital cover (2.6 %), Zn (2.4 %), Na (2.3 %), NH_4^+ (1.9 %)) accounting for ~90 % of the species richness variation, with B and Cu concentrations being the most significant (Table 4). Relationships fitted by the BRT models for the environmental variables and diatom species richness are highlighted in Fig. 2. Predicted species richness had an approximately sigmoid shaped (i.e. positive) relationship with B concentration, with high values occurring over a wide range characterised by low water depth, Zn, Mn and Fe, and high Cu and detrital cover. A more muted variation occurred in relation to water temperature,

Zn, Mn, detrital cover and Fe concentrations. (Fig. 2). A generally high diatom species richness was predicted for seasonal sites with high Cu, detrital cover and phosphates.

Discussion

The current study demonstrated that the Krom River diatom community was largely structured by sediment chemistry variables, particularly B, Cu and Fe which accounted for 67.5 % of the variation. This provides valuable information on factors influencing diatom composition within Mediterranean climate river systems in the Global South. Moreover, the monthly diatom composition differed due to underlying environmental variables, but also potentially as a result of the different fish species present in each zone and their respective feeding preferences. The present study therefore showed the existence of a strong spatial and temporal component in diatom community structuring, but with no distinctly different diatom diversities among river zones, as highlighted by our analyses. The sediment chemistry variables were generally low, as expected due to the near pristine environment where the river is located and the underlying area geology (Thamm and Johnson, 2006). A number of diatom species were common among zones and seasons, but there were differences in their frequencies, potentially due to fish community composition that could have exerted a top-down effect on the system. Our results are similar to do Nascimento Filho and do Nascimento Moura (2021), in the respect that rainbow trout, rockfish and sawfish had a strong predatory role affecting macroinvertebrates, which might have affected diatom community composition and abundances through cascade effects. A lack of differences among zones considering diversity indices suggests, however, that the effects of these alien and native fish species are consistent with regards to the richness and evenness of diatom communities.

Macroinvertebrates have been shown to efficiently reduce diatom species, especially in the

absence of top predators which consume them (Mermillod-Blondin et al., 2020; Vanni and Layne, 2021).

While diatom community structure differed significantly, certain similarities in diatom diversity could be due to differences in structure (i.e., species composition, but similar diversity levels therein). As a consequence, the indicator taxa for the near-natural streams proved to be type-specific. Approximately 110 diatom species were observed, which were dominated by weakly or poor-to-moderate electrolyte species, such as *Eunotia formica*, *Diatoma vulgare*, *Eunotia rhomboidea* and *Tabellaria flocculosa*. This was further supported by the weak ion and conductivity concentrations recorded within the study sites and several other studies (see Table 1; De Fabricius et al., 2003; Taylor et al., 2007). Diatom species diversity was relatively similar across river zones, but was significantly different between the study months, with lower species richness during the January month in particular. This could be attributed to changes water levels which caused changes in the water and sediment chemistry. Death et al. (2009), Boivie et al. (2010) and Dalu et al. (2016) highlighted that hydrological alterations accounted for ecosystem community composition, structure and functioning changes.

Diatom development depends on various environmental variables such as metals. However, when these metal micronutrients are present at an optimal level, diatoms exhibit a strong fitness and grow at their fastest rate (Masmoudi et al., 2013). Boosted regression tree results indicated a highly predictable relationship between diatom species richness and environmental (i.e., B, Cu, Fe, water depth, temperature) variables. Sediment B concentration was the strongest predictor of diatom species richness, with predicted richness peaking at a concentration $\sim 30 \text{ mg kg}^{-1}$. A positive association between diatom species richness and other

environmental variables (i.e., water depth, Mn and Zn concentration) was also observed, with maximum diatom richness being recorded at sites with low water depth, Zn, Fe and Mn concentrations, and high detrital cover, Cu and PO_4^{3-} concentrations. Lewin (1966) and Loomis and Durst (1992) highlighted that all diatoms require B, as do all higher plants, and it is thus considered as an essential nutrient for diatom growth (i.e., cell wall structure and function) and physiological needs. Thus, B has been shown to play an important role in the general structure and function of diatom plasma membranes, where it influences membrane domain stability through binding with the glycoproteins, glycolipids and phosphoinositides (Brown et al., 2002).

Similarly, diatoms require small amounts of metal ions, such as Ca, Cu, Fe, K, Mg, Mn and Zn, which are necessary for development and growth, and hence are crucial for cell physiology (Masmoudi et al., 2013). Studies have shown that *Cyclotella* spp. and *Fragilaria* spp. disappeared due to Cu pollution, whereas other species presented various deformations in response to Cu and Fe. In contrast, *Achnanthes* spp. were able to accumulate and tolerate Cu over time (Ruggiu et al., 1998; Lavoie et al., 2017; Moreno et al., 2020). Water temperature was found to be a further significant factor affecting diatom community structuring within the Krom River system across seasons and/or river sections. With monthly changes, there were significant zonal and seasonal differences in riparian cover and water temperature, with the riparian cover reducing in January and leading to increased water temperatures. Ingebrigtsen et al. (2016) and Xiao et al. (2018) highlighted that increased light intensity and water temperatures normally result in changes to diatom species composition, which could explain the differences in species richness observed here seasonally.

In conclusion, diatom species communities reflected environmental variables (i.e., sediment and water chemistry) in this Mediterranean climate river system, as sediment metals such as B, Cu and Fe were found to be important in structuring diatom communities. Biotic influences from fish communities had little effect on diversity, but shifted community structure. Therefore, the current study highlighted how Mediterranean river systems in the Global South have complex interactions that played an important role in determining the diatom species composition. More studies are required to fully understand how environmental variables and fish compositions shape diatom communities in these systems, with the aim to improve biomonitoring efforts.

Acknowledgements

We acknowledge the funding of the Water Research Commission Project K5–2538 for the execution of this project. We acknowledge the use of infrastructure and equipment provided by the SAIAB Research Platform and the funding channelled through the NRF–SAIAB Institutional Support system as well as support from the National Research Foundation – South African Research Chairs Initiative of the Department of Science and Technology (Grant No. 110507). TD and P.M.C acknowledge funding from the Wissenschaftskolleg zu Berlin Institute for Advanced Study/Stellenbosch Institute for Advanced Study and Alexander von Humboldt Foundation, respectively. Any opinion, finding and conclusion or recommendation expressed in this material is that of the author(s) and the NRF and WRC do not accept any liability in this regard. We thank Lubabalo Mofu, Dumisani Khosa, Casey Broom, Tomas Maiztegui and Jeff Hean for assisting with fieldwork.

Authors' contributions

TD: Conceptualization, Investigation, Methodology, Data curation, Formal analysis, Funding acquisition, Writing – original draft, review & editing.

RNC: Methodology, Writing – original draft, review & editing.

RJW: Conceptualization, Funding acquisition, Writing – original draft, review & editing.

OLFW: Supervision, Funding acquisition, Project administration.

Declarations and competing interests

All authors declare no conflict or financial interests exist for the manuscript.

Ethics approval

No ethical approval was required since research involved lower order plants i.e., algae (diatoms).

References

- Anderson M.J., Gorley R.N., Clarke, K.R., 2008. PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods. PRIMER-E, Plymouth, UK.
- Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, 26, 32–46.
- Anderson, M.J., Gorley, R.N., Clarke, K.R., 2008. PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods. PRIMER-E, Plymouth, UK.
- Boix, D., García-Berthou, E., Gascón, S., Benejam, L., Tornés, E., Sala, J., Benito, J., Munné, A., Solà, C., Sabater, S., 2010. Response of community structure to sustained drought in Mediterranean rivers. *Journal of Hydrology*, 383, 135–146.
- Boix, D., García-Berthou, E., Gascón, S., Benejam, L., Tornés, E., Sala, J., Benito, J., Munné, A., Solà, C., Sabater, S., 2010. Response of community structure to sustained drought in Mediterranean rivers. *Journal of Hydrology*, 383, 135–146.
- Bonada, N., Zamora-Muñoz, C., Rieradevall, M. and Prat, N., 2005. Ecological and historical filters constraining spatial caddisfly distribution in Mediterranean rivers. *Freshwater Biology*, 50, 781–797.
- Brown, P.H., Bellaloui, N., Wimmer, M.A., Bassil, E.S., Ruiz, J., Hu, H., Pfeffer, H., Dannel, F. and Römheld, V., 2002. Boron in plant biology. *Plant Biology*, 4, 205–223.
- Colin, N., Porte, C., Fernandes, D., Barata, C., Padrós, F., Carrassón, M., Monroy, M., Cano-Rocabayera, O., De Sostoa, A., Piña, B. and Maceda-Veiga, A., 2016. Ecological relevance of biomarkers in monitoring studies of macroinvertebrates and fish in Mediterranean rivers. *Science of the Total Environment*, 540, 307–323.
- Cowling, R.M., Holmes, P.M. (1992) Flora and vegetation. In: Cowling, R.M. (ed) *The Ecology of Fynbos: Nutrients, Fire and Diversity*. Oxford University Press, Cape Town.

- Dalu, T., Cuthbert, R.N., Taylor, J.C., Magoro, M.L., Weyl, O.L., Froneman, P.W., Wasserman, R.J., 2020c. Benthic diatom-based indices and isotopic biomonitoring of nitrogen pollution in a warm temperate Austral river system. *Science of the Total Environment*, 748, 142452.
- Dalu, T., Froneman, P.W., Bere, T., 2016. Assessment of water quality based on diatom indices in a small temperate river system, Kowie River, South Africa. *Water SA*, 42, pp.183–193.
- Dalu, T., Murudi, T., Dondofema, F., Wasserman, R.J., Cheri, L.D., Murungweni, F.M., Cuthbert, R.N., 2020b. Balloon milkweed *Gomphocarpus physocarpus* distribution and drivers in an internationally protected wetland. *BioInvasions Records*, 9, 627–641.
- Dalu, T., Tshivhase, R., Cuthbert, R.N., Murungweni, F.M. and Wasserman, R.J., 2020a. Metal distribution and sediment quality variation across sediment depths of a subtropical Ramsar declared wetland. *Water*, 12, 2779.
- Dalu, T., Wasserman, R.J., Tonkin, J.D., Alexander, M.E., Dalu, M.T., Motitsoe, S.N., Manungo, K.I., Bepe, O., Dube, T., 2017a. Assessing drivers of benthic macroinvertebrate community structure in African highland streams: an exploration using multivariate analysis. *Science of the Total Environment*, 601, 1340–1348.
- Dalu, T., Wasserman, R.J., Tonkin, J.D., Mwedzi, T., Magoro, M.L. and Weyl, O.L., 2017b. Water or sediment? Partitioning the role of water column and sediment chemistry as drivers of macroinvertebrate communities in an austral South African stream. *Science of The Total Environment*, 607, 317–325.
- De Fabricius, A.L.M., Maidana, N., Gómez, N., Sabater, S., 2003. Distribution patterns of benthic diatoms in a Pampean river exposed to seasonal floods: the Cuarto River (Argentina). *Biodiversity and Conservation*, 12, 2443–2454.

- Death, R.G., Dewson, Z.S., James, A.B.W., 2009. Is the structure or function a better measure of the effects of water abstraction on ecosystem integrity? *Freshwater Biology*, 54, 2037–2050.
- do Nascimento Filho, S.L., do Nascimento Moura, A., 2021. Strong top–down effects of omnivorous fish and macroinvertebrates on periphytic algae and macrophytes in a tropical reservoir. *Aquatic Ecology*, 55, 667–680.
- Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carre, G., Garcia Marquez, J.R., Gruber, B., Lafourcade, B., Leitao, P.J., Muenkemueler, N., McClean, C., Osborne, P.E., Reineking, B., Schroeder, B., Skidmore, A.K., Zureli, D., Lautenbach, S., 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography*, 36, 27–46.
- Dudgeon, D., Arthington, A.H., Gessner, M.O., Kawabata, Z.I., Knowler, D.J., Lévêque, C., Naiman, R.J., Prieur–Richard, A.H., Soto, D., Stiassny, M.L., Sullivan, C.A., 2006. Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews*, 81, 163–182.
- Dunham, J.B., Angermeier, P.L., Crausbay, S.D., Cravens, A.E., Gosnell, H., McEvoy, J., Moritz, M.A., Raheem, N., Sanford, T., 2018. Rivers are social–ecological systems: Time to integrate human dimensions into riverscape ecology and management. *Wiley Interdisciplinary Reviews: Water*, 5, e1291.
- Elith, J., Leathwick, J.R., Hastie, T., 2008. A working guide to boosted regression trees. *Journal of Animal Ecology*, 77, 802–813.
- Ellender, B.R., Weyl, O.L.F., 2014. A review of current knowledge, risk and ecological impacts associated with non–native freshwater fish introductions in South Africa. *Aquatic Invasions*, 9, 117–132.

- Feio, M.J., Aguiar, F.C., Almeida, S.F.P., Ferreira, J., Ferreira, M.T., Elias, C., Serra, S.R.Q., Buffagni, A., Cambra, J., Chauvin, C., Delmas, F., 2014. Least disturbed condition for European Mediterranean rivers. *Science of the Total Environment*, 476, 745–756.
- Gelis, M.M.N., Cochero, J., Donadelli, J., Gómez, N., 2020. Exploring the use of nuclear alterations, motility and ecological guilds in epipelagic diatoms as biomonitoring tools for water quality improvement in urban impacted lowland streams. *Ecological Indicators*, 110, 105951.
- Gotelli, N.J., Colwell, R.K., 2001. Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters*, 4, 379–391.
- Ingebrigtsen, R.A., Hansen, E., Andersen, J.H., Eilertsen, N.C., 2016. Light and temperature effects on bioactivity in diatoms. *Journal of Applied Phycology*, 28, 939–950.
- Lavoie, I., Hamilton, P.B., Morin, S., Tiam, S.K., Kahlert, M., Gonçalves, S., Falasco, E., Fortin, C., Gontero, B., Heudre, D., Kivaićinović–Sirinelli, M., 2017. Diatom teratologies as biomarkers of contamination: are all deformities ecologically meaningful? *Ecological Indicators*, 82, 539–550.
- Lewin, J., 1966. Boron as a growth requirement for diatoms. *Journal of Phycology*, 2, 160–163.
- Liang, J., Huang, C., Stevenson, M.A., Qiao, Q., Zeng, L., Chen, X., 2020. Changes in summer diatom composition and water quality in urban lakes within a metropolitan area in central China. *International Review of Hydrobiology*, 105, 94–105.
- Liu, B., Chen, S., Liu, H., Guan, Y., 2020. Changes in the ratio of benthic to planktonic diatoms to eutrophication status of Muskegon Lake through time: Implications for a valuable indicator on water quality. *Ecological Indicators*, 114, 106284.
- Loomis, W.D., Durst, R.W., 1992. Chemistry and biology of boron. *BioFactors*, 3, 229–239.

- Mangadze, T., Dalu, T., Froneman, P.W., 2019a. Biological monitoring in southern Africa: a review of the current status, challenges and future prospects. *Science of the Total Environment*, 648, 1492–1499.
- Mangadze, T., Taylor, J.C., Froneman, W.P., Dalu, T., 2019b. Water quality assessment in a small austral temperate river system (Bloukrans River system, South Africa): Application of multivariate analysis and diatom indices. *South African Journal of Botany*, 125, 353–359.
- Mangadze, T., Wasserman, R.J., Dalu, T., 2017. Use of diatom communities as indicators of conductivity and ionic composition in a small Austral temperate river system. *Water, Air, and Soil Pollution*, 228, 1–11.
- Marr, S.M., Bellingan, T.A., Hugo, S., Shelton, J.M., Impson, N.D., Gouws, J., Weyl, O.L.F., 2019. Rotenone policy support and capacity development. Part 2: Krom River base-line monitoring and capacity development. WRC Report TT780/2/19. Water Research Commission, Pretoria.
- Masmoudi, S., Nguyen-Deroche, N., Caruso, A., Ayadi, H., Morant-Manceau, A., Tremblin, G., 2013. Cadmium, copper, sodium and zinc effects on diatoms: from heaven to hell—a review. *Cryptogamie, Algologie*, 34, 185–225.
- Mermillod-Blondin, F., Marmonier, P., Tenaille, M., Lemoine, D.G., Lafont, M., Vander Vorste, R., Simon, L., Volatier, L., 2020. Bottom-up processes control benthic macroinvertebrate communities and food web structure of fishless artificial wetlands. *Aquatic Ecology*, 54, 575–589.
- Messenger, M.L., Lehner, B., Cockburn, C., Lamouroux, N., Pella, H., Snelder, T., Tockner, K., Trautmann, T., Watt, C., Datry, T., 2021. Global prevalence of non-perennial rivers and streams. *Nature*, 594, 391–397.

- Moreno, C.M., Gong, W., Cohen, N.R., DeLong, K., Marchetti, A., 2020. Interactive effects of iron and light limitation on the molecular physiology of the Southern Ocean diatom *Fragilariopsis kerguelensis*. *Limnology and Oceanography*, 65, 1511–1531.
- Mucina, L., Rutherford, M.C., 2006. *The Vegetation of South Africa, Lesotho and Swaziland*. South African National Biodiversity Institute, Pretoria.
- Pandey, L.K., Lavoie, I., Morin, S., Park, J., Lyu, J., Choi, S., Lee, H., Han, T., 2018. River water quality assessment based on a multi-descriptor approach including chemistry, diatom assemblage structure, and non-taxonomical diatom metrics. *Ecological Indicators*, 84, 140–151.
- Pappas, J.L., Stoermer, E.F., 1996. Formulation of a method to count number of individuals' representative of number of species in algal communities. *Journal of Phycology* 32, 693–196.
- Polis, G.A., Holt, R.D., Menge, B.A., Winemiller, K.O., 1996. Time, space and life history: influences on food webs. In: Polis, G.A., Winemiller, K.O. (eds). *Food Webs: Integration of Patterns and Dynamics*. Chapman and Hall, New York, New York.
- R Core Team., 2020. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Ridgeway, G., 2006. Generalized boosted regression models. Documentation on the R package 'gbm', version 1.5–7. Available at: <http://www.ipensieri.com/gregr/gbm.shtml>.
- Round, F.E., Crawford, R.M., Mann, D.G., 1990. *Diatoms: Biology and Morphology of the Genera*. Cambridge University Press, Cambridge.
- Ruggiu, D., Luglie, A., Cattaneo, A., Panzani, P., 1998. Paleoecological evidence for diatom response to metal pollution in Lake Orta (N. Italy). *Journal of Paleolimnology*, 20, 333–345.

- Schapiro, R.E., 2003. The boosting approach to machine learning: an overview. In: Denison, D.D., Hansen, M.H., Holmes, C.C., Mallick, B., Yu, B. (eds) *Nonlinear Estimation and Classification*. Springer Science + Business Media, LLC, New York.
- Skoulikidis, N.T., Sabater, S., Datry, T., Morais, M.M., Buffagni, A., Dörflinger, G., Zogaris, S., del Mar Sánchez–Montoya, M., Bonada, N., Kalogianni, E., Rosado, J., 2017. Non-perennial Mediterranean rivers in Europe: status, pressures, and challenges for research and management. *Science of the Total Environment*, 577, 1–18.
- Šmilauer, P., Lepš, J., 2014. *Multivariate Analysis of Ecological Data Using CANOCO 5*. Cambridge University Press, Cambridge.
- Sommer, U., Adrian, R., De Senerpont Domis, L., Elser, J.J., Gaedke, U., Ibelings, B., Jeppesen, E., Lüring, M., Molinero, J.C., Mooij, W.M., Van Donk, E., 2012. Beyond the Plankton Ecology Group (PEG) model: mechanisms driving plankton succession. *Annual review of ecology, evolution, and systematics*, 43, 429–448.
- Soria, M., Leigh, C., Datry, T., Bini, L.M., Bonada, N., 2017. Biodiversity in perennial and intermittent rivers: A meta-analysis. *Oikos*, 126, 1078–1089.
- SPSS Inc., 2007. *SPSS Release 16.0.0 for Windows*. Polar Engineering and Consulting. SPSS Inc., Chicago (IL).
- Stenger–Kovács, C., Lengyel, E., Crossetti, L.O., Üveges, V., Padisák, J., 2013. Diatom ecological guilds as indicators of temporally changing stressors and disturbances in the small Torna–stream, Hungary. *Ecological Indicators*, 24, 138–147.
- Stevenson, C.F., Demes, K.W., Salomon, A.K., 2016. Accounting for size–specific predation improves our ability to predict the strength of a trophic cascade. *Ecology and Evolution*, 6, 1041–1053.
- Stomp, M., Huisman, J., Mittelbach, G.G., Litchman, E., Klausmeier, C.A., 2011. Large-scale biodiversity patterns in freshwater phytoplankton. *Ecology*, 92, 2096–2107.

- Tan, X., Zhang, Q., Burford, M.A., Sheldon, F., Bunn, S.E., 2017. Benthic diatom based indices for water quality assessment in two subtropical streams. *Frontiers in Microbiology* 8, 601.
- Tan, X., Liu, Y., Burford, M.A., Zhang, Q., 2021. The performance of diatom indices in assessing temporal changes in water quality in a large lowland river ecosystem. *River Research and Applications*, 37, 423–432.
- Taylor, J.C., Harding, W.R., Archibald, C.G.M., 2005. A Methods Manual for the Collection, Preparation and Analysis of Diatom Samples. Version 1.0. WRC Report No. TT 281/07. Water Research Commission, Pretoria.
- Taylor, J.C., Harding, W.R., Archibald, C.G.M., 2007. An Illustrated Guide to some Common Diatom Species from South Africa. WRC Report No. TT 282/07. Water Research Commission, Pretoria.
- ter Braak, C.J.F., Šmilauer, P., 2002. *CANOCO Reference Manual and Cano Draw for Windows User's Guide: Software for Canonical Community Ordination (version4.5)*. Microcomputer Power, Ithaca.
- Van Dam, H., Mertens, A., Simenstad, J., 1994. A coded checklist and ecological indicator values of freshwater diatoms from the Netherlands. *Netherlands Journal of Aquatic Ecology*, 28, 117–152.
- Vanni, M.J., Layne, C.D., 1997. Nutrient recycling and herbivory as mechanisms in the “top-down” effect of fish on algae in lakes. *Ecology*, 78, 21–40.
- Vasselon, V., Rimet, F., Tapolczai, K., Bouchez, A., 2017. Assessing ecological status with diatoms DNA metabarcoding: Scaling-up on a WFD monitoring network (Mayotte island, France). *Ecological Indicators*, 82, 1–12.

Wasserman, R.J., Noyon, M., Avery, T.S., Froneman, P.W. 2013. Trophic level stability inducing effects of predaceous early juvenile fish in an estuarine mesocosms study. PLOS ONE 8, e61019. doi:10.1371/journal.pone.0061019

Wood, R.J., Mitrovic, S.M., Lim, R.P., Warne, M.S.J., Dunlop, J., Kefford, B.J., 2019.

Benthic diatoms as indicators of herbicide toxicity in rivers – A new SPEcies At Risk (SPEARherbicides) index. *Ecological Indicators*, 99, 203–213.

Xiao, W., Liu, X., Irwin, A.J., Laws, E.A., Wang, L., Chen, B., Zeng, Y., Huang, B., 2018.

Warming and eutrophication combine to restructure diatoms and dinoflagellates. *Water Research*, 128, 206–216.

Yoccoz, N.G., 1991. Use, overuse, and misuse of significance tests in evolutionary biology and ecology. *The Bulletin of the Ecological Society of America*, 72, 106–111.

Figures




Fig. 1. Location of the study sites in the Krom River, Cederberg Mountain region, South Africa




Fig. 2. Variation in diatom species richness predicted by a boosted regression trees (BRT) model for the first 9 environmental variables, and using environmental characteristics as predictors

Tables

Table 1. Mean (\pm standard deviation) of environmental variables measured from the KromRiver system, South Africa. Abbreviations: TDS – total dissolved solids, PO_4^{3-} – phosphates, NH_4^+ – ammonium

Variable	Units	October			January		
		Rainbow trout	Rock catfish	Sawfish	Rainbow trout	Rock catfish	Sawfish
<i>Water and habitat characteristics</i>							
Temperature	$^{\circ}\text{C}$	15.1 \pm 0.4	16.3 \pm 0.8	18.0 \pm 0.2	18.5 \pm 0.6	19.6 \pm 0.3	23.2 \pm 0.4
pH		6.8 \pm 0.2	6.7 \pm 0.6	6.3 \pm 0.3	6.1 \pm 0.5	6.9 \pm 0.3	6.5 \pm 0.1
Dissolved oxygen	mg L^{-1}	7.7 \pm 0.2	7.5 \pm 0.3	7.0 \pm 0.2	7.2 \pm 0.6	7.0 \pm 0.7	5.9 \pm 1.2
Conductivity	$\mu\text{S cm}^{-1}$	27.5 \pm 8.1	30.7 \pm 8.2	52.6 \pm 29.9	31.3 \pm 5.6	37.4 \pm 6.3	117.1 \pm 5.0
TDS	mg L^{-1}	19.5 \pm 5.6	23.1 \pm 5.3	37.4 \pm 21.2	19.7 \pm 3.4	26.2 \pm 3.9	81.4 \pm 3.6
Salinity	ppm	0.02 \pm 0.003	0.02 \pm 0.002	0.03 \pm 0.002	0.01 \pm 0.001	0.01 \pm 0.001	0.04 \pm 0.001
Turbidity	NTU	0.8 \pm 0.3	1.3 \pm 0.5	2.3 \pm 0.6	0.8 \pm 1.3	3.7 \pm 4.6	4.3 \pm 1.7
Water depth	cm	47.8 \pm 0.3	61.1 \pm 0.5	75.3 \pm 0.6	32.0 \pm 1.4	29.9 \pm 4.6	28.3 \pm 1.7
Riparian cover	%	27.4 \pm 31.5	37.0 \pm 32.4	45.3 \pm 31.2	20.6 \pm 14.0	15.0 \pm 16.2	15.0 \pm 9.0
Channel width	cm	312.2 \pm 101.3	271.1 \pm 102.8	279.2 \pm 97.2	301.0 \pm 144.9	388.3 \pm 89.4	563.6 \pm 135.6
Detrital cover	%	2.7 \pm 2.0	4.2 \pm 4.1	37.5 \pm 28.9	30.5 \pm 20.9	33.3 \pm 19.7	45.0 \pm 22.0
Substrate embeddedness		4.2 \pm 2.0	3.5 \pm 1.4	2.7 \pm 1.9	4.1 \pm 0.7	3.5 \pm 1.8	2.2 \pm 1.8
PO_4^{3-}	mg L^{-1}	0.6 \pm 0.9	0.3 \pm 0.7	0.3 \pm 0.7	1.3 \pm 1.5	0.9 \pm 0.5	1.1 \pm 0.4
NH_4^+	mg L^{-1}	0.6 \pm 1.8	0.1 \pm 0.1	0.4 \pm 0.5	0.2 \pm 0.3	0.2 \pm 0.2	0.5 \pm 0.3
<i>Sediment</i>							
N	%	0.03 \pm 0.01	0.03 \pm 0.004	0.04 \pm 0.01	0.02 \pm 0.01	0.04 \pm 0.03	0.03 \pm 0.01
K	%	0.01 \pm 0.01	0.02 \pm 0.0	0.02 \pm 0.01	0.02 \pm 0.01	0.10 \pm 0.09	0.03 \pm 0.03
Ca	%	0.001 \pm 0.003	0.002 \pm 0.004	0.01 \pm 0.02	0	0	0
Mg	%	0.001 \pm 0.003	0.003 \pm 0.01	0.01 \pm 0.01	0.0 \pm 0.0	0.003 \pm 0.01	0.01 \pm 0.01
Na	mg kg^{-1}	112.8 \pm 47.5	416.0 \pm 49.8	568.4 \pm 93.8	375.7 \pm 96.4	512.0 \pm 207.1	1009 \pm 156.0
Mn	mg kg^{-1}	5.7 \pm 3.4	4.1 \pm 2.6	23.0 \pm 40.0	17.1 \pm 16.5	17.8 \pm 22.5	21.8 \pm 21.8
Fe	mg kg^{-1}	1081 \pm 583.5	1169 \pm 795.8	1585 \pm 1141	1101 \pm 275.7	1333 \pm 734.6	816.0 \pm 411.5
Cu	mg kg^{-1}	0.6 \pm 0.4	2.1 \pm 1.5	6.5 \pm 3.3	451.3 \pm 981.5	40.5 \pm 40.2	88.8 \pm 170.3
Zn	mg kg^{-1}	2.8 \pm 1.3	3.9 \pm 3.0	15.0 \pm 11.1	3.2 \pm 3.5	2.5 \pm 1.9	96.0 \pm 144.0
B	mg kg^{-1}	30.5 \pm 11.9	29.3 \pm 16.9	38.1 \pm 25.6	0.2 \pm 0.6	0.2 \pm 0.4	2.0 \pm 4.9

Table 2. Kruskal-Wallis results of the measured environmental variables for the different fish zones and months in the Krom River system, South Africa

Variable	Zone		Season	
	H	p	H	p
<i>Water</i>				
Temperature	29.297	<0.001	45.273	<0.001
pH	36.117	<0.001	14.808	<0.001
Dissolved oxygen	24.772	<0.001	0.582	0.446
Conductivity	40.909	<0.001	5.149	0.023
TDS	53.158	<0.001	0.585	0.445
Salinity	44.657	<0.001	8.749	0.003
Turbidity	12.313	0.002	49.161	<0.001
Water depth	2.624	0.269	8.267	0.004
Riparian cover	6.815	0.033	5.423	0.020
Channel width	2.179	0.336	1.672	0.196
Detrital cover	9.065	0.011	14.597	<0.001
Substrate embeddedness	11.443	0.003	0.108	0.743
PO ₄ ³⁻	0.163	0.922	11.241	0.001
NH ₄ ⁺	6.184	0.045	1.930	0.165
<i>Sediment</i>				
N	8.188	0.017	3.664	0.056
K	2.097	0.350	4.302	0.038
Ca	3.214	0.200	5.500	0.019
Mg	6.963	0.031	0.478	0.489
Na	20.429	<0.001	0.372	0.542
Mn	2.329	0.312	7.677	0.006
Fe	0.850	0.652	0.005	0.944
Cu	2.275	0.321	30.715	<0.001
Zn	6.713	0.035	2.268	0.132
B	0.027	0.986	34.796	<0.001


Table 3. Dominant diatom species relative percentage abundances \pm standard deviation (species with $> 2\%$) collected in three river zones (rainbow trout zone, rock catfish zone, sawfish zone) between the two study months (October 2017, January 2018) along the Krom River system, South Africa

Species	October			January		
	Rainbow trout	Rock catfish	Sawfish	Rainbow trout	Rock catfish	Sawfish
<i>Cymbella cymbiformis</i>		2.1 \pm 4.3	2.0 \pm 3.2		1.4 \pm 1.4	0.2 \pm 0.4
<i>Cymbella tumida</i>	1.5 \pm 4.7	1.2 \pm 1.5	1.8 \pm 3.0	0.1 \pm 0.2	0.3 \pm 0.6	0.2 \pm 0.3
<i>Diatoma vulgare</i>	1.1 \pm 3.5	10.8 \pm 8.8	0.1 \pm 0.2	1.5 \pm 1.9	5.8 \pm 6.3	1.3 \pm 1.5
<i>Eunotia formica</i>	5.4 \pm 10.0	12.8 \pm 8.8	4.5 \pm 8.8	10.9 \pm 8.4	8.9 \pm 4.4	8.7 \pm 6.0
<i>Eunotia minor</i>	3.7 \pm 5.4	15.8 \pm 10.9	7.2 \pm 7.0	3.2 \pm 3.4	12.4 \pm 7.2	2.7 \pm 1.7
<i>Eunotia rhomboidea</i>	14.5 \pm 12.8	6.0 \pm 10.1	6.1 \pm 7.0	3.3 \pm 3.2	7.5 \pm 4.7	9.9 \pm 7.0
<i>Fragilaria sp.</i>	2.8 \pm 6.9		0.5 \pm 0.6	1.2 \pm 3.2	0.3 \pm 0.6	0.9 \pm 1.2
<i>Fragilaria ulna</i>	5.0 \pm 6.4	0.3 \pm 0.8	2.1 \pm 2.5	0.7 \pm 1.1	0.1 \pm 0.2	0.9 \pm 1.2
<i>Frustalia rostrata</i>	2.3 \pm 4.0	0.2 \pm 0.5		0.5 \pm 1.1	0.1 \pm 0.4	0.1 \pm 0.2
<i>Frustalia saxonica</i>	2.6 \pm 5.5	3.3 \pm 5.4	0.4 \pm 0.6	0.5 \pm 1.1	0.1 \pm 0.2	0.1 \pm 0.3
<i>Frustalia magaliesmontana</i>		0.6 \pm 1.1	3.1 \pm 5.7	0.7 \pm 1.5	0.2 \pm 0.5	1.1 \pm 2.0
<i>Gomphonema acuminatum</i>			0.1 \pm 0.2	3.4 \pm 5.8	1.3 \pm 3.2	0.4 \pm 0.9
<i>Gomphonema loticillum</i>				3.0 \pm 4.9	1.4 \pm 3.0	
<i>Gomphonema parvulum</i>	0.5 \pm 1.4		2.2 \pm 5.3	4.8 \pm 6.5	1.1 \pm 2.4	0.1 \pm 0.2
<i>Gomphonema venusta</i>	2.2 \pm 4.1			4.8 \pm 7.5	1.0 \pm 2.4	0.3 \pm 0.8
<i>Navicula heimansoides</i>	2.4 \pm 3.5	2.0 \pm 4.9	2.0 \pm 4.8	0.3 \pm 0.6	0.7 \pm 1.4	1.1 \pm 1.9
<i>Navicula ranomafanensis</i>	4.5 \pm 7.6	1.2 \pm 2.0	8.7 \pm 6.4		1.2 \pm 2.4	0.7 \pm 1.7
<i>Navicula radiosa</i>	10.0 \pm 7.5	8.5 \pm 8.4	6.3 \pm 4.6	5.1 \pm 5.7	4.8 \pm 4.6	1.8 \pm 2.1
<i>Navicula sp. 1</i>	2.5 \pm 3.4	0.1 \pm 0.2	2.5 \pm 2.7	0.5 \pm 1.0	0.1 \pm 0.1	0.9 \pm 1.2
<i>Navicula sp. 2</i>	0.4 \pm 1.1			2.4 \pm 5.5	0.1 \pm 0.2	
<i>Navicula subrhynchocephala</i>				2.0 \pm 3.6		0.4 \pm 0.9
<i>Navicula zanonii</i>	0.2 \pm 0.5	0.7 \pm 1.8		2.0 \pm 3.4	0.2 \pm 0.4	1.3 \pm 2.6
<i>Nitzschia sigma</i>	2.0 \pm 2.2	0.2 \pm 0.3	1.6 \pm 1.6	0.2 \pm 0.3	0.1 \pm 0.2	0.3 \pm 0.5
<i>Pinnularia acrosphaeria</i>				3.0 \pm 5.9	1.3 \pm 2.1	0.1 \pm 0.2
<i>Pinnularia divergens</i> var. <i>undulata</i>	2.1 \pm 4.4	0.7 \pm 0.8	1.6 \pm 3.5	0.6 \pm 1.7	0.6 \pm 1.3	0.3 \pm 0.5
<i>Pinnularia viridiformis</i>	2.8 \pm 3.5	3.2 \pm 3.8	4.8 \pm 5.5	2.1 \pm 4.4	2.0 \pm 2.8	2.6 \pm 2.8
<i>Staurosira elliptica</i>	2.4 \pm 6.1	2.9 \pm 7.1		0.5 \pm 1.2		0.1 \pm 0.3
<i>Surirella sp.</i>	0.1 \pm 0.3			3.8 \pm 11.6		0.1 \pm 0.3
<i>Tabellaria flocculosa</i>	5.0 \pm 11.5	7.9 \pm 12.3	10.3 \pm 13.2	4.7 \pm 6.1	8.7 \pm 10.5	25.5 \pm 17.6
<i>Tryblionella apiculata</i>	0.1 \pm 0.3			2.1 \pm 6.6	4.8 \pm 11.9	3.3 \pm 8.0
Taxa richness	27.1 \pm 5.4	27.2 \pm 4.1	28.2 \pm 5.7	13.4 \pm 2.7	13.8 \pm 2.2	16.2 \pm 3.3
Shannon-Weiner	2.6 \pm 0.3	2.6 \pm 0.2	2.4 \pm 0.3	2.2 \pm 0.3	2.1 \pm 0.2	2.3 \pm 0.2
Simpson	0.89 \pm 0.04	0.90 \pm 0.03	0.83 \pm 0.07	0.86 \pm 0.05	0.84 \pm 0.04	0.87 \pm 0.03
Evenness	0.50 \pm 0.08	0.52 \pm 0.10	0.42 \pm 0.06	0.71 \pm 0.08	0.61 \pm 0.10	0.63 \pm 0.08

Table 4. Predictor variable contribution to boosted regression tree models relating diatom species richness to environmental variables learning rate of 0.05 and tree complexity of 5.

Environmental variables are sorted in order of decreasing diatom richness contribution

Predictor	Relative contribution (%)	Predictor	Relative contribution (%)
B	43.3	Riparian cover	1.8
Cu	20.8	TDS	1.8
Fe	3.4	Turbidity	1.5
Water depth	3.2	Channel width	1.4
Temperature	3.0	Salinity	1.0
PO ₄ ³⁻	2.9	pH	1.0
Mn	2.6	Substrate embeddedness	1.0
Detrital cover	2.6	Conductivity	0.9
Zn	2.4	DO	0.9
Na	2.3	K	0.1
NH ₄ ⁺	1.9	N	0.1



Graphical Abstract

Journal Pre-proof

Highlights

- Sediment and water chemistry variables increased downstream in both events.
- 110 diatom species across the two study events identified.
- Significant differences observed in diatom community across river zones and events.
- Boron and copper most important variables in structuring diatoms.

Journal Pre-proof